

Biocontrol of Green Mould Disease of Oyster Mushroom (*Pleurotus ostreatus*) using *Bacillus amyloliquefaciens*

Mwangi, R. W.* Kariuki, S. T Wagara, I. N.
Department of Biological Sciences, Egerton University.

Abstract

The occurrence of *Trichoderma harzianum* and *T. asperellum* in cultivation of oyster mushroom (*Pleurotus ostreatus*) frequently results in serious crop losses and considerable inhibition of growth of mycelium and fruiting bodies of oyster mushroom thus lowering the yield substantially. *Bacillus amyloliquefaciens* strain isolated from groundnuts proved very effective in antagonizing the oyster mushroom pathogenic *T. harzianum* and *T. asperellum* without having a negative effect on *P. ostreatus* mycelia. The *Bacillus amyloliquefaciens* was found to produce diffusible and volatile organic compounds. This strain is a potential biocontrol candidate, in addition to the lack of antagonistic activity towards *P. ostreatus* mycelia. The present study, hence, provides a potential biocontrol agent for *Trichoderma* green mould. However, field studies of this isolate as substrate inoculant in oyster mushroom are required in order to establish its actual performance.

Keywords: *Bacillus amyloliquefaciens*, Green mould, Mushroom, Biocontrol

1. Introduction

Trichoderma green mould infection in edible mushrooms has a long history Sinden and Hauser, (1953). This green mould disease of cultivated oyster mushroom (*Pleurotus ostreatus*) has been reported in several countries where large scale production of oyster mushroom is practised Hatvani *et al.* (2008). Oyster mushroom is the third most important commercially grown basidiomycete worldwide; its cultivation has significantly increased in the world during the last few years. Among many pests and diseases in oyster mushroom cultivation, the most serious crop losses are due to *Trichoderma* green mould infections. Previously, the fungi responsible for the green mould disease of *P. ostreatus* was reported to be *T. aggressivum* Hatvani *et al.* (2007); Komon-Zelazowska *et al.* (2007). However, new species including *T. harzianum* and *T. asperellum* Park *et al.* (2006) have emerged and been reported.

Chen and Moy (2004) stated that parameters of mushroom cultivation, such as the sources of carbon and nitrogen, high relative humidity, warm temperatures, a fluctuation of these factors, and the absence of light during spawn run are ideal environmental conditions for moulds as well, which can easily lead to a contamination. In these favourable conditions moulds exhibit fast growth, thus competing for space and nutrients more successfully than the mushrooms. Additionally, they are able to produce extracellular enzymes, toxic secondary compounds as well as volatile organic compounds Williams *et al.* (2003), which can result in a substantial decrease in production or wiping out the entire crop. Pathogenic green moulds may colonize the substrate or grow on the surface of the emerging mushrooms, which become severely spotted and often distorted. Besides that, in serious outbreaks no fruiting bodies are produced. *Trichoderma* spp. produce whitish mycelia which are not easily distinguished from those of the mushrooms during spawn run, making it difficult to recognize the infection at an early stage Won (2000), Largeau-Mamoun *et al.* (2002).

The main symptom of green mould disease is the appearance of greenish mycelium in the compost, bagging layer or fruiting bodies of *P. ostreatus*, 2–5 weeks after the beginning of production cycle. The pathogen inhibits the growth of mushroom and in severe outbreaks the fruiting bodies are not produced. This severely affects the markets of the mushrooms as most of the oyster mushroom farms are affected by the *Trichoderma* green mould problem. Although the first flush of the production can be saved with very strict hygiene, *Trichoderma* green mould often reduces the yield of the second flush by 20-30% Nagy (2012). Prevention has therefore, to play a central role in green mould management; however, if the infection has already occurred at a farm, it has to be controlled.

Biological control offers an important alternative to synthetic chemicals. Biocontrol has been shown to be economical, environmentally friendly and an alternative to chemical fungicides for managing oyster mushroom diseases and contamination.

Bacillus is a genus of gram positive and rod shaped bacteria. They are capable of forming stable dormant structures called endospores in nutrient void and stressful environmental conditions. Spores are generally viable for a long period even under harsh conditions. The sporulation ability and easy cultivation of *Bacillus* species Ross *et al.* (2001); Tiago *et al.* (2004) are attractive for their practical use as biocontrol agents.

There have been no reports on the use of microorganisms for the biological control of mushroom green mold. It is considered that effective biological control of green mold relies both on the selection of antagonists that act specifically against the pathogens which cause the green mold, and that these same antagonists have no inhibition on the mycelial growth and fruiting bodies formation of mushrooms. Thus, this study was conducted

to select prospective antagonists for the biological control of green mold of mushrooms caused by *Trichoderma* spp.

The use of bacteria like *Bacillus* spp., have been investigated because of their ability to produce antifungal metabolites that protect plants from fungal infection Moita *et al.* (2005); Siddiqui *et al.* (2005); Nourozian *et al.* (2006). Microorganisms produce compounds with the following properties: high specificity against target plant pathogens, easy degradability and low cost for mass production. *Bacillus* spp. are widely distributed in soils and substrates, have high thermal tolerance, rapid growth in media culture and readily form resistant spores. They are considered safe biological agents and their potency is high Kim *et al.* (2003); Saharan and Nehra (2011). This study explored the potential of *B. amyloliquefaciens* as a biological control agent of green mold of oyster mushrooms.

2. Materials and Methods

2.1 Bacterial and fungal isolate

The *B. amyloliquefaciens* was isolated from healthy groundnuts and identified according to the morphological, biochemical and physiological tests as described by Collee *et al.* (1996). *Trichoderma* fungal isolates were isolated from the spent mushroom substrate and from infected wheat grain spawn. These included *T. harzianum*, and *T. asperellum*. The fungal isolates were characterized and identified based on their colonial morphology and microscopic characteristics using different identification keys and methods according to Eastburn and Butler, (1988). The bacterial isolate was maintained on nutrient agar (NA) slants while fungal pathogens were maintained on PDA slants. Slant cultures were stored at 4 °C in the refrigerator until use.

2.2 Antagonistic effect of *B. amyloliquefaciens* isolate (in vitro)

Bacillus amyloliquefaciens isolate was used in *in vitro* sensitivity experiments against the fungal isolates. Potato dextrose agar plates were inoculated with antagonistic isolate of *B. amyloliquefaciens* as two streak lines with a loop-full of 2 days-old culture on the periphery of the petri plate. The plates were doubly inoculated with mycelial disc (5 mm in diameter) of an actively growing culture of the pathogen placed 5 cm opposite to the other edge of the petri plate and incubated at 25 °C for 7 days Toure *et al.* (2004). (Plate 1A)



Plate 1A dual culture interaction between *B. amyloliquefaciens* and *T. harzianum*
Plates with *B. amyloliquefaciens* and the respective pathogens alone were used as checks. Inhibition zones (the distance between the edge of antagonistic bacterial growth and the edge of tested fungal isolates) were measured. The percentage inhibition of the growth of the pathogen were calculated with the help of the formula:

$$L = (C - T) / CX100,$$

Where:

L = inhibition of radial mycelial growth,

C = radial growth measurement of pathogen in control,

T = radial growth measurement of pathogen in the presence of antagonist.

Radial growth of pathogen was recorded and percentage inhibition calculated in relation with control according to Hajieghrari *et al.* (2008). The experiment was done twice and each test was done in five replicates. The percentage inhibition, L, was categorized on a growth inhibition category, GIC scale from 0-4:

Where:

0 = no growth inhibition,

1 = 1 - 25% growth inhibition,

2 = 26 - 50% growth inhibition,

3 = 51 - 75% growth inhibition

4 = 76 - 100% growth inhibition.

Mycelial growth of the pathogen was measured and observations were recorded on formation of inhibition zone, over growth and lysis of pathogen mycelium. The data obtained were statistically analyzed using the Statistical Analysis System (SAS).

2.3 Production of Volatile Antifungal Metabolites

Production of volatile metabolites by *B. amyloliquefaciens*, having antagonistic activity against *Trichoderma* pathogens was tested by paired plate technique as described by Fiddaman and Rossall (1993) with some slight modifications. A petri dish containing PDA medium was streak inoculated with a loopful of 48 hours old *B.*

amyloliquefaciens isolate. The top of the petri dish containing PDA was inoculated with a 5 mm plug of the actively growing *T. harzianum* and another set with *T. asperellum* separately, at the centre. Both half plates were sealed together and the paired plates were incubated at 25° C for seven days. This incubation ensured that both organisms were growing in the same culture conditions although they were physically separated. Control set of paired plates was designed with only the test pathogens on PDA half plate inverted over un streaked PDA half plate. The experiment was repeated twice and conducted in five replicates. After incubation period, the paired plates were observed for inhibition of fungal growth as compared to the control. The radial growth diameter of the pathogens was measured and compared with the control set. Percentage inhibition of radial growth of the pathogens was calculated as mentioned before.

2.4 Slide cultures

For slide cultures, a clean slide was placed on an L-shaped glass rod in a 9 cm diameter petri dish and autoclaved. Then a small amount of molten PDA was poured and evenly spread over the slide to make a thin agar film. One end of the slide was kept free of the medium to facilitate handling. Inocula from each *B. amyloliquefaciens* or *Trichoderma* isolates were placed separately on the slide 1 cm apart from each other. Two ml of sterile water was added to the petri dish to prevent drying, and the slide incubated at 25 °C for 3-5 days. *Trichoderma* species alone were used as controls. At the end of incubation period, regions where the *B. amyloliquefaciens* met the hyphae of the pathogen were observed under a light microscope for the presence of coil formation and penetration structures, or wall disintegration.

3. Results

3.1 Effect of *B. amyloliquefaciens* on the growth of the *Trichoderma* pathogens *in vitro*

Bacillus amyloliquefaciens grew faster than *T. harzianum* and *T. asperellum* on PDA media under the same culture conditions. *B. amyloliquefaciens* grew in all possible directions and came into contact with the pathogenic fungi on the fifth day after inoculation and started to suppress further growth of the pathogens. No inhibition zone was formed around the contact area between these species (Plate 1). Thus initially, *B. amyloliquefaciens* inhibited *T. harzianum* and *T. asperellum* by competing for space and nutrients. Later the mycelia of *T. harzianum* collapsed and died completely indicating that *B. amyloliquefaciens* produced antibiotics.




Plate 1: Antagonism of *B. amyloliquefaciens* (BA) against *T. asperellum* (TA) and *T. harzianum* (TH) ; a and c are treated while b and d are controls.

3.2 Production of volatile compounds by *B. amyloliquefaciens* against *Trichoderma* spp.

The antagonistic potential was noted to vary through volatile metabolites produced by *B. amyloliquefaciens*, and direct parasitism on the two pathogens. In addition, a change in mycelia colour which was different from the mycelium colour of the control, close to the colony was observed. A stronger antibiosis mechanism of antagonistic *B. amyloliquefaciens* and a higher pathogen inhibition through volatile metabolites were noted. Volatile toxic substances produced by antagonists were noted to spread easily and inhibit pathogens growth *in vitro* (Plate 2).

The volatile organic compounds (VOCs) produced by *B. amyloliquefaciens* reduced the mycelial growth of *T. harzianum* and *T. asperellum* (Plate 2, 1 & 2) in comparison with the control (Plate 2, 3 & 4). The VOCs decreased the length of fungal mycelia, and colonies seemed to be significantly reduced ($p = 0.05$). The inhibition of *T. harzianum* and *T. asperellum* by VOCs was about 71 and 72.8% (Figure 1) respectively compared with the control after seven days, suggesting that the bacterial VOCs were unable to completely kill these two pathogens but had a significantly inhibitory effect on fungal mycelia. The colour of the mycelia also changed from green to white indicating that there was no sporulation in the treated plates (Plate 2, 1 & 2).




Plate 1: Effects of volatile compounds of *B. amyloliquefaciens* against *Trichoderma* species




Figure 1: Percentage mean inhibition of *T. harzianum* and *T. asperellum* by volatile compound of *B. amyloliquefaciens*

4. Discussion

The production of antifungal compounds and siderophores is a primary mechanism in suppressing disease by *Bacillus* spp. Smitha *et al.* (2015). Peptide antibiotics and several other compounds which are toxic to plant pathogens have been recovered from several *Bacillus* strains Yu *et al.* (2002). Antagonism was evident in petri dishes through the different magnitudes of the *Trichoderma* suppression by the *B. amyloliquefaciens*. The control plates without *B. amyloliquefaciens*, were completely covered by pathogen mycelia showing no fungus growth inhibition (Plate 1, b & d). Sporulation was also inhibited completely compared to the control. The mean mycelium growth inhibition on the tested pathogens by this bacterial isolate revealed that the antagonist is highly effective.

A compound microscope was used to make observations, it was clear that from the mycelium collected from the interface region of *T. harzianum* and *T. asperellum* with *B. amyloliquefaciens* showed modifications in

the mycelium appearance. Some of the modifications included: mycelia colour changing from dark green to white, a coagulation of the fungal cytoplasm which could be observed up to the hypha, presence of small vesicles and the appearance of big vacuoles. In this case, the destructive effect of the *Trichoderma* spp by *B. amyloliquefaciens* was high, resulting in serious damage of the hyphae, associated with a series of degradation events.

The mycoparasitic potential of *Bacillus* spp. is well documented Johri *et al.* (2003); Saharan and Nehra (2011). Thus, this phenomenon has often been used as a means for *in vitro* screening of biocontrol agents Jat and Agalave, (2013). Similar conclusions have been reported by El Hassni *et al.* (2007) and Idris *et al.* (2007). They reported a modification of the fungal mycelium appearance, due to antifungal secondary metabolite production. Generally, biocontrol capacity through antagonistic bacteria involves either competition Wehner *et al.* (2010) or bacterial metabolite production. Bacterial metabolites produced for antagonism towards plant pathogens include siderophores, hydrogen cyanide, antibiotics or extracellular enzymes Kamilova *et al.* (2005); Sang *et al.* (2006). It has been reported that *Bacillus* spp. contains various biocontrol characteristics including secondary metabolites, the colonizing potential, and the production of competitors Yoshida *et al.* (2001); Schmidt *et al.* (2004).

According to the observations made in this study, production of diffusible and volatile organic compounds seems to be a primary source of inhibition of the tested fungal pathogens. This agrees with the work done by Prashar *et al.* (2013), who reported that isolate TNAM5 belonging to *Bacillus* spp. was found to be a strong producer of volatile and diffusible antifungal compounds, a character that has been previously established for various strains of *Bacillus* Wang *et al.* (2007); Dunlap *et al.* (2011).

Nonvolatile antibiotics, including lipopeptides, have strong antifungal activities. However, these nonvolatile antibiotics cannot spread over long distances, and only when these antagonists directly colonize the mushroom mycelia they can prevent pathogenic fungi from infecting the mushroom crop. Volatile organic compounds spread over long distances, producing fungistatic microenvironments around the pathogenic communities. In addition, the antifungal VOCs produced by bacteria can kill surviving spores in the mushroom substrate and limit both the production and the establishment of the green mould disease in spawn preparation and in mushroom houses where mushrooms are growing. Additionally, Munimbazi and Bullerman, (1998) reported that extracellular antifungal metabolites produced by *B. pumilus* inhibited mycelial growth of many species of *Aspergillus*, *Penicillium* and *Fusarium*.

5. Conclusion

The *B. amyloliquefaciens* used in this study is effective in suppressing mushroom pathogenic fungi, including *Trichoderma harzianum* and *T. asperellum*, the causative agents of green mould disease of oyster mushrooms. It exhibited broad-spectrum antifungal properties. It produced both volatile and nonvolatile organic compounds and showed good potential for biological control of green mould disease. This study has, therefore, provided a potential bacterial isolate suitable for controlling *Trichoderma* green mould. However, it is suggested that a detailed investigation must be carried out to evaluate this isolate for its field performance as a biocontrol.

References

- Collee, J.G., Fraser, A.G., Marmion, B.P. and Simmons, A. (1996). Practical Medical Microbiology. pp: 131-149. In 14th Edition. Churchill Livingstone, New York.
- Dunlap, C.A., Schisler, D.A., Price, N.P. and Vaughan, S.F. (2011). Cyclic Lipopeptide Profile of Three *Bacillus* Strains; Antagonist of *Fusarium* Head-blight. *Journal of Microbiology*, 49:603-609.
- Eastburn, D.M. and Butler, E.E. (1988). Microhabitat characterization of *Trichoderma harzianum* in natural soil: evaluation of factors affecting population density. *Soil Biology and Biochemistry*, 20:541-545.
- El-Hassni, M., El-Hadrami, A., Daayf, F., Cherif, M., Ait-Barka, E. and El-Hadrami, I. (2007). Biological control of bayoud disease in date palm: selection of microorganisms inhibiting the causal agent and inducing defense reactions. *Environmental and Experimental Botany*, 59 (2):224-234.
- Fiddaman, P.J. and Rossall, S. (1993). The Production of Antifungal Volatiles by *Bacillus subtilis*. *Journal of Applied Bacteriology*, 74:119-126.
- Hajieghrari, B., Torabi-Giglou, M., Mohammadi, M. R. and Davari, M. (2008). Biological potential of some Iranian *Trichoderma* isolates in control of soil borne plant pathogenic fungi. *African Journal of Biotechnology*, 7(8):967-972.
- Hatvani, L., Antal, Z., Manczinger, L., Szekeres, A., Druzhinina, I.S., Kubicek, C.P., Nagy, A., Nagy, E., Vagvolgyi, C. and Kredics, L. (2007). Green mold diseases of *Agaricus* and *Pleurotus* are caused by related but phylogenetically different *Trichoderma* species. *Phytopathology*, 97:532-537.
- Hatvani, L., Kocsube, S., Manczinger, L., Antal, Z., Szekeres, A., Druzhinina, I.S., Komo-Zelazowska, M., Kubicek, C.P., Nagy, A., Vagvolgyi, C. and Kredics, L. (2008). The green mould disease global threat to the cultivation of oyster mushroom (*Pleurotus ostreatus*): a review. *Mushroom Science*, 17:485-495.

- Idris, H.A., Labuschagne, N. and Korsten, L. (2007). Screening rhizobacteria for biological control of *Fusarium* root and crown rot of sorghum in Ethiopia. *Biological Control*, 40: 97–106.
- Jat, J. G., and Agalave, H.R. (2013). Antagonistic properties of *Trichoderma* species against oilseed-borne fungi. *Science Research Reporter*, 3(2), 171-174.
- Johri, B.N., Sharma, A. and Viridi, J.S. (2003). Rhizobacterial diversity in India and its influence on soil and plant health. *Advances in Biochemical Engineering/Biotechnology*, 84:49–89.
- Kamilova, F., Validov, S., Azarova, T., Mulders, I. and Lugtenberg, B. (2005). Enrichment for enhanced competitive plant root tip colonizers selects for a new class of biocontrol bacteria. *Environmental Microbiology*, 7:1809-1817.
- Kim, H.S., Park, J., Choi, S.W., Choi, K.H., Lee, G.L., Ban, S.J., Lee, C.H. and Kim, C.S. 2003. Isolation and characterization of *Bacillus* strains for biological control. *Journal of Microbiology*, 41(3):196-205.
- Komon-Zelazowska, M., Bissett, J., Zafari, D., Hatvani, L., Manczinger, L., Woo, S., Lorito, M., Kredics, L., Kubicek, C.P. and Druzhinina, I.S. (2007). Genetically closely related but phenotypically divergent *Trichoderma* species cause world-wide green mould disease in oyster mushroom farms. *Applied and Environmental Microbiology*, 73:7415-7426.
- Nagy, A., Laszlo, M., Dora, T., Lorant, H., Julia, G., Zsuzsanna, A., Enik, S., Csaba, V. and Laszlo, K. (2012). Biological control of oyster mushroom green mould disease by antagonistic *Bacillus* species. *Biological Control of Fungal and Bacterial Plant Pathogens*, 78:289-293.
- Nourozian J.H., Etebarian, R. and Khodakaramian, G. (2006). Biological control of *Fusarium graminearum* on wheat by antagonistic bacteria. *Nutraceutical and Functional Food*. 28:29-36.
- Park, M.S., Bae, K.S. and Yu, S.H. (2006). Two new species of *Trichoderma* associated with green mold of oyster mushroom cultivation in Korea. *Mycobiology*. 34:111-113.
- Prashar, P., Kapoor, N. and Sachdeva, S. (2013). Isolation and Characterization of *Bacillus* spp with *In-vitro* Antagonistic Activity against *Fusarium oxysporum* from Rhizosphere of Tomato. *Journal of Agriculture, Science and Technology*, 15:1501-1512.
- Ross, N., Villemur, R., Marcandella, E. and Deschenes, L. (2001). Assessment of Changes in Biodiversity when a Community of Ultra-microbacteria Isolated from Groundwater is Stimulated to Form a Biofilm. *Microbial Ecology*, 42:56-68.
- Saharan, B.S. and Nehra, V. (2011). Plant growth promoting rhizobacteria: a critical review. *Life Sciences and Medicine Research*, 21:23-31.
- Samuels, G.J., Dodd, S.L., Gams, W., Castlebury, L.A. and Petrini, O. (2002). *Trichoderma* species associated with the green mold epidemic of commercially grown *Agaricus bisporus*. *Mycologia*, 94:146-170.
- Sang, M.K., Chiang, M.H., Yi, E.S., Park, K.W. and Kim, K.D. (2006). Biocontrol of Korean ginseng root rot caused by *Phytophthora cactorum* using antagonistic bacterial strains ISE13 and KJ1R5. *Plant Pathology Journal*, 22:103-106.
- Schmidt, C.S., Agostini, F., Leifert, C., Killham, K. and Mullins, C.E. (2004). Influence of soil temperature and matric potential on sugar beet seedling colonization and suppression of *Pythium* damping-off by the antagonistic bacteria *Pseudomonas fluorescens* and *Bacillus subtilis*. *Phytopathology*, 94:351-363.
- Siddiqui, S., Siddiqui, Z.A. and Ahmad, I. (2005). Evaluation of fluorescent *Pseudomonas* and *Bacillus* isolates for the biocontrol of a wilt complex of pigeon pea. *World Journal of Microbiology and Biotechnology*, 21:729-739.
- Smitha, K.P., Mohan, R., Devadason, A. and Raguchander, T. (2015). Exploiting novel rhizosphere *Bacillus* species to suppress the root rot and wilt pathogens of chickpea. *African Journal of Microbiology Research*, 9(15), 1098-1104.
- Wehner, J., Antunes, P.M., Powell, J.R., Mazukatow, J., and Rillig, M.C. (2010). Plant pathogen protection by arbuscular mycorrhizas: a role for fungal diversity. *Pedobiologia*, 53(3), 197-201.
- Tiago, I., Teixeira, I., Silva, S., Chung, P., Verissimo, A. and Manaia, C.M. (2004). Metabolic and genetic diversity of mesophilic and thermophilic bacteria isolated from composted municipal sludge on poly-epsilon-caprolactones. *Current Microbiology*, 49:407-414.
- Wang, J., Liu, J., Chen, H. and Yao, J. (2007). Characterization of *Fusarium graminearum* Inhibitory Lipopeptide from *Bacillus subtilis* IB. *Applied Microbiology and Biotechnology*, 76:889-894.
- Yoshida, S., Hiradate, S., Tsukamoto, T., Hatakeda, K. and Shirata, A. (2001). Antimicrobial activity of culture filtrate of *Bacillus amyloliquefaciens* RC-2 isolated from mulberry leaves. *Phytopathology*, 91 (2):181-187.
- Yu, G.Y., Sinclair, J.B., Hartman, G.L. and Bertagnolli, B.L. (2002). Production of iturin A by *Bacillus amyloliquefaciens* suppressing *Rhizoctonia solani*. *Soil Biology and Biochemistry*, 34:955-963.