

## Environmentally benign un-catalyzed esterification of alcohols under solvent-free condition using acetyl chloride.

Dr. Sangita S Makone\*<sup>1</sup>

<sup>1</sup>Chemical Sciences Research Laboratory, Swami Ramanand Teerth Marathwada University, Vishnupuri, Nanded.

Maharashtra. India.

Pin code: 431606.

\*Email: [ss\\_makone@rediffmail.com](mailto:ss_makone@rediffmail.com).

Sandeep N Niwadange<sup>1</sup>

<sup>1</sup>Chemical Sciences Research Laboratory, Swami Ramanand Teerth Marathwada University, Vishnupuri, Nanded.

Maharashtra. India.

Pin code: 431606.

Email: [sandip.niwadange@gmail.com](mailto:sandip.niwadange@gmail.com)

### Abstract

A general and practical route for un-catalyzed esterification of alcohols using acetyl chloride is described under solvent free and energy efficient conditions. Excellent yields, wide applicability to various alcohols, mild reaction conditions, short reaction time, ease of operation makes this method economic and environmentally benign.

**Keywords:** Esterification, un-catalyzed, solvent-free, acetyl chloride, alcohol.

### 1. Introduction

Functional group protection and deprotection is an important in synthetic organic chemistry (Greene 1991). Amongst protecting groups for alcohols the esters are the most important with acetates being simplest and easiest of all. Ester protects the –OH groups particularly during oxidation, peptide coupling and glycosidation reaction (Greene 1991). Also ester groups are highly important in chemistry and can found in bulk chemicals, fine chemicals, natural products and polymers (Oetra 2003). Esters are prepared using transition metal catalyzed cross-coupling reactions which involved the carbonylation of aryl halides (Schoenberg 1974; Brennfuhrer 2009) but presence of halide anion and reaction conditions are not environmentally benign. Another way of synthesis of esters is an oxidative esterification of aldehydes with alcohols (Kovi 2008; Yoo 2007; Yoo 2006; Xu 2010; Wu 2009) using

stoichiometric amounts of metal salts like  $\text{KHSO}_5$ (oxome) or  $\text{MnO}_2$  as an oxidant (Travis 2003; Maki 2008). Many of the esterification reactions were reported using homogeneous basic catalyst such as amine(Zhdanov 1975), DMAP(Hofle 1978; Scriven 1983), tri-butyl phosphine(Vedejs 1993) or acidic catalyst like p-toluene sulfonic acid(Cope 1963),  $\text{COCl}_2$ (Iqbal 1992),  $\text{Sc}(\text{OTf})_3$ (Ishihara 1996) or  $\text{Bi}(\text{OTf})_3$ (Oriata 2000) with acid chloride or acid anhydride as the acetylating agent. However, the use of homogeneous catalyst possesses several problems such as difficulty in the separation and recovery of the catalyst, disposal of the spent catalyst, corrosion problems. Traditionally esters are prepared by the reaction of activated derivatives like acyl chlorides and acyl anhydride in presence of base(Muzler 1991; Trost 1995). Later modifications involve use of lewis acid (Izumi 1995; Vedejs 1996; Iranpoor 1998; Chandrasekhar 1998; Ganesan 1974) in combination with  $\text{Ac}_2\text{O}$  which is inherently wasteful since half of the every acid anhydride molecule loses as a carboxylic acid. Among the various base catalyst used for esterification of alcohols using acid chlorides, the most common are 1,4-diazobicyclo[2.2.2]-octane (DABCO)(Ishihara 1993; Hajipour 2002), N,N,N,N-tetramethylethylenediamine(TMEDA)(Sano 1999) and NaOH(Volker 1979). Few other catalysts are also reported for these syntheses are  $\text{Al}_2\text{O}_3$  (Yadav 2001; Paul 2003; Yadav 2004),  $\text{TiO}_2$  (Pasha 2008), ZnO (Tamaddon 2005; Sarvari 2005; Tayebbe 2010),  $\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$ (Ghosh 2005),  $\text{CeCl}_3$ (Torregiani 2005),  $\text{Ti}$  (Taylor 1968),  $\text{BiCl}_3$  (Ghosh 2004) and  $\text{BiFeO}_3$  (Farhadi 2009). A catalyst free reaction using acid halides and alcohols has been reported using dichloromethane solvent (Strazzolini 1994). Ranu et.al.(Ranu 2003), reported a solvent free esterification of different alcohols reacted with acetyl or propanoyl chloride. Recently catalyst free condensation of acid chlorides and alcohols has been reported by Christian V. Steven (Stevens 2012) using continuous flow for which microwave technology is required.

Some methods using eco-friendly catalysts i.e. commercial HY Zeolites (Ballini 1998) and montmorillonite K-10 or KSF clay (Li 1998) for the acylation of alcohols and phenols by acetic anhydride also reported.

However most of these methods still have limitations such as use of catalyst and solvent, high power microwave, harsh reaction conditions, long periods of reaction, use of large amount of organic solvents as the reaction medium and air sensitive nature of catalysts, critical product isolation process of high boiling solvents.

Nowadays, the reactions under catalyst and solvent-free conditions are considerably safe, nontoxic, environmentally friendly and inexpensive.

In continuation of our earlier work (Makone 2010) to the best of our knowledge which was first report on un-catalyzed esterification of alcohols under solvent free and energy efficient conditions starting from acid chloride, here we report novel, un-catalyzed esterification of alcohols using acetyl chloride as an acylating agent under solvent free and energy efficient conditions.

## 2. Experimental

### 2.1. General.

All alcohols and acetyl chloride used were commercially available (Sigma Aldrich). Melting point and boiling points were uncorrected and recorded on Veego melting/boiling point apparatus.  $^1\text{H}$  NMR spectra were recorded on a Bruker Avance II 400MHz spectrometer with TMS as internal standard. IR spectra were determined on a Perkin Elmer Spectrum RX FTIR spectrometer.

### 2.2. General procedure for esterification of alcohols.

In a 100 ml two-neck round bottom flask a magnetic stir bar and alcohol (10mmol) were placed. To this alcohol acetyl chloride (20mmol) was added slowly through side neck of round bottom flask. To this round bottom flask reflux condenser was attached through which continuously ice cold water is circulated to trap the fumes of HCl generated during the reaction, which gets condensed and dissolved in distilled water to form dil. HCl in the receiver. The reaction mixture was stirred continuously until corresponding esters are formed. For complete conversion of corresponding alcohols into corresponding esters appropriate proportion of acetyl chloride was added (Table 1). The progress of the reaction was monitored by using TLC. The product formed in the round bottom flask was collected by simple distillation and conversion of alcohols to esters was confirmed by taking following confirmative test of ester group.

A little quantity of formed product of ester was taken in a test tube in which 1-2 drops of phenolphthalein was added. To this mixture 5-6 drops of dil. NaOH was added. Then this mixture was heated in water bath. The pink color of the mixture slowly disappears. This confirms the presence of ester group in the product. The authenticity of the products was established by comparing their melting points and spectral data with the literature.

### 2.3. Spectral data information

#### *Ester of Trigol (entry m):*

IR in  $\text{cm}^{-1}$  : 1739  $\text{cm}^{-1}$  (for ester group).

$^1\text{H}$  NMR : 2.01(s, 6H); 3.45 (t, 4H); 3.65 (t, 4H); 4.2 (t, 4H).

#### *Ester of 1,6-Hexanediol(entry i):*

IR in  $\text{cm}^{-1}$  : 1740  $\text{cm}^{-1}$  (for ester group).

$^1\text{H}$  NMR : 1.30 (quintet, 4H); 1.60 (quintet, 4H); 2.01 (s, 6H); 4.08 (t, 4H).


*Ester of Glycerol (entry n):* IR in  $\text{cm}^{-1}$  :  $1747 \text{ cm}^{-1}$  (for ester group)

$^1\text{H NMR}$  : d 2.01 (s, 9H); 4.32 (d, 4H); 5.20 (quintet, 1H)

### 3. Result and Discussion

In continuation to our previous work (Makone 2010) as there was no earlier report on the application of acetyl chloride as an acetylating reagent for un-catalyzed esterification of alcohols under solvent free conditions, here we wish to report an efficient and environmentally benign method for the synthesis of an ester from alcohols using acetyl chloride.

In order to study the scope and generality of this method a variety of alcohols were subjected to the esterification using acetyl chloride as an acetylating agent. The reaction was carried out under catalyst free and solvent free conditions using following reaction setup (fig.1).



The representative reaction of un-substituted benzyl alcohol was studied to evaluate the results. In a 100 ml two-neck round bottom flask a magnetic stir bar and benzyl alcohol (10mmol) were placed. To this benzyl alcohol acetyl chloride (10mmol) was added slowly through side neck of round bottom flask. To this round bottom flask reflux condenser was attached through which continuously cold water is circulated to trap the fumes of HCl generated during the reaction. The reaction mixture was stirred for 1hr, completion of reaction was monitored by TLC. It was found that the reaction completes to produce 71% formation of corresponding product. For complete conversion of benzyl alcohol into benzyl acetate the percentage of acetyl chloride was increased (20mmol) and progress of reaction was evaluated. During reaction instead of cold water ice cold water was circulated continuously through reflux condenser (Scheme 1). The maximum conversion of benzyl alcohol in to benzyl acetate was observed when 20mmol of acetyl chloride was added and fumes of generated HCl during the reaction gets condensed when ice cold water was circulated continuously through reflux condenser. The reaction was completed in 1hr to produce 98% yield of the product (Table 1, entry a).

Subsequently, various aliphatic alcohols were reacted with acetyl chloride (Table 1, entry b, c, d) to produce corresponding esters with excellent yields. With increasing number of aliphatic carbon chain of various aliphatic alcohols such as heptanol, octanol, decanol were treated with acetyl chloride using 1:4 proportions under optimized reaction set-up conditions, the reaction proceeded successfully producing high yields of desired esters (Table 1, entry e, f, g). The applicability of this process was investigated using diol substrates such as 1,4-butanediol and 1,6-hexanediol, the desired esters were formed using acetyl chloride in 1:4 proportion (Scheme 2) producing high yields of desired esters (Table 1, entry h, i).

After successful application of esterification to various alcohols we tried to extend the newly developed protocol to long chain aliphatic diols. When 1,12-dodecanediol was introduced with acetyl chloride in 1:6 proportion desired ester was formed with 95% yield (Table 1, entry j). In addition we also investigate the reactions using trigol and glycerol as a substrates with acetyl chloride in 1:4 and 1:6 proportion to form the corresponding esters of trigol and glycerol with excellent yields (Table 1, entry m, n). The trigol and glycerol smoothly undergo the un-catalyzed esterification under solvent-free conditions.

With these results in hand, we set out to examine the scope of these reactions to phenolic –OH groups using simple un-substituted phenol. When phenol (10mmol) was stirred with acetyl chloride (20mmol) up to 1hr, no reaction was observed. The results are confirmed using 4-hydroxy benzyl alcohol substrate with acetyl chloride. To our surprise when 4-hydroxy benzyl alcohol (10mmol) reacted with acetyl chloride in 1:4 proportion at solvent free conditions only benzylic –OH group underwent conversion of ester and corresponding ester of 4-hydroxy benzyl alcohol is formed with 96% yield (Table 1, entry o), where as phenolic –OH remains intact. The results show that the generality of present protocol is effective for the alcoholic –OH groups present the alcoholic substrates.

#### 4. Conclusion

In summary we have developed an efficient un-catalyzed esterification of alcohols under solvent-free and energy efficient conditions using acetyl chloride. Method is applicable to various aromatic as well as aliphatic alcohols such as benzyl alcohol, diols, trigols, glycerols. This protocol also gives results to propargyl and allyl alcohols to give their corresponding esters in high to excellent yields. The advantages of this methodology, solvent free, catalyst free, energy efficient and environmentally benign conditions, simple work-up procedure and easy separation of the formed HCl will lead to this protocol a green technique for practical organic synthesis. However this method is limited to phenolic –OH group. Efforts are underway to elucidate the scope and limitations of this method to more variety of alcohols.

## 5. Acknowledgment

SNN thanks to University Grant Commission, New Delhi for providing Rajiv Gandhi Senior Research Fellowship for financial support.

## 6. References

- Ballini, R., Bosica, G., Carloni, S., Ciaralli, L., Maggi, R., Sartori, G. (1998), "Zeolite HSZ-360 as a new reusable catalyst for the direct acetylation of alcohols and phenols under solvent less conditions", *Tetrahedron Letter* **39**, 6049.
- Bernnufer, A., Neumann, H., Beller, M.(2009), "Palladiumkatalysierte Carbonylierungen von Arylhalogeniden und ähnlichen Substraten", *Angewandte Chemie* **121**, 4176-4196; "Palladium catalyzed carbonylation reactions of aryl halides and related compounds", *Angewandte Chemie International Edition* **48**, 4114-4133.
- Chandrasekhar, S., Ramachander, T., Takhi, M.(1998), "Acylation of alcohols with acetic anhydride catalyzed by TaCl<sub>5</sub>: some implications in kinetic resolution" *Tetrahedron Letter* **39**, 3263.
- Cope, A., Herrick, E. C.(1963), "Synthesis of Diethyl cis- $\Delta^4$ -tetrahydrophthalate", *Organic Synthesis* **4**, 304.
- Farhadi, S., Zaidi, M.(2009), "Bismuth ferrite (BiFeO<sub>3</sub>) nano-powder prepared by sucrose-assisted combustion method: A novel and reusable heterogeneous catalyst for acetylation of amines, alcohols and phenols under solvent free conditions", *Journal of Molecular Catalysis A: Chemistry* **299**, 18.
- Ganem, B., Small, V. R. Jr.(1974), "Ferric chloride in acetic anhydride, mild and versatile reagent for the cleavage of ethers", *Journal of Organic Chemistry* **39**, 3728.
- Ghosh, R., Maiti, S., Chakraborty, A.(2004), "Facile catalyzed acylation of heteroatoms using BaCl<sub>2</sub> generated in situ from the precatalyst BiOCl and acetyl chloride", *Tetrahedron Letter* **45**, 6775.
- Ghosh, R., Maiti, S., Chakraborty, A.(2005), "Facile catalyzed acylation of alcohols, phenols, amines and thiols based on ZrOCl<sub>2</sub> · H<sub>2</sub>O and acetyl chloride in solution and solvent free conditions", *Tetrahedron Letter* **46**, 147.
- Greene, T. W., Wuts, P. G.(1991), *Protective Groups in Organic Synthesis* (3<sup>rd</sup> edition). John Wiley & Sons Inc.: New York.
- Hajipour, A. R., Mazloumi, G.(2002), "An efficient and simple procedure for preparation of esters and anhydrides from acyl chlorides in the presence of 1,4-diazabicyclo[2.2.2]octane (DABCO) under solvent free conditions", *Synthetic Communication* **32**, 23.

- Hofle, G., Steglich, V., Vorbruggen, H. (1978), "4-Dialkylaminopyridines as a highly active acylation catalysts", *Angewandte Chemie International Edition* **17**, 569.
- Iqbal, J., Srivastava, R. R.(1992), "Cobalt(II) chloride catalyzed acylation of alcohols with acetic anhydride: scope and mechanism", *Journal of Organic Chemistry* **57**, 2001.
- Iranpoor, N., Firouzabadi, H., Zolfigol, M. A.(1998), "Selective acetylation of primary alcohols: acetyl and formyl transfer reactions with copper(II) salts", *Synthetic Communication* **28**, 1923.
- Ishihara, K., Kurihara, H., Yamamoto, H.(1993), " An extremely simple, convenient, and selective method for acetylating primary alcohols in the presence of secondary alcohols", *Journal of Organic Chemistry* **58**, 3791.
- Ishihara, K., Kubota, M., Kurihara, H., Yamamoto, H. (1996), "Scandium trifluorometahnesulfonate as an extremely active lewis acid catalyst in acylation of alcohols with acid anhydrides and mixed anhydrides", *Journal of Organic Chemistry* **61**, 4560.
- Izumi, J., Shiina, I., Mukiyama, T.(1995), "An efficient esterification reaction between equimolar amounts of free carboxylic acids and alcohols by the combined use of octamethylcyclotetrasiloxane and a catalytic amount of Titanium(IV) Chloride Tris(trifluoromethanesulfonate)", *Chemical Letters* **24**, 141-142.
- Kovi, K. E., Wolf, C.(2008), "One-pot oxidative esterification and amidation of aldehydes", *Chemistry: A European Journal* **14**, 6302-6315.
- Li, T. S., Li, A. X.(1998), "Montmorillonite clay catalysis. Part 10; K-10 and KSF catalyzed acylation of alcohols, phenols, and amines: scope and limitation", *Journal of Chemical Society Perkin Trans. I* **12**, 1913.
- Maki, B. E., Scheidt, K. A.(2008), "N-Heterocyclic carbene catalyzed oxidation of unactivated aldehydes to esters", *Organic Letters* **10**, 4331-4334.
- Makone, S. S., Niwadange, S. N. (2010), "Collection of abstract: Second asia pacific conference on Ionic liquids and Green process, September 7-10, Dalian, China", page 78.
- Muzler, J., Trost, B. M., Fleming, I.(1991), *Comprehensive Organic Synthesis, Eds. Oxford: Pergamon Press* **6**, 323.
- Oltra, J. (2003), *Esterification: Methods, Reactions and Applications*, Wiley-VCH, Weinheim.
- Orita, A., Tanashi, C., Kakuda, A., Otera.(2000), "Highly efficient and versatile acylation of alcohols with Bi(OTf)<sub>3</sub> as a catalyst", *Angewandte Chemie International Edition* **39**, 2877.

- Pasha, M. A., Manjula, K.(2008), "TiO<sub>2</sub>: A simple and an efficient catalyst for esterification of phenols under solvent free conditions", *Indian Journal of Chemistry* **47b**, 597.
- Paul, S., Nande, P., Gupta, R.(2003), "PhCOCl-Py/Basic alumina as a versatile reagents for benzylation in solvent free conditions", *Molecules* **8**, 374.
- Ranu, B. C., Dey, S. S., Hajra, A.(2003), "Highly efficient acylation of alcohols, amines and thiols under solvent free and catalyst free conditions", *Green Chemistry* **5**, 44.
- Sano, T., Ohashi, K., Oriyama, T.(1999), "Remarkably fast acylation of alcohols with benzoyl chloride promoted by TMEDA", *Synthesis* **7**, 1141.
- Sarvari, M. H., Sharghi, H.(2005), "Zinc oxide (ZnO) as a new, highly efficient and reusable catalyst for acylation of alcohols, phenols and amines under solvent free conditions", *Tetrahedron* **61**, 10903.
- Schoenberg, A., Heck, R. F.(1974), "Palladium catalyzed amidation of aryl, heterocyclic and vinylic halides", *Journal of Organic Chemistry* **39**, 3327-3331.
- Scriven, E. F. V.(1983), "4-Dialkylaminopyridines: super acylation and alkylation catalysts", *Chemical Society Review* **12**, 129.
- Stevens, C. V., Frederik, E. A., Waes, V., Drabowicz, J., Cukalovic, A.(2012), "Efficient and catalyst free condensation of acid chlorides and alcohols using continuous flow", *Green chemistry* **14**, 2776-2779.
- Strazzolini, P., Giumanini, A. G., Verardo, G.(1994), "The reaction between acyl halides and alcohols: alkyl halides vs. ester formation", *Tetrahedron* **50**, 217.
- Tamaddon, F., Amrollahi, M. A., Sharafat, L.(2005), " A green protocol for the chemoselective O-acylation in the presence of zinc oxide as a heterogeneous, reusable and eco-friendly catalyst", *Tetrahedron Letters* **46**, 7841.
- Tayebbe, R., Cheravi, F., Mirzaee, M., Amini, M. M.(2010), "Commercial zinc oxide (Zn<sup>2+</sup>) as an efficient and environmentally benign catalyst for homogeneous benzylation of hydroxyl functional groups", *Chinese Journal of Chemistry* **28**, 1247.
- Taylor, E. C., McLay, G. W., McKillop, A.(1968), "Thallium in organic synthesis. II. Acylation, aroylation and tosylation of phenols and carboxylic acids", *Journal of American Chemical Society* **90**, 2422.
- Torregiani, E., Seu, G., Minassi, A., Appendino, G.(2005), "Cerium (III) chloride promoted chemoselective esterification of phenolic alcohols", *Tetrahedron Letters* **46**, 2193.



Travis, B. R., Sivakumar, M., Hollist, G. O., Borhan, B.(2003), “Facile oxidation of aldehydes to acids and esters with oxone”, *Organic Letters* **5**, 1031-1034.

Trost, B. M.(1995), “Atom economy- A challenge for organic synthesis: Homogeneous catalysis leads the way”, *Angewandte Chemie International Edition England* **34**, 259.

Vedejs, E., Daugulis, J.(1996), “Dual activation in the esterification of hindered alcohols with anhydrides using  $MgBr_2$  and a tertiary amine” *Journal of Organic Chemistry* **61**, 5702-5703.

Vedejs, E., Diver, S. T.(1993), “Tributylphosphine: a remarkable acylation catalyst”, *Journal of American Chemical Society* **115**, 3358.

Wu, X. F., Darcel, C.(2009), “Iron catalyzed one pot oxidative esterification of aldehydes”, *European Journal of Organic Chemistry* **8**, 1144-1147.

Xu, B., Liu, X., Haubrich, J., Friend, C. M.(2010), “Vapour-phase gold surface mediated coupling of aldehydes with methanol”, *Nature Chemistry* **2**, 61-65.


Yadav, V. K., Babu, K. G.(2004), “Reactions on a Solid Surface. A simple, Economical and efficient acylation of alcohols and amines over  $Al_2O_3$ ”, *Journal of Organic Chemistry* **69**, 577.

Yadav, V. K., Babu, K. G., Mittal, M.(2001), “ $KF-Al_2O_3$  is an efficient solid support reagent for the acetylation of amines, alcohols, and phenols, impeding effect of solvent on the reaction rate”, *Tetrahedron* **57**,7047.

Yoo, W. J., Li, C. J.(2006), “Highly stereoselective oxidative esterification of aldehydes with  $\alpha$ -dicarbonyl compounds”, *Journal of Organic Chemistry* **71**, 6266-6268.


Yoo, W. J., Li, C. J.(2007), “Copper catalyzed oxidative esterification of alcohols with aldehydes activated by lewis acids”, *Tetrahedron Letters* **48**, 1033-1035.

Zhdanov, R. I., Zhenodarova, S. M.(1975), “Chemical methods of oligonucleotide synthesis”, *Synthesis* **4**, 222.



Where, R= A kyl, benzyl groups


Scheme 1



Where  $n$  = number of carbon atoms

Scheme 2

### Reaction mechanism of the esterification reaction



**Table 1.** Un-catalyzed esterification of alcohols under solvent free and energy efficient conditions.

Entry	R	Alcohols : Acetyl Chloride	Reaction time(hr)	Yield[%] <sup>a</sup>
a	$C_6H_5CH_2$	1:2	1	98
b	$CH_3CH_2CH_2$	1:2	1.5	95
c	$CH_3(CH_2)_4$	1:2	1.5	95
d	$CH_3(CH_2)_5$	1:2	1.5	97
e	$CH_3(CH_2)_6$	1:4	2	96
f	$CH_3(CH_2)_7$	1:4	2	96
g	$CH_3(CH_2)_9$	1:4	2.5	95
h	$CH_2(CH_2)_2CH_2$	1:4	2	97
i	$CH_2(CH_2)_4CH_2$	1:4	2	98
j	$CH_2(CH_2)_{10}CH_2$	1:6	2.5	95
k	$C_3H_3$	1:2	1	96
l	$C_3H_5$	1:2	1	97
m	$CH_2CH_2OCH_2CH_2OCH_2CH_2$	1:4	1.5	95
n	$CH_2CH(OH)CH_2$	1:6	2	97
o	$4OHC_6H_5CH_2$	1:4	4	96

<sup>a</sup>Yield refer to the pure isolated product.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

## CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

### IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

