

Antonie van Leeuwenhoek
DOI 10.1007/s10482-010-9547-9

ORIGINAL PAPER

Description of a novel actinobacterium *Kocuria assamensis* sp. nov., isolated from a water sample collected from the river Brahmaputra, Assam, India

Chandandeep Kaur · Ishwinder Kaur ·
Revti Raichand · Tarun Chandra Bora ·
Shanmugam Mayilraj

Received: 3 November 2010 / Accepted: 22 December 2010
© Springer Science+Business Media B.V. 2011

Abstract A Gram-positive, pale yellow pigmented actinobacterium, strain S9-65^T was isolated from a water sample collected from the river Brahmaputra, Assam, India and subjected to a polyphasic taxonomic study. The physiological and biochemical properties, major fatty acids (anteiso-C15:0 and anteiso-C17:0), estimated DNA G+C content (69.2 mol %) and 16S rRNA gene sequence analysis showed that strain S9-65^T belonged to the genus *Kocuria*. Strain S9-65^T exhibited highest 16S rRNA gene sequence similarity with *Kocuria palustris*

(99.1%); however, the DNA–DNA relatedness value between strain S9-65^T and *K. palustris* was 20.6%. On the basis of differential phenotypic characteristics and genotypic distinctiveness, strain S9-65^T should be classified as representative of a novel species *Kocuria*, for which the name *Kocuria assamensis* is proposed. The type strain is S9-65^T (=MTCC 10622^T = DSM 23999^T).

Keywords DNA–DNA hybridization · FAME · 16S rRNA gene sequence

Institute of Microbial Technology Chandigarh and North East Institute of Science & Technology—a constituent laboratory of Council of Scientific and Industrial Research (CSIR), Government of India

Chandandeep Kaur, Ishwinder Kaur, Both authors have contributed equally to the study.

The GenBank accession number for the 16S rDNA sequence of *Kocuria assamensis* strain S9-65^T is HQ018931.

Electronic supplementary material The online version of this article (doi:[10.1007/s10482-010-9547-9](https://doi.org/10.1007/s10482-010-9547-9)) contains supplementary material, which is available to authorized users.

C. Kaur · I. Kaur · R. Raichand · S. Mayilraj (✉)
Microbial Type Culture Collection (MTCC) & Gene
Bank, Institute of Microbial Technology,
Chandigarh 160 036, India
e-mail: mayil@imtech.res.in

T. C. Bora
Department of Biotechnology, North East Institute
of Science & Technology, Jorhat 785 005, India

Introduction

The genus *Kocuria* was proposed by Stackebrandt et al. (1995) to accommodate phylogenetically distinct Actinobacteria formerly classified in the genus *Micrococcus*. The type species of the genus is *Kocuria rosea*. Members of the genus have been isolated from different sources such as air, fermented sea food, mammalian skin, soil, the rhizosphere, freshwater or seawater, marine sediment and desert soil (Kloos et al. 1974; Stackebrandt et al. 1995; Kovacs et al. 1999; Reddy et al. 2003; Kim et al. 2004; Tvrtzová et al. 2005; Li et al. 2006; Mayilraj et al. 2006a, b; Zhou et al. 2008; Tang et al. 2009; Seo et al. 2009). At present the genus *Kocuria* consists of 17 species with validly published names (<http://www.bacterio.cict.fr/k/kocuria.html>). Here, we describe the taxonomic status of an actinobacterium,

strain S9-65^T, isolated from a surface water sample collected from the river Brahmaputra, Assam, India (93° 08'–98° 36'E and 26° 30'–20° 45'N) by using a polyphasic approach.

Materials and methods

Strains, cultivation and phenotypic characterization

The strain S9-65^T was isolated by the dilution plate technique on tryptic soy agar medium (TSA; HiMedia, India) and incubated for 4 days at 30°C. To study its phenotypic characteristics, the isolate was routinely cultivated on TSA medium at 30°C and maintained as glycerol stocks at –70°C. The reference type strain *Kocuria palustris* strain TAGA 27^T MTCC 10490^T (=DSM 11925^T) was obtained from Microbial Type Culture Collection & Gene Bank (MTCC), Institute of Microbial Technology, Chandigarh, India. Colony morphology, cell morphology, motility and Gram's reaction of the strain were determined by using standard methods (Barrow and Feltham 1993; Murray et al. 1994; Smibert and Krieg 1994). Phenotypic characterization was performed using TSA as basal medium and strains were incubated at their optimum growth temperatures. Physiological tests such as growth at different temperatures (between 10 and 45°C), pH (using biological buffers; Na₂HPO₄/NaH₂PO₄, Na₂CO₃/NaHCO₃ for pH below 8 and Na₂HPO₄/NaOH for pH above 8), NaCl concentrations and acid production from various carbohydrates and other biochemical tests were performed as described (Smibert and Krieg 1994). The API ZYM and API 20NE micro test strips were used as per the instructions of the manufacturer (bioMérieux). Sensitivity of the strain to antibiotics was tested by using antibiotic susceptibility discs (HiMedia, India) after incubation of 48 h.

Chemotaxonomic characterization

Freeze-dried cells for chemotaxonomic analysis (except for the fatty acids study) were prepared by harvesting the bacterial cells in the late exponential phase following their growth in Tryptic Soy Broth (TSB; HiMedia, India) at 30°C for 2 days. Isoprenoid quinones were extracted and purified as described by

Saha et al. (2005). The purified quinones were separated by reversed phase HPLC (SCL-10AVP, Shimadzu) using the solvent system of acetonitrile and isopropanol in a ratio of 65:35 with a flow rate of 1 ml/min and monitored at a wavelength of 269 nm. For cellular fatty acid analysis, the strains were grown on tryptic soy agar medium at 30°C for 36 h and the fatty acid methyl ester analysis was performed by using Sherlock Microbial Identification System (MIDI, USA) as described previously (Sasser 1990; Pandey et al. 2002). Extraction of polar lipids was done based on the modified protocol of Bligh and Dyer (1959). Two-dimensional TLC was run for identification of polar lipids according to procedures described by Komagata and Suzuki (1987). Lipid spots were detected using the following spray reagents: molybdotriphosphoric acid (5% w/v) in absolute ethanol, molybdenum blue spray reagent (1.3% Sigma), ninhydrin (0.2% w/v) in acetone and anisaldehyde reagent (Sigma) for detection of total lipids, phospholipids, aminolipids and glycolipids respectively. The peptidoglycan structure was determined by using a hydrolysate of purified cell walls according to Schleifer (1985). The amino acids and peptides were separated by two-dimensional ascending TLC as described by Schleifer and Kandler (1972), with the modification that TLC on cellulose sheets (Merck 5577) was used instead of paper chromatography. The G+C content of genomic DNA was determined spectrophotometrically (Lambda 35, Perkin Elmer, Waltham, MA, USA) using thermal denaturation method (Mandel and Marmur 1968).

Determination of 16S rRNA gene sequence, phylogenetic analysis and genomic relatedness

For 16S rRNA gene sequencing the genomic DNA extraction and amplification was performed as described previously (Mayilraj et al. 2006a, b). Identification of phylogenetic neighbours and the calculation of pairwise 16S rRNA gene sequence similarities were achieved using the EzTaxon server (<http://www.eztaxon.org/>; Chun et al. 2007). The 16S rRNA gene sequence of S9-65^T and representative closely related species were retrieved from the Eztaxon server and aligned using the MEGA version 4.0 (Tamura et al. 2007). Phylogenetic trees were constructed using the neighbour-joining as well as maximum parsimony algorithms. Bootstrap analysis

was performed to assess the confidence limits of the branching. The G+C content of genomic DNA was determined spectrophotometrically (Lambda 35, Perkin Elmer, Waltham, MA, USA) using thermal denaturation method (Mandel and Marmur 1968). DNA–DNA hybridization was performed each time with freshly isolated genomic DNA and was repeated three times by the membrane filter method (Tourova and Antonov 1987).

Results and discussion

Phenotypic characteristics

Growth of the strain on TSA produced a yellow pigment after incubation on TSA for 2 days. The detailed differential phenotypic properties are shown in Table 1 and also mentioned in species description. Phenotypic data presented in the table indicated that strain S9-65^T differed from the closely related species at least by 29 characters which includes acid production from carbohydrates like adonitol, raffinose, rhamnose, cellobiose, arbinose and xylose, casein hydrolysis, nitrate reduction, hydrogen sulphide production were negative for strain S9-65^T and positive for the closely related strain *K. palustris* TAGA 27^T. There were major differences in oxidation of different carbon sources using biolog. Strain S9-65^T was sensitive to antibiotics (µg disc⁻¹) such as nitrofurantion (300), norfloxacin (10), polymyxin B (300), cephalothin (30) and oxacillin (5) in comparison to the closely related strain *K. palustris* which showed resistance to all of the above mentioned antibiotics.

Both strain S9-65^T and *K. palustris* TAGA 27^T were positive for growth at pH 8.0–10.0; NaCl concentration up to 7%; temperature between 20–37°C. Both strains were positive for citrate utilization, gelatin liquefaction; for enzyme activities of (using API ZYM) alkaline phosphatase, esterase (C4), esterase lipase (C8), α-chymotrypsin, acid phosphatase, naphthol-AS-B1-phosphohydrolase, β-galactosidase and negative for lipase (C14), β-glucuronidase and α-fucosidase. Both the strains are positive (using API 20 NE) for hydrolysis of esculin and gelatin, ONPG, nitrate reduction, urease, D-glucose fermentation, assimilation of glucose, citrate, D-mannitol, D-mannose and negative for indole production, MR-VP, growth at pH 5 and 12, 15% NaCl; starch

Table 1 Characters that differentiate strain S9-65^T along with the closest species *K. palustris* (MTCC 10490^T): 1, strain S9-65^T and 2, *K. palustris*

Characteristics	1	2
Growth at pH 11	+	–
Growth at 12°C	–	+
Casein hydrolysis	–	+
Enzyme assayed for (API ZYM)		
Leucine arylamidase	–	+
Valine arylamidase	–	+
Cystine	–	+
Trypsin	–	+
α-galactosidase	–	+
α-glucosidase	–	+
β-glucosidase	–	+
N-acetyl-β-glucosaminidase	–	+
α-mannosidase	–	+
Assimilation of (API 20 NE)		
Arabinose	–	+
N-acetyl-glucosamine	–	+
Potassium gluconate	–	+
Malic acid	–	+
Hydrogen sulphide production	–	+
Nitrate reduction	–	+
Acid production from		
Adonitol	–	+
Raffinose	–	+
Rhamnose	–	+
Cellobiose	–	+
Arabinose	–	+
Xylose	–	+

+ positive; – negative. All the data from present study

hydrolysis; arginine dihydrolase, assimilation of capric acid, adipic acid and phenylacetic acid. Both the strains were sensitive to antibiotics (µg disc⁻¹) novobiocin (30), streptomycin (10), ampicillin (10), methicillin (5), penicillin G (10 U), neomycin (30), triple sulphas (300), sulfonamide (300), kanamycin (30), trimethoprim (5), colistin (10), rifampicin (2), lincomycin (2), cepoxitin (30), gentamycin (10) and bacitracin (8) and resistance to optachin and oxytetracycline (30). Both strains were positive for acid production from sucrose, trehalose, salicin, dextrose, maltose, sorbitol, mannitol, mannose, fructose and inositol; negative for melibiose, dulcitol, lactose and galactose.

Chemotaxonomic characterization

The chemotaxonomic properties of strain S9-65^T (presented in the species description) were typical of members of the genus *Kocuria*. The major fatty acids (anteiso-C15:0 and anteiso-C17:0) detected in the novel strain (presented in the species description) are consistently found in members of the genus *Kocuria*. The fatty acid compositions of the reference strain *K. palustris* assayed were qualitatively similar, but quantitatively varied from those of the novel strain (Table 2). The major polar lipids were phosphatidylglycerol (PG), diphosphatidylglycerol (DPG) and four unknown phospholipids (PL) (Supplementary Figure 1). The major menaquinones detected for the strain S9-65^T were MK-9(H₂), 77.4%, and MK-8(H₂), 8.4.; MK-7(H₂), 5.1%, and MK-6(H₂), 5.3% occur in minor amounts. The diagnostic diamino acid in cell wall hydrolyzates was lysine with peptidoglycan type Lys-Ala₃ (type A3 α). The DNA G+C content of strain S9-65^T was estimated to be 69.2 mol%, a value within the range of 66–75 mol% reported for members of the genus *Kocuria* (Stackebrandt et al. 1995; Rainey et al. 1997; Kovacs et al. 1999; Reddy et al. 2003; Kim et al. 2004; Trzová et al. 2005; Mayilraj et al. 2006a, b; Li et al. 2006; Zhou et al. 2008).

Table 2 Cellular fatty acid composition of strain S9-65^T along with the closest species *K. palustris* (MTCC 10490^T): 1 strain S9-65^T and 2 *K. palustris*

Fatty acid	1	2
Iso C _{14:0}	3.11	3.06
Iso C _{15:0}	1.83	ND
Iso C _{16:0}	2.44	2.76
C _{14:0}	2.03	3.06
C _{16:0}	1.4	2.77
ANTEISO-C _{15:0}	77.97	68.06
ANTEISO-C _{17:0}	8.37	16.92
C _{18:1} 2OH	1.40	3.38
Summed feature 4*	1.46	ND

Data from present study. ND not detected

Summed feature 4* consists of C17: 1 iso I/anteiso B, which could not be separated by MIDI

C17:1 iso I/ anteiso B: iso I/ anteiso B are monounsaturated iso/anteiso branched C17 fatty acids where the exact location of the double bond is not known with “I” / “B” indicating the probability of several isomers like A, B, C, etc

Phylogenetic analysis and genomic relatedness

The almost complete sequence (1425 bp) of the 16S rRNA gene of strain S9-65^T was determined (GenBank accession no. HQ018931) and compared with those of other closely related taxa retrieved from the EzTaxon database. Sequence analysis revealed that strain S9-65^T shared highest 16S rRNA gene sequence identity with *K. palustris* TAGA 27^T (99.1%) and the remaining species in the genus *Kocuria* were in the range between 95.6 and 96.5% 16S rRNA gene sequence similarity. Based on the 16S rRNA gene sequence identity, the strain could be assigned to the genus *Kocuria*. The neighbour-joining phylogenetic tree (Fig. 1) as well as maximum parsimony algorithms (data not shown) demonstrated that strain S9-65^T formed a separate lineage along with the closely related species *K. palustris*. However, the DNA–DNA relatedness value between strain S9-65^T and the closely related taxon *K. palustris* TAGA 27^T (MTCC 10490^T) was 20.6% ($\pm 0.6\%$), which was well below the 70% threshold value recommended for the delineation of bacterial species (Stackebrandt and Goebel 1994). DNA–DNA relatedness values between strain S9-65^T and the remaining type strains of the genus *Kocuria* were not determined, since organisms with more than 3% 16S rRNA gene sequence dissimilarity are considered to belong to different genomic species (Wayne et al. 1987).

Conclusion

Based on the phenotypic and genotypic results, strain S9-65^T has to be regarded as a new species of *Kocuria*. Table 1 shows the main features that distinguish strain S9-65^T from the closely related taxon *K. palustris* TAGA 27^T (MTCC 10490^T). Therefore, from the polyphasic evidence gathered in this study it is concluded that strain S9-65^T represents a novel species of the genus *Kocuria*, for which the name *Kocuria assamensis* sp. nov. is proposed.

Description of *Kocuria assamensis* sp. nov.

Kocuria assamensis (as.sam.en'sis. N.L. masc. adj. *assamensis*, pertaining to Assam, a north-eastern state in India, where the type strain was isolated).




Fig. 1 Phylogenetic neighbour-joining tree based on 16S rRNA gene sequences showing the relationship between *Kocuria assamensis* S9-65^T and related members of the genus *Kocuria*. *Micrococcus luteus* NCTC 2665^T (CP001628) was

used as an out-group. Bootstrap values (expressed as percentages of 1000 replications) greater than 50% are given at nodes. Bar 0.5% sequence variation. GenBank accession numbers are given in parentheses

Cells are Gram-positive, non-spore forming, aerobic, motile, catalase-positive, oxidase-positive, short rods (0.8–1.0 × 2.0–2.5 μm). The colonies are pale yellow pigmented, circular, opaque and convex with smooth margin on TSA. Growth occurs at 25–37°C, optimal temperature is 30°C, at a pH of 6.5–11.0, optimal pH is 7.0 and at 2–7% (w/v) NaCl. The strain does not grow on Simmon's citrate and MacConkey agar and do not reduce nitrate to nitrite. Detailed phenotypic characteristics are given in Table 1. The predominant fatty acids are anteiso C15:0 and anteiso-C17:0. The major polar lipids are phosphatidylglycerol (PG), diphosphatidylglycerol (DPG) and four unknown phospholipids (PL), as shown in Supplementary Figure 1. The major menaquinones detected for the strain S9-65^T were MK-9(H₂), 77.4%, and MK-8(H₂), 8.4%; MK-7(H₂), 5.09%, and MK-6(H₂), 5.33%, occur in minor amounts. The diagnostic diamino acid in cell wall hydrolyzates was lysine with peptidoglycan type Lys-Ala₃ (type A3α). The DNA G+C content is 69.2 mol%. The type

strain S9-65^T (MTCC 10622^T = DSM 23999^T) was isolated from a water sample collected from the river Brahmaputra, Assam, India.

Acknowledgments We thank Mr Malkit Singh for his excellent technical assistance. This work was supported by CSIR (net work project NWP006) Government of India. This is IMTECH communication number 073/2010.

References

- Barrow GI, Feltham RKA (1993) Cowan and steel's manual for the identification of medical bacteria, 3rd edn. Cambridge University Press, Cambridge
- Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. Can J Biochem Physiol 37:911–917
- Chun J, Lee JH, Jung Y, Kim M, Kim S, Kim BK, Lim YW (2007) EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. Int J Syst Evol Microbiol 57:2259–2261
- Kim SB, Nedashkovskaya OI, Mikhailov VV, Han SK, Kim KO, Rhee MS, Bae KS (2004) *Kocuria marina* sp. nov., a

- novel actinobacterium isolated from marine sediment. *Int J Syst Evol Microbiol* 54:1617–1620
- Kloos WE, Tornabene TG, Schleifer KH (1974) Isolation and characterization of micrococci from human skin, including two new species: *Micrococcus lylae* and *Micrococcus kristinae*. *Int J Syst Bacteriol* 24:79–101
- Komagata K, Suzuki K (1987) Lipid and cell-wall analysis in bacterial systematics. *Methods Microbiol* 19:161–207
- Kovacs G, Burghardt J, Pradella S, Schumann P, Stackebrandt E, Marialigeti K (1999) *Kocuria palustris* sp. nov. and *Kocuria rhizophila* sp. nov., isolated from the rhizoplane of the narrow-leaved cattail (*Typha angustifolia*). *Int J Syst Bacteriol* 49:167–173
- Li WJ, Zhang YQ, Schumann P, Chen HH, Hozzein WN, Tian XP, Xu LH, Jiang CL (2006) *Kocuria aegyptia* sp. nov., a novel actinobacterium isolated from a saline, alkaline desert soil in Egypt. *Int J Syst Evol Microbiol* 56:733–737
- Mandel M, Marmur J (1968) Use of ultraviolet absorbance temperature profile for determining the guanine plus cytosine content of DNA. *Methods Enzymol* 12B:195–206
- Mayilraj S, Kroppenstedt RM, Suresh K, Saini HS (2006a) *Kocuria himachalensis* sp. nov., an actinobacterium isolated from the Indian Himalayas. *Int J Syst Evol Microbiol* 56:1971–1975
- Mayilraj S, Saha P, Suresh K, Saini HS (2006b) *Ornithinimicrobium kibberense* sp. nov., isolated from the Himalayas, India. *Int J Syst Evol Microbiol* 56:1657–1661
- Murray RGE, Doetsch RN, Robinow CF (1994) Determinative and cytological light microscopy. In: Gerhard P, Murray RGE, Wood WA, Krieg NR (eds) *Methods for general and molecular bacteriology*. American Society for Microbiology, Washington, DC, pp 21–41
- Pandey KK, Mayilraj S, Chakraborti T (2002) *Pseudomonas indica* sp. nov., a novel butane-utilizing species. *Int J Syst Evol Microbiol* 52:1559–1567
- Rainey FA, Nobre MF, Schumann P, Stackebrandt E, Da Costa MS (1997) Phylogenetic diversity of the deinococci as determined by 16S ribosomal DNA sequence comparison. *Int J Syst Bacteriol* 47:510–514
- Reddy GSN, Prakash JSS, Prabakar V, Matsumoto GI, Stackebrandt E, Shivaji S (2003) *Kocuria polaris* sp. nov., an orange-pigmented psychrophilic bacterium isolated from an Antarctic cyanobacterial mat sample. *Int J Syst Evol Microbiol* 53:183–187
- Saha P, Mondal AK, Mayilraj S, Krishnamurthi S, Bhattacharya A, Chakraborti T (2005) *Paenibacillus assamensis* sp. nov., a novel bacterium isolated from a warm spring in Assam, India. *Int J Syst Evol Microbiol* 55:2577–2581
- Sasser M (1990) Identification of bacteria by gas chromatography of cellular fatty acids, MIDI Technical Note 101. MIDI Inc, Newark, DE
- Schleifer KH (1985) Analysis of the chemical composition and primary structure of murein. *Methods Microbiol* 18:123–156
- Schleifer KH, Kandler O (1972) Peptidoglycan types of bacterial cell wall and their taxonomic implications. *Bacteriol Rev* 36:407–477
- Seo YB, Kim DE, Kim GD, Kim HW, Nam SW, Kim YT, Lee JH (2009) *Kocuria gwangalliensis* sp. nov., an actinobacterium isolated from seawater. *Int J Syst Evol Microbiol* 59:2769–2772
- Smibert RM, Krieg NR (1994) Phenotypic characterization. In: Gerhard P, Murray RGE, Wood WA, Krieg NR (eds) *Methods for general and molecular bacteriology*. American Society for Microbiology, Washington, DC, pp 607–654
- Stackebrandt E, Goebel BM (1994) Taxonomic note: a place for DNA–DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* 44:846–849
- Stackebrandt E, Koch C, Gvozdiak O, Schumann P (1995) Taxonomic dissection of the genus *Micrococcus*: *Kocuria* gen. nov., *Nesterenkonia* gen. nov., *Kytococcus* gen. nov., *Dermacoccus* gen. nov., and *Micrococcus* Cohn 1872 gen. emend. *Int J Syst Bacteriol* 45:682–692
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596–1599
- Tang SK, Wang Y, Lou K, Mao PH, Xu LH, Jiang CL, Kim CJ, Li WJ (2009) *Kocuria halotolerans* sp. nov., a novel actinobacterium isolated from a saline soil in China. *Int J Syst Evol Microbiol* 59:1316–1320
- Tourova TP, Antonov AS (1987) Identification of microorganisms by rapid DNA–DNA hybridization. *Methods Microbiol* 19:333–355
- Tvrzová L, Schumann P, Sedláček I, Páková Z, Spröer C, Verburg S, Kroppenstedt RM (2005) Reclassification of strain CCM 132, previously classified as *Kocuria varians*, as *Kocuria carniphila* sp. nov. *Int J Syst Evol Microbiol* 55:139–142
- Wayne LG, Brenner DJ, Colwell RR, Grimont PAD, Kandler O, Krichevsky MI, Moore LH, Moore WEC, Murray RGE, Stackebrandt E, Starr MP, Truper HG (1987) Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* 37:463–464
- Zhou G, Luo X, Tang Y, Zhang L, Yang Q, Qio Y, Fang C (2008) *Kocuria flava* sp. nov. and *Kocuria turfanesis* sp. nov., airborne actinobacteria isolated from Xinjiang, China. *Int J Syst Evol Microbiol* 58:1304–1307