

Mahanta, J.J., Chutia, M., Sarma, T.C.

Study on weed flora and their influence on patchouli (*Pogostemon cablin* Benth.) oil and patchoulol (2007) *Journal of Plant Sciences*, 2 (1), pp. 96-101. Cited 1 time.

Regional Research Laboratory (CSIR), Jorhat-785 006, Assam, India

#### Abstract

Experiments were conducted to study the weed flora and its influence on the yield of oil and Patchoulol by co-distillation of fresh Patchouli leaves with weed biomass at different proportions (0, 5, 10, 15 and 30%) during 2004-05 and 2005-06. The survey of weeds commonly growing in patchouli plantation was made and a total of 17 weed species were recorded. *Alternanthera sessile*, *Cynodon dactylon* and *Oxalis corniculata* exhibited 100% frequency in both the years. *A. sessile*, *C. dactylon* and *O. corniculata* had highest density during 1st year and the density of most of the weed species increased during 2nd year except *C. dactylon* and *A. sessile*. Co-distillation of fresh Patchouli leaves with weeds at the rate of 0, 5, 10, 15 and 30% yielded 0.70, 0.67, 0.65, 0.50 and 0.43% oil, respectively. It was observed that the oil yield decreased gradually with the increase in weed biomass. However, the percentage of patchoulol showed a different behaviour. It decreased at 15% (53.7) and 30% (50.4) and increased at 5% (56.5) and 10% (63.8) treatments. The oil extracted with weed biomass imports a weedy odour, which may decrease its commercial value. © Academic Journals Inc.

#### Author Keywords

Co-distillation; Oil yield; Patchouli; Patchoulol; Weed; Weedy odour

#### References

- Akhila, A., Nigam, M.C.  
GC-MS analysis of the essential oil of *Pogostemon cablin* (Patchouli oil)  
(1984) *Fitoterapia*, 55, pp. 363-365.
- Akhila, A., Tewari, R.  
Chemistry of patchouli oil: A review  
(1984) *CROMAP*, 6, pp. 38-54.
- Chapman, B.  
Effect of weed in wheat. Top Document  
(2000) *Agricultural Food and Rural Development*, p. 25.
- Kaul, P.N., Rao, B.R.R., Bhattacharaya, A.K., Singh, K., Singh, C.P.  
Effect of partial shade on essential oils of three geranium (*Pelargonium* species) cultivar  
(1997) *Indian Perfumer*, 41, pp. 1-4.
- Lawrence, B.M.  
Progress in essential oils- Patchouli oil  
(1976) *Perfum. Flavor*, 1, p. 20.
- Lawrence, B.M.  
Progress in essential oils- Patchouli oil  
(1981) *Perfum. Flavor*, 6, pp. 73-76.
- Lawrence, B.M.

- Progress in essential oils- Patchouli oil  
(1990) *Perfum. Flavor*, 15, pp. 76-77.
- Lawrence, B.M.  
Progress in essential oils- Patchouli oil  
(1995) *Perfum. Flavor*, 20, pp. 72-73.
  - Maheshari, M.L., Vasantha, K.T., Sharma, N., Chandel, K.P.S.  
Patchouli-an Indian Perspective  
(1993) *Ind. Perfum*, 37, pp. 9-11.
  - Maillet, J., Lopez, G.C.  
What criteria are relevant for predicting the invasive capacity of a new agricultural weed? The case of invasive American species in France  
(2000) *Weed Research*, 40, pp. 11-26.
  - Nikiforov, A., Jirovetz, L., Buchbauer, G., Raverdino, V.  
GC-FTIR and GC-MS in odour analysis of essential oils  
(1988) *Mikrochim. Acta*, 2, pp. 193-198.
  - Nikiforov, A., Jirovetz, L., Buchbauer, G.  
Synthesis of tertiary alcohol carbamates  
(1989) *Liebigs Ann. Chem*, 5, pp. 489-491.
  - Odum, E.P.  
(1971) *Fundamentals of Ecology*, p. 574.  
W.B. Saunders Company, Philadelphia, pp
  - Powell, K., Jutsum, A.R.  
Technical and commercial aspects of bio control products  
(1993) *Pest. Sci*, 37, pp. 315-321.
  - Rao, B.R.R., Kaul, P.N., Singh, K., Mallavarapu, G.R., Ramesh, S.  
Influence of Co-distillation with weed Biomass on yield and chemical composition of Rose-Scented Geranium (Pelargonium species) Oil  
(2005) *J. Essential Oil Res*, 17, pp. 41-43.
  - Sarma, P.C., Kanjilal, P.B.  
Effect of planting time and row spacing on growth yield and quality of Patchouli (Pogostemon patchouli Benth)  
(2000) *Adv. Plant Sci*, 13, pp. 201-204.
  - Sarma, A., Sarma, T.C.  
Patchouli oil recovery and effect of leaf ageing  
(2003) *Indian Perfumer*, 47, pp. 151-154.
  - Sharma, N.  
Observation of Patchouli (Pogostemon patchouli) through in vitro methods  
(1999) *Ind. Perfum*, 43, pp. 19-22.
  - Sharma, R.P., Singh, P., Maliwal, P.L.  
Effect of weed management and phosphorous level on yield and quality of Indian mustard (Brassica juncea)  
(2002) *Ind. J. Agric. Sci*, 72, pp. 461-463.

- Sugimura, Y., Ichikawa, Y., Otsuji, K., Fujita, M., Toi, N., Kamata, N., Del Rosario, R.M., Taga, G.L.  
An- Cultivarietal comparison of patchouli plants in relation to essential oil production and quality (1990) *Flavour and Fragrance J*, 5, pp. 109-114.
- Wirjahadja, S.D., Pancho, J.Y.  
Weed survey method and vegetation analysis (1978) *Biotrop. Tech. Bull*, 4, p. 20.
- Yadav, R.L., Mohan, R., Singh, D.V.  
Requirement of weed free period for optimum herb and oil yield of Java citronell (1981) *Ind. Perfum*, 25, pp. 47-52.

Document Type: Article

Source: Scopus

Copyright © 2008 [Elsevier B.V.](#) All rights reserved. Scopus ® is a registered trademark of Elsevier B.V.