

Mathematical Theory and Modeling  
 ISSN 2224-5804 (Paper) ISSN 2225-0522 (Online)  
 Vol.2, No.3, 2012

[www.iiste.org](http://www.iiste.org)



## Common Fixed Points of Weakly Reciprocally Continuous Maps using a Gauge Function

R. K. Bisht\*, Jay Singh, AjayGairola, R. U. Joshi

Department of Mathematics, D. S. B. Campus, Kumaun University, Nainital-263002, India

\* E-mail of the corresponding author: [ravindra.bisht@yahoo.com](mailto:ravindra.bisht@yahoo.com)

### Abstract

The aim of the present paper is to obtain a common fixed point theorem by employing the recently introduced notion of weak reciprocal continuity. We demonstrate that weak reciprocal continuity ensures the existence of fixed points under contractive conditions which otherwise do not ensure the existence of fixed points. Our result generalize and extend several well-known fixed point theorems due to Boyd and Wong (1969), Jungck(1976), Pant (1994) and Pathak et al (1997).

**Keywords:** Fixed point theorems, compatible maps, A-compatible maps, T-compatible maps, reciprocal continuity, weak reciprocal continuity

### 1. Introduction

The question of continuity of contractive maps in general and of continuity at fixed points in particular emerged with the publication of two research papers by Kannan (1968, 1969) in 1968 and 1969 respectively. These two papers generated unprecedented interest in the fixed point theory of contractive maps which, in turn, resulted in vigorous research activity on the existence of fixed points of contractive maps and the question of continuity of contractive maps at their fixed points turned into an open question.

In (1998), 30 years after Kannan's celebrated papers, Pant (1998) introduced the notion of reciprocal continuity for a pair of mappings and as an application of this concept obtained the first fixed point theorem, in which the common fixed point was a point of discontinuity.

#### Definition 1.1

Two self mappings  $A$  and  $T$  of a metric space  $(X, d)$  are defined to be reciprocally continuous iff  $\lim_n ATx_n = At$  and  $\lim_n TAx_n = Tt$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_n Ax_n = \lim_n Tx_n = t$  for some  $t$  in  $X$ .

The notion of reciprocal continuity has been employed by many researchers in diverse settings to establish fixed point theorems which admit discontinuity at the fixed point. Imdad *et al* (2009) used this concept in the setting of non-self mappings. Singh and Mishra (2002) have used reciprocal continuity to establish general fixed point theorems for hybrid pairs of single valued and multi-valued maps. P.Balasubramaniam *et al* (2002) extended the study of reciprocal continuity to fuzzy metric spaces. Suneel Kumar *et al* (2008) studied this concept in the setting of probabilistic metric space. S. Murlishankar *et al* (2009) established a common fixed point theorem in an intuitionistic fuzzy metric space using contractive condition of integral type. Chugh *et al* (2003) and Kumar *et al* (2002) have, in the setting of metric spaces, obtained interesting fixed point theorems which do not force the map to be continuous at the fixed point.

The notion of reciprocal continuity is mainly applicable to compatible mapping satisfying contractive conditions. To widen the scope of the study of fixed points from the class of compatible mappings

satisfying contractive conditions to a wider class including compatible as well as noncompatible mappings satisfying contractive, nonexpansive or Lipschitz type condition Pant *et al*(2011) generalized the notion of reciprocal continuity by introducing the new concept of weak reciprocal continuity as follows:

*Definition 1.2*

Two self mappings  $A$  and  $T$  of a metric space  $(X, d)$  are defined to be weak reciprocally continuous iff  $\lim_n ATx_n = At$  or  $\lim_n TAx_n = Tt$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_n Ax_n = \lim_n Tx_n = t$  for some  $t$  in  $X$ .

Jungck(1986) generalized the notion of weakly commuting maps by introducing the concept of compatible maps.

*Definition 1.3*

Two self mappings  $A$  and  $T$  of a metric space  $(X, d)$  are compatible iff  $\lim_n d(ATx_n, TAx_n) = 0$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_n Ax_n = \lim_n Tx_n = t$  for some  $t$  in  $X$ .

In (1993), Jungck et al (1993) further generalized the concept of weakly commuting mappings by introducing the notion of compatible of type (A).

*Definition 1.4*

Two self mappings  $A$  and  $T$  of a metric space  $(X, d)$  are compatible of type (A) iff  $\lim_n d(AAx_n, TAx_n) = 0$  and  $\lim_n d(ATx_n, TTx_n) = 0$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_n Ax_n = \lim_n Tx_n = t$  for some  $t$  in  $X$ .

In (1998), Pathak et al (1998) generalized the notion of compatibility of type (A) by introducing the two analogous notions of compatibility, i.e., A-compatible and T-compatible.

*Definition 1.5*

Two self mappings  $A$  and  $T$  of a metric space  $(X, d)$  are A-compatible iff  $\lim_n d(ATx_n, TTx_n) = 0$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_n Ax_n = \lim_n Tx_n = t$  for some  $t$  in  $X$ .

*Definition 1.6*

Two self mappings  $A$  and  $T$  of a metric space  $(X, d)$  are T-compatible iff  $\lim_n d(AAx_n, TAx_n) = 0$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_n Ax_n = \lim_n Tx_n = t$  for some  $t$  in  $X$ .

It may be noted that the notions compatible and A-compatible (or T-compatible, compatible mappings of type (A)) are independent to each other. If both  $A$  and  $T$  are continuous, then all the analogous notions of compatibility including compatibility are equivalent to each other.

As an application of weak reciprocal continuity we prove a common fixed point theorem for a  $\Phi$ -contractive condition that extend the scope of the study of common fixed point theorems from the class of compatible or analogous compatible continuous mappings to a wider class of mappings which also includes discontinuous mappings.

**1. Main Results**

**Theorem 2.1**

Let  $A$  and  $T$  be weakly reciprocally continuous self mappings of a complete metric space  $(X, d)$  satisfying

(i)  $AX \subset TX$

(ii)  $d(Ax, Ay) \leq \Phi (\max\{ d(Tx, Ty), d(Ax, Tx), d(Ay, Ty), [d(Ax, Ty) + d(Ay, Tx)]/2 \})$ ,

where  $\Phi: R_+ \rightarrow R_+$  denotes an upper semi continuous function such that  $\Phi(t) < t$  for each  $t > 0$ .

If  $A$  and  $T$  are either compatible or A-compatible or T-compatible then  $A$  and  $T$  have a unique common fixed point.

**Proof.**

Let  $x_0$  be any point in  $X$ . Define sequences  $\{x_n\}$  and  $\{y_n\}$  in  $X$  given by the rule

$$y_n = Ax_n = Tx_{n+1}.$$

This can be done since  $AX \subset TX$ . Then using (ii) we obtain

$$\begin{aligned} d(y_n, y_{n+1}) = d(Ax_n, Ax_{n+1}) &\leq \Phi(\max\{d(Tx_n, Tx_{n+1}), d(Ax_n, Tx_n), d(Ax_{n+1}, Tx_{n+1}), \\ &\quad [d(Ax_n, Tx_{n+1}) + d(Ax_{n+1}, Tx_n)]/2\}) \\ &= \Phi(d(y_{n-1}, y_n) < d(y_{n-1}, y_n)). \end{aligned}$$

Thus  $d(y_n, y_{n+1}) \leq (d(y_{n-1}, y_n) < d(y_{n-1}, y_n))$ .

$$(2.1)$$

Similarly,  $d(y_{n-1}, y_n) \leq (d(y_{n-2}, y_{n-1}) < d(y_{n-2}, y_{n-1}))$ .

$$(2.2)$$

Thus we see that  $\{d(y_n, y_{n+1})\}$  is strictly decreasing sequence of positive numbers and hence tends to a limit  $r \geq 0$ .

Suppose  $r > 0$ , then relation (2.1) on making  $n \rightarrow \infty$  and in view of upper semi continuity of  $\Phi$  yields  $r \leq \Phi(r) < r$ , a contradiction. Hence  $r = \lim_{n \rightarrow \infty} d(y_n, y_{n+1}) = 0$ . We claim  $\{y_n\}$  is a Cauchy sequence.

Suppose it is not. Then there exists an  $\varepsilon > 0$  and a subsequence  $\{y_{n_i}\}$  of  $\{y_n\}$  such that  $d(y_{n_i}, y_{n_i+1}) > 2\varepsilon$ .

Since  $\lim_n d(y_n, y_{n+1}) = 0$ , there exists integers  $m_i$  satisfying  $n_i < m_i < n_{i+1}$  such that

$d(y_{n_i}, y_{m_i}) \geq \varepsilon$ . If not then

$$d(y_{n_i}, y_{m_i+1}) \leq d(y_{n_i}, y_{n_i+1}) + d(y_{n_i+1}, y_{m_i+1}) < \varepsilon + d(y_{n_i+1}, y_{m_i+1}) < 2\varepsilon,$$

a contradiction. If  $m_i$  be the smallest integer such that  $d(y_{n_i}, y_{m_i}) > \varepsilon$ , then

$$\begin{aligned} \varepsilon &\leq d(y_{n_i}, y_{m_i}) < d(y_{n_i}, y_{m_i-2}) + d(y_{m_i-2}, y_{m_i-1}) + d(y_{m_i-1}, y_{m_i}) \\ &< \varepsilon + d(y_{m_i-2}, y_{m_i-1}) + d(y_{m_i-1}, y_{m_i}), \end{aligned}$$

that is, there exists integer  $m_i$  satisfying  $n_i < m_i < n_{i+1}$  such that

$$d(y_{n_i}, y_{m_i}) \geq \varepsilon \text{ and } \lim_n d(y_{n_i}, y_{m_i}) = \varepsilon \quad (2.3)$$

without loss of generality we can assume that  $n_i$  is odd and  $m_i$  is even. Now by virtue of (1), we have

$$d(y_{n_i+1}, y_{m_i+1}) \leq \Phi((d(y_{n_i}, y_{m_i}) + d(y_{n_i}, y_{n_i+1})).$$

Now on letting  $n_i \rightarrow \infty$  and in view of (2.3) and upper semi continuity of  $\Phi$ , the above relation yields  $\varepsilon \leq \Phi(\varepsilon) < \varepsilon$ , a contradiction. Hence  $\{y_n\}$  is a Cauchy sequence. Since  $X$  is complete, there exists a point  $u$  in  $X$  such that  $y_n \rightarrow u$  as  $n \rightarrow \infty$ . Moreover,

$$y_n = Ax_n = Tx_{n+1} \rightarrow u. \quad (2.4)$$

Suppose that  $A$  and  $T$  are compatible mappings. Now, weak reciprocal continuity of  $A$  and  $T$  implies that  $\lim_n ATx_n = Au$  or  $\lim_n TAx_n = Tu$ . Let  $\lim_n TAx_n = Tu$ . Then compatibility of  $A$  and  $T$  yields  $\lim_n d(ATx_n, TAx_n) = 0$ , i.e.,  $\lim_n ATx_n = Tu$ . By virtue of (2.4) this yields  $\lim_n ATx_{n+1} = \lim_n AAx_n = Tu$ . If  $Au \neq Tu$  then using (ii) we get  $d(Au, AAx_n) \leq \Phi(\max\{d(Tu, TAx_n), d(Au, Tu), d(AAx_n, TAx_n), [d(Au, TAx_n) + d(AAx_n, Tu)]/2\})$ . On letting  $n \rightarrow \infty$  we get  $d(Au, Tu) \leq \Phi(d(Au, Tu)) < d(Au, Tu)$ , a contradiction. Hence  $Au = Tu$ . Again compatibility of  $A$  and  $T$  implies commutativity at coincidence points. Hence  $ATu = T Au = AAu = TTu$ . Further, if  $Au \neq AAu$  then in view of (ii), we get  $d(Au, AAu) \leq \Phi(\max\{d(Tu, T Au), d(Au, Tu), d(AAu, T Au), [d(Au, T Au) + d(AAu, Tu)]/2\}) = \Phi(d(Au, AAu)) < d(Au, AAu)$ , a contradiction. Hence  $Au = AAu = T Au$ . Therefore,  $Au = Tu$  is a common fixed point of  $A$  and  $T$ .

Next suppose that  $\lim_n ATx_n = Au$ . Then  $AX \subset TX$  implies that  $Au = Tv$  for some  $v$  in  $X$  and  $\lim_n ATx_n = Au = Tv$ . Compatibility of  $A$  and  $T$  implies,  $\lim_n TAx_n = Tv$ . By virtue of (2.4) this also yields  $\lim_n ATx_{n+1} = \lim_n AAx_n = Tv$ . If  $Av \neq Tv$  then using (ii) we get  $d(Av, AAx_n) \leq \Phi(\max\{d(Tv, TAx_n), d(Av, Tv), d(AAx_n, TAx_n), [d(Av, TAx_n) + d(AAx_n, Tv)]/2\})$ . On letting  $n \rightarrow \infty$  we get  $d(Av, Tv) \leq \Phi(d(Av, Tv)) < d(Av, Tv)$ , a contradiction. Hence  $Av = Tv$ . Compatibility of  $A$  and  $T$  implies commutativity at coincidence points. Hence  $ATv = T Av = AA v = TT v$ . Further, if  $Av \neq AA v$  then in view of (ii), we get  $d(Av, AA v) \leq \Phi(d(Av, AA v)) < d(Av, AA v)$ , a contradiction. Hence  $Av = AA v = T Av$ . Therefore,  $Av = Tv$  is a common fixed point of  $A$  and  $T$ .

Now, suppose that  $A$  and  $T$  are  $A$ -compatible mappings. Weak reciprocal continuity of  $A$  and  $T$  implies that  $\lim_n ATx_n = Au$  or  $\lim_n TAx_n = Tu$ . Let  $\lim_n TAx_n = Tu$ . Then  $A$ -compatibility of  $A$  and  $T$  yields  $\lim_n d(ATx_n, TTx_n) = 0$ . By virtue of  $\lim_n TTx_{n+1} = \lim_n TAx_n = Tu$  this yields  $\lim_n ATx_n = Tu$ . If  $Au \neq Tu$  then using (ii) we get  $d(Au, ATx_n) \leq \Phi(\max\{d(Tu, TTx_n), d(Au, Tu), d(ATx_n, TTx_n), [d(Au, TTx_n) + d(ATx_n, Tu)]/2\})$ . On letting  $n \rightarrow \infty$  we get  $d(Au, Tu) \leq \Phi(d(Au, Tu)) < d(Au, Tu)$ , a contradiction. Hence  $Au = Tu$ . Again  $A$ -compatibility implies commutativity at coincidence points. Hence  $ATu = TAU = AAu = TTu$ . Further, if  $Au \neq AAu$  then in view of (ii), we get  $d(Au, AAu) \leq \Phi(\max\{d(Tu, TAU), d(Au, Tu), d(AAu, TAU), [d(Au, TAU) + d(AAu, Tu)]/2\}) = \Phi(d(Au, AAu)) < d(Au, AAu)$ , a contradiction. Hence  $Au = AAu = TAU$ . Therefore,  $Au = Tu$  is a common fixed point of  $A$  and  $T$ .

Next suppose that  $\lim_n ATx_n = Au$ . Then  $AX \subset TX$  implies that  $Au = Tv$  for some  $v$  in  $X$  and  $\lim_n ATx_n = Au = Tv$ .  $A$ -Compatibility of  $A$  and  $T$  implies,  $\lim_n TTx_n = Tv$ . If  $Av \neq Tv$  then using (ii) we get  $d(Av, ATx_n) \leq \Phi(\max\{d(Tv, TTx_n), d(Av, Tv), d(ATx_n, TTx_n), [d(Av, TTx_n) + d(ATx_n, Tv)]/2\})$ . On letting  $n \rightarrow \infty$  we get  $d(Av, Tv) \leq \Phi(d(Av, Tv)) < d(Av, Tv)$ , a contradiction. Hence  $Av = Tv$ .  $A$ -Compatibility implies commutativity at coincidence points. Hence  $ATv = TAV = AAu = TTv$ . Further, if  $Av \neq AAu$  then in view of (ii), we get  $d(Av, AAu) \leq \Phi(d(Av, AAu)) < d(Av, AAu)$ , a contradiction. Hence  $Av = AAu = TAV$ . Therefore,  $Av = Tv$  is a common fixed point of  $A$  and  $T$ .

Finally, suppose that  $A$  and  $T$  are  $T$ -compatible mappings. Now, weak reciprocal continuity of  $A$  and  $T$  implies that  $\lim_n ATx_n = Au$  or  $\lim_n TAx_n = Tu$ . Let  $\lim_n TAx_n = Tu$ . Then  $T$ -compatibility of  $A$  and  $T$  yields  $\lim_n d(TAx_n, AAx_n) = 0$ , i.e.,  $\lim_n AAx_n = Tu$ . If  $Au \neq Tu$  then using (ii) we get  $d(Au, AAx_n) \leq \Phi(\max\{d(Tu, TAx_n), d(Au, Tu), d(AAx_n, TAx_n), [d(Au, TAx_n) + d(AAx_n, Tu)]/2\})$ . On letting  $n \rightarrow \infty$  we get  $d(Au, Tu) \leq \Phi(d(Au, Tu)) < d(Au, Tu)$ , a contradiction. Hence  $Au = Tu$ . Again  $T$ -compatibility implies commutativity at coincidence points. Hence  $ATu = TAU = AAu = TTu$ . Further, if  $Au \neq AAu$  then in view of (ii), we get  $d(Au, AAu) \leq \Phi(\max\{d(Tu, TAU), d(Au, Tu), d(AAu, TAU), [d(Au, TAU) + d(AAu, Tu)]/2\}) = \Phi(d(Au, AAu)) < d(Au, AAu)$ , a contradiction. Hence  $Au = AAu = TAU$ . Therefore,  $Au = Tu$  is a common fixed point of  $A$  and  $T$ .

Next suppose that  $\lim_n ATx_n = Au$ . Then  $AX \subset TX$  implies that  $Au = Tv$  for some  $v$  in  $X$  and  $\lim_n ATx_n = Au = Tv$ . By virtue of (2.4) this yields  $\lim_n ATx_{n+1} = \lim_n AAx_n = Au = Tv$ .  $T$ -Compatibility of  $A$  and  $T$  implies,  $\lim_n TAx_n = Tv$ . If  $Av \neq Tv$  then using (ii) we get  $d(Av, AAx_n) \leq \Phi(\max\{d(Tv, TAx_n), d(Av, Tv), d(AAx_n, TAx_n), [d(Av, TAx_n) + d(AAx_n, Tv)]/2\})$ . On letting  $n \rightarrow \infty$  we get  $d(Av, Tv) \leq \Phi(d(Av, Tv)) < d(Av, Tv)$ , a contradiction. Hence  $Av = Tv$ .  $T$ -Compatibility implies commutativity at coincidence points. Hence  $ATv = TAV = AAu = TTv$ . Further, if  $Av \neq AAu$  then in view of (ii), we get  $d(Av, AAu) \leq \Phi(d(Av, AAu)) < d(Av, AAu)$ , a contradiction. Hence  $Av = AAu = TAV$ . Therefore,  $Av = Tv$  is a common fixed point of  $A$  and  $T$ . Uniqueness of the common fixed point theorem follows easily in each of the three cases.

We now furnish an example to illustrate Theorems 2.1.

**Example 2.1**

Let  $X = [2, 20]$  and  $d$  be the usual metric on  $X$ . Define  $A$  and  $T: X \rightarrow X$  as follows

$$Ax = 2 \text{ if } x = 2 \text{ or } x > 5, Ax = 6 \text{ if } 2 < x \leq 5,$$

$$Tx = 2, Tx = 12 \text{ if } 2 < x \leq 5, Tx = x - 3 \text{ if } x > 5.$$

Then  $A$  and  $T$  satisfy all the conditions of Theorem 2.1 and have a common fixed point at  $x = 2$ . It can be verified in this example that the mappings  $A$  and  $T$  are  $T$ -compatible. It can also be noted that  $A$  and  $T$  are weakly reciprocally continuous. To see this, let  $\{x_n\}$  be a sequence in  $X$  such that  $fx_n \rightarrow t, gx_n \rightarrow t$  for some  $t$ . Then  $t=2$  and either  $\{x_n\}=2$  for each  $n$  or  $\{x_n\}=5+ \epsilon_n$  where  $\epsilon \rightarrow 0$  as  $n \rightarrow \infty$ . If  $x_n=2$  for each  $n$ ,  $ATx_n \rightarrow 2 = A2$  and  $TAx_n \rightarrow 2 = T2$ . If  $x_n=5+ \epsilon_n$ , then  $Ax_n \rightarrow 2, Tx_n \rightarrow 2, ATx_n \rightarrow 6 \neq A2$  and  $TAx_n \rightarrow 2 = T2$ . Thus,  $\lim_n ATx_n = T2$  but  $\lim_n ATx_n \neq A2$ . Hence  $A$  and  $T$  are weakly reciprocally continuous. It is also obvious that  $A$  and  $T$  are not reciprocally continuous mappings.

**Remark 2.1**

Theorem 2.1 contains proper generalizations of many important fixed point theorems, we mention only those due to Boyd and Wong (1969), Jungck (1976), Pant (1994) and Pathak et al (1997).

As a direct consequence of the above theorem we get the following corollary.

*Corollary 2.1*

Let  $A$  and  $T$  be reciprocally continuous self mappings of a complete metric space  $(X, d)$  satisfying

(i)  $AX \subset TX$

(ii)  $d(Ax, Ay) \leq \Phi (\max\{ d(Tx, Ty), d(Ax, Tx), d(Ay, Ty), [d(Ax, Ty) + d(Ay, Tx)]/2 \})$ ,

where,  $\Phi: R_+ \rightarrow R_+$  denotes an upper semi continuous function such that  $\Phi(t) < t$  for each  $t > 0$ .

If  $A$  and  $T$  are either compatible or  $A$ -compatible or  $T$ -compatible then  $A$  and  $T$  have a unique common fixed point.

If we let  $\Phi(t) = kt$ ,  $0 \leq k < 1$ , then we get the following corollaries:

*Corollary 2.2*

Let  $A$  and  $T$  be weakly reciprocally continuous self mappings of a complete metric space  $(X, d)$  satisfying

(i)  $AX \subset TX$

(ii)  $d(Ax, Ay) \leq k(\max\{ d(Tx, Ty), d(Ax, Tx), d(Ay, Ty), [d(Ax, Ty) + d(Ay, Tx)]/2 \})$ ,  $0 \leq k < 1$ ,

If  $A$  and  $T$  are either compatible or  $A$ -compatible or  $T$ -compatible then  $A$  and  $T$  have a unique common fixed point.

*Corollary 2.3*

Let  $A$  and  $T$  be reciprocally continuous self mappings of a complete metric space  $(X, d)$  satisfying

(i)  $AX \subset TX$

(ii)  $d(Ax, Ay) \leq k(\max\{ d(Tx, Ty), d(Ax, Tx), d(Ay, Ty), [d(Ax, Ty) + d(Ay, Tx)]/2 \})$ ,  $0 \leq k < 1$ ,

If  $A$  and  $T$  are either compatible or  $A$ -compatible or  $T$ -compatible then  $A$  and  $T$  have a unique common fixed point.

**References**

- Boyd D. W. and Wong J. S.(1969), "On nonlinear contractions", *Proc. Amer. Math. Soc.***20**, 458-464.  
 Balasubramanian P., Murlishankar S. M. and Pant R. P.(2002), "Common fixed points of four mappings in a fuzzy metric space", *J. Fuzzy Math.*, **10-2**,379-384.  
 ChughRenu and Kumar S.(2003), "Minimal commutativity and common fixed points", *J. Indian Math. Society*,**70**,169-177.  
 Imdad M., and Ali Javid(2009), "Reciprocal continuity and common fixed points of nonselfmappings", *Taiwanese Journal of Mathematics*, **13(5)**, 1457-1473.  
 Jungck, G.(1976), "Commuting mappings and fixed point", *Amer. Math. Monthly*,**83**, 261-263.  
 Jungck, G.(1976), "Compatible mappings and common fixed points", *Internat. J. Math. Sci.* **9**, 771-779.  
 Kannan R.(1968), "Some results on fixed points", *Bull. Cal. Math. Soc.***60**, 71-76.  
 Kannan R.(1969), "Some results on fixed points – II", *Amer. Math. Monthly***76**, 405-408.  
 Kumar Sanjay and ChughRenu (2002), "Common fixed point theorems using minimal commutativity and reciprocal continuity in metric space", *Scientiae Math. Japonicae*, **56**, 269-275.  
 Kumar Suneel and Pant B. D.(2008), "A common fixed point theorem in probabilistic metric space using implicit relation", *Filomat*, Vol. **22(2)**, 43-52.  
 Muralisankar S. and Kalpana G.(2009), "Common fixed point theorem in intuitionistic fuzzy metric space using general contractive condition of integral type", *Int. J. Contemp. Math. Sciences*, **4(11)**, 505 – 518.  
 Pant R. P. (1998), "Common fixed points of four mappings", *Bull. Cal. Math. Soc.***90**, 281-286.  
 Pant R. P. (1999), "A common fixed point theorem under a new condition", *Indian J. pure appl. Math.*,**30(2)**, 147-152.  
 Pant R. P.(1994), "Common fixed points of noncommuting mappings:.", *J. Math. Anal. Appl.***188**, 436-440.  
 Pant R. P.(1999), "Discontinuity and fixed points", *J. Math. Anal. Appl.***240**, 284-289.  
 Pant R. P., Bisht R. K. and Arora D.(2011), "Weak reciprocal continuity and fixed point theorems", *Ann*



*Univ Ferrara (Springer)*, **57**, 181–190.

Pathak H. K., Cho Y. J. and Kang S. M. (1997), “Remarks on R-weakly commuting mappings and common fixed point theorems”, *Bull. Korean Math. Soc.* **34**, 247-257.

Pathak H. K. and Khan M. S. (1997), “A comparison of various types of compatible maps and common fixed points”, *Indian J. Pure Appl. Math.* **28(4)**, 477–485.

Singh S. L. and Mishra S. N. (2002), “Coincidence and fixed points of reciprocally continuous and compatible hybrid maps”, *Internat. J. Math. and Math. Sci.* **30**, 627-635.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

### **IISTE Knowledge Sharing Partners**

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

