

Some Special Artex Spaces Over Bi-monoids

K.Muthukumaran (corresponding author)

Assistant Professor

PG and Research Department Of Mathematics, Saraswathi Narayanan College, Perungudi

Madurai-625022,Tamil Nadu, India

Mobile Phone : 9486727007 E-mail : nkmuthukumaran@gmail.com

M.Kamaraj

Associate Professor

Department Of Mathematics, Government Arts College, Melur, Madurai-625106,

Tamil Nadu, India.

E-mail : kamarajm17366@rediffmail.co.in

ABSTRACT

We introduce a Special Artex Space over a bi-monoid namely Distributive Artex space over a bi-monoid. We give some examples of Distributive Artex space over bi-monoids. We prove the Cartesian product of any two Distributive Artex Spaces over a bi-monoid is a Distributive Artex Space over the bi-monoid. Also we prove the Cartesian product of a finite number of Distributive Artex Spaces over a bi-monoid is a Distributive Artex Space over the bi-monoid. We prove under the Artex space homomorphism $f : A \rightarrow B$, the homomorphic image of a SubArtex Space of an Artex space A over a bi-monoid is a SubArtex space of B . We prove the homomorphic image of a Distributive Artex Space over a bi-monoid is a Distributive Artex Space over the bi-monoid. We prove a SubArtex space of a Distributive Artex space over a bi-monoid is a Distributive Artex space over the bi-monoid. We solve three problems on Bounded Artex spaces over bi-monoids. 1. A SubArtex space of a Lower Bounded Artex space over a bi-monoid need not be a Lower Bounded Artex space over the bi-monoid. 2. A SubArtex space of an Upper Bounded Artex space over a bi-monoid need not be an Upper Bounded Artex space over the bi-monoid and 3. A SubArtex space of a Bounded Artex space over a bi-monoid need not be a Bounded Artex space over the bi-monoid.

Keywords : Distributive Artex space, Bounded Artex space, Homomorphic image

1.INTRODUCTION

The theory of Groups is one of the richest branches of abstract algebra. Groups of transformations play an important role in geometry. A more general concept than that of a group is that of a semi-group. The aim of considering semi-groups is to provide an introduction to the theory of rings. Two binary operations are defined for a ring. With respect to the first operation written in the order, it is an abelian group and with respect to the second one, it is enough to be a semi-group. We study many examples of rings and we have so many results in rings. However, there are many spaces or sets which are monoids with respect to two or more operations. This motivated us to define a more general concept bi-monoid than that of a ring. So we can have many algebraic systems which are bi-monoids, but not rings. With these concepts in mind we enter into Lattices. In Discrete Mathematics Lattices and Boolean algebra have important applications in the theory and design of computers. There are many other areas such as engineering and science to which Boolean algebra is applied. Boolean Algebra was introduced by George Boole in 1854. A more general algebraic system is the lattice. A Boolean Algebra is introduced as a special lattice. The study of Lattices and Boolean algebra together with our bi-monoid

motivated us to bring our papers titled “Artex Spaces over Bi-monoids”, “Subartex Spaces Of an Artex Space Over a Bi-monoid”, and “Bounded Artex Spaces Over Bi-monoids and Artex Space Homomorphisms”. As an another development of it now, we introduce a Special Artex Space over a bi-monoid namely Distributive Artex space over a bi-monoid. These Distributive Artex spaces over bi-monoids and the Propositions, we prove here, are important results for further development. The examples of Distributive Artex spaces over bi-monoids are interesting and useful. The main part under consideration deals with Distributive Artex spaces over bi-monoids, SubArtex spaces of Bounded Artx spaces over bi-monoids. The principal concepts that we consider here are those of homomorphism and, epimorphism. As a result, the theory of Artex spaces over bi-monoids like the theory of Lattices and Boolean Algebra, will, in future, play a good role in many fields especially in science and engineering and in computer fields.

2.Preliminaries

2.1.1 Definition : Distributive Lattice : A lattice (L, \wedge, \vee) is said to be a distributive lattice if

for any $a, b, c \in L$, (i) $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$ (ii) $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$

2.1.2 Definition : Bi-monoid : An algebraic system $(M, +, \cdot)$ is called a Bi-monoid if

1. $(M, +)$ is a monoid 2. (M, \cdot) is a monoid

and 3 (i) $a \cdot (b+c) = a \cdot b + a \cdot c$ and (ii) $(a+b) \cdot c = a \cdot c + b \cdot c$, for all $a, b, c \in M$.

Note : The identity elements of $(M, +, \cdot)$ with respect to $+$ and \cdot are denoted by 0 and 1 respectively.

2.1.3 Definition : Artex Space Over a Bi-monoid : A non-empty set A is said to be an Artex Space Over a Bi-monoid $(M, +, \cdot)$ if 1. (A, \wedge, \vee) is a lattice and

2. for each $m \in M$, $m \neq 0$, and $a \in A$, there exists an element $ma \in A$ satisfying the following conditions :

$$(i) \quad m(a \wedge b) = ma \wedge mb$$

$$(ii) \quad m(a \vee b) = ma \vee mb$$

$$(iii) \quad ma \wedge na \leq (m+n)a \quad \text{and} \quad ma \vee na \leq (m+n)a$$

$$(iv) \quad (mn)a = m(na), \text{ for all } m, n \in M, m \neq 0, n \neq 0, \text{ and } a, b \in A$$

$$(v) \quad 1.a = a, \quad \text{for all } a \in A$$

Here, \leq is the partial order relation corresponding to the lattice (A, \wedge, \vee)

The multiplication ma is called a bi-monoid multiplication with an artex element or simply bi-monoid multiplication in A .

Unless otherwise stated A remains as an Artex space with the partial ordering \leq which need not be “less than or equal to” and M as a bi-monoid with the binary operations $+$ and \cdot need not be the usual addition and usual multiplication.

Proposition 2.1.4 : If A and B are any two Artex spaces over a bi-monoid M and if \leq_1 and \leq_2 are the partial ordering on A and B respectively, then AXB is also an Artex Space over M , where the partial ordering \leq on AXB and the bi-monoid multiplication in AXB are defined by the following :

For $x, y \in AXB$, where $x=(a_1, b_1)$ and $y=(a_2, b_2)$, $x \leq y$ means $a_1 \leq_1 a_2$ and $b_1 \leq_2 b_2$

For $m \in M$, $m \neq 0$, and $x \in AXB$, where $x=(a,b)$, the bi-monoid multiplication in AXB is defined by

$$mx = m(a,b) = (ma,mb), \text{ where } ma \text{ and } mb \text{ are the bi-monoid multiplications in } A \text{ and } B \text{ respectively.}$$

In other words if \wedge_1 and \vee_1 are the cap, cup of A and \wedge_2 and \vee_2 are the cap, cup of B , then the cap, cup of AXB denoted by Λ and V are defined by $x \Lambda y = (a_1,b_1) \Lambda (a_2,b_2) = (a_1 \wedge_1 a_2, b_1 \wedge_2 b_2)$

$$x V y = (a_1,b_1) V (a_2,b_2) = (a_1 \vee_1 a_2, b_1 \vee_2 b_2).$$

Corollary 2.1.5 : If $A_1, A_2, A_3, \dots, A_n$ are Artex spaces over a bi-monoid M , then $A_1 X A_2 X A_3 X \dots X A_n$ is also an Artex space over M .

2.1.6 Definition : Complete Artex Space over a bi-monoid : An Artex space A over a bi-monoid M is said to be a Complete Artex Space if as a lattice, A is a complete lattice, that is each nonempty subset of A has a least upper bound and a greatest lower bound.

2.1.7 Remark : Every Complete Artex space must have a least element and a greatest element. The least and the greatest elements, if they exist, are called the bounds or units of the Artex space and are denoted by 0 and 1 respectively.

Note : The identity elements of $(M, +, \cdot)$ with respect to $+$ and \cdot are also denoted by 0 and 1 respectively.

2.1.8 Definition : Lower Bounded Artex Space over a bi-monoid : An Artex space A over a bi-monoid M is said to be a Lower Bounded Artex Space over M if as a lattice, A has the least element 0 .

2.1.9 Definition : Upper Bounded Artex Space over a bi-monoid : An Artex space A over a bi-monoid M is said to be an Upper Bounded Artex Space over M if as a lattice, A has the greatest element 1 .

2.1.10 Definition : Artex Space Homomorphism : Let A and B be two Artex spaces over a bi-monoid M , where \wedge_1 and \vee_1 are the cap, cup of A and \wedge_2 and \vee_2 are the cap, cup of B . A mapping $f : A \rightarrow B$ is said to be an Artex space homomorphism if

1. $f(a \wedge_1 b) = f(a) \wedge_2 f(b)$
2. $f(a \vee_1 b) = f(a) \vee_2 f(b)$
3. $f(ma) = mf(a)$, for all $m \in M$, $m \neq 0$ and $a, b \in A$.

2.1.11 Definition : Artex Space Epimorphism : Let A and B be two Artex spaces over a bi-monoid M . An Artex space homomorphism $f : A \rightarrow B$ is said to be an Artex space epimorphism if the mapping $f : A \rightarrow B$ is onto.

2.1.12 Definition : Artex Space Monomorphism : Let A and B be two Artex Spaces over a bi-monoid M . An Artex space homomorphism $f : A \rightarrow B$ is said to be an Artex Space monomorphism if the mapping $f : A \rightarrow B$ is one-one.

2.1.13 Definition : Artex Space Isomorphism : Let A and B be two Artex spaces over a bi-monoid M . An Artex Space homomorphism $f : A \rightarrow B$ is said to be an Artex Space Isomorphism if the mapping $f : A \rightarrow B$ is both one-one and onto, ie, f is bijective.

2.1.14 Definition : Isomorphic Artex Spaces : Two Artex spaces A and B over a bi-monoid M are said to be isomorphic if there exists an isomorphism from A onto B or from B onto A .

2.1.15 Definition : SubArtex Space : Let (A, \wedge, \vee) be an Artex space over a bi-monoid $(M, +, \cdot)$

Let S be a nonempty subset of A . Then S is said to be a Subartex space of A if (S, \wedge, \vee) itself is an Artex space over M .

Proposition 2.1.16 : Let (A, \wedge, \vee) be an Artex space over a bi-monoid $(M, +, \cdot)$

Then a nonempty subset S of A is a SubArtex space of A if and only if for each $m, n \in M$, $m \neq 0$, $n \neq 0$, and

$$a, b \in S, ma \wedge nb \in S \text{ and } ma \vee nb \in S$$

3 Homomorphic image of a SubArtex Space of an Artex space over a bi-monoid

Proposition 3.1.1 : Let A and B be Artex spaces over a bi-monoid M . Let $f : A \rightarrow B$ be an Artex space homomorphism. Let S be a SubArtex space of A . Then $f(S)$ is a SubArtex Space of B . In other words, under the Artex space homomorphism $f : A \rightarrow B$, the homomorphic image of a SubArtex Space of the Artex space A over M is a SubArtex space of B over M .

Proof : Let A and B be Artex spaces over a bi-monoid M .

Let S be a SubArtex space of A .

Let \leq_1 and \leq_2 be the partial orderings of A and B respectively.

Let \wedge_1 and \vee_1 be the cap and cup of A and let \wedge_2 and \vee_2 be the cap and cup of B .

Suppose $f : A \rightarrow B$ is an Artex space homomorphism.

To show $f(S)$ is a SubArtex Space of B .

Let $a', b' \in f(S)$ and $m, n \in M$, where $m \neq 0$, and $n \neq 0$.

Then there exists a and b in A such that $f(a) = a'$ and $f(b) = b'$

$$ma' \wedge_2 nb' = mf(a) \wedge_2 n f(b) = f(ma) \wedge_2 f(nb) = f(ma \wedge_1 nb) \quad (\text{since } f \text{ is a homomorphism})$$

Since S is a subArtex space of A , $ma \wedge_1 nb \in A$

Therefore, $f(ma \wedge_1 nb) \in f(S)$

$$ma' \wedge_2 nb' \in f(S)$$

$$ma' \vee_2 nb' = mf(a) \vee_2 n f(b) = f(ma) \vee_2 f(nb) = f(ma \vee_1 nb).$$

Since S is a subArtex space of A , $ma \vee_1 nb \in A$

Therefore, $f(ma \vee_1 nb) \in f(S)$

$$ma' \vee_2 nb' \in f(S)$$

Hence by the Proposition 2.1.16, $f(S)$ is a SubArtex space of B .

4 Distributive Artex Space over a bi-monoid : An Artex space A over a bi-monoid M is said to be a Distributive Artex Space over the bi-monoid M if as a lattice, A is a distributive lattice.

In other words, an Artex space A over a bi-monoid M is said to be a Distributive Artex Space over the bi-monoid M if for any $a, b, c \in A$, (i) $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$ (ii) $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$

4.1.1 Example : Let A be the set of all sequences (x_n) in Z , where Z is the set of all integers

and let $W = \{0,1,2,3,\dots\}$.

Define \leq' , an order relation, on A by for $(x_n), (y_n)$ in A , $(x_n) \leq' (y_n)$ means $x_n \leq y_n$, for each n ,

where \leq is the usual relation “less than or equal to”

Let $x \in A$, where $x = (x_n)$

Clearly $x_n \leq y_n$, for each n

So, $(x_n) \leq' (x_n)$

Therefore, \leq' is reflexive.

Let $x, y \in A$, where $x = (x_n)$ and $y = (y_n)$ be such that $x \leq' y$ and $y \leq' x$, that is, $(x_n) \leq' (y_n)$ and

$(y_n) \leq' (x_n)$.

Then $(x_n) \leq' (y_n)$ implies $x_n \leq y_n$, for each n

$(y_n) \leq' (x_n)$ implies $y_n \leq x_n$, for each n

Now, $x_n \leq y_n$, for each n , and $y_n \leq x_n$, for each n , implies $x_n = y_n$, for each n .

Therefore, $(x_n) = (y_n)$, that is $x = y$

Therefore, \leq' is anti-symmetric.

Let $x, y, z \in A$, where $x = (x_n)$, $y = (y_n)$ and $z = (z_n)$ be such that $x \leq' y$ and $y \leq' z$,

That is, $(x_n) \leq' (y_n)$

and $(y_n) \leq' (z_n)$.

Then $(x_n) \leq' (y_n)$ implies $x_n \leq y_n$, for each n

$(y_n) \leq' (z_n)$ implies $y_n \leq z_n$, for each n

Now, $x_n \leq y_n$, for each n , and $y_n \leq z_n$, for each n , implies $x_n \leq z_n$ for each n .

Therefore, $(x_n) \leq' (z_n)$

Therefore, \leq' is transitive.

Hence, \leq' is a partial order relation on A

Now the cap, cup operations are defined by the following :

$(x_n) \wedge (y_n) = (u_n)$, where $u_n = \min\{x_n, y_n\}$, for each n .

$(x_n) \vee (y_n) = (v_n)$, where $v_n = \max\{x_n, y_n\}$, for each n .

Clearly (A, \leq') is a lattice.

The bi-monoid multiplication in A is defined by the following :

For each $m \in W, m \neq 0$, and $x \in A$, where $x = (x_n)$, mx is defined by $mx = m(x_n) = (mx_n)$.

Clearly $(mx_n) \in A$

Let $x, y \in A$, where $x = (x_n)$, $y = (y_n)$ and let $m \in W, m \neq 0$

Then, it is clear that (i) $m(x \wedge y) = mx \wedge my$

(ii) $m(x \vee y) = mx \vee my$

(iii) $mx \wedge nx \leq (m+n)x$ and $mx \vee nx \leq (m+n)x$

(iv) $(mn)x = m(nx)$

(v) $1.x = x$, for all $m, n \in W, m \neq 0, n \neq 0$, and $x, y \in A$

Therefore, A is an Artex space over W .

Clearly, $(x_n) \wedge [(y_n) \vee (z_n)] = [(x_n) \wedge (y_n)] \vee [(x_n) \wedge (z_n)]$

and $(x_n) \vee [(y_n) \wedge (z_n)] = [(x_n) \vee (y_n)] \wedge [(x_n) \vee (z_n)]$.

Hence, A is a Distributive Artex Space over the bi-monoid W .

4.1.2 Example : If B is the set of all sequences (x_n) in Q , where Q is the set of all rational numbers, then as in Example 4.1.1, B is a Distributive Artex Space over the bi-monoid W .

4.1.3 Example : If D is the set of all sequences (x_n) in R , where R is the set of all real numbers, then as in Example 4.1.1, D is a Distributive Artex Space over the bi-monoid W .

Proposition 4.2.1: A SubArtex space S of a Distributive Artex space A over a bi-monoid M is a Distributive Artex space over M .

Proof : Let A be a Distributive Artex space over a bi-monoid M .

Let S be SubArtex space of A .

As a SubArtex space of A , by the definition of a SubArtex space, $S \neq \emptyset$ and S itself is an Artex space over M .

Let $a, b, c \in S$

Since S is a subset of A , $a, b, c \in A$

Since A is a Distributive Artex space over M , (i) $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$

and (ii) $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$

Hence, S is a Distributive Artex space over M .

Proposition 4.2.2 : If D and D' are any two Distributive Artex spaces over a bi-monoid M , then DXD' is also a Distributive Artex Space over M , If \leq_1 and \leq_2 are the partial orderings on D and D' respectively, then the partial ordering \leq on DXD' and the bi-monoid multiplication in DXD' are defined by the following :

For $x, y \in DXD'$, where $x = (a_1, b_1)$ and $y = (a_2, b_2)$, $x \leq y$ means $a_1 \leq_1 a_2$ and $b_1 \leq_2 b_2$

For $m \in M, m \neq 0$, and $x \in DXD'$, where $x = (a, b)$, the bi-monoid multiplication in DXD' is defined by $mx = m(a, b) = (ma, mb)$, where ma and mb are the bi-monoid multiplications in D and D' respectively.

In other words if \wedge_1 and \vee_1 are the cap, cup of D and \wedge_2 and \vee_2 are the cap, cup of D' , then the cap, cup of DXD' denoted by \wedge and \vee are defined by $x \wedge y = (a_1, b_1) \wedge (a_2, b_2) = (a_1 \wedge_1 a_2, b_1 \wedge_2 b_2)$

and $x \vee y = (a_1, b_1) \vee (a_2, b_2) = (a_1 \vee_1 a_2, b_1 \vee_2 b_2)$.

Proof : Let $A = DXD'$

We know that if (D, \leq_1) and (D', \leq_2) are any two Artex spaces over a bi-monoid M , then by the Proposition 2.1.4 DXD' is an Artex space over the bi-monoid M . Therefore, it is enough to prove that $A = DXD'$ is Distributive.

Let $x, y, z \in A = DXD'$, where $x = (b, b')$, $y = (c, c')$, $z = (d, d')$

$$\begin{aligned} \text{Now, } x \wedge (y \vee z) &= (b, b') \wedge ((c, c') \vee (d, d')) \\ &= (b, b') \wedge (c \vee_1 d, c' \vee_2 d') \\ &= (b \wedge_1 (c \vee_1 d), b' \wedge_2 (c' \vee_2 d')) \\ &= ((b \wedge_1 c) \vee_1 (b \wedge_1 d), (b' \wedge_2 c') \vee_2 (b' \wedge_2 d')) \\ &= (b \wedge_1 c, b' \wedge_2 c') \vee (b \wedge_1 d, b' \wedge_2 d') \\ &= ((b, b') \wedge (c, c')) \vee ((b, b') \wedge (d, d')) \\ &= (x \wedge y) \vee (x \wedge z) \end{aligned}$$

To show $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$

$$\begin{aligned} \text{Now, } x \vee (y \wedge z) &= (b, b') \vee ((c, c') \wedge (d, d')) \\ &= (b, b') \vee (c \wedge_1 d, c' \wedge_2 d') \\ &= (b \vee_1 (c \wedge_1 d), b' \vee_2 (c' \wedge_2 d')) \\ &= ((b \vee_1 c) \wedge_1 (b \vee_1 d), (b' \vee_2 c') \wedge_2 (b' \vee_2 d')) \\ &= (b \vee_1 c, b' \vee_2 c') \wedge (b \vee_1 d, b' \vee_2 d') \\ &= ((b, b') \vee (c, c')) \wedge ((b, b') \vee (d, d')) \\ &= (x \vee y) \wedge (x \vee z) \end{aligned}$$

Hence, $A = DXD'$ is a Distributive Artex space over M .

Corollary 4.2.3 : If $D_1, D_2, D_3, \dots, D_n$ are Distributive Artex spaces over a bi-monoid M , then

$D_1 \times D_2 \times D_3 \times \dots \times D_n$ is also a Distributive Artex space over M .

Proof : The proof is by induction on n

When $n=2$, by the Proposition $B_1 \times B_2$ is a Distributive Artex space over M

Assume that $B_1 \times B_2 \times B_3 \times \dots \times B_{n-1}$ is a Distributive Artex space over M

Consider $B_1 \times B_2 \times B_3 \times \dots \times B_n$

Let $B = B_1 \times B_2 \times B_3 \times \dots \times B_{n-1}$

$$\begin{aligned} \text{Then } B_1 \times B_2 \times B_3 \times \dots \times B_n &= (B_1 \times B_2 \times B_3 \times \dots \times B_{n-1}) \times B_n \\ &= B \times B_n \end{aligned}$$

By assumption B is a Distributive Artex space over M .

Again by the Proposition $B \times B_n$ is a Distributive Artex space over M

Hence, $B_1 \times B_2 \times B_3 \times \dots \times B_n$ is a Distributive Artex space over M

Proposition 4.2.4 : Let A be a Distributive Artex space over a bi-monoid M and let B be an Artex space over M . Let $f : A \rightarrow B$ be an Artex space epimorphism of A onto B . Then B is a Distributive Artex Space over M . In other words, the homomorphic image of a Distributive Artex Space over a bi-monoid is a Distributive Artex space over the bimonoid.

Proof : Let A be a Distributive Artex Space over a bi-monoid M and let B be an Artex space over M .

Let \leq_1 and \leq_2 be the partial orderings of A and B respectively.

Let \wedge_1 and \vee_1 be the cap and cup of A and let \wedge_2 and \vee_2 be the cap and cup of B .

Suppose $f : A \rightarrow B$ is an Artex space epimorphism of A onto B .

To show that $B = f(A)$ is a Distributive Artex Space over the bi-monoid M .

Let $x', y', z' \in B$

To show (i) $x' \wedge_2 (y' \vee_2 z') = (x' \wedge_2 y') \vee_2 (x' \wedge_2 z')$ and (ii) $x' \vee_2 (y' \wedge_2 z') = (x' \vee_2 y') \wedge_2 (x' \vee_2 z')$.

Since $f : A \rightarrow B$ is an Artex space epimorphism of A onto B , there exist elements $x, y, z \in A$ such that

$f(x) = x', f(y) = y',$ and $f(z) = z'$.

To show (i) $x' \wedge_2 (y' \vee_2 z') = (x' \wedge_2 y') \vee_2 (x' \wedge_2 z')$

$$\begin{aligned} \text{Now, } x' \wedge_2 (y' \vee_2 z') &= f(x) \wedge_2 (f(y) \vee_2 f(z)) \\ &= f(x) \wedge_2 f(y \vee_1 z) && \text{(since } f \text{ is an Artex space homomorphism)} \\ &= f(x \wedge_1 (y \vee_1 z)) && \text{(since } f \text{ is an Artex space homomorphism)} \\ &= f((x \wedge_1 y) \vee_1 (x \wedge_1 z)) && \text{(since } A \text{ is a Distributive Artex Space)} \\ &= f(x \wedge_1 y) \vee_2 f(x \wedge_1 z) && \text{(since } f \text{ is an Artex space homomor.)} \\ &= (f(x) \wedge_2 f(y)) \vee_2 (f(x) \wedge_2 f(z)) && \text{(since } f \text{ is an Artex space homomor.)} \\ &= (x' \wedge_2 y') \vee_2 (x' \wedge_2 z'). \end{aligned}$$

To show (ii) $x' \vee_2 (y' \wedge_2 z') = (x' \vee_2 y') \wedge_2 (x' \vee_2 z')$

$$\begin{aligned} \text{Now, } x' \vee_2 (y' \wedge_2 z') &= f(x) \vee_2 (f(y) \wedge_2 f(z)) \\ &= f(x) \vee_2 f(y \wedge_1 z) && \text{(since } f \text{ is an Artex space homomorphism)} \end{aligned}$$

$$\begin{aligned}
 &= f(x \vee_1 (y \wedge_1 z)) \quad (\text{since } f \text{ is an Artex space homomorphism}) \\
 &= f((x \vee_1 y) \wedge_1 (x \vee_1 z)) \quad (\text{since } A \text{ is a Distributive Artex Space}) \\
 &= f(x \vee_1 y) \wedge_2 f(x \vee_1 z) \quad (\text{since } f \text{ is an Artex space homomorphism}) \\
 &= (f(x) \vee_2 f(y)) \wedge_2 (f(x) \vee_2 f(z)) \quad (\text{since } f \text{ is an Artex space homomorphism}) \\
 &= (x' \vee_2 y') \wedge_2 (x' \vee_2 z').
 \end{aligned}$$

Hence, B is a Distributive Artex Space over M.

SubArtex Spaces Of Bounded Artex Spaces Over Bi-monoids

Problem 1 : A SubArtex space of a Lower Bounded Artex space over a bi-monoid need not be a Lower Bounded Artex space over the bi-monoid.

Proof : Let A be the set of all constant sequences (x_n) in $[0, \infty)$ and let $W = \{0, 1, 2, 3, \dots\}$.

Define \leq' , an order relation, on A by for $(x_n), (y_n)$ in A, $(x_n) \leq' (y_n)$ means $x_n \leq y_n$, for each n

Where \leq is the usual relation “less than or equal to”

Since the sequences in A are all constant sequences, $x_n \leq y_n$, for some n implies $x_n \leq y_n$, for each n

Therefore, $x_n \leq y_n$, for each n and $x_n \leq y_n$, for some n in this problem are the same.

Let $x \in A$, where $x = (x_n)$

Clearly $x_n \leq x_n$, for each n

So, $(x_n) \leq' (x_n)$

Therefore, \leq' is reflexive.

Let $x, y \in A$, where $x = (x_n)$ and $y = (y_n)$ be such that $x \leq' y$ and $y \leq' x$, that is, $(x_n) \leq' (y_n)$ and $(y_n) \leq' (x_n)$.

Then $(x_n) \leq' (y_n)$ implies $x_n \leq y_n$, for each n

and $(y_n) \leq' (x_n)$ implies $y_n \leq x_n$, for each n

Now, $x_n \leq y_n$, for each n, and $y_n \leq x_n$, for each n, implies $x_n = y_n$, for each n.

Therefore, $(x_n) = (y_n)$, that is $x = y$

Therefore, \leq' is anti-symmetric.

Let $x, y, z \in A$, where $x = (x_n)$, $y = (y_n)$ and $z = (z_n)$ be such that $x \leq' y$ and $y \leq' z$, that is, $(x_n) \leq' (y_n)$

and $(y_n) \leq' (z_n)$.

Then $(x_n) \leq' (y_n)$ implies $x_n \leq y_n$, for each n

$(y_n) \leq' (z_n)$ implies $y_n \leq z_n$, for each n

Now, $x_n \leq y_n$, for each n, and $y_n \leq z_n$, for each n, implies $x_n \leq z_n$ for each n.

Therefore, $(x_n) \leq' (z_n)$

Therefore, \leq' is transitive.

Hence, \leq' is a partial order relation on A

Now the cap ,cup operations are defined by the following :

$$(x_n) \wedge (y_n) = (u_n), \text{ where } u_n = \min\{x_n, y_n\}, \text{ for each } n.$$

$$(x_n) \vee (y_n) = (v_n), \text{ where } v_n = \max\{x_n, y_n\}, \text{ for each } n.$$

Clearly (A, \leq') is a lattice.

The bi-monoid multiplication in A is defined by the following :

For each $m \in W, m \neq 0$, and $x \in A$, where $x = (x_n)$, mx is defined by $mx = m(x_n) = (mx_n)$.

Since (x_n) is a constant sequence belonging to A , (mx_n) is also a constant sequence belonging to A.

Therefore $(mx_n) \in A$

Let $x, y \in A$, where $x = (x_n)$, $y = (y_n)$ and let $m \in W, m \neq 0$

Then, it is clear that (i) $m(x \wedge y) = mx \wedge my$

$$(ii) \quad m(x \vee y) = mx \vee my$$

$$(iii) \quad mx \wedge nx \leq (m+n)x \quad \text{and} \quad mx \vee nx \leq (m+n)x$$

$$(iv) \quad (mn)x = m(nx) , \text{ for all } m, n \in W, m \neq 0, n \neq 0, \text{ and } x, y \in A$$

$$(v) \quad 1.x = x , \text{ for all } x \in A$$

Therefore, A is an Artex space over W.

The sequence (0_n) , where 0_n is 0 for all n, is a constant sequence belonging to A

Also $(0_n) \leq' (x_n)$, for all the sequences (x_n) belonging to A

Therefore, (0_n) is the least element of A.

That is, the sequence $0, 0, 0, \dots$ is the least element of A

Hence A is a Lower Bounded Artex space over W.

Now let S be the set of all constant sequences (x_n) in $(0, \infty)$.

Clearly S is a SubArtex Space A.

But S has no least element.

Therefore, S is not a Lower Bounded Artex Space over W.

Hence, a SubArtex space of a Lower Bounded Artex space over a bi-monoid need not be a Lower Bounded Artex space over the bi-monoid.

Problem 2 : A SubArtex space of an Upper Bounded Artex space over a bi-monoid need not be an Upper Bounded Artex space over the bi-monoid.

Proof : Let A be the set of all constant sequences (x_n) in $(-\infty, 0]$ and let $W = \{0, 1, 2, 3, \dots\}$.

Define \leq' , an order relation, on A by for $(x_n), (y_n)$ in A , $(x_n) \leq' (y_n)$ means $x_n \leq y_n$, for $n = 1, 2, 3, \dots$

where \leq is the usual relation “less than or equal to”

Then as in Example 3.3.1, A is an Artex space over W .

Now, the sequence (1_n) , where 1_n is 0, for all n , is a constant sequence belonging to A

Also $(x_n) \leq' (1_n)$, for all the sequences (x_n) in A

Therefore, (1_n) is the greatest element of A .

That is, the sequence $(0, 0, 0, \dots)$ is the greatest element of A

Hence A is an Upper Bounded Artex Space over W .

Now let S be the set of all constant sequences (x_n) in $(-\infty, 0)$.

Clearly S is a SubArtex Space A .

But S has no greatest element.

Therefore, S is not an Upper Bounded Artex Space over W .

Hence, a SubArtex space of an Upper Bounded Artex space over a bi-monoid need not be an

Upper Bounded Artex space over the bi-monoid.

Problem 3 : A SubArtex space of a Bounded Artex space over a bi-monoid need not be a Bounded Artex space over the bi-monoid.

Proof : The proof is clear from the Problems 1 and 2

In other way, if A is taken as the set of all constant sequences in the extended real line and S is taken as the set of all sequences in the real line, then as in the Problems 1 and 2 the proof follows. Here by convention the symbols $-\infty$ and ∞ are considered to be the least and the greatest elements of the extended real line, but in the real line there exist no such elements. Therefore S is not a Bounded Artex space over W .

References : 1. K.Muthukumar and M.Kamaraj, ”Artex Spaces Over Bi-monoids”, Research Journal Of Pure Algebra,2(5),May 2012, Pages 135-140.

2. K.Muthukumar and M.Kamaraj, “Subartex Spaces Of an Artex Space Over a Bi-monoid”,(accepted) Mathematical Theory and Modeling ,an IISTE Journal.

3. K.Muthukumar and M.Kamaraj, “Bounded Artex Spaces Over Bi-monoids and Artex Space Homomorphisms”,(accepted) Research Journal Of Pure Algebra.

4. J.P.Tremblay and R.Manohar, Discrete Mathematical Structures with Applications to Computer Science, Tata McGraw-Hill Publishing Company Limited, New Delhi, 1997.

5. John T. Moore, The University of Florida /The University of Western Ontario, Elements of Abstract Algebra, Second Edition, The Macmillan Company, Collier-Macmillan Limited, London, 1967.
6. Garrett Birkhoff & Thomas C. Bartee, Modern Applied Algebra, CBS Publishers & Distributors, 1987.
7. J. Eldon Whitesitt, Boolean Algebra And Its Applications, Addison-Wesley Publishing Company, Inc., U.S.A., 1961.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

