

On Continuity of Complex Fuzzy Functions

Pishtiwani O. Sabir

Department of Mathematics, Faculty of Science and Science Education, University of Sulaimani, Iraq
pishtiwani.sabir@gmail.com

Abstract

In this paper, Some important theorems on fuzzy type I-continuous and II-continuous of complex fuzzy functions mapping generalized rectangular valued bounded closed complex complement normalized fuzzy numbers into itself are proved.

Keywords: Fuzzy Complex Numbers, Fuzzy Complex Functions, Fuzzy Continuity.

1. Introduction

It is well known that fuzzy complex numbers and fuzzy complex analysis were first introduced by (Buckley, 1989; Buckley and Qu, 1991, 1992). Scholars did series research about the properties of fuzzy complex number from various aspects (Quan, 1996; Ma *et al.*, 2009; Zheng and Ha, 2009). But these achievements were very abstract, and it did not consummate until today. In view of (Buckley, 1989), Guangquan (1992) discussed the limit theory of the sequence of fuzzy complex numbers in detail, giving a series of results about limit theory, which are the counterparts of well-known results valid for real numbers in classical mathematics analysis. Buckley (1989) suggested that introducing a metric on the space of fuzzy complex numbers provide to study convergence, continuity and differentiation of fuzzy complex function (Chun and Ma, 1998; Qiu *et al.*, 2000, 2001; Ousmane and Congxin, 2003; Shengquan, 2006; Cai, 2009; Sabir, 2012). On the basis of Buckley's work, some authors continued research and have extensively studied the theory of fuzzy complex numbers and fuzzy complex analysis (Wu and Qiu, 1999; Zengtai and Shengquan, 2006; Qiu and Shu, 2008; Sun and Guo, 2010; Sabir *et al.*, 2012b). Sabir *et al.* (2012a) giving the definitions of the complement normalized fuzzy numbers (CNFNs), bounded closed complex CNFNs (BCCCNFNs), generalized rectangular valued BCCCNFNs (GRVBCCCNFNs) and discussed some of their basic properties. In section two, we first review the definitions and characterizations related to fuzzy complex sets. We will also present the notations needed in the rest of the paper. In the last section, some theorems on the continuity of complex fuzzy functions are proved.

2. Preliminaries

A fuzzy set \tilde{A} defined on the universal set X is a function $\mu(\tilde{A}, x) : X \rightarrow [0,1]$. Frequently, we will write $\mu_{\tilde{A}}(x)$ instead of $\mu(\tilde{A}, x)$. The family of all fuzzy sets in X is denoted by $\mathcal{F}(X)$. The α -level of a fuzzy set \tilde{A} , denoted by $\alpha^+ \tilde{A}$, is the non-fuzzy set of all elements of the universal set that belongs to the fuzzy set \tilde{A} at least to the degree $\alpha \in [0,1]$. The weak α -level $\alpha^- \tilde{A}$ of a fuzzy set $\tilde{A} \in \mathcal{F}(X)$ is the crisp set that contains all elements of the universal set whose membership grades in the given set are greater than but do not include the specified value of α . The largest value of α for which the α -level is not empty is called the height of a fuzzy set \tilde{A} denoted $\alpha_{\tilde{A}}^{max}$. The core of a fuzzy set \tilde{A} is the non-fuzzy set of all points in the universal set X at which $\sup_x \mu_{\tilde{A}}(x)$ is essentially attained.

Let $\tilde{A}_i \in \mathcal{F}(X)$. Then the union of fuzzy sets \tilde{A}_i , denoted $\cup_i \tilde{A}_i$, is defined by $\mu_{\cup_i \tilde{A}_i}(x) = \sup_x \mu_{\tilde{A}_i}(x) = \vee_x \mu_{\tilde{A}_i}(x)$, the intersection of fuzzy sets \tilde{A}_i , denoted $\cap_i \tilde{A}_i$, is defined by $\mu_{\cap_i \tilde{A}_i}(x) = \inf_x \mu_{\tilde{A}_i}(x) = \wedge_x \mu_{\tilde{A}_i}(x)$, and the complement of \tilde{A}_i , denoted $\neg \tilde{A}_i$, is defined by $\mu_{\neg \tilde{A}_i}(x) = 1 - \mu_{\tilde{A}_i}(x)$, for all x in the universal set X .

A fuzzy number \tilde{a} is a fuzzy set defined on the set of real numbers R^1 characterized by means of a membership function $\mu_{\tilde{a}}(x) : R^1 \rightarrow [0,1]$, which satisfies: (1) \tilde{a} is upper semicontinuous, (2) $\mu_{\tilde{a}}(x) = 0$ outside some interval $[c, d]$, (3) There are real numbers a, b such that $c \leq a \leq b \leq d$ and $\mu_{\tilde{a}}(x)$ is increasing on $[c, a]$, $\mu_{\tilde{a}}(x)$ is decreasing on $[b, d]$, $\mu_{\tilde{a}}(x) = 1$, $a \leq x \leq b$. We denote the set of all fuzzy numbers by \mathcal{F}^* . A fuzzy complex number \tilde{Z} is defined by its membership functions $\mu_{\tilde{Z}}(z)$ which is a mapping from the set of ordinary complex numbers into $[0,1]$ if and only if $\mu_{\tilde{Z}}(z)$ is continuous; $\alpha^- \tilde{Z}$ is open, bounded, and connected; and ${}^{1+} \tilde{Z}$ is non-empty, compact, and arcwise connected. We use \mathcal{F}^{**} to the set of all fuzzy complex numbers.

Let $f(z', z'') = w$ be any mapping from $\mathbb{C} \times \mathbb{C}$ into \mathbb{C} . Buckley (1989) extend f to $\mathcal{F}^{**} \times \mathcal{F}^{**}$ into \mathcal{F}^{**} and write $f(\tilde{Z}', \tilde{Z}'') = \tilde{W}$ if $\mu_{\tilde{W}}(w) = \vee_{f(z', z'')=w} (\mu_{\tilde{Z}'}(z') \wedge \mu_{\tilde{Z}''}(z''))$. One obtains $\tilde{W} = \tilde{Z}' \oplus \tilde{Z}''$ or $\tilde{W} = \tilde{Z}' \odot \tilde{Z}''$ by using $f(\tilde{Z}', \tilde{Z}'') = \tilde{Z}' \oplus \tilde{Z}''$ or $f(\tilde{Z}', \tilde{Z}'') = \tilde{Z}' \odot \tilde{Z}''$, respectively.

3. Properties of Continuous Complex Fuzzy Functions

In this section, we give the continuity of complex fuzzy function mapping GRVBCCCNFNs into itself. Most results, definitions and standard notations on fuzzy complex analysis which are used in this section can be found

in Sabir *et al.* (2012a). Some of the results in this section are without proofs owing to the simplicities.

Definition 3.1. Let $\tilde{\mathcal{F}}^* \subseteq [\tilde{\mathcal{F}}_{-N}^{**}]$, and \tilde{F} be a mapping from $\tilde{\mathcal{F}}^*$ to the set of all GRVBCCCNFNs. If for arbitrary $[Z] \in \tilde{\mathcal{F}}^*$, there exists unique $[\tilde{W}] \in [\tilde{\mathcal{F}}_{-N}^{**}]$, make $\tilde{F}([Z]) = [\tilde{W}]$, we call \tilde{F} a complex fuzzy function defined on $\tilde{\mathcal{F}}^*$. Let $[\tilde{W}] \in [\tilde{\mathcal{F}}_{-N}^{**}]$, we say $\tilde{F}([Z])$ is fuzzy continuous at $[\tilde{W}]$ if for all $\bar{\varepsilon} > \bar{0}$ there exists $\bar{\delta} > \bar{0}$ such that $\tilde{r}(\tilde{F}([Z]), \tilde{F}([\tilde{W}])) < \bar{\varepsilon}$ as $\tilde{r}([Z], [\tilde{W}]) < \bar{\delta}$.

Definition 3.2. We say $\tilde{F}([Z])$ is fuzzy type I-continuous (resp. II-continuous) at $[\tilde{W}] \in [\tilde{\mathcal{F}}_{-N}^{**}]$ if for each $\bar{\varepsilon} > \bar{0}$ there is $\bar{\delta} > \bar{0}$ such that $\tilde{F}([Z]) < \tilde{F}([\tilde{W}])[+]\bar{\varepsilon}$ (resp. $\tilde{F}([\tilde{W}]) < \tilde{F}([Z])[+]\bar{\varepsilon}$) whenever $[Z][-][\tilde{W}] < \bar{\delta}$.

Theorem 3.3. Let \tilde{F} and \tilde{G} both are fuzzy continuous at GRVBCCCNFN $[\tilde{W}]$ then so is $\tilde{F}[*]\tilde{G}$ for $[*] \in \{[+], [-], [\cdot], [/], [\vee], [\wedge]\}$.

Proof: We only prove for $[*] = [+]$, the proof of the rest are similar. By hypothesis, for any $\bar{\varepsilon} > \bar{0}$, there exists $\bar{\delta} > \bar{0}$, when $\tilde{r}([Z], [\tilde{W}]) < \bar{\delta}$, $\tilde{r}(\tilde{F}([Z]), \tilde{F}([\tilde{W}])) < \bar{\varepsilon}/2$ and $\tilde{r}(\tilde{G}([Z]), \tilde{G}([\tilde{W}])) < \bar{\varepsilon}/2$. Therefore, we have

$$\begin{aligned} & \tilde{r}(\tilde{F}([+]\tilde{G})([Z]), \tilde{F}([\tilde{W}])[+]\tilde{G}([\tilde{W}])) \\ &= \tilde{r}(\tilde{F}([Z]), \tilde{F}([\tilde{W}])[+]\tilde{G}([\tilde{W}])[-]\tilde{G}([Z])) \\ &\leq \tilde{r}(\tilde{F}([Z]), \tilde{F}([\tilde{W}])) [+]\tilde{r}(\tilde{F}([\tilde{W}]), \tilde{F}([\tilde{W}])[+]\tilde{G}([\tilde{W}])[-]\tilde{G}([Z])) \\ &= \tilde{r}(\tilde{F}([Z]), \tilde{F}([\tilde{W}])) [+]\tilde{r}(\tilde{G}([\tilde{W}]), \tilde{G}([Z])) < \bar{\varepsilon}. \end{aligned}$$

Theorem 3.4. Let \tilde{F} be fuzzy continuous at $[\tilde{W}] \in [\tilde{\mathcal{F}}_{-N}^{**}]$. Then there exist GRVBCCCNFNs $[\tilde{Z}_0] = [\tilde{\mu}_0][+][\tilde{\mu}_1]$ and $[\tilde{Z}_1] = [\tilde{\lambda}_0][+][\tilde{\lambda}_1]$ satisfy $\gamma^+[\tilde{\mu}_0] < \gamma^+(\text{Re}\tilde{F}([\tilde{Z}])) < \gamma^+[\tilde{\lambda}_0]$ and $\gamma^+[\tilde{\mu}_1] < \gamma^+(\text{Im}\tilde{F}([\tilde{Z}])) < \gamma^+[\tilde{\lambda}_1]$ when $\tilde{r}([Z], [\tilde{W}]) < \bar{\delta}$ for $\bar{\delta} > \bar{0}$.

Theorem 3.5. Let \tilde{F} be fuzzy continuous function at $[\tilde{W}] \in [\tilde{\mathcal{F}}_{-N}^{**}]$, $[\tilde{Z}_0] \in [\tilde{\mathcal{F}}_{-N}^{**}]$, and there exists $\bar{\delta} > \bar{0}$ such that $\gamma^+(\text{Re}\tilde{F}([\tilde{Z}])) < \gamma^+(\text{Re}[\tilde{Z}_0])$ (resp. $\gamma^+(\text{Re}\tilde{F}([\tilde{Z}])) > \gamma^+(\text{Re}[\tilde{Z}_0])$) and $\gamma^+(\text{Im}\tilde{F}([\tilde{Z}])) < \gamma^+(\text{Im}[\tilde{Z}_0])$ (resp. $\gamma^+(\text{Im}\tilde{F}([\tilde{Z}])) > \gamma^+(\text{Im}[\tilde{Z}_0])$) when $\tilde{r}([Z], [\tilde{W}]) < \bar{\delta}$ then $\gamma^+(\text{Re}\tilde{F}([\tilde{W}])) < \gamma^+(\text{Re}[\tilde{Z}_0])$ (resp. $\gamma^+(\text{Re}\tilde{F}([\tilde{W}])) > \gamma^+(\text{Re}[\tilde{Z}_0])$) and $\gamma^+(\text{Im}\tilde{F}([\tilde{W}])) < \gamma^+(\text{Im}[\tilde{Z}_0])$ (resp. $\gamma^+(\text{Im}\tilde{F}([\tilde{W}])) > \gamma^+(\text{Im}[\tilde{Z}_0])$).

Theorem 3.6. Let \tilde{F} and \tilde{G} are both fuzzy type I-continuous (resp. II-continuous) and $[*] \in \{[\vee], [\wedge], [\cdot]\}$. Then

- $\tilde{F}[*]\tilde{G}$ is also fuzzy type I-continuous (resp. II-continuous), such that $\gamma^+(\text{Re}\tilde{F}([\tilde{Z}]))^l$, $\gamma^+(\text{Im}\tilde{F}([\tilde{Z}]))^l$, $\gamma^+(\text{Re}\tilde{G}([\tilde{Z}]))^u$, $\gamma^+(\text{Im}\tilde{G}([\tilde{Z}]))^u$, $\gamma^+(\text{Re}\tilde{F}([\tilde{Z}]))^u$, $\gamma^+(\text{Im}\tilde{F}([\tilde{Z}]))^u$, $\gamma^+(\text{Re}\tilde{G}([\tilde{Z}]))^l$, $\gamma^+(\text{Im}\tilde{G}([\tilde{Z}]))^l$ are all greater than zero for all $\gamma \in I_{[0]}^1$.
- $\tilde{F}[+]\tilde{G}$ is also fuzzy type I-continuous (resp. II-continuous).
- $[-]\tilde{G}$ is fuzzy type II-continuous (resp. I-continuous).
- $\tilde{I}[/math>]/ \tilde{G} is also fuzzy type I-continuous (resp. II-continuous) such that, $\gamma^+(\text{Re}\tilde{G}([\tilde{Z}]))^{l-}$, $\gamma^+(\text{Re}\tilde{G}([\tilde{Z}]))^{l+}$, $\gamma^+(\text{Re}\tilde{G}([\tilde{Z}]))^{u+}$, $\gamma^+(\text{Im}\tilde{G}([\tilde{Z}]))^{u-}$, $\gamma^+(\text{Im}\tilde{G}([\tilde{Z}]))^{l-}$, $\gamma^+(\text{Im}\tilde{G}([\tilde{Z}]))^{l+}$, $\gamma^+(\text{Re}\tilde{G}([\tilde{Z}]))^{u-}$, $\gamma^+(\text{Im}\tilde{G}([\tilde{Z}]))^{u+}$ are all positive for any $\gamma \in I_{[0]}^1$.$

Proof: By hypothesis, for any $\bar{\varepsilon} > \bar{0}$ there is a $\bar{\delta}' > \bar{0}$, $\bar{\delta}'' > \bar{0}$ such that $\tilde{F}([Z]) < \tilde{F}([\tilde{W}])[+]\bar{\varepsilon}$ (resp. $\tilde{F}([\tilde{W}]) < \tilde{F}([Z])[+]\bar{\varepsilon}$) whenever $[Z][-][\tilde{W}] < \bar{\delta}'$ and $\tilde{G}([Z]) < \tilde{G}([\tilde{W}])[+]\bar{\varepsilon}$ (resp. $\tilde{G}([\tilde{W}]) < \tilde{G}([Z])[+]\bar{\varepsilon}$) whenever $[Z][-][\tilde{W}] < \bar{\delta}''$.

- Let $\bar{\delta} = \bar{\delta}' \wedge \bar{\delta}''$ and $\gamma \in I_{[0]}^1$, we have
- $$\begin{aligned} \gamma^+(\text{Re}\tilde{F}([\tilde{Z}]))^{l-} &< \gamma^+(\text{Re}\tilde{F}([\tilde{W}]))^{l-} + \varepsilon \left(\text{resp. } \gamma^+(\text{Re}\tilde{F}([\tilde{W}]))^{l-} < \gamma^+(\text{Re}\tilde{F}([\tilde{Z}]))^{l-} + \varepsilon \right). \\ \gamma^+(\text{Re}\tilde{F}([\tilde{Z}]))^{l+} &< \gamma^+(\text{Re}\tilde{F}([\tilde{W}]))^{l+} + \varepsilon \left(\text{resp. } \gamma^+(\text{Re}\tilde{F}([\tilde{W}]))^{l+} < \gamma^+(\text{Re}\tilde{F}([\tilde{Z}]))^{l+} + \varepsilon \right). \end{aligned}$$

$$\begin{aligned}
 & \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{l^-} < \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{l^-} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{l^-} < \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{l^-} + \varepsilon \right). \\
 & \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{l^+} < \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{l^+} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{l^+} < \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{l^+} + \varepsilon \right). \\
 & \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{u^-} < \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{u^-} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{u^-} < \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{u^-} + \varepsilon \right). \\
 & \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{u^+} < \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{u^+} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{u^+} < \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{u^+} + \varepsilon \right). \\
 & \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{u^-} < \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^-} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^-} < \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{u^-} + \varepsilon \right). \\
 & \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{u^+} < \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^+} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^+} < \gamma^+ (\text{Im}\tilde{F}([\tilde{Z}]))^{u^+} + \varepsilon \right). \\
 & \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} < \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} < \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} + \varepsilon \right). \\
 & \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^+} < \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^+} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^+} < \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^+} + \varepsilon \right). \\
 & \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{l^-} < \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^-} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^-} < \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{l^-} + \varepsilon \right). \\
 & \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{l^+} < \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^+} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^+} < \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{l^+} + \varepsilon \right). \\
 & \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{u^-} < \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^-} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^-} < \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{u^-} + \varepsilon \right). \\
 & \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{u^+} < \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^+} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^+} < \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{u^+} + \varepsilon \right). \\
 & \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{u^-} < \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^-} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^-} < \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{u^-} + \varepsilon \right). \\
 & \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{u^+} < \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^+} + \varepsilon \left(\text{resp. } \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^+} < \gamma^+ (\text{Im}\tilde{G}([\tilde{Z}]))^{u^+} + \varepsilon \right). \text{ Hence,} \\
 & \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} \\
 & \quad < \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} + \left(\gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{l^-} + \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} + \varepsilon \right) * \varepsilon \\
 & \quad < \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} + \left({}^{0+} (\text{Re}\tilde{F}([\tilde{W}]))^{l^-} + {}^{0+} (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} + \varepsilon \right) * \varepsilon \\
 & \quad = \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} + \varepsilon' \\
 & \left(\text{resp. } \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} \right. \\
 & \quad < \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} + \left(\gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{l^-} + \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} + \varepsilon \right) * \varepsilon \\
 & \quad < \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} + \left({}^{0+} (\text{Re}\tilde{F}([\tilde{Z}]))^{l^-} + {}^{0+} (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} + \varepsilon \right) * \varepsilon \\
 & \quad = \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{l^-} * \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^-} + \varepsilon'' \left. \right). \\
 & \gamma^+ (\text{Re}\tilde{F}([\tilde{Z}]))^{l^+} * \gamma^+ (\text{Re}\tilde{G}([\tilde{Z}]))^{l^+} < \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{l^+} * \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^+} + \varepsilon' \left(\text{resp. } \gamma^+ (\text{Re}\tilde{F}([\tilde{W}]))^{l^+} * \right. \\
 & \gamma^+ \text{Re}\tilde{G}([\tilde{W}]))^{l^+} < \gamma^+ \text{Re}\tilde{F}([\tilde{Z}]))^{l^+} * \gamma^+ \text{Re}\tilde{G}([\tilde{Z}]))^{l^+} + \varepsilon''.
 \end{aligned}$$

$$\begin{aligned} & \gamma^+ (\text{Im}\tilde{F}([Z]))^{u^-} + \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^-} < \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^-} + \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^-} + 2\varepsilon \\ & \left(\text{resp. } \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^-} + \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^-} < \gamma^+ (\text{Im}\tilde{F}([Z]))^{u^-} + \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^-} + 2\varepsilon \right). \\ & \gamma^+ (\text{Im}\tilde{F}([Z]))^{u^+} + \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^+} < \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^+} + \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^+} + 2\varepsilon \\ & \left(\text{resp. } \gamma^+ (\text{Im}\tilde{F}([\tilde{W}]))^{u^+} + \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^+} < \gamma^+ (\text{Im}\tilde{F}([Z]))^{u^+} + \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^+} + 2\varepsilon \right). \end{aligned}$$

3. For any $\gamma \in I_{[0]}^1$, we have

$$\begin{aligned} & - \gamma^+ (\text{Re}\tilde{G}([Z]))^{l^-} > - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^-} > - \gamma^+ (\text{Re}\tilde{G}([Z]))^{l^-} + \varepsilon \right). \\ & - \gamma^+ (\text{Re}\tilde{G}([Z]))^{l^+} > - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^+} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{l^+} > - \gamma^+ (\text{Re}\tilde{G}([Z]))^{l^+} + \varepsilon \right). \\ & - \gamma^+ (\text{Im}\tilde{G}([Z]))^{l^-} > - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^-} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^-} > - \gamma^+ (\text{Im}\tilde{G}([Z]))^{l^-} + \varepsilon \right). \\ & - \gamma^+ (\text{Im}\tilde{G}([Z]))^{l^+} > - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^+} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{l^+} > - \gamma^+ (\text{Im}\tilde{G}([Z]))^{l^+} + \varepsilon \right). \\ & - \gamma^+ (\text{Re}\tilde{G}([Z]))^{u^-} > - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^-} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^-} > - \gamma^+ (\text{Re}\tilde{G}([Z]))^{u^-} + \varepsilon \right). \\ & - \gamma^+ (\text{Re}\tilde{G}([Z]))^{u^+} > - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^+} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Re}\tilde{G}([\tilde{W}]))^{u^+} > - \gamma^+ (\text{Re}\tilde{G}([Z]))^{u^+} + \varepsilon \right). \\ & - \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^-} > - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^-} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^-} > - \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^-} + \varepsilon \right). \\ & - \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^+} > - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^+} + \varepsilon \left(\text{resp. } - \gamma^+ (\text{Im}\tilde{G}([\tilde{W}]))^{u^+} > - \gamma^+ (\text{Im}\tilde{G}([Z]))^{u^+} + \varepsilon \right). \end{aligned}$$

4. Obvious.

Theorem 3.7. If \tilde{F} and \tilde{G} are a fuzzy type I-continuous and II-continuous such that $\gamma^+ (\text{Re}\tilde{F}([Z])) \leq \gamma^+ (\text{Re}\tilde{G}([Z]))$ and $\gamma^+ (\text{Im}\tilde{F}([Z])) \leq \gamma^+ (\text{Im}\tilde{G}([Z]))$ for every $[Z] \in [\tilde{\mathcal{F}}_{-N}^{**}]$ and $\gamma \in I_{[0]}^1$, then there is a fuzzy continuous function \tilde{H} satisfy $\gamma^+ (\text{Re}\tilde{F}([Z])) \leq \gamma^+ (\text{Re}\tilde{H}([Z])) \leq \gamma^+ (\text{Re}\tilde{G}([Z]))$ and $\gamma^+ (\text{Im}\tilde{F}([Z])) \leq \gamma^+ (\text{Im}\tilde{H}([Z])) \leq \gamma^+ (\text{Im}\tilde{G}([Z]))$.

Here, an open problem is presented for further investigations: One can study that \tilde{F} is both fuzzy type I-continuous and II-continuous if and only if \tilde{F} is fuzzy continuous.

References

- Buckley, J.J. and Qu, Y. (1991); (1992), "Fuzzy complex analysis I: Differentiation; II: Integration", *Fuzzy Sets and Systems* 41; 49, Elsevier, 269–284; 71–179.
- Buckley, J.J. (1989), "Fuzzy complex numbers", *Fuzzy Sets and Systems* 33, Elsevier, 333–345.
- Cai, Q.P. (2009), "The continuity of complex fuzzy function", *Adv. in Int. and Soft Computing* 2, 695-704.
- Chun, C. and Ma, S. (1998), "The differentiation of complex fuzzy functions", *Proc. of the 9th NCFMFS*, Baoding, Hebei U. Press, 162-166.
- Guangquan, Z. (1992), "Fuzzy limit theory of fuzzy complex numbers", *Fuzzy Sets and Systems* 46, Elsevier, 227–235.
- Ma, S.Q., Cai, Q.P., and Peng, D.J. (2009), "Some correlative conception and properties of bounded closed fuzzy complex number set", *Adv. in Int. and Soft Computing* 62, 705-715.
- Ousmane, M. and Congxin, W. (2003), "Semi continuity of complex fuzzy functions", *Tsinghua Science and Technology* 8, 65-70.
- Qiu, D. and Shu, L. (2008), "Notes on "On the restudy of fuzzy complex analysis: Part I and Part II", *Fuzzy Sets and Systems* 159, Elsevier, 2185–2189.
- Qiu, J., Wu, C. and Li, F. (2000); (2001), "On the restudy of fuzzy complex analysis: Part I. The sequence and series of fuzzy complex numbers and their convergences; Part II. The continuity and differentiation of fuzzy complex functions", *Fuzzy Sets and Systems* 115, Elsevier, 445–450; 120 517–521.

- Quan, M.S. (1996), "Fuzzy complex numbers and some operational properties", *J. of Lanzhou University* 32, NSE, 643-645.
- Sabir, P.O. (2012), "On fuzzy complex analysis", *Ph.D. Thesis*, University of Sulaimani, Sulaimani.
- Sabir, P.O., Adil, K.J., Munir, A.A. (2012)^a, "The continuity and differentiation of complex fuzzy functions for new fuzzy quantities", *Asian Transactions on Science & Technology* 2(4), 1-11.
- Sabir, P.O., Adil, K.J., Munir, A.A. (2012)^b, "On fuzzy complex integrals", *Asian Transactions on Science & Technology* 2(4), 12-17.
- Shengquan, M. (2006), "The series of complex fuzzy valued and its convergence", *Journal of Fuzzy Mathematics* 5, 200-211.
- Sun, J. and Guo, S. (2010), "The solution algorithm of complex fuzzy valued function integral by fuzzy element", *ACFIE* 78, 55-63.
- Wu, C. and Qiu, J. (1999), "Some remarks for fuzzy complex analysis", *Fuzzy Sets and Systems* 106, Elsevier, 231-238.
- Zengtai, G. and Shengquan, M. (2006), "The research advances in fuzzy complex analysis", *Math. in Prac. and Theory* 36, 200-211.
- Zheng, L. and Ha, M. (2009), "Further discussions on rectangular fuzzy complex numbers", *Proc. of the 8th ICMLC*, Baoding, 642-646.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

