

Common Unique Fixed Point Theorem for Random Operators in Hilbert Space

Anupama Gupta

Asst. Prof. BUIT (BU) BHOPAL

Abstract: The object of this paper is to obtain a common fixed point theorem for four continuous random operators by considering a sequence of measurable functions satisfying certain contractive condition in separable Hilbert space.

Keywords: Separable Hilbert space, random operators, common random fixed point, rational inequality.

Introduction: The study of the random fixed point theorems in abstract spaces is initiated by Hans [3] and is the stochastic generalizations of the classical fixed point theorems in separable Banach spaces. The research along this line gained momentum after the publication of the paper by Bharucha – Reid [1] and since then several random fixed point theorems have been proved in the literature. Random operator theory is needed for the study of various classes of random equations. Now this theory has become the full fledged research area and various ideas associated with random fixed point theory are used to obtain the solution of non linear random system [7]. The study of the random fixed point theory has attracted much attention in recent years [2, 5, 6]. In this paper we construct a sequence of measurable functions and consider its convergence to the common unique random fixed point of four continuous random operators defined on a non – empty closed subset of a separable Hilbert space. For the purpose of obtaining the random fixed point of the four continuous random operators, we have used a rational inequality and the parallogram law.

Preliminaries:

Throughout this paper (Ω, Σ) denotes a measurable space consisting of a set Ω and sigma algebra Σ of subsets of Ω . H stands for a separable Hilbert space, and C is a non – empty closed subset of H .

Definition 2.1: A function $f: \Omega \rightarrow C$ is said to be measurable if $f^{-1}(B \cap C) \in \Sigma$ for every Borel subset B of H .

Definition 2.2: A function $F: \Omega \times C \rightarrow C$ is said to be random operator if $F(., x): \Omega \rightarrow C$ is measurable for every $x \in C$.

Definition 2.3: A measurable function $g: \Omega \rightarrow C$ is said to be random fixed point of the random operator $F: \Omega \times C \rightarrow C$ if $F(t, g(t)) = g(t)$ for all $t \in \Omega$.

Definition 2.4: A random operator $F: \Omega \times C \rightarrow C$ is said to be continuous if for fixed $t \in \Omega$, $F(t, .): C \rightarrow C$ is continuous.

Main Result:

Theorem 1: Let C be a non – empty closed subset of a separable Hilbert space H . Let E, F, S, T be four continuous random operators defined on C such that for $t \in \Omega$, $E(t, .), F(t, .), T(t, .), S(t, .): C \rightarrow C$ satisfy condition.

$$\begin{aligned}
 \|Ex - Fy\|^2 &\leq \beta_1 \frac{\|y - Ty\|^2 [1 + \|x - Sx\|^2]}{1 + \|x - y\|^2} + \beta_2 [\|x - Sx\|^2 + \|y - Ty\|^2] \\
 &+ \beta_3 \frac{\|Sx - Ex\|^2 [\|Ty - Fy\|^2 + \|Ex - Ty\|^2]}{\|Sx - Ty\|^2 + \|Ex - Ty\|^2} + \beta_4 \frac{\|Ex - Ty\|^2 [\|Sx - Ex\|^2 + \|Ty - Fy\|^2]}{\|Sx - Ty\|^2 + \|Ex - Ty\|^2} \\
 &+ \beta_5 \frac{\|Sx - Ex\|^2 \|Ty - Fy\|^2}{\|Sx - Ty\|^2} + \beta_6 \frac{\|Ty - Fy\|^2 [1 + \|Sx - Ex\|^2]}{1 + \|Sx - Ty\|^2} \\
 &+ \beta_7 \frac{\|Sx - Fy\|^2 [\|Sx - Ex\|^2 + \|Ty - Fy\|^2]}{\|Sx - Fy\|^2 + \|Ex - Ty\|^2} \tag{1}
 \end{aligned}$$

for each $x, y \in C$, $Sx \neq Ty$ and $\|Sx - Ty\|^2 + \|Ex - Ty\|^2 \neq 0$ and $\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7$ being positive real number. Then E, F, T, S have a common unique random fixed point in C.

Proof: Let the function $g_0: \Omega \rightarrow C$ be arbitrary measurable function. By (1), there exists a function $g_1: \Omega \rightarrow C$ such that $T(t, g_1(t)) = E(t, g_0(t))$ for $t \in \Omega$ and for this function $g_1: \Omega \rightarrow C$, we can choose another function $g_2: \Omega \rightarrow C$ such that $F(t, g_1(t)) = S(t, g_2(t))$ for $t \in \Omega$, and so on. Inductively, we can define a sequence of functions for $t \in \Omega$, $\{y_n(t)\}$ such that

$$y_{2n}(t) = T(t, g_{2n+1}(t)) = E(t, g_{2n}(t)) \text{ and } y_{2n+1}(t) = S(t, g_{2n+2}(t)) = F(t, g_{2n+1}(t)) \text{ for } t \in \Omega \text{ and } n = 0, 1, 2, 3 \dots \tag{2}$$

From (1). Now consider for $t \in \Omega$

$$\begin{aligned}
 \|y_{2n}(t) - y_{2n+1}(t)\|^2 &= \|E(t, g_{2n}(t)) - F(t, g_{2n+1}(t))\|^2 \leq \\
 &\beta_1 \frac{\|S(t, g_{2n}(t)) - E(t, g_{2n}(t))\|^2 [\|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2 + \|E(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2 + \|S(t, g_{2n}(t)) - F(t, g_{2n+1}(t))\|^2]}{\|S(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2 + \|E(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ \beta_2 \frac{\|E(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2 [\|S(t, g_{2n}(t)) - E(t, g_{2n}(t))\|^2 + \|E(t, g_{2n}(t)) - F(t, g_{2n+1}(t))\|^2 + \|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2]}{\|S(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2 + \|E(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2 + \|E(t, g_{2n}(t)) - F(t, g_{2n+1}(t))\|^2} \\
 &+ \beta_3 \frac{\|S(t, g_{2n}(t)) - E(t, g_{2n}(t))\|^2 \|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2}{\|S(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ \beta_4 \frac{\|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2 [1 + \|S(t, g_{2n}(t)) - E(t, g_{2n}(t))\|^2]}{1 + \|S(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2} + \beta_5 \frac{\|S(t, g_{2n}(t)) - E(t, g_{2n}(t))\|^2 \|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2}{\|S(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ \beta_6 \frac{\|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2 [1 + \|S(t, g_{2n}(t)) - E(t, g_{2n}(t))\|^2]}{1 + \|S(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ \beta_7 \frac{\|S(t, g_{2n}(t)) - F(t, g_{2n+1}(t))\|^2 [\|S(t, g_{2n}(t)) - E(t, g_{2n}(t))\|^2 + \|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2]}{\|S(t, g_{2n}(t)) - F(t, g_{2n+1}(t))\|^2 + \|E(t, g_{2n}(t)) - T(t, g_{2n+1}(t))\|^2} \\
 \|y_{2n}(t) - y_{2n+1}(t)\|^2 &\leq \beta_1 \frac{\|y_{2n-1}(t) - y_{2n}(t)\|^2 [\|y_{2n}(t) - y_{2n+1}(t)\|^2 + \|y_{2n}(t) - y_{2n}(t)\|^2 + \|y_{2n-1}(t) - y_{2n+1}(t)\|^2]}{\|y_{2n-1}(t) - y_{2n}(t)\|^2 + \|y_{2n}(t) - y_{2n}(t)\|^2} \\
 &+ \beta_2 \frac{\|y_{2n}(t) - y_{2n}(t)\|^2 [\|y_{2n-1}(t) - y_{2n}(t)\|^2 + \|y_{2n}(t) - y_{2n+1}(t)\|^2 + \|y_{2n}(t) - y_{2n+1}(t)\|^2]}{\|y_{2n-1}(t) - y_{2n}(t)\|^2 + \|y_{2n}(t) - y_{2n}(t)\|^2 + \|y_{2n}(t) - y_{2n+1}(t)\|^2} \\
 &+ \beta_3 \frac{\|y_{2n-1}(t) - y_{2n}(t)\|^2 \|y_{2n}(t) - y_{2n+1}(t)\|^2}{\|y_{2n-1}(t) - y_{2n}(t)\|^2} + \beta_4 \frac{\|y_{2n}(t) - y_{2n+1}(t)\|^2 [1 + \|y_{2n-1}(t) - y_{2n}(t)\|^2]}{1 + \|y_{2n-1}(t) - y_{2n}(t)\|^2} \\
 &+ \beta_5 \frac{\|y_{2n-1}(t) - y_{2n}(t)\|^2 \|y_{2n}(t) - y_{2n+1}(t)\|^2}{\|y_{2n-1}(t) - y_{2n}(t)\|^2} + \beta_6 \frac{\|y_{2n}(t) - y_{2n+1}(t)\|^2 [1 + \|y_{2n-1}(t) - y_{2n}(t)\|^2]}{1 + \|y_{2n-1}(t) - y_{2n}(t)\|^2}
 \end{aligned}$$

$$\begin{aligned}
 & + \beta_7 \frac{\|y_{2n-1}(t) - y_{2n+1}(t)\|^2 [\|y_{2n-1}(t) - y_{2n}(t)\|^2 + \|y_{2n}(t) - y_{2n+1}(t)\|^2]}{\|y_{2n-1}(t) - y_{2n+1}(t)\|^2 + \|y_{2n}(t) - y_{2n+1}(t)\|^2} \\
 & = \beta_1 [\|y_{2n}(t) - y_{2n+1}(t)\|^2 + \|y_{2n-1}(t) - y_{2n+1}(t)\|^2] + \beta_3 \|y_{2n}(t) - y_{2n+1}(t)\|^2 \\
 & \quad + \beta_4 \|y_{2n-1}(t) - y_{2n}(t)\|^2 + \beta_5 \|y_{2n}(t) - y_{2n+1}(t)\|^2 + \beta_6 \|y_{2n}(t) - y_{2n+1}(t)\|^2 \\
 & \quad + \beta_7 [\|y_{2n-1}(t) - y_{2n}(t)\|^2 + \|y_{2n}(t) - y_{2n+1}(t)\|^2]
 \end{aligned}$$

$$(1 - 2\beta_1 - \beta_3 - \beta_5 - \beta_6 - \beta_7) \|y_{2n}(t) - y_{2n+1}(t)\|^2 \leq (\beta_1 + \beta_4 + \beta_7) \|y_{2n-1}(t) - y_{2n}(t)\|^2$$

$$\|y_{2n}(t) - y_{2n+1}(t)\|^2 \leq \left(\frac{\beta_1 + \beta_4 + \beta_7}{(1 - 2\beta_1 - \beta_3 - \beta_5 - \beta_6 - \beta_7)} \right) \|y_{2n-1}(t) - y_{2n}(t)\|^2$$

$$\|y_{2n}(t) - y_{2n+1}(t)\| \leq k \|y_{2n-1}(t) - y_{2n}(t)\| \text{ where } k = \left(\frac{\beta_1 + \beta_4 + \beta_7}{(1 - 2\beta_1 - \beta_3 - \beta_5 - \beta_6 - \beta_7)} \right)^{1/2}. \text{ Replacing } 2n \text{ by } n \text{ we get}$$

$$\|y_n(t) - y_{n+1}(t)\| \leq k \|y_{n-1}(t) - y_n(t)\|$$

$$\text{on further reducing } \|y_n(t) - y_{n+1}(t)\| \leq K^n \|y_0(t) - y_1(t)\| \text{ for all } t \in \Omega. \quad (3)$$

Now we shall prove for $t \in \Omega$, $\{y_n(t)\}$ is a Cauchy sequence. For this for every positive integer p we have, $t \in \Omega$.

$$\begin{aligned}
 \|y_n(t) - y_{n+p}(t)\| & = \|y_n(t) - y_{n+1}(t) + y_{n+1}(t) - y_{n+2}(t) + \dots + y_{n+p-1}(t) - y_{n+p}(t)\| \\
 & \leq \|y_n(t) - y_{n+1}(t)\| + \|y_{n+1}(t) - y_{n+2}(t)\| + \dots + \|y_{n+p-1}(t) - y_{n+p}(t)\| \\
 & \leq [K^n + K^{n+1} + \dots + K^{n+p-1}] \|y_0(t) - y_1(t)\| \\
 & = K^n [1 + k + K^2 + \dots + K^{p-1}] \|y_0(t) - y_1(t)\| \\
 & \leq \frac{K^n}{(1-k)} \|y_0(t) - y_1(t)\| \text{ for all } t \in \Omega.
 \end{aligned}$$

$$\|y_n(t) - y_{n+p}(t)\| \rightarrow 0 \text{ as } n \rightarrow \infty \text{ for } t \in \Omega. \quad (4)$$

From eq (4), it follows that for $t \in \Omega$, $\{y_n(t)\}$ is a Cauchy sequence and hence is convergent in closed subset C of Hilbert space H .

$$\text{For } t \in \Omega, \text{ let } \{y_n(t)\} \rightarrow y(t) \text{ as } n \rightarrow \infty. \quad (5)$$

Since C is closed, g is a function from C to C and consequently the subsequence $\{E(t, g_{2n}(t))\}$, $\{F(t, g_{2n+1}(t))\}$, $\{T(t, g_{2n+1}(t))\}$ and $\{S(t, g_{2n+2}(t))\}$ of $\{y_n(t)\}$ for $t \in \Omega$ also converges to the $\{y(t)\}$ and continuity of E, F, T, S gives

$$\begin{aligned}
 \{E(t, S(t, g_n(t)))\} & \rightarrow E(t, y(t)), \{S(t, E(t, g_n(t)))\} \rightarrow S(t, y(t)), \\
 \{F(t, T(t, g_n(t)))\} & \rightarrow F(t, y(t)) \text{ and } \{T(t, S(t, g_n(t)))\} \rightarrow T(t, y(t)) \\
 E(t, y(t)) & = S(t, y(t)), F(t, y(t)) = T(t, y(t)) \text{ for } t \in \Omega \text{ from (1)} \quad (6)
 \end{aligned}$$

Consider for $t \in \Omega$

$$\begin{aligned}
 \|E(t, y(t)) - y(t)\|^2 & = \|E(t, y(t)) - y_{2n+1}(t) + y_{2n+1}(t) - y(t)\|^2 \\
 & \leq 2\|E(t, y(t)) - y_{2n+1}(t)\|^2 + 2\|y_{2n+1}(t) - y(t)\|^2 \\
 & \leq 2\|E(t, y(t)) - F(t, g_{2n+1}(t))\|^2 + 2\|y_{2n+1}(t) - y(t)\|^2 \text{ by (4)}
 \end{aligned}$$

[By Parallelogram law $\|x + y\|^2 \leq 2\|x\|^2 + \|y\|^2$]

$$\begin{aligned}
 \|Ex - Fy\|^2 &\leq 2\beta_1 \frac{\|(t, g_{2n+1}(t)) - T(t, g_{2n+1}(t))\|^2 [1 + \|(t, y(t)) - S(t, y(t))\|^2]}{1 + \|(t, y(t)) - (t, g_{2n+1}(t))y\|^2} \\
 &+ 2\beta_2 [\|(t, y(t)) - S(t, y(t))\|^2 + \|(t, g_{2n+1}(t)) - T(t, g_{2n+1}(t))\|^2] \\
 &+ 2\beta_3 \frac{\|S(t, y(t)) - E(t, y(t))\|^2 [\|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2 + \|E(t, y(t)) - T(t, g_{2n+1}(t))\|^2]}{\|S(t, y(t)) - T(t, g_{2n+1}(t))\|^2 + \|E(t, y(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ 2\beta_4 \frac{\|E(t, y(t)) - T(t, g_{2n+1}(t))\|^2 [\|S(t, y(t)) - E(t, y(t))\|^2 + \|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2]}{\|S(t, y(t)) - T(t, g_{2n+1}(t))\|^2 + \|E(t, y(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ 2\beta_5 \frac{\|S(t, y(t)) - E(t, y(t))\|^2 \|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2}{\|S(t, y(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ 2\beta_6 \frac{\|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2 [1 + \|S(t, y(t)) - E(t, y(t))\|^2]}{1 + \|S(t, y(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ 2\beta_7 \frac{\|S(t, y(t)) - F(t, g_{2n+1}(t))\|^2 [\|S(t, y(t)) - E(t, y(t))\|^2 + \|T(t, g_{2n+1}(t)) - F(t, g_{2n+1}(t))\|^2]}{\|S(t, y(t)) - F(t, g_{2n+1}(t))\|^2 + \|E(t, y(t)) - T(t, g_{2n+1}(t))\|^2} \\
 &+ 2\|y_{2n+1}(t) - y(t)\|^2 \text{ by (2)} \\
 &\leq 2\beta_1 \frac{\|y(t) - y(t)\|^2 [1 + \|y(t) - S(t, y(t))\|^2]}{1 + \|y(t) - y(t)\|^2} \\
 &+ 2\beta_2 [\|y(t) - S(t, y(t))\|^2 + \|y(t) - y(t)\|^2] \\
 &+ 2\beta_3 \frac{\|S(t, y(t)) - E(t, y(t))\|^2 [\|y(t) - y(t)\|^2 + \|E(t, y(t)) - y(t)\|^2]}{\|S(t, y(t)) - y(t)\|^2 + \|E(t, y(t)) - y(t)\|^2} \\
 &+ 2\beta_4 \frac{\|E(t, y(t)) - y(t)\|^2 [\|S(t, y(t)) - E(t, y(t))\|^2 + \|y(t) - y(t)\|^2]}{\|S(t, y(t)) - y(t)\|^2 + \|E(t, y(t)) - y(t)\|^2} \\
 &+ 2\beta_5 \frac{\|S(t, y(t)) - E(t, y(t))\|^2 \|y(t) - y(t)\|^2}{\|S(t, y(t)) - y(t)\|^2} \\
 &+ 2\beta_6 \frac{\|y(t) - y(t)\|^2 [1 + \|S(t, y(t)) - E(t, y(t))\|^2]}{1 + \|S(t, y(t)) - y(t)\|^2} \\
 &+ 2\beta_7 \frac{\|S(t, y(t)) - y(t)\|^2 [\|S(t, y(t)) - E(t, y(t))\|^2 + \|y(t) - y(t)\|^2]}{\|S(t, y(t)) - y(t)\|^2 + \|E(t, y(t)) - y(t)\|^2} + 2\|y(t) - y(t)\|^2 \\
 &\|E(t, y(t)) - y(t)\|^2 \leq (2\beta_2 + 2\beta_7) [\|y(t) - S(t, y(t))\|^2 + (2\beta_3 + 2\beta_4) \|E(t, y(t)) - y(t)\|^2] \\
 &(1 - (2\beta_3 + 2\beta_4)) \|E(t, y(t)) - y(t)\|^2 \leq (2\beta_2 + 2\beta_7) \|y(t) - S(t, y(t))\|^2 \\
 &(1 - 2(\beta_2 + \beta_3 + \beta_4 + \beta_7)) \|E(t, y(t)) - y(t)\|^2 \leq 0 \\
 &\|E(t, y(t)) - y(t)\|^2 = 0
 \end{aligned}$$

$$2(\beta_2 + \beta_3 + \beta_4 + \beta_7) = 1$$

$$E(t, y(t)) = y(t) \text{ for } t \in \Omega \tag{7}$$

From eq (6) and eq (7)

$$E(t, y(t)) = y(t) = S(t, y(t)), \tag{8}$$

In an exactly similar way we can prove for all $t \in \Omega$.

$$F(t, y(t)) = y(t) = T(t, y(t)) \tag{9}$$

Again, if $A: \Omega \times C \rightarrow C$ is a continuous random operator on a non – empty subset C of a separable Hilbert space H then for any measurable function $f: \Omega \rightarrow C$, the function $h(t) = A(t, f(t))$ is also measurable [8]. It follows from the construction of $\{y_n(t)\}$ by (4) and the above consideration that $\{y_n(t)\}$ is a sequence of measurable functions. From (5), it follows that $y(t)$ for $t \in \Omega$ is also a measurable function. This fact along with (8) and (9) shows that $g: \Omega \rightarrow C$ is a common random fixed point of E, F, S, T .

Uniqueness: Let $h: \Omega \rightarrow C$ be another random fixed point common to E, F, T and S that is for $t \in \Omega$, $E(t, g(t)) = g(t)$, $F(t, h(t)) = h(t)$, $T(t, h(t)) = h(t)$, $S(t, g(t)) = g(t)$. Then for $t \in \Omega$.

$$\begin{aligned} \|g(t) - h(t)\|^2 &= \|E(t, g(t)) - F(t, h(t))\|^2 \\ &\leq \beta_1 \frac{\|h(t) - T(t, h(t))\|^2 [1 + \|g(t) - S(t, g(t))\|^2]}{1 + \|g(t) - h(t)\|^2} \\ &\quad + \beta_2 [\|g(t) - S(t, g(t))\|^2 + \|h(t) - T(t, h(t))\|^2] \\ &\quad + \beta_3 \frac{\|S(t, g(t)) - E(t, g(t))\|^2 [\|T(t, h(t)) - F(t, h(t))\|^2 + \|E(t, g(t)) - T(t, h(t))\|^2]}{\|S(t, g(t)) - T(t, h(t))\|^2 + \|E(t, g(t)) - T(t, h(t))\|^2} \\ &\quad + \beta_4 \frac{\|E(t, g(t)) - T(t, h(t))\|^2 [\|S(t, g(t)) - E(t, g(t))\|^2 + \|T(t, h(t)) - F(t, h(t))\|^2]}{\|S(t, g(t)) - T(t, h(t))\|^2 + \|E(t, g(t)) - T(t, h(t))\|^2} \\ &\quad + \beta_5 \frac{\|S(t, g(t)) - E(t, g(t))\|^2 \|T(t, h(t)) - F(t, h(t))\|^2}{\|S(t, g(t)) - T(t, h(t))\|^2} \\ &\quad + \beta_6 \frac{\|T(t, h(t)) - F(t, h(t))\|^2 [1 + \|S(t, g(t)) - E(t, g(t))\|^2]}{1 + \|S(t, g(t)) - T(t, h(t))\|^2} \\ &\quad + \beta_7 \frac{\|S(t, g(t)) - F(t, h(t))\|^2 [\|S(t, g(t)) - E(t, g(t))\|^2 + \|T(t, h(t)) - F(t, h(t))\|^2]}{\|S(t, g(t)) - F(t, h(t))\|^2 + \|E(t, g(t)) - T(t, h(t))\|^2} \\ \|g(t) - h(t)\|^2 &\leq \beta_1 \frac{\|h(t) - h(t)\|^2 [1 + \|g(t) - g(t)\|^2]}{1 + \|g(t) - h(t)\|^2} + \beta_2 [\|g(t) - g(t)\|^2 + \|h(t) - h(t)\|^2] \\ &\quad + \beta_3 \frac{\|g(t) - g(t)\|^2 [\|h(t) - h(t)\|^2 + \|g(t) - h(t)\|^2]}{\|g(t) - h(t)\|^2 + \|g(t) - h(t)\|^2} \\ &\quad + \beta_4 \frac{\|g(t) - h(t)\|^2 [\|g(t) - g(t)\|^2 + \|h(t) - h(t)\|^2]}{\|g(t) - h(t)\|^2 + \|g(t) - h(t)\|^2} \\ &\quad + \beta_5 \frac{\|g(t) - g(t)\|^2 \|h(t) - h(t)\|^2}{\|g(t) - h(t)\|^2} + \beta_6 \frac{\|h(t) - h(t)\|^2 [1 + \|g(t) - g(t)\|^2]}{1 + \|g(t) - h(t)\|^2} \\ &\quad + \beta_7 \frac{\|g(t) - h(t)\|^2 [\|g(t) - g(t)\|^2 + \|h(t) - h(t)\|^2]}{\|g(t) - h(t)\|^2 + \|g(t) - h(t)\|^2} \end{aligned}$$

$$\|g(t) - h(t)\|^2 \leq 0$$

$\Rightarrow g(t) = h(t)$ for all $t \in \Omega$.

References:

- [1] Bharucha – Reid, A. T., Fixed Point Theorems in Probabilistic analysis, Bull. Amer. Math. Soc., 82, 611 – 645 (1996).
- [2] Choudhary, B.: A common unique fixed point theorem for two random operators in Hilbert space. IJMMS, 32(3), 177 – 182 (2002).
- [3] Hans, P. : Random fixed point theorems, Transactions of the Prague Conference on Information Theory, Statistical Decision Functions, Random Process, pp. 105 – 125, (1957).

- [4] Himmelberg, C. J. Measurable relations, *Fund Math*, 87, 53 – 72,(1975).
- [5] Nair, S. and shrivastava, S. : Fixed point theorem for Hilbert spac, *Jour. Pure Math.* 22, 33 – 37, (2005).
- [6] Nashin, H.: Existence of common random fixed point and random best approximation for non – commuting random operators, *Bulletin if the Institute of Mathematics* vol. 5 No. 1, pp 25 – 40, (2010).
- [7] Regan, D. O.: Fixed Points and Random Fixed Points for Weakly inward Approximable Maps. *Proceeding of the American and Mathematical Society* 126 No. 10, 3045 – 3053, (1998).

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

