

Sexual Behavior and Risk Perception of HIV Infection Among Youths in Bahirdar Town

Teferi Beyene (MA)
Bahir Dar University, Bahir Dar, PO.Box 407

ABSTRACT

Ethiopia is a developing country with a demographic profile dominated by a young population. Due to biological, psychological, socio cultural and economic factors, young people, particularly those aged 15-29 years, are generally at a high risk of HIV/AIDS and other reproductive health problems. This paper presented results of a cross-sectional descriptive study conducted in Bahir Dar Town, northwest Ethiopia, to assess sexual behavior and risk perception of HIV infection among youths. Both quantitative and qualitative method of data-collection methods were employed to conduct the study. For quantitative data collection, a household questioner survey was conducted among 232 youth, aged 15-29 years, within the 4kebeles (villages) of the town. Qualitative data were collected by conducting focus-group discussions and in-depth interviews with 17 participants. Descriptive statistics was used to characterize socio-economic, demographic and behavioral variables and the level of risk perception of HIV. Chi-square was used to examine the association of socio-economic, demographic and behavioral variables to HIV risk perception. Logistic regression model was used further examined to identify the prediction independent variables to risk perception. Data obtained by interview and focus group discussion were qualitatively analyzed. The minimum mean age at first sexual commencement was 16.71(± 1.45) years and the maximum is 26.25(± 1.18). Socio-economic and demographic variables such as Job($\chi^2=4.7151$, $p=0.03$), alcohol use($\chi^2=16.8405$, $p=0.001$), monthly income($\chi^2=12.769$, $p=0.026$), gender($\chi^2=9.4788$, $p=0.002$) and education status($\chi^2=11.8883$, $p=0.003$) were significantly associated with risk perception of HIV among the youth. Behavioral variables such as sex ever had ($\chi^2=10.1561$, $p=0.001$), age at first sex ($\chi^2=7.524$, $p=0.023$), no of sexual partners($\chi^2=7.2156$, $p=0.002$) and knowledge of HIV status($\chi^2=16.0624$, $p=0.000$) were also significantly associated with risk perception of HIV among the youths. IN logistic regression model, age ($z=-2.13$, $p=0.033$), education status ($z=-4.36$, $p=0.000$), marital status ($z=-2.48$, $p=0.013$), alcohol use ($z=4.88$, $p=0.000$), and knowledge of HIV status ($z=-3.69$, $p=0.000$) were significantly and independently predicted HIV risk perception. In conclusion, further research should be conducted to better understand the nature of association between the above socio-economic and demographic, and sexual behavioral variables with risk perception of HIV/AIDS Staking behaviors.

Keywords: sexual behavior, risk perceptions

1. Introduction

HIV and AIDS, since it appeared in 1981, has become one of the most devastating epidemics in human history. Since then more than 20 million people have lost their lives, and more 80 million people are projected to die of it by 2010. Currently there are about 33 million carriers across the worldwide. HIV/AIDS have become a serious threat with its adverse consequences on people and all sectors of economic development (UNAIDS, 2007).

Among the HIV/AIDS infected people, more than 90% of them live in the developing countries. Of the 33 million people living with the virus worldwide, Africa hosts about 22 million and Sub-Saharan Africa, with over 10 % of the world's population, is a home to two - thirds of HIV/AIDS carriers. When the spread of AIDS is seen on gender bases, the prevalence is high among women. Usually they are being infected at an earlier age than men. The main reasons behind this difference are wide spread poverty among women, gender inequality, and social and political inertia. Especially, in developing countries these factors have widened the gap of AIDS prevalence between both sexes (UNFPA, 2008).

There are many ways by which AIDS can spread from one person to the other. But of all, unsafe sex has been identified to be the leading cause of HIV infection globally. From this, it is obvious that the epidemic affects more the youth group than the other segment of society. Studies indicate that young people under the age of 25 are estimated to account for half of all new HIV infections worldwide, with an average of two young people being infected every hour of a day (Doll LS, Beeker, 1996).

HIV/AIDS is rapidly spreading across the world, and has claimed millions of lives and billions of dollars to humanity. Studies have shown that despite biological factors like sexual relationships, drug injection, blood transfusion, that intensify the spread of this disease there are also other underlying psychological factors like anxiety, stress, depression, lack of communication, posttraumatic stress disorder, negative effect, domestic problems, and others that make someone vulnerable to HIV infection. Keeping this in mind, contemporary researchers have considered some more relevant psychological factors that seem more influential in expanding HIV infection. These are HIV risk perception, peer pressure and drug abuse behaviors (Fisher, Byrne, White, 1988).

Ethiopia, the third largest country in Sub-Saharan Africa, is located in the horn neighboring countries like Sudan, Eritrea, Djibouti, Somalia and Kenya. It has a total area of 1.13 million sq km with an estimated 80 million population. It is one of the Sub-Saharan African countries that are seriously hit by HIV/AIDS. In this country too it has indiscriminately affected all groups of the community, especially the young, children, and women.

The first evidence of HIV/AIDS infection was found in 1984 while the first two cases were reported in 1986 from Addis Ababa hospitals to the Ministry Of Health /MOH/. Since then, for more than two decades, the epidemic has continued to spread at a rapid pace both in urban and rural areas (UNAIDS, 2007). This time, Ethiopia is the 16th country with highest HIV and AIDS prevalence in the world, and the third largest with numbers of HIV/AIDS carriers in Africa.

HIV/AIDS pandemic has hit the country with catastrophic consequences. The average national prevalence rate among the adult population is estimated to be 3.51 % (10.5 % urban and 1.94 % rural) with a steadily rise in the rural areas. Nationally, the estimated prevalence among males is 3.8 %. The number of people living with HIV/AIDS in 2006 is estimated at 1.3 million, of which 96,000 are children. In 2006 some 207,270 new People living with HIV/AIDS were in need of antiretroviral (ART) throughout the country (UN AIDS Rport-2008).

The Amhara regional state is one of the nine regions of Ethiopia in which the spread of HIV/AIDS spread in this region is one of the worst in the country with high prevalence in urban areas in particular. The AIDS prevalence in this region was estimated at 6.1%, 6.5%, 6.7%, and 7% for 2003, 2004, 2005 and 2006 respectively.

In this region, while the spread of the epidemic seems to be stabilizing in urban areas since 1996/97, it shows a rising trend in its rural part. Hence, the overall trend of AIDS prevalence in the region shows an increasing rate; where in 2006 it was estimated at 15% and 5.9% for urban and rural areas respectively (HAPCO, 2007).

Currently, the number of people living with HIV/AIDS in this region is estimated to be 673,488. Even though, there is no study showing reasons for the wider variation between the national and the regional figures of HIV carriers, there are some speculations like high rate of poverty, drought and famine, illiteracy, unemployment, rural to urban migration, silence about HIV/AIDS among family members, harmful traditional practices like early marriage and female genital mutilation, and the polygamous culture in the region to be the possible causes (ADA, 2008).

The HIV/AIDS epidemic shows great variation between urban and rural areas of the Amhara region. The prevalence is at least five times higher in urban areas than in rural. In 2005, the infection level was ranging from 0.5 percent in rural areas to 20 percent in urban areas such as in the town of Bahir Dar (ADA, 2008).

As stated above, though the prevalence and spread of the disease is high, studies conducted on assessing the behavioral risk factors of its transmission and prevention among the various segments of the society in Ethiopia are so scarce. But it is obvious that in order to plan intervention measures, it is important to assess people's knowledge of HIV infection, sexual behaviors, perceived risks of infection, and attitude on VCT. Therefore, this research paper will be conducted in view of these basic issues by targeting the youth group in Bahir Dar Town.

2. Review of Related Literature

2.1 Sexuality in adolescents

Attitudes towards particular aspects of sexuality are always changing. For instance, since the recent times, Masturbation and premarital sex have become more usual. Not only this, but also there is a greater openness about sexual orientations, alternative behavior, and gender identities, with a significant degree of debate about their acceptability (Kelly Gary F, 1998). This seems something related to an increasing human awareness on sexuality. According to Kelly (1998), children move from a generalized awareness of their sexual nature to more specific experiences of sexual feelings. Likewise, adolescents explore their sexuality through relationships with others and there is evidence that they have become sexually active at increasingly younger ages. And he came to the conclusion that sexuality is a universal phenomenon in all-young people.

2.2 Socio-Economic and Demographic factors affecting risk perception

Young people are not a homogenous group. They have different needs owing to lots of factors like gender, age, marital status, income or employment status, cognitive development stage, educational level, access to media, and cultural norms. Despite the various factors for youth heterogeneity, different studies have shown that there are consistent trend towards sex outside marriage among the youth. There are also other common observations including weakening of social ties, late marriage, and changing sexual norms. In many instances, economic pressures and social norms are found to be among the major factors that force young people into sexual activity (Hughes and McCauley, 1998).

2.3 Risk perception of HIV/AIDS and STDs

Various literatures written on health related behavior consider the perception of being at risk of infection to be one of the necessary conditions for behavioral change (Lavra , 2002). Moreover, the degree of the perceived risk seems to affect individual actual control in adopting preventive measures. Individual risk perception is dependent on the perception of other members in a given network. Individual risk perception as well as individual knowledge is likely to be influenced by a social environment as long as social interaction allows information exchange, facilitates common evaluation and definition of the meaning and its validity.

2.4 Sexual Behavior and Risk Perception

Several studies have found that perception of risk is strongly related to the self protecting behavior of individuals (Adih and Alexander, 1999; Diamond and Madise, 2001). This is largely because of the adoption of protective behaviors, which is unlikely to occur unless the person is well aware of the risk of HIV infection. Studies show that people can judge their risk of HIV infection (Maharaj, 2004). However, sometimes those who are at risk may not perceive their risk and are less motivated to protect themselves (Varga, 2001).

2.5 Knowledge about HIV/AIDS and Sexual Behavior

Individual's knowledge of HIV transmission and accurate assessment of their own risk seem to be among the key factors in adoption of safer sexual practice (UNAIDS 2001). Knowledge of HIV/AIDS and related sexual behavior among the youths within the age range of 15-29 is of particular interest because the period between sexual initiation and marriage is a time of sexual experimentation that may involve high-risk behaviors. As indicated in EDHS (2005) only around one –fifth of women and one-third of men within the age of 15-24 know all of the basic facts about this disease, and the level of knowledge about it does not vary greatly among the young people.

2.6 Knowledge of HIV Status

Knowledge of HIV status helps HIV-negative individuals to make specific decisions to reduce risk and practice safer sex. For those who are HIV positive, knowledge of their status allows them to take action to protect their sexual partner, access treatment and plan for their future life. A study in South Africa depicts that young men who had been tested for HIV were more likely to have used condom than those who had not been tested ($p=0.001$). Testing for HIV/AIDS during pregnancy is especially important to prevent a mother to child transmission. However, in Ethiopia, as revealed in the EDHS (2005), knowledge of HIV status among the community is very low, where only 6% of men and 2% of women have been tested.

2.7 The role of non-sexual risk behaviors for HIV infection

A study conducted on adolescent reproductive health indicates that having ever used alcohol and drugs was a risk factor for ever having had sex, having more sexual partners over life time, and having more than one partner during the last three months (Adolescent Reproductive Health Task Force, August 2003). Taking alcohol and some drugs is a common practice among the youths both in the urban and rural areas of Ethiopia. The effect of taking alcohol and using chat regularly was observed on sexual behavior. Amongst the youth who were reported to have had risky sex in a previous 12 months, 44% of them had used alcohol and chat regularly (HIV/AIDS Behavioral Surveillance Survey (BSS), Round one, Ethiopia, 2002). A study conducted on casual sex-debuts among female adolescents in Addis Ababa also shows that 'alcohol' and 'chat' use have strong links with the incidence of 'rape' as a factor contributing to early sex initiation (Fekadu .,2001).

2.8 Misperception about HIV/AIDS

The routes of HIV transmission are well documented by scientists, but health officials and concerned social behavioral scientists continually grapple with the public's unfounded fears concerning the potential for HIV

2.9 Condom use

According to some studies, although the majority of young people have heard of AIDS, many of them do not know how it is transmitted and do not believe that they are at risk. Those who know something of HIV often do not protect themselves because they lack the knowledge, the support or the means to adopt safe behavior (Young people and HIV/AIDS opportunity in crisis, 2002). For instance, young girls and women are regularly and repeatedly denied information about and access to condoms.

3. The Methods

3.1 Sampling Method

The population of this study covers youths living and working in Bahir Dar City administrations which are divided into 9 kebeles. After grouping the kebeles in to three strata, the first strata which represent the central zone of the town was taken using purposive sampling based on the points mentioned above. Then the sample

size was distributed proportionally to the four kebeles under these strata and samples were selected using systematic random sampling. Eventually, the quantitative data was gathered by way of survey questionnaire.

In addition to distributing questionnaire to 247 young people, key informant interview and group discussion with politicians, community leaders, parents, and other stakeholders were also conducted to get adequate information about the problem. In this case also availability and snowball sampling was employed to draw the participants in these techniques. Accordingly, the 4 kebeles administrators, one representative from the regional HIV/AIDS prevention and control secretariat, four youth group leaders, 4 parents from both sexes, and two representatives from NGOs working on this area were interviewed for a more insight into the complex pattern of sexual behavior and risk perception of the youth about HIV/AIDS in the study area.

3.2 Data Collection Tools

The data required for this study was collected using both qualitative and quantitative methods. Preparing and distributing a structured questionnaire was the instrument for collecting the quantitative data.

Survey questionnaire: a well organized and structured questionnaire was prepared for the data collection process. The questionnaire was first developed in English, translated to Amharic and translated back into English so that accuracy and consistency in the wording are ensured. The questionnaire contained three main parts. The first Part of the questionnaire was contained demographic and socio-economic characteristics of participants; the second part was dealt about sexual behaviors of participants; the third part was about the knowledge of the respondents concerning HIV/AIDS and STIs, knowledge and use of condoms, and the level of risk perception of HIV infection among the youth.

Focus group discussion (FGD): was conducted to generate qualitative data that can Supplement the results of the questionnaire. And because of the sensitiveness and personal nature of the issue under investigation, conducting FGDs play irreplaceable role. There was one group include representatives from at each kebele. The participants were selected from those young adults, politicians, community leaders, and representatives from governmental and nongovernmental organizations who were willing to participate on discussion. Six FGDs with a group of 8-10 discussants participated in the discussion. The principal investigator was the facilitator for the FGDs. Unstructured questionnaire with open ended questions were also distributed to some stakeholders in order to receive their views and understanding on the problem. Most discussion guides are open ended to give more opportunity for discussion. Discussions were held in local language- Amharic, recorded by a tape recorder and later translated into English and analyzed by the researcher. **In-depth interviews** with selected key informants were also made to substantiate and crosscheck the information obtained from the questionnaire.

3.3 Data Collection Procedures

Four enumerators who have diploma or above educational level, with better understanding of the issue and the study area, who can speak the local language, and were interested to participate in this study was selected and trained to administer the survey questioner. Training manual and working guideline had been prepared for this purpose. The questionnaires were prepared in English but were translated into the local language, Amharic, so that the enumerators can understand it better and explain fully to their interviewees.

One supervisor from the town health office was selected and trained to control the overall data collection process. The writer was supervise everything daily and would give orientation and supervision to all the enumerators and supervisors on how the work was going on. The key informant interviews and group discussion was conducted by the researcher himself based on the action plan of the study. In this process interview checklists were used to go through all the relevant points and exploit the views and opinions of the participants.

The questionnaires were pre-tested in one kebele prior to the actual data collection on 15 respondents those who will not be included in the main survey. The result of the pre-test was used to revise the questionnaires based on the responses from the interviewees. During the actual data collection times the principal investigator and the supervisor was check whether the questionnaires were filled correctly or not. And accordingly, some advises was given to the enumerators so that they continue filling the questionnaires without any problem.

3.4 Variables Treated in the Study

The variables for HIV/AIDS risk perception; sexual behavior and awareness were defined. The information was collected on variables such as score of socio-economic and demographic characteristics, sexual behavior, and risk perception of the participants. The variables are basically of two types, dependent and independent variables.

Dependent Variable

The dependent variable in this study was individual's level of risk perception of HIV infection. In order to measure the risk perception of getting infected by HIV, respondents were asked a key question '*do you consider your chance of getting HIV to be high, medium, low or no chance at all?*' For the purpose of analysis, respondents were divided into high risk, medium risk and low or no risk categories depending on their responses.

Independent variables

The explanatory variables that influence perceived risk of HIV infection were selected based on the literature review. These factors were categorized into socio-economic, demographic and behavioral factors. The socio-economic variables included level of education, monthly income, chat chewing, smoking and alcohol use. The demographic factors included in this study were age, sex, marital status. And the behavioral factors included were age at first sex, number of life time sex partners, contact with CSW, condom use, ever having an STDs, knowledge of HIV status, and age difference with sex partner. Most of the variables mentioned above are self explanatory variables.

3.5 Methods of Data Analysis

Following the data collection in the field using various instruments, the data was analyzed by using Stata version 10 computer software packages. Analysis of frequencies of different variables and chi- squared test for some selected variables were done. Logistic regression was used in the analysis to examine the relationship between risky sexual behavior and perception of HIV risk in categorical variables based on gender while controlling for other factors that are likely to influence sexual behavior.

4. Results and Discussions

4.1. Results

In this study the sample size was distributed based on selected demographic and socio-economic characteristics which include sex, age, level of education, monthly income, alcohol drink, smoking, and chat chewing. Hence, 247 individuals were drawn as samples from the total population and they were addressed using questionnaires. Of the 247 questionnaires distributed to gather information from the determined sample size the researcher collected 232 fully completed questionnaires. And the remaining 15 questionnaires were missing or inappropriately filled and hence rejected by the researcher.

The participants considered in this study were both males 134 (57.75%) & females 98 (38.79%). As can be seen from table 4.1 below, 93 (40.09 %) of the respondents were in the age group of 15-19, 93 (40.09%) of them were between 20 and 24, and the remaining 46 (19.83%) were in the age group of 25-29. With regard to their educational status, 45 of them (19.40%) were of grade 1-8, 113 (48.71%) grade 9-12, 74 of them (31.90%) were diploma holders and above. The sample distribution based on their religion shows that 156 (67.24%) were Orthodox Christians, 61 (26.29%) were Muslims, whereas the proportion of Protestants and followers of other religions were 11(4.74%) and 4(1.72%) respectively.

Speaking about their source of income, the participants of this study reported that they have various forms of income generating activities. Accordingly, 62 (26.72%) of the respondents earn a monthly income of below 300 Birr, 85 (36.64%) of them earn between 301 and 500, 48(20.69%) of them earn 501-1000, 24(10.34%) earn 1001-1500, 10(4.31%) earn 1501-2000 and the rest 3(1.29%) earn a monthly income of 2000 birr and above. It is also found out that 138(59.48%), 38(16.38%) and 60(25.86%) of the participants drink alcohol, smoke cigarette and chew chat respectively.

Table1. Percentage distribution of respondents by selected background characteristics

Respondents Characteristics		Sex of respondents				total	
		Male		female			
		frequency	%	frequency	%	frequency	%
Age	15-19	56	41.79	37	37.76	93	40.09
	20-24	52	38.81	41	41.84	93	40.09
	25-29	26	19.40	20	20.41	46	19.83
Marital status	Single	106	79.10	58	59.18	164	70.69
	Married	26	19.40	27	27.55	53	22.84
	Divorced	2	1.49	9	9.18	11	4.74
	widowed	—	—	4	4.08	4	1.72
Have a job	Yes	68	50.75	60	61.22	128	55.17
	No	66	49.25	38	38.78	104	44.83
Level of education	Can't read & write	—	—	—	—	—	—
	Can read & write	—	—	—	—	—	—
	Grade 1-8	30	22.39	15	15.31	45	19.40
	Grade 9-12	65	48.51	48	48.98	113	48.71
	Diploma & above	39	29.10	35	35.75	74	31.90
Monthly income	Below 300 birr	33	24.63	29	29.59	62	26.72
	301-600	35	26.12	50	51.02	85	36.64
	601-1000	34	25.37	14	14.29	48	20.69
	1001-1500	21	15.67	3	3.06	24	10.34
	1501-2000	8	5.97	2	2.04	10	4.31
	Above 2000	3	2.24	—	—	3	1.29

The assessment of family profile of the respondents indicate that 79 (34.05%) and Sixty five (28.08%) of their mothers and fathers respectively were able to read and write. Likewise, the mothers of 25.43% of the participants were householders while the fathers of 58 (25.43%) of the respondents were found to be civil servants.

4.2. Sexual behavior of participants

This part shows the percentage distribution of some risk related factors which will further be investigated using chi-square test and then binary logistic regression to discuss the net effect of individual factors on risk perception of HIV infection. The sexual behavior of participants has been shown using various methods. Table 2. below shows that 208(89.66%) of the respondents reported to have ever had sex where the percentage was slightly higher among males (90.3%) than among females (88.78%).

Table 2. Distribution of respondents by variables related to risky sexual behavior

Respondents characteristics		Sex of respondents				total	
		male		female			
		frequency	%	frequency	%	frequency	%
Ever had sex	Yes	121	90.30	87	88.78	208	89.66
	No	13	9.70	11	11.22	24	10.34
Age at first sexual intercourse	15-19 years	78	64.46	51	52.04	129	62.02
	20-24 years	35	28.93	28	28.57	63	30.29
	25-29 years	8	6.61	8	8.16	16	7.69
Reason to have sex	Fell in love	41	33.88	28	32.18	69	33.17
	Sexual desire	51	42.15	35	40.23	86	41.35
	Marriage	11	9.09	8	9.20	19	9.13
	Rap	-	-	-	-	-	-
	Peer pressure	1	0.83	4	4.60	5	2.40
	Get drunk	11	9.09	6	6.90	17	8.17
	Others	6	4.96	6	6.90	12	5.77
First sexual relationship	Lover	58	47.93	42	48.28	100	48.08
	Causal partner	49	40.50	34	39.08	83	39.90
	Partner	12	9.92	10	11.49	22	10.58
	others	2	1.65	1	1.15	3	1.44
No of sexual partners ever had	1	43	35.54	30	34.48	73	35.10
	2	21	17.36	17	19.54	38	18.27
	3	30	24.79	24	27.59	54	25.96
	4	10	8.26	8	9.20	18	8.65
	Above 4	17	14.05	8	9.20	25	12.02

In this study the minimum and maximum ages of first sex have been reported to be 14 and 26 respectively. Among those 232 respondents who ever had sex, 73(35.10%) of them had at least 1 sexual partner, 38 respondents(18.27%) reported to have two sexual partners, 54 of them (25.96%) had exactly 3 partners, 18 respondents(8.65%) had four, and the rest 25 (12.02%) respondents had four and above sex partners.

Among the respondents who ever had sex, 100 of them (48.08%) did their first sex with their lover, 83 respondents (39.90%) with causal partner, while 22 (10.58%) and 3(1.44%) of the sexually active respondents did their first sex with partners and others respectively. Assessment of the major reasons for beginning sex among the respondents show that falling in love (33.17%), sexual desire (41.35%), marriage (9.13%), peer pressure (2.40%), get drunk (8.17%) and others (5.77%) respectively, while the other 24(10.34%) respondents have never started sex so far.

4.3 Knowledge about HIV/AIDS and Other STDs

Knowledge about sexually transmitted diseases is vital to protect oneself from risky sexual behavior. In this study assessment has been done to find out the status and level of knowledge about HIV/AIDS and other STDs among the youths in the study area. Based on the finding in this study almost 100 percent of the respondents have information and knowledge about HIV/AIDS.

Regarding the HIV prevention methods more than 13% of the respondents have an understanding that abstinence from sex and 18.97% faithfulness to one sex partner prevents from being infected by HIV. Although most of the respondents also believe that using condom is one method of protecting the infection, it was found out that only about 59% of them use it consistently and 7.76% claim to prevent AIDS transmission by other methods.

Table 3. HIV/AIDS related preventive practice perceived by the respondents

knowledge about HIV/AIDS	frequency	Percent
HIV/Aids can be cured/prevented		
1.Yes	232	100
2.No	0	0
Preventive method		
1. Abstinence	32	13.79
2. being faithful to one partner	44	18.97
3. condom use	138	59.48
4. others	18	7.76

Due to the presence of a highly organized national and local HIV risk reduction campaign and the existence of more than two decades history of the epidemic, only 214(92.24 %) of the respondents reported that it is difficult to identify HIV carriers by looking at some one. Moreover 224(96.55%) of the respondents knew that the presence of STDs can increase the risk towards HIV infection, while about 55% believe that HIV can be transmitted in the first time of sexual contact. Furthermore almost 100% of them know that HIV/AIDS can be treated, and no one responded that the pandemic has no cure at all.

The participants of this study were asked if they have ever faced STDs. Accordingly, 8 respondents (3.45%) said that they were hit at least once with 138 (59.48%) participants believing that condom use is a practical protective option against HIV/AIDS and other STDs. Although use of condom is popular among the youths, 46 (19.83%) respondents believe that using condom is a sign of mistrusting partners while 47 (20.26%) respondents said that discussing about condom with young people could promote promiscuity (please see table 4.5). Moreover, 175(75.43%) respondents agreed that alcohol consumption and drug use can lead to HIV acquisition while 220 (94.83%) of the participants reported that multiple sexual contact leads to this problem.

Table 4. Awareness and beliefs towards risks of sexual activities and HIV/AIDS among adolescents

Variable		Male		Female		Total	
		Frequency	%	Frequency	%	Frequency	%
1 Person can get HIV when he/she has sex the first time	yes	79	58.96	48	48.98	127	54.74
	No	36	26.87	30	30.61	66	28.45
	I don't know	19	14.18	20	20.41	39	16.81
2 By looking carefully you can know if she/he has HIV	Yes	11	8.25	7	7.14	18	7.76
	No	123	91.79	91	92.86	214	92.24
	I don't know						
3 Using condom is a sign of not trusting your partner	Agree	27	20.15	19	19.39	46	19.83
	Disagree	93	69.40	71	72.45	164	70.69
	Not sure	14	10.45	8	8.16	22	9.48
4 Discussing condom or contraception with young people promotes promiscuity	Agree	85	63.43	70	71.43	155	66.81
	Disagree	34	25.37	13	13.27	47	20.26
	Not sure	15	11.19	15	15.31	30	12.93
5 Is AIDS curable	Yes	0	0	0	0	0	0
	No	134	100	98	100	232	100
	I don't know	0	0	0	0	0	0
6 Do you believe having multiple sexual contact leads to HIV acquisition	Yes	126	94.03	94	95.92	220	94.83
	No	8	5.97	4	4.08	12	5.17
7 Do you believe alcohol consumption and drug use can predispose to HIV acquisition	Yes	98	73.13	77	78.57	175	75.43
	No	24	17.91	21	21.43	45	19.40
	Don't know	12	8.96	--	--	12	5.17

4.4. Condom use

Using condom is one of the methods of preventing STDs including HIV/AIDS. Condom, which is ubiquitous and easy to use, is especially one of the three methods being used to safeguard sexually active youths from the risk of HIV infection. Use of condom, as a means of safer sex practice, has gained greater significance especially at a time when the spread of HIV/AIDS is increasing rapidly. In this context, it is imperative for men and women to be aware of the importance of condom use. This study has tried to assess the attitude of respondents towards condom use its effectiveness in preventing HIV transmission.

As can be seen from table 4.6 below, among the sexually active respondents 164(78.85%) had ever used condom with only 35(44.9%) respondents claiming to have used it consistently during the last 12 months. on the other hand 46 (54.8%) respondents reported that they have used condom only in their first sexual intercourse and 64(76.2%) of them have used condom in their last sexual intercourse. Out of those who used condom 59(70.2%) respondents used it while doing sex with friends, 15(17.9%) with casual partners and 4(4.8%) with commercial sex workers. When asked why they use condom during sexual intercourse, they replied to avoid HIV/AIDS (48.17%), prevent pregnancy (1.22%), and for both reasons (50.61%).

Table 5. Condom use among adolescents by sex

Variables	Male		Female		Total	
	frequency	percent	frequency	percent	frequency	Percent
Ever used condom						
1.yes	103	85.12	61	70.11	164	78.85
2.No	18	14.88	26	29.89	44	21.15
Condom use in the last 12 months						
1.alwys	55	53.40	24	39.34	79	48.17
2.most of the time	26	25.24	30	49.18	56	34.15
3.sometimes	22	21.36	7	11.48	29	17.68
Condom use during the first sexual intercourse						
1.yes	7	6.80	1	1.64	8	4.88
2.No	96	93.20	60	98.36	156	98.12
Condom use during the last sexual intercourse						
1.yes	59	57.28	28	45.90	87	53.05
2.No	44	42.72	33	54.10	77	46.95
Reason for condom use						
1.avoid HIV/AIDS & STD	78	75.73	1	1.64	79	48.17
2.avoid pregnancy	-	-	2	3.28	2	1.22
3.Both	25	24.27	58	95.08	83	50.61

4.5 Risk perception of respondents

In the assessment done to know the risk perception of respondents, this study has found out that 48(21.64%) of the male participants and 26(15.31%) of female participants have perceived themselves to be at high risk of HIV infection, while 34(25.37%) of the males and 40(40.82%) of females have perceived a medium chance of getting HIV. Similarly there were respondents who perceived themselves to be at a less or no risk of getting infected with this epidemic.

This study shows that 23(17.16%) of the male and 17(17.35%) of the female respondents believe to have low chance of acquiring the virus while 48(35.82%) of the male and 15(15.31%) of the female respondents replied that they have no chance of getting HIV. In general it was found out that the male respondents were considering themselves to be at a higher risk of HIV infection than the female respondents (Table: 6)

Table 6. Percent distributions of respondents by HIV risk perception

variable	male		female		Total	
	frequency	percent	frequency	percent	frequency	Percent
Chance of acquiring HIV:						
None	48	35.82	15	15.31	63	27.16
Low	23	17.16	17	17.35	40	17.24
Medium	34	25.37	40	40.82	74	31.90
High	29	21.64	26	26.53	55	23.71

4.6 Demographic and socio-economic characteristics and perceived level of HIV risk among the youths

As partly this study is aimed at highlighting the relationship of variables and interpreting the implication, Chi square method has been used to examine the association between demographic and socio-economic characteristics and the risk perception of HIV among the youths in the study area.

The table below shows that there are various levels of association between demographic and socio-economic characteristics and risk perception of HIV among the youths. Accordingly, age level was not found to have a significant association with risk perception of HIV among the youths ($\chi^2= 7.0930$, $P= 0.29$). A better association was observed between the job variable and risk perception. In this study job was found to be significantly associated with risk perception of HIV among the youths ($\chi^2=4.7151$, $P= 0.03$). This means that youths who had job had high risk perception as compared to the jobless ones.

Alcohol use was the other variable which has shown a significant association with risk perception of HIV among the youths ($\chi^2= 16.8405$, $P=0.001$). The alcoholic youths were found to have high risk perception than those who did not use. The level of monthly income was significantly associated with risk perception of HIV ($\chi^2=12.7690$, $P=0.026$) where the statistics show that youths with high monthly income had a relatively better risk perception.

The same is true with the sex variable which was significantly associated with risk perception of HIV among the youths ($\chi^2= 9.4788$, $P= 0.002$). Hence the level of risk perception among males was found to be

higher than the female respondents. A significant association has also been observed between youth level of education and risk perception of HIV ($\chi^2 = 11.8883$, $P=0.003$), where the level of risk perception was better among those with higher educational level than those who are otherwise.

Table 7. Demographic and socio-economic characteristics and perceived level of HIV risk among the youths

Variables		Perceived chance of getting HIV		χ^2	P-value
		Lower chance (low risk +no risk)	Higher chance (high +medium risk)		
Age	15-19	32	61	7.0930	0.29
	20-24	50	43		
	25-29	21	25		
Job	Yes	65	63	4.7151*	0.03
	No	38	66		
Drinking alcohol	No	57	40	16.8405**	0.001
	Once &twice	4	3		
	Sometimes	28	66		
	daily	14	20		
Monthly income	Below 300	27	35	12.7690*	0.026
	301-500	31	54		
	501-1000	22	26		
	1001-1500	18	6		
	1501-2000	3	7		
	Above 2000	2	1		
Sex	Male	71	63	9.4788**	0.002
	Female	32	66		
Level of education	1-8	17	28	11.8883**	0.003
	9-12	41	72		
	Above diploma	45	29		

P<0.05 p<0.01

4.7 Behavioral factors and Risk Perception among the Youth

The chi square method used to examine the association between behavioral factors and risk perception of HIV among the youths has shown various results. As seen in the table below, the behavioral factors have shown some level of association with risk perception of HIV. The results indicate that sexual practice was significantly associated to risk perception ($t=10.1561$, $P= 0.001$), where the risk perception of people who had ever practiced sex was greater than those who didn't.

Table 8. Association of behavioral factors and risk perception of HIV infection

Variables		Perceived chance of getting HIV		χ^2	p-value
		Low risk	High risk		
Ever had sex	Yes	85	123	10.1561**	0.001
	No	18	6		
No of sexual partners ever had	1	43	31	17.2151**	0.002
	2	14	24		
	3	19	34		
	4	3	15		
	Above 4	6	19		
Age at first sex	15-19	60	69	7.5240*	0.023
	20-24	23	40		
	25-29	2	14		
Condom use	Yes	68	96	0.1147	0.735
	No	17	27		
Ever had STDs	Yes	3	5	0.1596	0.689
	No	100	124		
Knowledge of HIV status	Tested	51	16	16.0624**	0.000
	Not tested	78	87		

P* < 0.05 p< 0.01**

Similarly, the number of sexual partners someone ever had was found to be significantly associated with risk perception of HIV among the youths ($\chi^2=17.2151$, $P= 0.002$). This means that the risk perception of people who had multiple sexual partners was found to be lower as compared to those who didn't. Likewise, the age of

to the participants, the main reason for early sex are early marriage, peer pressure, illegal video house (watching pornography film), rape and abduction, alcohol, khat and economic problems. The participants also stated that due to early sex females are exposed to unwanted pregnancy, abortion and other complications like fistula. Generally, the participants agreed that early sex predisposes to HIV/AIDS and finally death. Majority of the participants stated that sexual intercourse should be started after marriage and if possible after 18 years and after getting their own income. One of the factors facilitating the spread of HIV is having multiple sexual partners. To be able to assess this, the participants were asked whether people had many sexual partners and reasons for this attitude. Different reasons were given for many sexual partners in the era of AIDS. Peer pressures, experimentation by the youth and intensive sexual urge were the reasons for many sexual partners. A male discussant also said "*absence of recreation place for the youth is the main reason for youth sexuality as sex is one method of recreation for the youth*". Another issue discussed by the participants as a cause of youth sexuality is unemployment and excess free time. Majority of participants agreed on the fact that multiple sexual partners predispose to HIV/AIDS except a young female discussant, who strongly argued that multiple sexual partners does not predispose to HIV, if safe sex is practiced.

Risk perception of HIV/AIDS

The discussants in the focus groups were asked about what they perceive to be the risk of contracting HIV/AIDS among people who had multiple sexual partners and how do the youth perceive risk of HIV/AIDS. Almost all participants in the groups perceived that people with multiple sexual partners are at high risk of contracting HIV/AIDS. According to the participants, all the youth seemed to fear HIV/AIDS, but still chose to participate in unprotected sex. A 17 year old male said "*Some of the youth perceive there is no infection of HIV after 10 pm.*" Most of the participants stated that the reasons why youth do not perceive themselves at risk of HIV are due to over indulgence in alcohol and khat. Alcohol and khat make the youth not to think of risk perception, not to be open in the sexual matter, feel hopelessness and the nature of the disease not causing sudden death. According to participants, even though youth have high knowledge on HIV/AIDS they don't bring behavior change because of peer pressure, poverty, absence of recreation area, khat and drug abuse, no vision, unemployment and generally they don't give attention due to their ages.

Condom use

Majority of the group didn't consider the use of condom as acceptable means of prevention because of perceived reduction in sexual pleasure. According to the participants, some of the reasons for non-use of condom are perceived reduction in sexual pleasure, shame to buy, create tension, feeling that condom may contain the virus and over indulgence in alcohol. Explaining the situation a young boy said, "*Using condom during sexual intercourse is as if having banana with its cover.*" Another male participant also said "*Using condom is walking in the rain with umbrella, because the rain may wet some part of the body.*" Another issue raised in the discussion was the difference of price for different types of condom that influence the youth understanding on the protective capacity of condom. Regarding where condoms were obtained from, all groups reported the sources of condom distribution were from shop, health institution, hotels and pharmacy. Majority of participants agreed that condom should be distributed in recreation area, meeting places, schools, public offices, and kebele associations and bus stations in addition to previous distribution.

Willingness to undergo VCT

The study obtained information on various aspects of HIV testing including perception of the youth about HIV testing and knowledge of VCT. Majority of the discussants have knowledge on the importance of VCT. Almost all groups indicate that, even though youth have knowledge on VCT, they are afraid to be tested. According to the participants, voluntary counseling and testing is necessary to plan for the future, to know status, for marriage, for DV and to prevent transmission of HIV.

Generally almost all participants of in the focused-group discussions and in-depth interviews agreed that the pattern of HIV/AIDS epidemic is getting worse over time, and is seriously affecting the young which is the productive population group in the town.

All the participants in the focus-group discussions and in-depth interviews agreed that the key risk factor putting youths at risk of HIV/AIDS in the central part of the Town is the increasing number of places for consuming alcohol and *chat*. The participants in these methods have also witnessed that there is a trend of taking too much alcohol among youths aiming at avoiding the problem of sleep disturbance and anxiety that usually follow chewing of *chat*. And it is obvious that taking of too much alcohol, in turn, results in practicing unprotected sex.

4.2 Discussions

This study has examined the association of variables using the chi-square test which has shown variables like sexual practice, multiple sexual partners, age at first sex, and knowledge of HIV status were found to have significant association with risk perception of HIV of among the youths. A similar study conducted in Uganda has revealed that the risk perception on HIV/AIDS is associated with condom use, educational attainment,

marital status, and number of sexual partners (Ahimbisibwe, Odwee and Ayiga, 2003).

In this study some variables like job and alcoholism have shown significantly association with risk perception where people who have jobs and do not consume alcohol were found to have high risk perception than those who do otherwise. This finding similarity and consistency with previous studies conducted on this area. For instance, various studies conducted on adolescent reproductive health indicate that use of alcohol and drugs was a risk factor for having had sex, and having multiple sexual partners over life time (Adolescent Reproductive Health Task Force, August 2003). A study conducted on this area in 2002 backs this finding. Among youths who were reported to have had risky sex during 12 months time, 44% of them had used alcohol and chat regularly (HIV/AIDS Behavioral Surveillance Survey (BSS), Round one, Ethiopia, 2002). Similarly, Fekadu (2001) reported that 'alcohol' and 'chat' use have strong links with the incidence of 'rape' as which is again a factor contributing to early sex initiation among female adolescents in Addis Ababa.

Various literatures written on alcohol use related behavior have considered the perception of being at risk of infection to be one of the necessary conditions for behavioral change (Lavra, 2002). This study has come up with a finding that shows alcohol use is a predictor that explained variation in the likelihood of risk perception of HIV infection. As it is observed from the logistic regression model, respondents' who consume more alcohol were 1.701 times likely to be at a higher risk of being infected by HIV than those who consume less. Another study has indicated that alcohol consumption has shown an effect on risky sexual practices among school anti-AIDS club member and non member youths in Jima and Agaro town of Ethiopia (Yazachew, 2003). Those who are engaged in drug use (including alcohol) may become more vulnerable to sex related infections including HIV/AIDS (UNAIDS 1997). Similarly, Langer, et al., (2001) has identified alcohol intake to be a significant predictor of risky sexual practices using regression analysis.

In addition to this, monthly income was found to be significantly associated with risk perception of HIV among the youths, where respondents with low monthly income had low risk perception of HIV than those with high income youths. Likewise, gender status was found to be significantly associated with risk perception of HIV youths, and that the level of risk perception of males was much higher than female respondents. This finding is somehow similar to previous findings which revealed that the level of HIV risk perception of males was higher than females. According to various studies, the main reason behind this difference are wide spread poverty among women, gender inequality, and social and political inertia. Especially, in developing countries these factors have widened the gap of AIDS prevalence between both sexes (UNFPA, 2008).

Level of education was also found to have a significant association with youths' risk perception of HIV/AIDS. The assessment in this regard shows that people with low level of education had low risk perception of HIV than those with a better educational level, which goes in harmony with previous studies conducted in this area. For example, as revealed in EDHS (2005), knowledge about HIV/AIDS among the youths rises with the level of education, but the level of risk perception declines. In this study educational level was found to be an independent and significant predictor of risk perception towards HIV/AIDS among the youths. As it has been observed from the logistic regression model, respondents' with higher educational status were 1.914 times at a higher risk than those who had lower educational level. Similarly, as revealed in EDHS (2005) the spread of HIV/AIDS rises with the level of education; and youths in the lowest educational level were found to be at a lower risk. Marital status was also found to be a predictor in explaining variation in the likelihood of risk perception of HIV infection in this study. As can be observed from the logistic regression model, married respondents had 0.857 times higher risk perception than the unmarried people who were likely at risk of being infected by HIV. A study conducted by Langer, et al., (2001) in Uganda that revealed marital status to be a significant predictor of risky sexual perception backs this finding. In this study sex experience was also found to be significantly associated with risk perception of HIV among people. The assessment in this regard shows that people who ever practiced sex had low risk perception than those who didn't. A study by Macintyre (2004) has come up with the same finding where he found that Sexually inexperienced youth may perceive such a risk if, for example, they doubt their potential to maintain consistent condom use or to identify partners who are at low risk of infection once they become sexually active. The other variable that has shown a sort of association with risk perception is the number of sexual partners. In this study it has been found that the risk perception towards HIV/AIDS was higher among those who had limited sexual partners than those with multiple partners. And several studies show that this trend can lead to a high AIDS prevalence. For instance, Encarta (2006) stated that the potential negative consequence of practicing unsafe sex with multiple partners is high risk of contracting HIV/AIDS.

According to the finding in this study, the age at first sexual practice of youths was significantly associated with risk perception of HIV, where the youths who had experienced sex at their early age had less level of risk perception than those who practiced sex at their later ages. There are also other studies conducted on this area that support this finding. A study conducted on high-risk sexual behavior among youths in Tanzania has revealed that the largest group, 55% of girls and 45% of boys, had their first sexual intercourse experience between the ages of 14 and 17 (Ikamba and Ovedraogo, 2003). Another study also shows that 30% of respondents were

sexually experienced at an average age of 12.7 (Family Planning perspectives 2000; 32(1)). These studies indicate that as people start sexual intercourse at their earlier ages it is highly likely to have less risk perception towards AIDS. And this is the reason why in many countries 60 percent of all new HIV infections are among people of 15-24 years old. That means the highest rates of STDs are usually found in the age range of 20-24 years, followed by 15-19 years (UNAIDS, 1997). Generally as stated by Langer *et al*, 2001, the age level of first sex practice determines individual risk perception. Similarly, in this study age was found to be a predictor that explains variation in the likelihood of risk perception of HIV infection. As it is observed from the logistic regression model, respondents' age is 0.9280 times likely to perceive higher risk of being infected by HIV. Langer, et al., (2001) has come up with a finding which identified age as a significant predictor of risky sexual practices.

Knowledge of HIV status was also found to be significantly associated with risk perception of HIV among youths. This means that the risk perception of persons who know of their HIV status was higher than those who did not undergo VCT. A study conducted on high-risk sexual behavior among youth who had the knowledge of HIV status in Tanzania revealed that 11.7% of the participants felt they were at high risk of getting HIV/AIDS and STDS, 25% felt that they had a very low risk, while 53.1% felt that they were not at risk at all (Ikamba and Ovedraogo, 2003). Similarly, knowledge of HIV was found to be a predictor variable that explained variation in the likelihood of risk perception of HIV infection. As it is observed from the logistic regression model, respondents' Knowledge of HIV is 1.591 times likely to predict higher risk than lower risk of being infected by HIV, and a research conducted by Ellen et al (2006) among young adults in South Africa have revealed the same finding. According to her, young men who had been tested for HIV were significantly more likely to have used condom during their recent sexual intercourse.

On the other hand, in this study, condom use was not significantly associated with risk perception of HIV among youths. Likewise, experience of STDs was not significantly associated with risk perception of HIV among youths. Contrary to this finding, different studies revealed that condom use was significantly associated with risk perception of HIV. Therefore, it seems that this new outcome would open a room for further research. Generally speaking, the logistic regression in this study reveals that socio demographic and behavioral factors like age, educational level, marital status, alcohol use and Knowledge of AIDS and of HIV status were significant and independent predictors of risk perception. However variables like sex, employment status, monthly income and condom use were not found to be significant and independent predictors of risk perception.

5. Conclusions and Implications

5.1 Conclusions

As it has been mentioned somewhere in the introductory part of this thesis, the main purpose of the study is to investigate the risk perception of HIV among youths in Bahir Dar town. And to address this research problem the following research questions were entertained:

1. Do socio-economic, demographic and behavioral variables predict the level of risk Perception of HIV infection among young adults?
2. How do youths perceive the risks associated with sexual activity?
3. How aware are youths to prevent such risks?

Participants of the study were a sample of 232 youths living in Bahir Dar town of Ethiopia. In addition, 17 concerned officials drawn from government offices, non-governmental organizations, civic societies, and community leaders were participated in this study by way of focused group discussion and key informant interviews. A self administered questionnaire that was designed in a way that includes socio-economic, demographic and behavioral characteristics served to collect data from the youths. On top of this, observation by the researcher has had a vital contribution in substantiating the data collected using the various methods.

While descriptive statistics has been used to characterize socio-economic, demographic and behavioral variables and the level of risk perception of HIV/AIDS, chi square was used to examine the association of socio-economic, demographic and behavioral variables with HIV risk perception. These variables were further examined to identify their independent prediction using the logistic regression model. Finally, the data gathered using interview and focus group discussion was qualitatively analyzed.

Analysis of the quantitative and qualitative data collected for this study has led to generate the following finding.

1. Socio-economic and demographic variables such as Job, alcohol use, monthly income, and gender and educational status were significantly associated with risk perception of HIV among the youths. And behavioral variables like sex experience, age at first sex, no of sexual partners and knowledge of HIV status were significantly associated with risk perception of HIV among the youths.
2. Age, education status, marital status, alcohol use and knowledge of HIV status were significantly and independently predicted HIV risk perception.

6.2 IMPLICATIONS

1. This study provides important information about socio-economic, demographic and behavioral predictors of risk perception of HIV among youths. It also provides important information for further research, as well as implications for preventive strategies and intervention on the sexual behavior of youths to reduce the prevalence of HIV/ AIDS among youths. Being the young are at risk of facing socio-economic, demographic and behavioral related and other problems, it is important to understand the mechanism behind the relationships among these variables. This would definitely help to design appropriate prevention strategies tailored towards this group.
2. This study is primarily conducted based on cross-sectional data collection and analysis; hence it is not possible to make any causal connections between sexual behavior and risk perceptions of HIV or know the direction in their relationship. Furthermore, the measure of risky sexual behavior used in this study is based only on an individual's behavior in the past preceding the study yet actual risk of exposure to HIV is a combination of several factors such as the type and number of lifetime sexual partners; use of condoms by the individual and/or the partner currently and in the past; and the partner's past and current sexual behavior.

REFERENCE

- Abate, S. (1999). Determinants of high- risk sexual behavior for HIV/AIDS among out of- school youth in Addis Ababa, Ethiopia,(thesis). Department of Community Health, Faculty of Medicine, Addis Ababa University.
- Adamu, R.Samuel, M. and Ingidushet, S. (2003). Patterns and correlates of sexual initiation, sexual risk behaviors, and condom use among secondary school students in Ethiopia. *Journal of Health Development*, 41(2): 163-177
- Adih, W. and Alexander, C. (1999). Determinants of condom use to prevent HIV infection among youth in Ghana, *Journal of Adolescent Health* ,24(1).
- Ahimbisibwe, E.Odwee, J.and Ayiga, N. (2003). Risk perception and condom use in Uganda. *African population studies* ,vol. 18 (1): 68-80
- AIDS in Ethiopia technical report, (2007)
- Caldwell, J.C, Orubuloye, I.O, and Caldwell, P. (1999). Obstacles to Behavioral Change to lessen the risk of HIV infection in the African AIDS epidemic
- Campbell, C. (1997). Migrancy, masculine identities and AIDS: the psycho-social context of HIV transmission on the south African gold mines; *Social Science and Medicine* 45(2)
- Catherine M. and Catherine C. (2001). Condom use coming adolescents and young people in a southern African township. *Social Science and Medicine* 52(11): 1613-1627
- Central Statistical Authority. (2006). Ethiopian demographic and health survey (EDHS) report. Addis Ababa: Central Statistical Authority: 179-212.
- Dawud, A. (2002). Perception of the risks of sexual activities among out-of-school Adolescents in south Gondar administrative zone, Amhara Region,(thesis) Department of Community Health, Faculty of Medicine, Addis Ababa University.
- Doll LS, Beeker C. (1996). Male bisexual behavior and HIV risk in the United States: Synthesis of research with implications for behavioral interventions. *AIDS Educ Prev*; 8:205-25.
- Encarta. (2006). Microsoft Encarta Encyclopedia standard
- EPHA. (2003). Adolescent reproductive health global and national initiatives and lessons learned Adolescent Reproductive Health Task Force.
- Eshetu F Zakus D and Kebede D. (1997). The attitudes of students, parents and teachers 53 towards the promotion and provision of condoms for adolescents in Addis Ababa.Ethiopia. *Journal of Health Development*, 11(1): 7-16
- Ethiopia: HIV/AIDS causes lower population growth in Amhara region, 2008 Ethiopia. Ministry Culture and Youth Affairs. Ethio pian youth basic challenges and prospects. Addis Ababa: Children and Youth Affairs Organization, Ministry of Culture and Youth Affairs, Government of Ethiopia, 1995:40-6. (Monograph).
- Fantahun Mulu. and Chala F. (1996). Sexual behaviour, and knowledge and attitude towards HIV/AIDS among out-of-school youth in Bahir Dar Town, Northwest Ethiopia. *Medical Journal*, 34(4): 233-242
- Fekadu Zewedu. (2001).Casual sex debuts among female adolescents in Addis Ababa, Ethiopia. *Journal of Health Development* . , 15(2): 109-116
- Fisher WA, Byrne D, White LA. (1988). Erotophobia-erotophilia as a dimension of personality. *Journal of Sex Research*, 25:123-51.
- Fitaw, Yala. and Worku Ada.(2002). High-risk sexual behavior and pattern of condom utilization of the Gondar College of Medical Science(GCMS) students, North-west Ethiopia. *Ethiop. Journal of Health Development*. 16(3): 335-339.

- Govindasamy P, Kidanu A, Bantayerga H. (2002). Youth reproductive health in Ethiopia. Calverton, MD: ORC Macro, 99 p.
- Has HIV/AIDS Epidemic Changed Sexual Behaviour of High Risk Groups in Uganda. A paper to be presented to the conference of the International Union for scientific study of population, to be held in Salvador, Brazil 18-24 August 2001
- HIV/AIDS Behavioural Surveillance Survey (BSS). Round one. Ethiopia, 2002
- HIV/AIDS and work: global estimates, impact and response (ILO) ., 2005.
- Ikamba L. and Ovedraogo B. (2003). High-risk sexual behaviour: knowledge, attitudes and practice among youth at Kichangan ward, Tanga, Tanzania. Action research reports. Retrieved May 10, 2010 from <http://11-www.fhs.usyd.edu.au/arow/arer/018.htm>,
- Ismail S, Bitsuamlak H, and Alemu, D. (1997). High-risk sexual behaviors for STD/HIV pregnancies and contraception among high school students in a rural town Northwestern Ethiopia. *Journal of Health Development*, (1): 29-36.
- Jemmott, L. S., & Jemmott, J. B. (1992). Family structure, parental strictness, and Sexual Behavior among inner-city Black male adolescents. *Journal of Adolescent Research*, 7(2), 192-207.
- Kapiga S. and Lugalla J. (2003). Male condom use in Tanzania: Results from a National survey. *East African medical journal*, 18 (4): 181-190.
- Karra A. and Haile M. (1999). Sexual behavior and level of awareness on reproductive health among youths: evidence from Harar, Eastern Ethiopia. *Journal of Health Development*. 13(2): 107-113
- Kelly Gary F. (1998). Sexuality today the human perspective 6th edition. United States.
- Kermyt G Anderson, Ann M. Beutel and Brendan Maughan-Brown. (2007). HIV Risk Perceptions and First Sexual Intercourse Among Youth in Cape Town South Africa : International Family Planning Perspectives vol.33, no.3.
- Langer, L. M., Warheit, G. J., and McDonald, L. P. (2001). Correlates and Predictors of Risky Sexual Practices among a multi-racial/ethnic sample of university students. *Sexual Behavior and Personality*, 29(2), 133-144.
- Lavra B. (2002) Determinant of individual AIDS risk perception: knowledge, behavioral control and social influence. MPIDR working paper. Retrieved June 02, 2010 <http://www.Demogr.mpg.de/papers/working/wp-2002-029> .
- Lemma. E. (2000). Predictors of HIV/AIDS related sexual behaviour of high-school adolescents based on the classical health behaviour models, Jimma Town, Southwest Ethiopia, (thesis) Department of Community Health, Faculty of Medicine, Addis Ababa University.
- Maharaj, P. (2004). Perception of Risk of HIV Infection in Marital and Cohabiting partnerships; *African journal of AIDS research* 3(2)
- Mahesh, P. (2002). Sexual Risk Behavior and Risk Perception of Unwanted Pregnancy and Sexually Transmitted Infection among young factory workers in Nepal
- Macintyre, K. (2004). Understanding Perception of HIV risk among Adolescents in Kwazulu-Natal, AIDS and behavior, 8(3)
- Macintyre, K. Brown, L. and Soster, S. (2001). "It is not what you know, but who you know", Examining the Relationship between Behavior Change and AIDS Mortality in Africa, *AIDS Education Prevention* 13(2).
- Masulanya, E. Moji, K I Hariguchi, et.al. (1999). Knowledge, risk perception of AIDS and reported AIDS and reported sexual behavior among students in secondary schools and colleges in Tanzania. *Health education research theory and practice*. 14(2): 185 –196.
- Melanie, G. and Eleanor, M.T. (2004). School-based HIV prevention programmes for African youth. *Social science and medicine*. 58(7): 1337-135.
- Miller, K. S., Forehand, R., and Kotchick, B. A. (2000). Adolescent Sexual Behavior in two ethnic minority groups: A multi-system perspective, *Adolescence*, 35(138):313-333
- Ministry of Health. (2007). AIDS in Ethiopia, 4th edition. Disease Prevention and Control Department,
- Ndola, P. Leo, M. Elizio, M. Farnaz, V., and Mrk, S. (2006). Relationships between HIV risk perception and condom use: evidence from a population based survey in Mozambique; *International Family Planning Perspectives*, 32(4).
- Petros, B. Belayneh, S. and Mekonnen, Y. (1997). AIDS and college students in Addis Ababa: A study of knowledge, attitude and behavior. *Journal of Health Development*. 1997; 11(2): 115-123
- Position statement on condom and HIV prevention. (2004). Retrieved January 23, 2010 from <http://www.Who.int/3by5/en/condom.04/08/04>
- Policy on HIV/AIDS of the Federal Democratic Republic of Ethiopia. August 1998.
- Prata, Ndola et al. (2006). Relationship between HIV risk perception and condom use: evidence from a population-based survey in Mozambique ; *International Family Planning Perspective*, 32(4).
- Robert, J. Ali, M. Lisa, A. Katherine, C. Musonda, L. and Gwendolyn, T. (2000). Reproductive Health Risk and

- Protective Factors among Youth in Lusaka, Zambia. Social factors play major role in making young people sexual risk takers. *Family Planning perspectives* 2000; 32(1).
- Stoner, B. WLH Whittington Aral S, et.al. (2003). Avoiding risky sex, partner Perception of partners' risks vs partners' self reported risks. *Sex transm infect*, 79:197-201.
- Tadesse, E. Gudunfa, A. and Mengistu ,G. (1996).A survey of adolescent reproductive health in the city of Addis Ababa,Ethiopia. *Journal of. Health Development*. 10(1): 35-39
- Taffa, N. (1998).Sexuality of out-of-school youth, and their knowledge and Attitude about STDs and HIV/AIDS in Southern Ethiopia *Journal of. Health Development*. . 12(1): 17-22.
- Taffa, N. Johanne, S. Carolo ,H. and Gunnar, B. (2002).HIV prevalence and socio cultural contexts of sexuality among young youth in Addis Ababa, Ethiopia. *Journal of. Health Development*. 16(2): 139-145
- UNAIDS. (2001). Children and Young people in a world of AIDS.
- UNAIDS. (2001). The global strategy framework on HIV/AIDS.
- UNAIDS. (2004). Report on the global AIDS epidemic, fourth report.
- UNAIDS. (2006). Report on the global HIV/AIDS epidemic.
- UNAIDS. (2007). Report on the global HIV/AIDS epidemic.
- UNAIDS. (2008). Report on the global HIV/AIDS epidemic.
- UNAIDS. (2008). A review of household and community responses to the HIV/AIDS.
- UNFPA. (2006). Preventing HIV infections in young people. Retrieved December 5, 2010 from <http://www.unfpa.org/aids/index.htm>
- UNFPA. (2008). Preventing HIV infections in young people.
- United Nations Children Fund. (2002). Young people and HIV/AIDS opportunity in crisis, 55 Joint United Nations programme on HIV/AIDS and World Health Organization.
- Verga, C. (2001). coping with HIV/AIDS in Durban's commercial sex industry. *AIDS care* 13(3)
- WHO. (2004). The World Health Report. Changing history.
- World Health Organization. (1998). Youth and HIV/AIDS: forces for change (report). Geneva: Joint United Nations Programme on HIV/AIDS, 22-40
- Yazachew Mekuria. (2003). Assessment of sexual risk behaviour for HIV infection between out of school anti-AIDS club members and non-club member youth in Jima and Agaro, (thesis) Department of Community Health, Faculty of Medicine, Addis Ababa University.