

## Efficacy of Insecticides on Rice Crop Against Rice Leaf Folder, *Cnaphalocrocis Medinalis* Guen. (Pyralidae: Lepidoptera)

SAJJAD ALI KHUHRO\* ABDUL WAHEED SOLANGI, ABDUL GHANI LANJAR

Department of Entomology, Sindh Agriculture University, Tandojam.

\*Corresponding author: SAJJAD ALI KHUHRO Email: [sajjadento.sau@gmail.com](mailto:sajjadento.sau@gmail.com)

### ABSTRACT

A field study was carried out during 2010 at experimental area Sindh Agriculture University Tandojam in RCBD to examine the reduction percentage in the population of *C. medinalis* on rice crop with three insecticides viz; Deltaphos, Tracer, Thioluxan and impact of these insecticides was compared with control plot (untreated). The data showed the maximum reduction (76.19%) in the population of *C. medinalis* was recorded in the plots treated with Deltaphos followed by Tracer (65.93%) and Thioluxan (46.92%) after 1<sup>st</sup> spray on rice crop. In the second spray maximum reduction (71.08%) was also recorded in the plots treated with Deltaphos followed by Tracer (57.40%) and Thioluxan (30.15%). Similarly, after 3<sup>rd</sup> spray the maximum reduction (78.99%) was recorded in the plots treated with Deltaphos followed by Thioluxan (55.88%) and Tracer (54.34%) on rice crop. While, in fourth spray, the maximum reduction (71.92%) was recorded in plots treated with Deltaphos followed by Tracer (39.36%) and Thioluxan (14.36%) on rice crop. The results further indicated that the maximum yield (4783.00kg/ha<sup>-1</sup>) was recorded in plots treated with Deltaphos followed by Tracer (4486.46kg/ha<sup>-1</sup>) and Thioluxan (4033.66kg/ha<sup>-1</sup>) as compared control (3066.50kg/ha<sup>-1</sup>). Highly significantly ( $P < 0.01$ ) difference in the efficacy of insecticides was recorded during all sprays carried out in cropping season of rice crop.

**Key word:** *Oryza sativa*, *Cnaphalocrocis medinalis*. Management and Insecticides.

### INTRODUCTION

Rice (*Oryza sativa* L.) is cultivated and consumed by 2500 million people in almost 112 countries. About 90% of the world's rice is grown and consumed in Asia (Ahmed et al., 2005). Rice is an important cereal crop which plays multifarious role in the economy of Pakistan (Sabir et al., 2006). There are large numbers of insect pests, which damage rice crop right from nursery sowing to the harvest causing considerably high yield losses. About 128, different species of insect pests have been reported to attack rice crop in Pakistan (Ahmed, 2005). Insect pests damage rice crop at different stages of its growth. Among that leaf feeding insect pests are of major importance because of their ability to defoliate or to remove the chlorophyll content of the leaves leading to considerable reduction in yield. Rice leaf folder, *Cnaphalocrocis medinalis* (Guen. Pyralidae Lepidoptera) was considered as pests of minor importance have increased in abundance in late 1980's and have become major pests in many parts world (Ahmed et al., 2010). Paddy leaf folder is one of the most important insect pests (Gunathilangaraj et al., 1986). Out of the eight species of leaf folder, the most widespread and important one is *Cnaphalocrocis medinalis* (Guenee) (Bhatti et al., 1995). *C. medinalis* (LF) has been reported to attain the major pest status in some important paddy growing areas (Maragesan et al., 1987). Second instars Leaf folder larvae glues the growing paddy leaves longitudinally for accommodation and feeds on green foliage voraciously which results in papery dry leaves (Khan et al., 1989). Loss incurred to the growing paddy crop is insurmountable (Ahmed et al., 2010). Feeding often results in stunting, curling or yellowing of plant green foliage (Alvi et al., 2003). The extent of loss may extend up to 63 to 80 percent depending on agro-ecological situations as reported by (Rajendran et al., 1986). The control of rice insect pests has often relied on extensive use of insecticides, which disrupt the beneficial insects and other insect fauna and also cause environmental contamination (Heong, 2005). The heavy use of insecticides and high fertilizer rates seem to favour leaf-folder population outbreaks (Gottfried and Fallil 1986). For the control of this pest in Asia, more than 25 per cent of applied pesticides are aimed to this pest in a year (Heong et al., 1997). In Sri Lanka, LF affects roughly 20 per cent of the total paddy field (Nugaliyadde et al., 1997). Application of Lorsban, Sumithion, Methyl Parathion, Denital and Thiodan gave more than 90% mortality of the insect larvae and were statistically at par in controlling the rice leaf-folder. Ramasubbaiah et al. (1980) tested fenitrothion, phosphamidon, fenitrothion, endosulphon dimethoate, quinalphos, diazinon and carbaryl against *C. medinalis* and reported that all insecticides gave effective control. Saroja and Raju (1982) studied the effect of some foliar insecticides against leaf folder and found that cypermethrin and fanvalerate provided effect control and increased the yield of rice significantly. Hence, this study was taken up and the results of which will be useful for the scientists to work out for an Eco-friendly Integrated Pest Management, thus it will support the ultimate beneficiary to the farming community (Ahmed et al., 2010).

## MATERIAL AND METHOD

Experiment was conducted at Sindh Agriculture University Tandojam during 2010. Rice crop (IRRI-6) was grown to determine the population dynamics of *C. medinalis*. The experiment was laid out in a Randomized Complete Block Design with four treatments and three replications, in a field measuring about 1245m<sup>2</sup> (12 ghunta) with sub plot size 207m<sup>2</sup> (2 ghunta). The crop was transplanted on 1<sup>st</sup> week of July in 2010. All the agronomic practices were carried out accordingly. The population was recorded by counting the number of insect larvae from randomly selected 10 plants calculating average number of larvae per hills. Three insecticides were evaluated against rice leaf folder on rice crop.

The detail of treatments is given as under;

Trade name and formulation	Common name	Dose (acre)	Dose ml/sub plot
Deltaphos 10.5 EC	Trizophos	600 ml	30 ml
Thioluxan 10.5 EC	Endosulfan	1000 ml	50 ml
Tracer 480SL	Spinosad	100 ml	5 ml
Control	Without pesticides		

The insecticides were sprayed with the help of knapsack sprayer machine and sprays was done in the early morning. Four sprays was done during the morning hours, the 1<sup>st</sup> spray was done on 026-07-2010., while second, third and fourth sprays was applied at 15 days interval, respectively. Observations at each spray were taken one day before spray (Pre treatment observation) and three observations after spray (Post treatment observations) at the interval of 1 day, 3 day and after one week. To see the efficacy of pesticides reduction percentage was calculated as per the standard formula of Hinderson and Titten (1955).

$$\text{Percent mortality} = 1 - \frac{T_a}{T_b} \times \frac{C_a}{C_b} \times 100$$

Where

Tb = Number of pest in treated plots before treatment.

Ta = Number of pest in treated plots after treatment.

Cb = Pest population in the control plots before treatment.

Ca = Pest population in the control plots after treatment.

Yield of the crop from each treatment was ascertained to compare the input of each treatment.

## RESULTS

The experiment was carried out on the effect of some chemical insecticides on rice leaf folder, *Cnaphalocrocis medinalis* Guen. (Pyralidae : Lepidoptera) at Sindh Agriculture University Tandojam under field conditions during 2010. The results of present studies are given below:


The result on reduction in the population of *C. medinalis* was recorded after one, three and seven days after 1<sup>st</sup> spray of insecticides. Table1 revealed that the different pesticides reduce *C. medinalis* population progressively. All insecticides showed maximum reduction %age after 7<sup>th</sup> day of 1<sup>st</sup> sprays except Thioluxan, which reduced maximum population on 3<sup>rd</sup> day. The maximum reduction percentage was recorded in the plots treated with Deltophos (76.19%), followed by the plots treated with Tracer (65.93%). The maximum reduction (46.92%) brought by Thioluxan 3<sup>rd</sup> day of spray after that it reduced its efficacy and rice leaf folder enhanced it activates on the crop.

**Table 1 Mean population and reduction %age of different pesticides against *C. medinalis* on rice crop after 1<sup>st</sup> spray by the application of insecticides.**

Treatments	Pre Treatment population	Reduction percentage in population		
		1 DAS	3 DAS	7 DAS
Deltaphos	3.70	64.85%	70.24%	76.19%
Tracer	3.17	51.66%	62.40%	65.93%
Thioluxan	3.07	43.91%	46.92%	40.67%

**DAS= Day After Spry**

Figure 1 indicated that the total reduction %age of *C. medinalis* of different treatments during 1<sup>st</sup> spray. The data showed the maximum total mean reduction %age (76.19%) was recorded in the plots treated with Deltaphos followed by Tracer (65.93%). However, the minimum reduction %age was recorded in the plots treated with Thioluxan (46.92%) after 1<sup>st</sup> spray on rice crop.



**Figure 1 Total reduction %age in the population of *C. medinalis* after different time intervals of 1<sup>st</sup> spray by insecticides on rice crop.**


The result of second spray showed that all insecticides behaved very similarly against rice leaf folder as record during the 1<sup>st</sup> spray. However, the efficacy with regard to time interval was changed a 1<sup>st</sup> during 2<sup>nd</sup> spray both Tracer and Thioluxan lost then efficacy after 3<sup>rd</sup> day. Table 2 showed that the maximum reduction (71.08%) was recorded in the plots treated with Deltaphos followed, by Trace (57.40%) and (30.15%) in Thioluxan respectively.

**Table 2 Mean population and reduction %age of different pesticides against *C. medinalis* on rice crop after 2<sup>nd</sup> spray by the application of insecticides.**

Treatments	Pre Treatment population	Reduction percentage in population		
		1 DAS	3 DAS	7 DAS
<b>Deltaphos</b>	1.93	49.04%	52.92%	71.08%
<b>Tracer</b>	2.03	35.09%	57.40%	53.38%
<b>Thioluxan</b>	2.07	28.00%	30.15%	17.27%

DAS= Day After Spry

Figure 2 indicated that the total mean reduction %age of *C. medinalis* of different treatments of 2<sup>nd</sup> spray. The data showed the maximum total mean reduction %age (71.08%) was recorded in the plots treated with Deltaphos followed by in the plots treated with Tracer (57.40%). However, the minimum reduction %age was recorded in the plots treated with Thioluxan (30.15%) after 2<sup>nd</sup> spray on rice crop.



**Figure 2 Total reduction %age in the population of *C. medinalis* after different time intervals of 2<sup>nd</sup> spray by insecticides on rice crop.**


During 3<sup>rd</sup> spray the behavior of insecticides was recorded in contrast of the results of 1<sup>st</sup> spray. During 3<sup>rd</sup> spray all insecticides lost the efficacy after 3rd day of spray. The data in Table 3 showed that the maximum reduction percentage was recorded in the plots treated with Deltaphos, (78.99%) followed by Thioluxan (55.88%) and Tracer (54.88%). However no significantly difference was recorded in the efficacy of Tracer and Thioluxan against rice leaf roller on rice crop.

**Table 3 Mean population and reduction %age of different pesticides against *C. medinalis* on rice crop after 3<sup>rd</sup> spray by the application of insecticides.**

Treatments	Pre Treatment population	Reduction percentage in population		
		1 DAS	3 DAS	7 DAS
<b>Deltaphos</b>	1.95	74.23%	78.99%	59.46%
<b>Tracer</b>	2.225	39.95%	54.34%	51.48%
<b>Thioluxan</b>	2.375	39.99%	55.88%	48.53%

DAS= Day After Spry

Figure 3 indicated that the total mean reduction %age in the population of *C. medinalis* by the application of different treatments during 3<sup>rd</sup> spray. The data showed the maximum reduction (78.99%) was recorded in the plots treated with Deltaphos followed by the plots treated with thioluxan (55.88%) and the plots treated with Tracer (54.34%).



**Figure 3 Total reduction %age in the population of *C. medinalis* after different time intervals of 3<sup>rd</sup> spray by insecticides on rice crop.**


The data in Table 4 revealed that the performance of the insecticides during 4<sup>th</sup> spray the maximum reduction was recorded in the plots treated with Deltaphos (71.92%), followed by the plots treated with Tracer (39.36%), and in the plots treated with Thioluxan (14.36%) at 1<sup>st</sup> day interval respectively.

**Table 4 Mean population and reduction %age of different pesticides against *C. medinalis* on rice crop after 4<sup>th</sup> spray by the application of insecticides.**

Treatments	Pre Treatment population	Reduction percentage in population		
		1 DAS	3 DAS	7 DAS
<b>Deltaphos</b>	1.00	71.92%	64.06%	35.15%
<b>Tracer</b>	1.25	39.36%	33.30%	19.21%
<b>Thioluxan</b>	1.25	14.36%	6.29%	6.35%

DAS= Day After Spry

However, Figure 4 showed that the maximum reduction %age in *C. medinalis* population that is recorded in the plots treated with Deltaphos (71.92%) followed by in plots treated with Tracer (39.36%). However, the minimum reduction %age was recorded in the plots treated with Thioluxan (14.36%) after 4<sup>th</sup> spray on rice crop.



**Figure 4 Total reduction %age in the population of *C. medinalis* after different time intervals of 4<sup>th</sup> spray by insecticides on rice crop.**

All insecticidal treatments gave the significantly more yield as compared untreated plots. Table 5 showed that the average paddy yield obtained after harvesting. The highest average paddy yield was obtained in the plots treated with Deltaphos (4783.00kg/ha<sup>-1</sup>), followed by the plots treated with Tracer (4486.46kg/ha<sup>-1</sup>) and the plots treated with Thioluxan (4033.66kg/ha<sup>-1</sup>) as compared control (3066.50kg/ ha<sup>-1</sup>). Increase in average yield over control (1716.50 kg/ ha<sup>-1</sup>) was recorded in plots treated with Deltophos. The average yield obtained in the treated plots with the application of Thioluxan (967.16 kg/ha<sup>-1</sup>) was statistically lower than the other former insecticides.

**Table 5 Average yield of rice crop after control operations of rice leaf-folder, *C. medinalis* at Sindh Agriculture University Tandojam during 2010.**

TREATMENTS	Average yield (kg/ ha <sup>-1</sup> )	Increase in yield over control(kg/ ha <sup>-1</sup> )
Deltaphos	4783.00	1716.50
Tracer	4486.46	1419.96
Thioluxan	4033.66	967.16
Control	3066.50	

## DISCUSSION

Present results partly corroborates to those of Misra *et al.* (2012) and Kaushik and Deb (2011) who have noted that the population suppression capacity of monocrotophos and cypermethrin was essentially prudent in some regions of India. The rice leaf folder population before and after application of insecticides were averagely counted. When the rice leaf folder was appeared on the rice field sprays with different chemical insecticides against rice leaf folder. The rice field was sprayed till sizeable reduction was brought by the different chemical products which were testes against rice leaf folder. In the same pattenen Ramasubbaiah *et al.* (1980) noted that fenthion, phosphamidin, fenitrothion, endosulphon dimethoate, quinalphos, diazinon and carbaryl could effectively suppress LF menace. The rice field was sprayed six times since the appearance of rice leaf folder onto them the 1<sup>st</sup> spray was done. The data reveal that maximum reduction numbers in rice leaf folder population was recorded with one week after spray in all treatments during all six sprays. Saroja and Raju (1982) have viewed that cypermethrin and fanvalerate are best suitable pesticide to suppress rice leaf folder population and accordingly to maximize paddy yield. Our findings are in agreement with their of Bhanu *et al.* (2008) who have noted considerable variations in the efficacy of pesticides in field condition. Wakil *et al.* (2001) from Pakistan have reported that not all the pesticides were equally effective to check leaf folder attack.

Data further reveled that the maximum reduction percentage of rice leaf folder population was recorded as (73.22%) in the plots treated with Deltaphos followed by in the plots treated with Tracer (64.17%). However, the minimum reduction %age was recorded in the plots treated with Thioluxan (43.80%) during all for applications of insecticides on rice crop. The plots treated with Deltaphos gave the highest reduction in the population of rice leaf folder followed by Tracer and Thioluxan. The reduction in the population caused by these insecticides was statistically at par. Thioluxan significantly lower mortality than the former pesticides. Our results are in partial agreement with those of Mishra *et al.* (1998) and Kaushik and Deb (2011) who observed that monocrotophos and cypermethrin gave good control of rice leaf-folder and were at par statistically. All insecticidal treatments significantly out yielded the untreated plots. The highest average paddy yield was obtained in plots treated with the application of Deltaphos followed by Tracer, Thioluxan and Control. Our results are almost similar to the Saroja and Raju (1982) who obtained similar increase in yield by controlling *C. medinalis* damage by the application of synthetic pyrethroids.

## Conclusion

The Deltophos pesticide proved better to control rice leaf folder as compared Tracer in rice field . Maximum reduction in the population of *C. medinalis* was recorded very next day of application. All insecticides lost their effect within in a week.

## Recommendations

The insecticide Daltaphos should be used an excellent and rapid tool to diminish rice leaf folder population in rice field. Spraying of Deltaphos should be kept continue at week interval until a sizeable reduction in the population of RLF is achieved.

## LITURECURE CITED

- Ahmad M., I. U. Haq, M.S. Wains, M. Anwer and M. Ahmad, 2005. Screening of advanced breeding materials for resistance to rice leaf folder under field conditions. *Proceedings of the International seminar on Rice crop*. Oct. 2-3, 2005, Rice Research Institute, Kalashahkaku. pp 293-296.
- Ahmed, H., R.B.Khan, D. Sharma, V.V.S. Jamwal, and S. Gupta, 2010. Seasonal incidence, infestation and trap catches of *Cnaphalocrocis medinalis* (Guenee) in rice. *Annals Pl. Prot. Sci.* 18(2):38-383.
- Alvi, S.M., Ali, M.A., Chaudhary, S. and S. Iqbal, 2003. Population trends and chemical control of rice leaf folder, *Cnaphalocrocis medinalis* on rice crop. *Inter. J. Agric. Biol.* 5: 615–617.
- Bhanu, K. V. and P.S. Reddy, 2008. Field evaluation of certain newer insecticides against rice insect pests. *J. Appl. Zool. Res.* 19 (1):11-14.
- Bhatti, M.N. 1995. Rice leaf folder (*Cnaphalocrocis medinalis*): A review. *Pak. Entomol.* 17: 126– 131.
- Gottfried F., F. Fallil, 1980. The spinning (Stitching) behaviour of the rice leaf folder, *Cnaphalocrocis medinalis*. *Entomologia Experimentalis et Applicata*. vol: 29 Issue (2) pp 138-146.
- Gunathilagaraj, K. and M.Gopalan, 1986. Rice Leaf folder Complex in Madurai, TN, India. *Intl. Rice Res. Notes* 11(6): 24.
- Heong, K.L., 2005. Rice leaf folder: are they serious pests? In: Hu, G.W., Guo, Y.J., Li, S.W. (Eds.), *Research on Rice Leaf Folder Management in /china*, Proceeding of the International Workshop on Economic threshold level of for rice leaf folder in China, March 4-6, 1992, Beijing. China Agricultural Science and Technology Publisher, Beijing, China, pp.8-11.
- Heong, K.L. and M.M.Escalanda, 1997. A comparative analysis of pest management practices of rice farmers in Asia. In *pest management of rice farmers of Asia*. Eds. K.L. Heong and M.M. Escalanda. Pp. 227-245. International Rice Research Institute, Philippines.
- Kaushik, C. and D. C. Deb, 2011. Extent of suppression of leaf folder, *cnaphalocrocis medinalis*, guen. population by some selected insecticides in the field of scented local paddy cultivar tulaipanji at Raiganj, Uttar Dinajpur, West Bengal, India, *International Journal of Plant, Animal and Environmental Sciences*, 1 (3): 142-149.
- Khan, M.R., M. Ahmad and S. Ahmad, 1989. Some studies on biology, chemical control and varietal preference of rice leaf folder, *Cnaphalocrocis medinalis*. *Pak. J. Agric. Sci.* 26: 253–63.
- Maragesan, S. and S. Chellish, 1987. Yield losses and economic injury by rice leaf folder. *Indian J. Agric. Sci.* 56: 282–85.
- Mishra, B.K., P. R. Mishra and S. M. A. Mandal, 1998. Screening of rice cultivars for resistance to leaf folder. *Indian J. Entomol.*, 64(1):68-72.
- Misra H P., 2012. Bioefficacy of Fenprothrin 30% EC for the control of the rice leaf folder, *Cnaphalocrocis medinalis* (Guen). *An International Journal on Rice*. Vol: 49, Issue : 4 pp 284 - 287.
- Nugaliyadde L., T. Hidaka, and M.P. Dhanapala. 1997. Pest management practices of rice farmers in Sri Lanka. In *pest management of rice farmers in Asia*. Eds. K.L. Heong and M.M. Escalanda. Pp 171- 184. International Rice Research Institute . Phillipines.
- Rajendran, R., S. Rajendran and P.C. Sandra, 1986. Varietals resistance of rice of leaf folder. *Intl. Rice Res. News* 11: 17.
- Ramasubbaiah, K.; P.S. Rao and A.G. Rao, 1980. Nature of damager and control of rice leaf folder. *Indian J. Entomol.* 42: 214–7
- Sabir, A. M., S. Ahmad, M. Hassan and A. Qadir, 2006. Pest weather interaction of major insect pests in rice ecosystem. *SAARC J. Agric.*, 4: 203-212.
- Saroja, R. and N. Raju, 1982. An effect of foliar insecticides on rice stem borers and leaf folders. *Intl. Rice Res. News* 7:14.
- Wakil Waqas, Mansoor-ul-Hasan, Rizwan Akbar & Assam Gulzar 200 I. Evaluation of different insecticides against rice stem borer and rice leaf folder. *Pak. 1. Agri. se: Val' 38 (3-4)*,



The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:  
<http://www.iiste.org>

## CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

**Prospective authors of journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

## MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

## IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

