

# Incidence of aflatoxins, fumonisins, trichothecenes and ochratoxins in Nigerian foods and possible intervention strategies

Adejumo\*, T. O. and Adejoro, D. O.

Dept. of Microbiology, Adekunle Ajasin University, P.M.B. 01, Akungba-Akoko, Ondo State, Nigeria.

\*Corresponding Author: [toadejumo@yahoo.com](mailto:toadejumo@yahoo.com)

## Abstract

Mycotoxins are toxic secondary metabolites produced by some species of fungi. Aflatoxins, fumonisins, trichothecenes and ochratoxins are the common mycotoxins in Nigeria. Aflatoxin is the most frequently reported in literatures, with trichothecenes being the least, they cause yield loss to farmers as well as constituting major health risk to humans. The occurrence of mycotoxins in food is a serious problem that Nigeria is facing presently, as it continues to pose threat to feed and food safety of animals and humans. There is the need to seek for approaches that would lead to reduction in their toxicity. The practice of good sanitary measures right from the farm to storage, creation of awareness campaign to indicate the toxic effects associated with mycotoxin poisonings in humans and livestock, and proper evaluation of food crops for its presence can go a long way in achieving the target reduction in incidence of mycotoxins in Nigeria.

## 1.0 Introduction

Mycotoxins are secondary metabolites of fungal origin which produce toxic responses when ingested by animals or humans. The word 'mycotoxin' is a combination of a Greek word, 'mykes' meaning 'fungus' and a Latin word 'toxicum' meaning 'by poison' (Rai *et al.*, 2012). Mycotoxicosis is a term used to denote the diseases that result from the ingestion of mycotoxin by animals and humans (Nierman *et al.*, 2008). The acuteness of a specific mycotoxicosis may vary in different victims. Various factors determine the severity of a mycotoxin; among them are the level of toxicity, dosage, age, nutritional and immunity status of the victim (Peraica *et al.*, 1999).

The general attraction to the study of mycotoxins started in 1960 when more than 100,000 turkeys died in London, England after consuming contaminated groundnut meal that was imported from Brazil (Blount, 1961). The disease that resulted was called 'Turkey X' disease. *Aspergillus flavus* was later isolated from the groundnut meal and the toxic agents named 'aflatoxin' (*A. flavus* toxin) accordingly (Splensley, 1963; Kensler *et al.*, 2011). The word 'mycotoxin' was then coined in 1962 after the Turkey X incident.

Most mycotoxins of concern are produced by three genera of fungi, namely, *Aspergillus*, *Penicillium*, and *Fusarium*. These are the predominant fungal genera associated with food grains during storage (CAST, 2003). These genera are considered significant because of their ubiquity in the environment and their ability to produce various types of mycotoxins like aflatoxins, fumonisins and ochratoxins (Juan *et al.*, 2010). However, the presence of these fungi does not necessarily imply the presence of toxins.

The mycotoxigenic fungi can be classified as either field or storage fungi. It should be noted that a fungus may produce several mycotoxins and a mycotoxin may be produced by different fungi (Fernandez-Cruz *et al.*, 2010). There are intrinsic and extrinsic factors which influence the production of mycotoxins. The intrinsic factors include pH, water activity ( $a_w$ ) and redox potential of the substrate while the extrinsic factors are the relative humidity, temperature and oxygen availability of the environment where the commodities are produced, stored or processed (Whitlow and Hagler, 2004; Nagwa *et al.*, 2013).

Over 400 types of mycotoxins have been reported (Bhat *et al.*, 2010), but only four (aflatoxins, fumonisins, trichothecenes and ochratoxins) will be discussed in this review. While the other mycotoxins are important for studies, aflatoxins, fumonisins, trichothecenes and ochratoxins are of greater interests because of their prevalence in agricultural produce and the harmful effects they exert when they are consumed by animals and humans (Richard, 2007).

The Food and Agriculture Organization (FAO) estimated that about 25% of the world's agricultural produce is contaminated with mycotoxins which cause huge losses for farmers (Wu, 2007). These losses are measured in reduced crop yields, lower quality, reduced animal performance and reproductive capabilities, and increased disease incidence.

The biochemical properties of mycotoxins are diverse, and their toxic effects are exceedingly variable. Mycotoxins are carcinogenic, tremorogenic, haemorrhagic, genotoxic, teratogenic, nephrotoxic, hepatotoxic and

immunotoxic (Refai, 1988; Hosseini and Bagheri, 2012). Mycotoxins usually enter the body via ingestion of contaminated foods, but inhalation of toxigenic spores and direct dermal contact are also important routes (Omar, 2013).

Nigeria enjoys the humid tropical climate type and experiences consistent high temperatures all year round. The tropical climate is characterized by the hot and wet conditions associated with the movement of the Inter-Tropical Convergence Zone (ITCZ) north and south of the equator. This produces two seasons: the rainy or wet season that lasts from mid-March to November in the southern parts of the country and from May to October in the northern parts of the country; and the dry season that fills up the rest of the year (Oyenuga, 1967).

Iloeje (2001) classified Nigerian agroecological zones into two broad categories: forest and savannah zones. Forest is the ecological zone in which trees are the dominant species while savannah is a grassland area with no forest cover. The important agroecological zones for mycotoxin studies in Nigeria are humid forest, derived savannah and Guinea savannah zones. The humid forest zone is characterized by two growing seasons, starting from April to November with an annual rainfall of 1,500 mm and 2,000 mm, average annual temperature of 24.5 to 27.5°C, and mean relative humidity of 78 to 100% (Afolabi *et al.*, 2013). Cash crops such as oil palm, cocoa and kolanut, and staple crops such as maize, rice, beans and groundnut are found in this region. Derived Savannah is the transition between the rainforest and guinea savannah zones. It has an annual rainfall range of 1,200mm to 1,700 mm, average annual relative humidity range of 66 to 78% and mean temperatures of 26 to 27 °C; maize, cassava, yam and rice are the common crops grown in this region. The Guinea Savannah is the most extensive ecological zone in Nigeria, occupying almost half the area of the country. The zone is sometimes subdivided into the Southern and Northern Guinea savannah. It has a wet season which lasts from May to October, with an annual rainfall of 1,000mm to 1,200 mm, relative humidity of 57 to 66%, and average annual temperature of 25.5 to 26.5 °C (Afolabi *et al.*, 2013). Crops in areas like this are more prone to contamination with mycotoxin-producing fungi because of the high temperature and humidity (Sherif *et al.*, 2009).

In Nigeria, mycotoxin contamination of cereals (rice), grains (maize) and seeds (cocoa) has raised a lot of concern for food safety (Kumar *et al.*, 2008) as these foods, especially rice and maize are not only eaten directly, but are also used in the production of various forms of indigenous foods like *ogi*, *eko*, *tuwo*, *kunu*, *donkwa*, and *masa* among others. Bankole and Adebajo (2003) noted that the mycotoxin that has received the most attention in scientific literature in food products from West African sub-region is aflatoxin, while there are few researches carried out on other mycotoxins like fumonisin and ochratoxin A. However, there have been some recent studies on the mycotoxins present in food products from Nigeria, especially maize, rice, groundnuts, guinea corn, sorghum, cocoa and cocoa-based beverages (Adejumo *et al.*, 2007a, 2007b; Ayejuyo *et al.*, 2008; Dongo *et al.*, 2008; Amadi and Adeniyi, 2009; Makun *et al.*, 2010; Olayiwola *et al.*, 2013).

## 2.0 Mycotoxins

### 2.1 Aflatoxin

Aflatoxins are a group of toxic and carcinogenic secondary metabolites of fungal origin. They are produced by strains of *Aspergillus flavus*, *A. parasiticus* and, in rare cases, *A. nominus* and *A. pseudotamari* (Mazaheri, 2009). Among the 18 different types of aflatoxins identified the four naturally occurring ones are Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>), Aflatoxin B<sub>2</sub> (AFB<sub>2</sub>), Aflatoxin G<sub>1</sub> (AFG<sub>1</sub>) and Aflatoxin G<sub>2</sub> (AFG<sub>2</sub>) (Filazi and Sireli, 2013). Aflatoxin B<sub>1</sub> and B<sub>2</sub> were named because of the strong blue fluorescence under ultraviolet light, hence the 'B', while Aflatoxin G<sub>1</sub> and G<sub>2</sub> are so named because of their greenish yellow fluorescence (Kensler *et al.*, 2011). The subscript designates relative chromatographic mobility during thin-layer chromatography, '1' and '2' indicate major and minor compounds respectively (Zain, 2011).

Apart from the four major aflatoxins, there are two other metabolites, designated as M<sub>1</sub> and M<sub>2</sub>. Aflatoxin M<sub>1</sub> and M<sub>2</sub> are metabolites of Aflatoxin B<sub>1</sub> and B<sub>2</sub> respectively and they may be present in milk, eggs or blood of animals that are fed with AFB<sub>1</sub> and AFB<sub>2</sub>-contaminated feed (IARC, 2002; CAST, 2003). Toxigenic *A. flavus* generally only produces AFB<sub>1</sub> and AFB<sub>2</sub> while




Figure 1. Chemical structures of AFB<sub>1</sub>, AFB<sub>2</sub>, AFG<sub>1</sub>, AFG<sub>2</sub>, AFM<sub>1</sub> and AFM<sub>2</sub> (Zain, 2011).

*A. parasiticus* (the most toxigenic species) produce AFB<sub>1</sub>, AFB<sub>2</sub>, AFG<sub>1</sub> and AFG<sub>2</sub> (Davis and Diener, 1983; Van Egmond, 1989). The order of toxicity in these aflatoxins is

AFB<sub>1</sub>>AFG<sub>1</sub>>AFB<sub>2</sub>>AFG<sub>2</sub>; AFB<sub>2</sub> and AFG<sub>2</sub> are typically nontoxic except they are metabolized into AFB<sub>1</sub> and AFG<sub>1</sub> inside the cells (Kensler *et al.*, 2011; Filazi and Sireli, 2013). AFB<sub>1</sub> is always genotoxic *in vitro* and *in vivo* (EFSA, 2007).

Aflatoxins are among the most powerful carcinogenic, mutagenic and teratogenic compounds that occur naturally (Jackson and Al-Taher, 2008). In view of this, the International Agency for Research on Cancer (IARC) classified aflatoxin as a Class I human carcinogen (IARC, 1993).

Aflatoxicosis is the disease associated with the ingestion of aflatoxins. Outbreaks of aflatoxicosis have been reported in Kenya, India and Thailand, but none has been reported so far in Nigeria. The outbreak which occurred in Kenya is one of the largest and most acute outbreaks of aflatoxicosis ever documented (CAST, 2003), with 317 cases reported and 125 deaths recorded (CDC, 2004). Investigations revealed that the outbreak was due to the consumption of maize contaminated with extremely toxic AFB<sub>1</sub>. The case-control studies carried out by Lewis *et al.* (2005) showed that aflatoxicosis in the affected area was associated with eating home-grown maize stored under humid conditions. During the outbreak, individual daily exposure of AFB<sub>1</sub> was estimated to be 50mg/day (Probst *et al.*, 2007).

Bankole and Adebajo (2003b) reported that many studies on aflatoxin had been carried out in West Africa. This includes investigation to know the levels of aflatoxin in rice, maize, groundnuts, guinea corn, sorghum and cocoa, as well as impacts of aflatoxin on human beings. McDonald (1964) reported that groundnuts cultivated in the northern parts of Nigeria were contaminated with aflatoxin levels of up to 2000µg/kg. Salifu (1981) studied fungal and mycotoxin contamination of sorghum in Nigeria, and reported that aflatoxin levels ranged from 10 to 80 µg/kg<sup>-1</sup>. In a recent report by Ezekiel *et al.* (2013), about 90% (26/29) of peanut cake samples had aflatoxin concentrations exceeding the maximum limit of 20 mg/kg set by USDA (FAO, 2004) (Table 1).

### 2.1.1 Health Risk of Aflatoxins to Humans and Animals

The primary disease associated with the ingestion of aflatoxins is hepatocellular carcinoma (liver cancer) (Williams *et al.*, 2004), which according to WHO (2008) is the third-leading cause of cancer worldwide, with about 600,000 fresh cases each year. Aflatoxin contamination has been shown to reduce feed intake, increase liver and kidney weights of farm animals, as well as induce immunosuppression and hepatitis in them; all of which contribute to increased mortality in farm animals (Hussein and Brussel, 2001; Zain, 2011).

Smith *et al.* (1992) reported that feeding broilers with a combined high level (3.5mg/kg) feed of aflatoxin blend (AFB<sub>1</sub>: 79%; AFB<sub>2</sub>: 4%; AFG<sub>1</sub>: 16% and AFG<sub>2</sub>: 1%) caused a reduction in body weight, and an increase in weights of liver and kidney. In cattle, increasing aflatoxin to levels such as 10, 26, 56.4, 81.1 and 108.5µg/kg has been shown to reduce feed intake depending on the dosage (Choudhary *et al.*, 1998). The quality of milk produced by these cattle is also affected by the aflatoxins (Zain, 2011).

Report on aflatoxin-related human health problems in Nigeria is limited. Aflatoxin M<sub>1</sub> has been reported in breast milk of lactating mothers in Ogun State, with 85% of the breast milk screened

**Table 1. Incidence of Aflatoxin-contamination of Common Nigerian Foods**

Commodities	Region	Type of Aflatoxin	Incidence Rate	Range µg/kg	References
<b>Rice</b>	Niger State, Nigeria	AFB <sub>1</sub> AFB <sub>2</sub> AFG <sub>1</sub> AFG <sub>2</sub> Total AF	100% 21/21	4.1-30.9 1.3 – 24.2 5.5-76.8 3.6-44.4 28-372	Makun et al., 2011.
<b>Dried Yam Chips</b>	South-western Nigeria	AFB <sub>1</sub> AFG <sub>1</sub> AFB <sub>2</sub>	54.2% 5.2% 32.3%	4-186 4-18 2-55	Bankole and Adebajo, 2003a.
<b>White Yam Flour</b>	South-western Nigeria	AFB <sub>1</sub> AFG <sub>1</sub>	57% 21%	0.02-3.2 0.05-3.5	Somarin et al., 2012
<b>Water Yam Flour</b>	South-western Nigeria	AFB <sub>1</sub>	32%	0.02-0.6	Somarin et al., 2012
<b>Kulikuli</b>	Oyo State, Nigeria.	AFB <sub>1</sub>	90% (18/20) 100% (29/29)	20-455 13-2824	Akano and Atanda, 1990; Ezekiel et al., 2012.
<b>Dried Mushrooms</b>	Lagos State, Nigeria.	ND*			Ezekiel et al., 2013.
<b>Tiger nuts</b>	Nigeria		35%	10-120	Bankole and Esegbe, 1996
<b>Beans Cassava flour Semovita Yam Wheat meal "Gari"</b>	Ogun State, Nigeria	AFB <sub>1</sub>		nd-0.89 nd-0.07 nd-0.17 nd-0.27 nd-0.06 nd-0.69	Adejumo et al., 2013

**Table 1. Incidence of Aflatoxin-contamination of Common Nigerian Foods Contd.**

Commodities	Region	Type of Aflatoxin	Incidence Rate	Range µg/kg	References
<b>Maize</b>	Niger State, Nigeria	AFB <sub>1</sub> AFB <sub>2</sub>		0-1874 0-608	Atehnkeng et al., 2008
<b>Palm oil</b>	Ibadan, Nigeria	AFB <sub>1</sub>	ND*	ND*	Olorunfemi et al., 2014
<b>Wheat Millet Guinea corn Breadfruit Groundnut</b>	Niger Delta Region, Nigeria.	AFB <sub>1</sub>	100% 100% 100% 100%	17.01-20.53 34.00– 40.30 27.22-36.13 40.06-48.59 74.03-82.12	Odoemelam and Osu, 2009
<b>Yam chips</b>	Ibadan, Oyo State.	AFB <sub>1</sub>	97.5%	11-196	Abiala et al., 2011

<b>Fresh milk</b>	Bida, Niger State	AFM <sub>1</sub>	100%	0.407-0.952µg/l	Okeke et al., 2012
<b>Nono</b>			100%	0.248-2.510µg/l	
<b>Kindirmo</b>			100%	0.139-1.238µg/l	
<b>Powdered milk</b>	Lagos, Lagos State	AFM <sub>1</sub>	19%	0.02 - 0.41µg/l	Makun et al., 2010
<b>Beans</b>	Minna, Niger State	AFB <sub>1</sub>	58%	63.5 - 106.2	Makun et al., 2010
<b>Wheat</b>			54%	102.9 - 198.4	
<b>Breast milk</b>	Ogun State	AFM <sub>1</sub>	85%	3.49-35ng/l	Adejumo et al., 2013

nd – Not Determined

\*ND – Not Detected

contaminated with AFM<sub>1</sub> (3.49-35 ng/l) and 16% of the breast milk samples exceeding the European Union (EU) limit of 25 ng/l (Adejumo *et al.*, 2013).

Uriah *et al.* (2001) reported that the blood and semen in infertile men attending the University of Benin Teaching Hospital had aflatoxin levels which ranged from 700 to 1392ng/ml and 60 to 148ng/ml respectively and these values were significantly higher than the concentrations of the toxin in fertile men. Twenty-two out of the 60 samples (37%) of serum and semen of infertile men had aflatoxin B<sub>1</sub> ranging from 60ng/ml to 148ng/ml. Although, their studies indicated that aflatoxins might have a negative effect on fertility, more studies are needed to ascertain the universality of this connection.

Aflatoxins were reported in the autopsy kidney specimens of 58% of children with kwashiorkor at Obafemi Awolowo University Teaching Hospital, Ile-Ife (Oyelami *et al.*, 1998). In a similar study by Onyemelukwe *et al.* (2012), aflatoxins were found in both the sera and the blood of children with kwashiorkor. Aflatoxins were present in the sera of 88.9% of the children with a range of 5.3 to 7646µg/l, and in the urine of 84.6% of the children with a range of 15 to 361µg/l. The study data indicate that higher aflatoxins were more frequently detected in sera and urine of patients with kwashiorkor, as compared to the healthy control within a similar age-group (Hendrickse, 1983).

## 2.2 Fumonisin

Fumonisin are fungal metabolites produced by various *Fusarium* species (Senyuva *et al.*, 2010), primarily by *Fusarium verticillioides*, formerly *F. moniliforme* (Seifert *et al.*, 2003), and *Fusarium proliferatum*, which are global contaminants of maize and maize products (Adejumo *et al.*, 2007a; Somorin *et al.*, 2012). Recently, it was discovered that fumonisins are also produced by *Aspergillus niger* (Soares *et al.*, 2013). The fumonisins were first isolated from cultures of *F. moniliforme* strain MRC at the South African Medical Research Council by Gelderblom *et al.* (1988). The structures of Fumonisin B<sub>1</sub> and B<sub>2</sub> (Figure 2) were first elucidated by Bezuidenhout *et al.* (1988).

Fumonisin, primarily FB<sub>1</sub>, FB<sub>2</sub> and FB<sub>3</sub>, are fluorescent cancer-promoting metabolites that have a long-chain hydrocarbon unit which is similar to that of sphingosine and sphinganine, which plays a role in their toxicity (Marin *et al.*, 2013). Maize is the major commodity affected by fumonisins, although some occurrences have been reported in sorghum and rice (Hosseini and Bagheri, 2012). Fumonisin are formed in maize prior to harvest or during the early stage of storage. Except under extreme conditions, their concentrations do not increase during storage (Marin *et al.*, 2013).






Figure 2. Chemical Structures of Fumonisin B<sub>1</sub> and B<sub>2</sub> (Zain, 2011; Marin *et al.*, 2013)

Among fumonisins, FB<sub>1</sub> is the most toxic, and has been proven to stimulate tumour growth in rats, cause leukoencephalomalacia in horses, pulmonary oedema in pigs and brain haemorrhage in rabbits (Gelderblom *et al.*, 1988; Marasas, 1995; Hussein and Brasel, 2001). They have also been linked with high incidence of human oesophageal cancer in some regions of South Africa and China (Sydenham *et al.*, 1990; Sun *et al.*, 2007).

Missmer *et al.* (2006) reported that fumonisin exposure increases the risk of neural tube defects (with respect to dose) up to a threshold level, at which point foetal death may be more likely to occur. In Nigeria, Bankole *et al.* (2003) detected fumonisin B<sub>1</sub> in 51% (55 out of 104 samples) of maize samples analysed with concentration range of 65 to 1830 µg/kg, and mean value of 390 µg/kg (Table 2). In a similar study on the natural occurrence of fumonisins in pre-harvest maize in south western Nigeria, Bankole and Mabekoje (2004) reported *F. verticillioides* occurring in 89.3% of samples. Fumonisin B<sub>1</sub> was reported to be the predominant toxin (frequency at 76.8%, concentration between 70 and 1780 µg/kg, and mean of 495 µg/kg), while Fumonisin B<sub>2</sub> was detected in 66% of the samples with a mean of 114 µg/kg.

Avantaggio *et al.* (2002) reported that insect damage was a good predictor of *Fusarium* mycotoxin contamination. The insect damage to maturing maize ears allows the strains of *Fusarium* to enter the ears and kernels. *Fusarium* species are always present in the ears and kernels of maize and fumonisins have even been shown to be present in symptomless kernels of maize in Nigeria (Thomas and Buddenhagen, 1980). Adejumo *et al.* (2007b) reported FB<sub>1</sub> in 73% of maize samples from four South-western states of Nigeria (mean 117 µg/kg; range 10 to 760 µg/kg). In a later study, the authors also investigated 80 maize samples collected from South-western Nigeria for fumonisin B<sub>1</sub> content (Adejumo *et al.*, 2009), and reported FB<sub>1</sub> in 55 (68.7%) of the total samples (range: 10 to 126.7 µg/kg; mean: 98.5 µg/kg). Somorin *et al.* (2012) reported that low levels of fumonisins were detected in yam flours which were being used in making *amala* in south western Nigeria, FB<sub>1</sub> was in 32% of white yam flour samples (range: 0.5 to 90 µg/kg; mean 5 µg/kg), and 5% of water yam samples (range 0.5 to 2 µg/kg). Makun *et al.* (2011) reported FB<sub>1</sub> and FB<sub>2</sub> at frequencies of 14.3% and 4.8% respectively in rice samples from Niger State.

Most of these occurrences of fumonisins in Nigerian commodities are still within the acceptable limit of <1000 µg/kg for fumonisins (CAST, 2003), although there is still the need for frequent monitoring of foods for fumonisin contamination.

### 2.3 Trichothecenes

Trichothecenes are a group of structurally related compounds with a common tetracyclic sesquiterpenoid 12, 13-epoxytrichothec-9-ene ring system. The trichothecenes are chemically the most diverse of all the mycotoxins with over 200 identified (Grove, 2007). Some trichothecenes are among the most toxic non-nitrogenous compounds known to man. They are powerful inhibitors of eukaryotic protein synthesis, insecticidal, phytotoxic, and toxic to animals (Grove, 2007).


**Table 2. Incidence of fumonisin-contamination of common Nigerian foods and feeds**

Commodity	Region	Type	Incidence	Range (µg/kg)	Mean (µg/kg)	References
<b>Rice</b>	Niger state	FB1	14.3%	0.4-4.4		Makun et al., 2011
		FB2	4.8%	132.5-132.5		
<b>White yam flour</b>	South western Nigeria	FB1	32%	0.5-91	5	Somorin et al., 2012
		FB2	12%	1-32	ND	
<b>Water Yam Flour</b>	South western Nigeria	FB1	5%	0.5-2		Somorin et al., 2012
<b>Maize</b>	South	FB1	51%	65-1830	390	Bankole et al., 2003
	western	FB <sub>1</sub>	73%	10-760	117	Adejumo et al., 2007b
	Nigeria	FB <sub>1</sub>	68.7%	10-714	98.5	Adejumo et al., 2009
<b>Maize (Pre-harvest)</b>	South western, Nigeria	FB <sub>1</sub>	78.6%	70-1780	495	Bankole and Mabekoje, 2004
		FB <sub>2</sub>	66%	70-1780	114	
<b>Dried Meat (Tinko)</b>	Ibadan, Nigeria	FB <sub>1</sub>	10%	0-1.91	ND	Oladejo and Adebajo, 2011
		FB <sub>2</sub>	10%	0-1.04	ND	
<b>Animal feed</b>	Nigeria	FB <sub>1</sub>	89%	20-2860	1092	Rodrigues et al., 2011
		FB <sub>2</sub>	89%	20-855	338	

ND - Not Determined


The trichothecenes are grouped into four types (A, B, C and D), named according to their characteristic functional groups with types A and B trichothecenes being the most common (Krska *et al.*, 2007). Examples of type A trichothecenes are HT-2 toxin (HT2), T-2 toxin (T2) 4,15-diacetoxyscirpenol, 15-monoacetoxyscirpenol, T-2 triol, T-2 tetraol, neosolaniol, 4,15-diacetylverrucarol, and verrucarol; while examples of type B are deoxynivalenol (DON), 3-acetyl-DON, 15-acetyl-DON, nivalenol (NIV) and fusarenon-X (FUS-X) (Krska *et al.*, 2007);

**Deoxynivalenol (Type B)**



**T-2 Toxin (Type A)**

**Diacetoxyscirpenol**



**Satratoxin G (Type D)**

Figure 3. Chemical structures of deoxynivalenol, diacetoxyscirpenol, T-2 toxin and satratoxin G (Hussein and Brasel, 2001; Gottschalk *et al.*, 2009)

Gottschalk *et al.*, 2009; Marin *et al.*, 2013). The C trichothecenes include crotoxin and baccharin, while type D trichothecenes are verrucarin, satratoxin G and H, and roridin A (Sudakin, 2003). However, type C and D trichothecenes are of lesser importance (Marin *et al.*, 2013).

The simple type A and B trichothecenes are produced by a variety of different *Fusarium* species (Thrane, 2001), although a number of other genera including *Trichoderma*, *Stachybotrys*, *Myrothecium*, and *Trichothecium* are also known to produce them (Sweeny and Dobson, 1998). However, the most important T-2 and HT-2-producing species are *F. sporotrichioides*, *F. langsethiae*, *F. acuminatum*, and *F. poae*. The main producers of DON are *F. graminearum*, *F. culmorum*, and *F. cerealis* (Liddell, 2003; Marin *et al.*, 2013). Type D trichothecenes are

**Table 3. Trichothecenes-producing fungi in food commodities**

Trichothecenes	Fungi	Reference
<b>Type A and B trichothecenes</b>	<i>Fusarium</i> species, <i>Trichoderma</i> , <i>Stachybotrys</i> , <i>Myrothecium</i> and <i>Trichothecium</i> .	Sweeny and Dobson, 1998; Thrane, 2001
<b>T-2 Toxin (Type A)</b>	<i>F. sporotrichioides</i> , <i>F. langsethiae</i> , <i>F. acuminatum</i> and <i>F. poae</i> .	Liddell, 2003; Marin <i>et al.</i> , 2013
<b>HT-2 (Type A)</b>	<i>F. sporotrichioides</i> , <i>F. langsethiae</i> , <i>F. acuminatum</i> and <i>F. poae</i> .	Liddell, 2003; Marin <i>et al.</i> , 2013
<b>DON (Type B)</b>	<i>F. graminearum</i> , <i>F. culmorum</i> and <i>F. cerealis</i> .	Liddell, 2003; Marin <i>et al.</i> , 2013
<b>Roridin A and verrucarin A (Type D)</b>	<i>Myrothecium</i> spp.	Ueno, 1983
<b>Satratoxin G and H (Type D)</b>	<i>Stachybotrys</i> spp.	Ueno, 1983

metabolites of *Myrothecium* spp. (e.g. roridin A, verrucarin A) and *Stachybotrys* spp. (e.g. satratoxin G and H) (Ueno, 1983).

There is paucity of information on contamination of Nigerian commodities trichothecenes. A few notable studies include the one conducted by Bankole *et al.* (2010) between 2005 and 2006 on Nigerian maize, with a reported 15-monoacetoxyscirpenol (15MAS) in one out of 32 samples at concentration of 4 µg/kg and T-2 teraol in two out of 32 samples at levels of 73 and 280 µg/kg. The authors concluded from the study that trichothecenes toxins do not appear to be major contaminants of Nigerian maize. However, Adejumo *et al.* (2007a) reported the screening of 180 maize samples meant for human consumption for twelve major trichothecenes (deoxynivalenol (DON), 3, mono-acetyldeoxynivalenol 93-AcDON), 15, mono-acetyldeoxynivalenol (15-AcDON), nivalenol (NIV), HT-2 toxin (HT-2), neosolaniol (NEO), T-2 toxin (T-2), T-2 tetraol and T-2 triol, diacetoxyscirpenol (DAS), MAS-monoacetoxyscirpenol (MAS) and fusarenone-X), and showed that sixty-six (36.3%) samples were contaminated with trichothecenes -DON, 3-AcDON, and DAS at different ranges. Authors of the report claimed that it was the first comprehensive report of the natural occurrence of trichothecenes in maize for direct human consumption in Nigeria.

A similar study was conducted on rice by Makun *et al.* (2011) who detected DON in 23.8% of rice sampled from Niger State, at a mean concentration of 18.9µg/kg and range of 0 to 112.2µg/kg. The reports on maize by Adejumo *et al.* (2007a) and Bankole *et al.* (2010) were followed up by Afolabi *et al.* (2013), who reported the detection of T-2 toxin in 36% of total maize samples; 29µg/kg, 13µg/kg and 12µg/kg were the maximum, mean and median values for T-2 toxin respectively.

## 2.4 Ochratoxins

Ochratoxins are toxic fungal metabolites produced by various *Aspergillus* species, *Penicillium verrucosum* and *P. nordicum* (Frisvad *et al.*, 2004). They are named after *A. ochraceus* from which the compound was first isolated (van der Merwe *et al.*, 1965). Several ochratoxins occur in nature, namely Ochratoxin A (OTA), Ochratoxin B (dechlorinated OTA), and Ochratoxin C (ethylated OTA) which are often co-produced (Reddy and Bhoola, 2010). However, OTA is the most toxic and most prevalent member of the ochratoxins (Marin *et al.*, 2013). Ochratoxin A is nephropathic (Kumar *et al.*, 2008), immunosuppressive, teratogenic, genotoxic, hepatotoxic and



**Table 4. Incidence of Trichothecene Contamination of Nigerian food commodities**

Commodity	Region	Type	Frequency	Mean and Range	Reference
<b>Maize</b>	Nigeria	15 MAS T-2 teraol	1/32 2/32	Mean:4µg/kg 73 µg/kg and 280µg/kg	Bankole et al., 2010
<b>Maize</b>	Southwestern Nigeria	DON	22%	Mean: 226.2µg/kg Range: 9.6-745.1µg/kg	Adejumo et al., 2007a
		3-AcDON DAS	17% 9%	Mean: 17.3µg/kg Range: 0.7-72.4µg/kg Mean: 16.0µg/kg Range: 1.0-51.0µg/kg	
<b>Maize</b>	Nigeria	T-2	36%	Mean:13µg/kg Range:0-29µg/kg	Afolabi et al., 2013
<b>Rice</b>	Niger State.	DON	23.8%	Mean: 18.9µg/kg Range:0-112µg/kg	Makun et al., 2011
		T-2	4.76%	ND	
<b>Processed Cocoa samples</b>	South western Nigeria.	DON	59%	Mean:3.3µg/kg Range:0.2-8.5µg/kg	Egbuta et al., 2013
<b>Cocoa-Based Powder Beverage</b>	South western Nigeria.	DON	32%	Mean:2.2µg/kg Range:0.1-7.6µg/kg	Egbuta et al., 2013
<b>Animal feed</b>	Nigeria	DON	56%	Mean: 181 µg/kg Range: 50- 451 µg/kg	Rodrigues et al., 2011
		NIV	2%	Mean: 4 µg/kg Range: 50-186 µg/kg	
		AcDON	4%	Mean: 9 µg/kg Range: 50-239 µg/kg	

ND – Not Determined

mutagenic (Qi *et al.*, 2014). Presently, there is no conclusive evidence of its carcinogenicity in human beings, but there is sufficient evidence in experimental animals. In view of this, the International Agency for Research on Cancer (IARC) has classified OTA in Group 2B as a possible carcinogen to humans (IARC, 1993).

OTA is found as natural contaminants of many foodstuffs including cereals, dried fruits, cocoa and cocoa products, cassava flour, grains, wine, poultry eggs, and milk ( Bankole and Adebajo, 2003b; Belli *et al.*, 2004; Gollucke *et al.*, 2004; Dongo *et al.*, 2008). These toxins are generally associated with grains stored in temperate climates of Europe and North America and it is not considered a major problem under Nigeria's tropical climate, although it has been reported in some food products (Adebajo, 1993; Steyn, 1995; Bankole and Adebajo, 2003b). The origin of OTA in warm temperate and tropical zones is commonly associated with *A. ochraceus* and the black Aspergilli (Accensi *et al.*, 2004). OTA production by *A. ochraceus* is usually at a temperature ranging from 12°C to 37°C with an optimum of 31°C and water activity level of 0.80 (Adebajo *et al.*, 1994).

Ochratoxin A is fairly stable to heat; in cereal products, up to 35% of the toxin survives autoclaving for up to 3 hours (IARC, 1976), hence OTA is not easily destroyed by common food preparation methods (Boudra *et al.*, 1995). The European Food Safety Authority (EFSA) has, on a request from European Commission (EC) derived a tolerable weekly intake (TWI) of 120ng/kg body weight (EFSA, 2006).

High frequency of OTA (66.7%) was reported in rice from Niger State, with a mean concentration of 141.9µg/kg (Makun *et al.*, 2011). The occurrence of OTA in the rice samples exceeded the maximum acceptable levels (CAST, 2003; EC 2006) in cereals for human consumption. Co-occurrence of OTA with AFB<sub>1</sub> has been reported to increase the mutagenicity of the latter (Sedmikova, *et al.*, 2001). Dried mushroom (Ezekiel *et al.*, 2013) and yam flour (Somorin *et al.*, 2012) from Nigeria were screened for the presence of Ochratoxin A, but they were not detected in any of the samples. This shows that OTA is not a likely contaminant of these food products. It has been observed that whenever OTA is detected in high levels in human breast milk, AFB<sub>1</sub> is absent or present at very low levels and vice versa. This relationship has also been reported in peanut cakes (Banu and Muthumary, 2008). These suggest some kind of competition between the toxins at either the production level in the substrate or their rate of absorption in the human gastrointestinal tract (Zain, 2011).




Figure 5. Chemical Structure of Ochratoxin A (Zain, 2011)

### 3.0 Intervention Strategies

It is important to know that complete elimination of mycotoxin-contaminated products/commodities is not feasible. In view of this, good agricultural practices (GAP) represents a primary line of defense against contamination of cereals with mycotoxins, followed by implementation of good manufacturing practices (GMP) during the handling, storage, processing, and distribution of cereals for human food and animal feed (Codex Alimentarius Commission, 2003; Atanda *et al.*, 2013). In addition, employing food safety practices like Hazard Analysis and Critical Control Point (HACCP) system can be useful in preventing and reducing mycotoxin contamination of agricultural produce (Adegoke and Letuma, 2013). Many intervention strategies to the problem of mycotoxins abound, out of which some of them itemized below can be found suitable to Nigeria's situation.

#### 3.1 Good Agricultural Practice

##### 3.1.1 Crop Rotation

A crop rotation routine should be developed and sustained to ensure that the same crop is not planted in the same field for two consecutive years. Crops that have been identified to be susceptible to toxigenic fungi should not be used in rotation with each other. For example, wheat and maize have been identified to be susceptible to toxigenic *Fusarium* spp, and should not be used in rotation with each other (CAC, 2003; Munkvold, 2003).

Table 5. Incidence of Ochratoxin in Nigerian foods

Commodities	Region	Type	Incidence Rate	Range µg/kg	Mean (µg/kg)	References
Tiger nuts	Nigeria	OTA	44.4%	10-80		Adebajo, 1993
Rice	Niger State, Nigeria	OTA	66.7%	0-341.3	141.7	Makun et al., 2011.
Cocoa seeds extract	South western Nigeria	OTA	88.2%	0.7-525.4	88.8	Egbuta et al., 2013.
Cocoa based powder beverage	South western Nigeria	OTA	56%	0.3-884.8	80.8	Egbuta et al., 2013.
Cocoa beans	Cross River, Edo and Ondo States	OTA	90%	1.0-277.5	37.7	Dongo et al., 2008
Cocoa beans	South	OTA	34.4%	0.05-	0.31	Aroyeun et al.,

	western Nigeria			2.70ng/ml		2007.
<b>Peanut cake</b>	Five Nigerian states	OTA	13.8%	-	4.0	Ezekiel et al., 2012.
<b>'Tinko'</b>	Ibadan, Nigeria	OTA	13.3%	0-2.13	ND	Oladejo and Adebayo (2011).

**Table 5. Incidence of Ochratoxin in Nigerian foods contd.**

Commodities	Region	Type	Incidence Rate	Range µg/kg	Mean (µg/kg)	References
<b>White Kolanut</b>	Ibadan, Oyo State	OTA	96%	13.9-65.3	30.2	Dongo et al., 2007.
<b>Red Kolanut</b>	Ibadan, Oyo State	OTA	100%	0.8–19.1	11.3	Dongo et al., 2007.
<b>Poultry feeds</b>	South western, Nigeria	OTA	98%	2.0-14.2	11.22	Ezekiel et al., 2011.
<b>Maize</b>	Minna, Niger State	OTA	98.2%	0-139.2	-	Makun et al., 2013.
<b>Millet</b>				10.20- 46.57		
<b>Sorghum</b>				0-29.50		
<b>Sesame</b>				1.90-15.66		
<b>Acha</b>				1.38-23.90		
<b>Garri</b>				3.28-22.75		

ND – Not Determined

### 3.1.2 Sanitation

Basic sanitation measures such as removal and destruction of debris from previous harvest would help in minimizing infection and infestation of produce in the field. Cleaning of the stores before loading new produce has been shown to be correlated with reduced aflatoxin levels (Hell *et al.*, 2000).

### 3.1.3 Insect Management

The level of insect damage influences the extent of mycotoxins contamination (Bankole and Adebajo, 2003b). Insect damage of maize has been found to be good predictor of *Fusarium* mycotoxins contamination (Avantaggio *et al.*, 2002). Insects carry spores of mycotoxigenic fungi from plant surfaces to the interior of the stalk, kernels or create infection wounds through their feeding habits, and promote fungal growth (Munkvold, 2003; Ariyo *et al.*, 2013). Therefore, proper management of insect pests through any appropriate control strategy would reduce mycotoxins contamination.

### 3.1.4 Proper Irrigation and Fertilization

Proper irrigation and fertilization should be carried out to reduce plants' stress which is sometimes responsible for mycotoxin development. The need to apply fertilizer and other soil conditioners to assure adequate soil pH and plant nutrition is critical (CAC RC, 2003; Munkvold, 2003).

### 3.1.5 Early harvesting

Early harvesting reduces fungal infection of crops in the field, as well as contamination of the harvested produce. In Nigeria, unpredictable weather, lack of storage space, labour constraint, need for cash, threat of thieves, rodents and other animals compel farmers to harvest at inappropriate times (Amyot, 1983). Rachaputi *et al.* (2002) reported that early harvesting and threshing of groundnuts resulted in lower aflatoxin levels and higher gross returns of 27% than in delayed harvesting.

### 3.1.6 Proper drying

Rapid drying of agricultural products to low moisture level is critical as it creates less favourable conditions for fungal growth and proliferation, insect infestation and helps produce keep longer (Lanyasunya *et al.*, 2005). In order to reduce or prevent the production of most mycotoxins, drying should take place soon after harvest, and as rapidly as possible. The critical water content for safe storage corresponds to water activity ( $a_w$ ) of about 0.7. Maintenance of feeds below 0.7  $a_w$  is an effective technique used throughout the world for controlling fungal spoilage and mycotoxin production in foods (Ariyo *et al.*, 2013). A community-based intervention trial in Guinea, West Africa focused on thorough drying and proper storage of groundnuts in subsistence farm villages and achieved a 60% reduction in mean aflatoxin levels in intervention villages (Turner *et al.*, 2005). During storage, transportation and marketing, maintenance of low moisture levels should be maintained by avoiding leaking roofs and condensation arising from inadequate ventilation (Bankole and Adebajo, 2003b).

### 3.1.7 Physical Separation

Physical treatments implemented through sorting, winnowing, washing, crushing and de-hulling are effective in reducing mycotoxin concentration according to Fandohan *et al.* (2005).

## 3.2 Biological Control

Biological control of mycotoxins may be a product of many factors which include competition for space and nutrients, production of anti-mycotoxigenic metabolites by co-existing microorganisms, changes in pH of the substratum, or a combination of these factors (Bianchini and Bullerman, 2010). In the presence of cell free supernatant of *Bacillus pumilus*, Munibabazi and Bullerman (1998) reported that 98% inhibition in aflatoxin production due to *A. parasiticus* and 53% inhibition of mould growth.

One of the biocontrol management strategies that had been developed is that which make use of competitive exclusion mechanism. The principle is utilized by the field application of atoxigenic strains that outcompete the toxigenic ones upon introduction, thereby shifting the fungal population from toxigenic to atoxigenic. Dorner and Cole (2002) reported a field application of non-toxigenic strains of *A. flavus* and *A. parasiticus* that reduced post-harvest aflatoxin contamination by 95.9%. This control method has also been successfully used in the USA and Nigeria as claimed by the authors (Atanda *et al.*, 2013, Bandyopadhyay and Cotty, 2013).

In Nigeria, the International Institute of Tropical Agriculture (IITA) has pioneered this technique with the production of Aflasafe™. There are single strain and multistrain kinds of this product, with the multistrain Aflasafe™ having the advantage of displaying both immediate and long term efficacy in various environments (Probst *et al.*, 2011). The multistrain Aflasafe™ contains a mixture of four native atoxigenic strains specifically targeted for a particular country or agroecosystem. The authors reported 50-99% reduction in aflatoxin contamination in maize field trials (Bandyopadhyay and Cotty, 2013).

## 3.3 Chemical Control

The application of fungicides and insecticides controls transmission of spores of mycotoxigenic fungi, fungal growth and minimize insect infestation of crops (Munkvold, 2003). However, poor education background of farmers often leads to misuse of pesticides. Hundreds of people have died in Nigeria as a result of consumption of cowpea treated with inappropriate pesticides (Bankole and Adebajo, 2003b). In 2008, 120 students of Government Girls' Secondary School, Dome, Gombe State were rushed to the hospital after consuming a meal of beans that was suspected to have been preserved with poisonous chemicals, 10 of the students were reported to be in critical condition (Gwary *et al.*, 2012). Analysis of the cooked and uncooked samples of the beans revealed that they contained high levels of Lindane, a banned pesticide (NAFDAC, 2004). In 2011, it was reported that 112 people were hospitalized in Calabar and 2 children confirmed dead, while 20 people were hospitalized in Oshogbo and 10 people died after eating a bean delicacy (Gbadeyan, 2010). Analyses revealed that the beans became poisonous because more pesticide was added than normal. In view of this growing

concern for food safety and potential health and environmental hazards associated with these chemicals, their use is being discouraged (Zain, 2011). Adejumo (2012) tested five botanicals against the maize mycotoxigenic fungus, *Fusarium verticillioides* and reported that *Piper guineense*, *Garcinia kola* and *Aframomum melegueta* had the potentials to be successfully used as sustainable alternative biopesticides, as a result of their cheap price, effectiveness, availability and environmental-friendliness.

### 3.4 Breeding for Resistance

This is one of the most promising intervention strategies to mycotoxin menace in Nigeria. Munkvold (2003) reported that sources of resistance to *A. flavus* and *Fusarium* spp., particularly *F. verticillioides* had been identified and had already been incorporated into public and private breeding programs. Genetic resistance is another approach. The genes for resistance to *Aspergillus* and *Fusarium* have been identified in cereals and peanuts, with prototypes of these crops containing genes encoding fungal growth inhibitors for reducing fungal infection. Scientists at United States Department of Agriculture have identified two maize lines that are resistant to *A. flavus* and *F. verticillioides* (Hamiton, 2000).

### 3.5 Detoxification and Decontamination

Detoxification involves the systemic degradation of mycotoxins into less toxic products. It was reported by Duvick *et al.* (2006) that the fungus *Expoliata spinnifera* was able to grow on fumonisin B<sub>1</sub> as a sole of carbon source (The hydrolysis of fumonisin B<sub>1</sub> yields free tricarboxylic acid and aminopental, the intermediate aminopental undergo oxidative deamination). In similar circumstances, *Agrobacterium-Rhizobium* strain E3-39 isolated from soil samples effectively converted deoxynivalenol into 3- keto-deoxynivalenol (which is less toxic) under aerobic conditions (Shima *et al.*, 1997). Zhou *et al.* (2008) isolated a bacterium strain Barpee from soil samples enhanced with *F. graminearum* infected maize. The strain showed strong DON transforming activity, transforming DON into two major products: a stereo-isomer of DON and 3-keto-DON, under aerobic conditions.

Another detoxification technique that is currently under investigation is the use of enterosorption, which is based on the introduction of harmless phyllosilicate clay (Hydrated Sodium Calcium Alumino Silicates (HASCAS)), which is widely used as an anticaking agent in animal feed, into animal diets. (Phillips, 1997). HSCAS is capable of tightly and selectively absorbing aflatoxins *in vitro* and *in vivo* (Bankole and Adebajo, 2003). There are different adsorption agents and their efficacies in preventing mycotoxicosis vary. In enterosorption, there is a risk that non-specific adsorption agents may prevent uptake of micronutrients from the food (Zain, 2011).

Decontamination as an intervention strategy seeks to reduce or eliminate contamination by mycotoxigenic fungi. The essential oils and aqueous extracts of *Aframomum danielli* have been reported to reduce ochratoxin A by between 64% and 95% in spiked cocoa powder. Although, ochratoxin molecule is stable, the authors stated that 40 to 90% of OTA got destroyed during roasting of coffee beans (Aroyeun and Adegoke, 2007).

Irradiation is also a useful tool in inactivating some mycotoxins. Herzallah *et al.* (2008) showed that exposure to sunlight (solar radiation),  $\gamma$ -radiation (<sup>60</sup>Co), and microwave heating were efficient in decontaminating aflatoxin residues in poultry feeds. The effect of irradiation depends on the fungus, application dose, moisture content, composition of food and storage conditions (Kabak and Var, 2005; Aziz *et al.*, 2006). There is a theoretical risk, however, of mycotoxins increasing after irradiation. One theory is that the higher radio-resistance of moulds and yeasts compared with bacteria results in a loss of competitive inhibition of mould and yeast growth. Any mould surviving under treatment with irradiation may be expected to grow more rapidly in the absence of competitors and eventually dominate the mycoflora (Temur and Tiryaki, 2013), although their potential to produce mycotoxins is the major concern.

### 3.6 Modification of Diet

One of the factors that impair food safety in Africa is the reliance on a single crop, especially cereals such as maize (Adebayo *et al.*, 2010). With high daily intake of the cereal, only moderate mycotoxin contamination levels are required to exceed recommended tolerable intake, thereby predisposing people to mycotoxicosis (Bryden, 2007). Bandyopadhyay *et al.* (2007) reported that Nigerian maize was more heavily colonised by aflatoxigenic strains of *Aspergillus* spp. than sorghum and millet, with corresponding higher aflatoxin levels. Consumption of risky foods such as maize was then discouraged by the authors in favour of less risky ones such as sorghum and millet.

### 3.7 Education and Awareness Creation

Due to the fact that the risks posed by mycotoxins are largely known to the scientific researchers and not to the general populace (Bankole and Adebajo, 2003b), there is the need to properly inform people, especially farmers, on the economic importance of mycotoxins and the dangers they pose to human health. Due to limited time and resources, the entire populace cannot be realistically reached; people who are trained and given



orientation about mycotoxins should therefore be advised to train others by sharing their knowledge with them. This effectively creates an advantageous ripple effect. The national bodies and societies should sensitize people on mycotoxins. For instance, the National Agency for Food and Drug Administration and Control (NAFDAC) could partner with a society like Mycotoxicology Society of Nigeria (MYCOTOXSON) on ways to educate the general public on mycotoxins. This can be carried out on media outlets such as television and radio stations, newspapers and on internet blogs and social media. WHO (2006) reported that field projects, strengthening surveillance and awareness raising and educating consumers on matters related to mycotoxins were part of WHO's agenda for Africa.

### 3.8 Legislation

In controlling mycotoxin contamination of food and feeds, certain legislation needs to be put in place. In most countries, regulations are established to control the contaminants in foodstuffs to protect human health. These regulations may include specific maximum limits for several contaminants for different foods and a reference to the sampling methods and performance criteria of analysis to be used (Kubo, 2012). In the Codex Alimentarius Standard, aflatoxin limits are 0.05 $\mu\text{g/l}$  for milk and milk products, 4 to 5 $\mu\text{g/kg}$  for beans, 10 $\mu\text{g/kg}$  in nuts such as peanuts and almonds, and 20 $\mu\text{g/kg}$  for cereals. The acceptable limit for fumonisins is <1000  $\mu\text{g/kg}$  (CAST, 2003). These limits are adopted in Nigeria. The EU set a limit of 0.5 $\mu\text{g/kg}$  OTA in raw cereals, with a tolerable weekly intake (TWI) of 0.12 $\mu\text{g/kg}$ , while maximum DON limits are set between 200 $\mu\text{g/kg}$  for processed cereal-based foods and baby foods for infants and young children, and 1750 $\mu\text{g/kg}$  for unprocessed wheat, oats and maize (EC, 2006). However, there is presently no maximum tolerable limit for trichothecenes in Nigeria. The responsible authority for these regulations in Nigeria is the National Agency for Food and Drug Administration and Control (NAFDAC). Official regulations concerning maximum acceptable limits for mycotoxins in Nigerian foods are urgently needed.

In addition to the intervention strategies discussed above, Atanda *et al.* (2013) suggested the following methods for the control of the hazards posed by mycotoxins in Nigeria: This include collection of a database of predominant fungi and mycotoxin in Nigeria, establishment of a mycotoxin occurrence map to know the areas prone to mycotoxin contamination and the establishment of a permanent culture collection media.

### 4.0 CONCLUSION

A considerable number of studies have been done on the problems of mycotoxins in Nigeria and the ways by which they can be solved or managed. Some techniques for management of mycotoxin contamination may be in usual practice in developed countries, in order to reduce or eliminate contamination of agricultural products meant for export and local use, and to protect Nigerians from the harmful effects that can result from consuming mycotoxin-contaminated food, we can adopt some of the methods described in this review. It is very important to properly educate Nigerian farmers on the implementation of good methods of cultivation and storage of agricultural products that would not promote fungal contamination. This review has shown that there are food products in Nigeria with mycotoxin levels above the maximum acceptable limits discussed earlier. Unfortunately, majority of the populace do not know the inherent dangers of consuming mouldy products (with possible contamination by mycotoxigenic fungi) because of their lack of knowledge on mycotoxin. In view of this, the general public needs to be educated on the economic and health hazards posed by mycotoxins. Control measures such as physical selection, proper washing and cooking practices of food commodities should be emphasized. There is a large pool of information on aflatoxins and ochratoxin A contamination in food commodities, but few on other major mycotoxins (fumonisins, trichothecenes and zearalenone) occurring naturally in food commodities. It is therefore advised that wider range of agricultural produce and food commodities be screened for all major mycotoxins to improve food safety in Nigeria.

### References

- Abiala M.A., Ezekiel, C.N., Chukwura, N.I. and Odebode, A.C. (2011). Toxigenic *Aspergillus* section *flavi* and aflatoxins in dried yam chips in Oyo state, Nigeria. *Acad. Arena* 3(5): 42-49.
- Accensi, F., Abarca, M.L., and Cabañes, F.J. (2004). Occurrence of *Aspergillus* species in mixed feeds and component rawmaterials and their ability to produce ochratoxin A," *Food Microbiol.* 21(5):623-627.
- Adebajo, L.O. (1993). Survey of aflatoxins and ochratoxin A in stored tubers of *Cyperus esculentus*. *Mycopathologia* 124: 41-46.
- Adebajo, L.O., Idowu, A.A. and Adesanya, O.O. (1994). Mycoflora and mycotoxins production in Nigerian corn and corn based snacks. *Mycopathologia* 126:183-192.

- Adebayo, G.B., Otunola, G.A. and Ajao, T.A. (2010). Physico-chemical, microbiological and sensory characteristics of kunu prepared from millet, maize and guinea corn and stored at selected temperature. *Adv. J. Food. Sci. Tech.* 2(1): 41-46.
- Adegoke, G.O. and Letuma, P. (2013). Strategies for the Prevention and Reduction of Mycotoxins in Developing Countries In: H.A. Makun (Ed) *Mycotoxin and Food Safety in Developing Countries*. InTech, Rijeka, Croatia, pp123-136.
- Adejumo, O., Atanda, O., Raiola, A., Bandyopadhyay, R., Somorin, Y. and Ritieni, A. (2013). Correlation between Aflatoxin M1 Content of Breast Milk, Dietary Exposure to Aflatoxin B1 and Socioeconomic Status of Lactating Mothers in Ogun State, Nigeria. *Food Chem. Toxicol.* 56:171-177.
- Adejumo, T.O. (2012). Evaluation of botanicals as biopesticides on the growth of *Fusarium verticillioides* causing rot diseases and fumonisin production of maize. *J. Microbiol. Antimicrob.* 4(1):23-31.
- Adejumo, T.O., Hettwer, U. and Karlovsky, P. (2007a). Occurrence of *Fusarium* species and trichothecenes in Nigerian maize. *Int. J. Food Microbiol.* 116: 350-357.
- Adejumo, T.O., Hettwer, U. and Karlovsky, P. (2007b). Survey of maize from south-western Nigeria for zearalenone,  $\alpha$ - and  $\beta$ -zearalenols, fumonisin B1 and enniatins produced by *Fusarium* species. *Food Add. Cont.* 24: 993-1000.
- Adejumo, T.O., Hettwer, U., Nutz, S. and Karlovsky, P. (2009). Real-time PCR and agar plating method to predict *Fusarium verticillioides* and fumonisin B1 content in Nigerian maize. *J. Plant Prot. Res.* 49(4):399-404.
- Afolabi, C.G., Ekpo, E. J. A. and Bandyopadhyay, R. (2013). Maize contamination by zearalenone and T-2 toxin and human exposure in Nigeria *Mycotoxins* 63(2):143-149
- Akano, D. A. and Atanda, O. O. (1990). The present Level of Aflatoxin in Nigeria Groundnut Cake ('Kulikuli'). *Letters in App. Microbiol.* 10: 187-18.
- Amadi, J. E. and Adeniyi, D. O. (2009). Mycotoxin production by fungi isolated from stored grains. *A. J. Biotechnol.* 8(7):1219-1221.
- Amyot, J. (1983). Social and Economic Aspects of Dryer Use for Paddy and Other Agricultural Produce in Thailand. Chulalongkorn University Social Research Institute and International Development Research Center.
- Ariyo, A.L., Peter, A.O., Abdulazeez, S.N. and Olatunde, O.H. (2013). Avian Mycotoxicosis in Developing Countries. In: H.A. Makun (Ed) *Mycotoxin and Food Safety in Developing Countries*. InTech, Rijeka, Croatia, pp 93-119.
- Aroyeun, S.O. and Adegoke, G.O. (2007). Reduction of ochratoxin A (OTA) in spiked cocoa powder and beverage using aqueous extracts and essential oils of *Aframomum danielli*. *Afr. J. Biotechnol.* 6:612-616.
- Aroyeun, S.O., Adegoke, G.O., Varga, J., Kocsubé, S., Pál, K. and Vágvolgyi, C. (2007). Effect of fermentation and storage on mycotoxigenic fungi, ochratoxin A and aflatoxin B1 in cocoa beans from south western Nigeria. *Malaysian Cocoa J.* 3: 35-46.
- Atanda, O., Makun, H.A., Ogara, I.M., Edema, M., Idahor, K.O., Eshiett, M.E. and Oluwabamiwo, B.F. (2013). Fungal and mycotoxin contamination of Nigerian foods and feeds Countries. In: H.A. Makun (Ed) *Mycotoxin and Food Safety in Developing Countries*. InTech, Rijeka, Croatia pp3-38.
- Atehnkeng, J., Ojiambo, P.S., Donner, M., Ikotun, T., Richard, A. Sikora, R.A., Cotty, P.J. and Bandyopadhyay, R. (2008). Distribution and toxigenicity of *Aspergillus* species isolated from maize kernels from three agro-ecological zones in Nigeria. *Int. J. Food Microbiol.* 122:74-84.
- Avantaggio, G., Quaranta F., Desidero, E. and Visconti, A. (2002). Fumonisin contamination of maize hybrids visibly damaged by *Sesamia*. *J. Sci. Food Agric.* 83: 13-18.
- Ayejuyo, O.O., Williams, A.B. and Imafidon, T.F. (2008). Ochratoxin A burdens in rice from Lagos markets, Nigeria. *J. Environ. Sci. Technol.* 1: 80-84.
- Aziz, N.H., Souzan, R.M. and Azza, A.S. (2006). Effect of gamma irradiation on the occurrence of pathogenic microorganism and nutritive value of four principal cereal grains. *Appl. Radiat. Isot.* 64:1555-1562.
- Bankole, S.A. and Adebajo, A. (2003a). Aflatoxin contamination of dried yam chips marketed in Nigeria. *Trop. Sci.* 43: 3-4.
- Bankole, S.A. and Adebajo, A. (2003b). Mycotoxins in food in West Africa: current situation and possibilities of controlling it. *A. J. Biotechnol.* 2 (9):254-263.
- Bankole, S.A. and Eseigbe, D.A. (1996). Occurrence of mycoflora and aflatoxins in marketed tiger nut. *Crop Res.* 11: 219-223.
- Bankole, S.A. and Mabekoje, O.O. (2004). Occurrence of aflatoxins and fumonisins in preharvest maize from south-western Nigeria. *Food Addit Contam.* 21(3):251-255.
- Bankole, S.A., Mabekoje, O.O. and Enikuomhin, O.A. (2003). *Fusarium moniliforme* and fumonisin B1 in stored maize from Ogun State, Nigeria. *Trop. Sci.* 43: 76-79.

- Bankole, S.A., Schollenberger, M. and Drochner, W. (2010). Survey of ergosterol, zearalenone and trichothecenes contamination in maize from Nigeria. *J. Food Comp. Anal.* 23(8):837-842.
- Banu, N. and Muthumary, J. (2008). Screening of commercial feed samples for the presence of multimycotoxins. *I. J. Sci. Technol.* 1(4), 1e4.
- Belli, N., Marin, S., Dauaigües, A., Ramos, A.J. and Sanchis, V. (2004). Ochratoxin A in wines, musts and gape juices from Spain. *J. Sci. Food Agric.* 84: 591-594.
- Bezuidenhout, S. C., Gelderblom, W.C.A., Gorst-Allman, C.P., Horak, R.M., Marasas, W.F.O., Spiteller, G. and Vlegaar, R. (1988). Structure elucidation of the fumonisins, mycotoxins from *Fusarium moniliforme*. *J. Chem. Soc. Chem. Commun.* 1988:743-745.
- Bhat, R., Rai, R. V., and Karim, A. A. (2010). Mycotoxins in Food and Feed: Present Status and Future Concerns. *Comp. Rev. Food Sci. Food Safety.* 9-57.
- Bianchini, A. and Bullerman, L.B. (2010). Biological Control of Molds and Mycotoxins in Foods In Mycotoxin Prevention and Control in Agriculture; Eds Appell, M., et al.; ACS Symposium Series 1031; American Chemical Society: Washington, DC.
- Blount, W. P. (1961). Turkey 'X' disease. *J. Brit. Turkey Federation* 9:52-61.
- Boudra, H., Le Bars, P. and Le Bars, J. (1995). Thermostability of ochratoxin A in wheat under two moisture conditions. *Appl. Environ. Microbiol.* 61:1156-1158.
- Bryden, W.L. (2007). Mycotoxins in the food chain: human health implications. *Asian Pac. J. Clin. Nutr.* 16(1): 93-101.
- Centers for Disease Control and Prevention (CDC). (2004). Outbreak of aflatoxin poisoning—eastern and central provinces, Kenya, January–July, 2004. *MMWR Morb Mortal Wkly Rep* 53:790–792. Available: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5334a4.htm> [Accessed March 17, 2014].
- Choudhary, P.L., Sharma, R.S., Borkhataria, V.N. and Desai, M.C. (1998). Effect of feeding aflatoxin B1 on feed consumption through naturally contaminated feeds. *I. J. Anim. Sci.* 68:400-401.
- Codex Alimentarius Commission. (2003). Code of practice for the prevention and reduction of mycotoxin contamination in cereals, including annexes on ochratoxin a, zearalenone, fumonisins and trichothecenes. Available online at [http://www.fao.org/fileadmin/user\\_upload/agns/pdf/CXS\\_051e.pdf](http://www.fao.org/fileadmin/user_upload/agns/pdf/CXS_051e.pdf). [Accessed August 10, 2014].
- Council for Agricultural Science and Technology (CAST) (2003). Mycotoxins: risks in plant, animal, and human systems. *Ames, IA*. Task Force Report. 139.
- Davis, N. D. and Diener, U. L. (1983). Some characteristics of toxigenic and nontoxigenic isolates of *Aspergillus flavus* and *A. parasiticus*. In: Diener, U.L., Asquith, R.I., Dickens, J.W., eds. Aflatoxin and *A. flavus* in corn. *Southern Coop Series Bull.* 279:1-5.
- Dongo, L. N., Manjula, K. and Orisajo, S. B. (2007). Occurrence of ochratoxin A in Nigerian kola nuts. *A. Crop Sci. Conference Proceedings* 8: 2133-2135.
- Dongo, L., Kumar, M., Ojiambo, P.S. and Bandyopadhyay, R. (2008). Occurrence of ochratoxin A in Nigerian ready for sale cocoa beans. *Agric. J.* 3:4-9.
- Dorner, J.W. and Cole, R.J. (2002). Effect of application of nontoxigenic strain of *Aspergillus flavus* and *A. parasiticus* on subsequent aflatoxin contamination of peanuts in storage. *J. Stored Prod. Res.* 38: 329-339.
- Dors, G. C., Caldas, S. S., Feddern, V., Bemvenuti, R. H., Hackbart, H. C. S., Souza, M.M., Oliveira, M. S., Garda-Bufferon, J., Primel, E. G. and Badiale-Furlong, E. (2011). Aflatoxins: Contamination, Analysis and Control. In: Guevara-González, R.G. (Ed), *Aflatoxins- Biochemistry and Molecular Biology*, InTech, pp415-438.
- Duvick J., Rood, T.A., Maddox, J.R. and Gilliam, J. (2006). Advances in ochratoxin A biosynthesis. Book of Abstracts, International Conference on “Advances on genomics, biodiversity and rapid systems for detection of toxigenic fungi and mycotoxins” September 26-29, 2006. Monopoli (Bari), Italy Pp.39 Retrieved from <http://www.Ispa.onr.n/mycoglobe-2006> [Accessed August 10, 2014].
- Egbuta, M.A., Chilaka, C.A., Phoku, J.Z., Mwanza, M. and Dutton, M.F. (2013). Co-contamination of Nigerian cocoa and cocoa-based powder beverages destined for human consumption by mycotoxins. *Ethno Med.* 7(3): 187-194.
- European Commission (2006). Commission Regulation (EC) No 1881/2006 of 19 December 2006 Setting maximum levels for certain contaminants in foodstuffs (text with EEA relevance). *Off. J. Eur. Union* 364:5-24.
- European Food Safety Authority (EFSA) (2006). Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to ochratoxin A in food. *The EFSA J.* 365:1-56.
- European Food Safety Authority (EFSA) (2007). Opinion of the scientific panel on contaminants in the food chain on a request from the commission related to the potential increase of consumer health risk by a

- possible increase of the existing maximum levels for aflatoxins in almonds, hazelnuts and pistachios and derived products. *The EFSA J.* 446:1-127.
- Ezekiel, C.N., Odebode, A.C., Fapohunda, S.O., Tayo, G.O., Olawuyi, O.J., Olaoye, O.B., Olarinmoye, A.O. and Adeyemi, O.O. (2011). Toxigenic potential of co-occurring aflatoxin and ochratoxin a detected in poultry feed on *Clarias gariepinus* larvae. *Nature and Sci.* 9(5): 186-192.
- Ezekiel, C. N., Sulyok, M., Warth, B., Odebode, A.C. and Krska, R. (2012). Natural occurrence of Mycotoxins in peanut cake from Nigeria. *Food control.* 27:338-342.
- Ezekiel, C.N., Sulyok, M., Frisvad, J.C., Somorin, Y.M., Warth, B., Houbraken, J., Samson, Krska, R. and Odebode, A.C. (2013). Fungal and mycotoxin assessment of dried edible mushroom in Nigeria. *Int. J. Food Microbiol.* 162:231–236.
- Fernandez-Cruz, M.L., Mansilla, M.L. and Tadeo, J.L. (2010). Mycotoxins in fruits and their processed products: Analysis, occurrence and health implications. *J. Advanced Res.* 1:113–122.
- Filazi, A. and Sireli, U.T. (2013). Occurrence of Aflatoxins in Food. In: Mehdi Razzaghi-Abyaneh (Ed) *Aflatoxins - Recent Advances and Future Prospects*, InTech, pp143-170.
- Food and Agriculture Organization (FAO) (2004). Worldwide regulations for mycotoxins in food and feed in 2003. Rome: FAO. Food and Nutrition Papers No. 81.
- Frisvad, J.C., Frank, J.M., Houbraken, J.A., Kuijpers, A.F. and Samson, R.A. (2004). New ochratoxin A producing species of *Aspergillus* section *Circumdati*. *Stud. Mycol.* 50:23-43.
- Gbadayan, R.A. (2010). De-marketing of harmful products in Nigeria. *Pakistan J. Social Sci.* 7(6):449-455.
- Gelderblom, W.C., Jaskiewicz, K., Marasas, W.F., Thiel, P.G., Horak, R.M., Vleggaar, R. and Kriek, N.P.J. (1988). Fumonisin - novel mycotoxins with cancer-promoting activity produced by *Fusarium moniliforme*. *Appl. Environ. Microbiol.* 54:1806-1811.
- Gottschalk, C., Barthel, J., Engelhardt, G., Bauer, J. and Meyer, K. (2009). Simultaneous determination of type A, B, and D trichothecenes and their occurrence in cereals and cereal products. *Food Add. Cont.* 26(9): 1273-1289.
- Gollucke, A.P.B., Taniwaki, M.H. and Tavares, D.Q. (2004). Survey on ochratoxin A in Brazilian green coffee destined for exports. *Cienc. Tecnol. Aliment. Campinas.* 24(4): 641-645.
- Grove, J.F. (2007). The trichothecenes and their biosynthesis. *Prog. Chem. Org. Nat. Prod.* 88:63–130.
- Gwary, O.M., Hati, S.S., Dimari, G.A. and Ogugbuaja, V.O. (2012). Pesticide residues in bean samples from Northeastern Nigeria. *ARPJ J. Sci. Technol.* 2(2): 79-84.
- Hamilton, D. (2000). Toxic Fungus Threatens Health of Consumers. [www.agnic.org/pmp/2000/ama0826.htm](http://www.agnic.org/pmp/2000/ama0826.htm)
- Hell, K., Cardwell, K.F., Setamou, M. and Poehling, H.M. (2000). The influence of storage practices on aflatoxin contamination in maize in four agroecological zones of Benin, West Africa. *J. Stored Prod. Res.* 36, 365–382.
- Hendrickse, R.G. (1983). Aflatoxin and kwashiorkor: epidemiology and clinical studies in Sudanese children and findings in autopsy liver sample from Nigeria and South Africa. *Bull. Soc. Pathol. Exotique* 76:559-566.
- Herzallah, S., Alshawabkeh, K. and Al Fataftah, A.A. (2008). Aflatoxin decontamination of artificially contaminated feeds by sunlight,  $\gamma$ -Radiation, and microwave heating. *J. Appl. Poult. Res.* 17:515–521.
- Hosseini, S.S. and Bagheri, R. (2012). Some major mycotoxin and their mycotoxicoses in nuts and dried fruits. *Int. J. Agro. Plant Prod.* Vol., 3 (5):179-184.
- Hussein, H. S. and Brasel, J. M., (2001). Toxicity, metabolism and impact of mycotoxins on humans and animals. *Toxicology* 167: 101-134.
- Iloje, N.P. (2001). A new geography of Nigeria. New Revised Edition. Longman Nigeria PLC. 200 p.
- International Agency for Research on Cancer (IARC) (1976). Ochratoxin A. In *Some Naturally Occurring Substances*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 10. Lyon, France: International Agency for Research on Cancer. pp. 191-197.
- International Agency for Research on Cancer (IARC) (1993). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins. *IARC Monogr Eval Carcinog Risks Hum*, 56: 489–521.
- International Agency for Research on Cancer (IARC) (2002). Some traditional herbal medicines, some mycotoxins, naphthalene and styrene. *IARC Monogr Eval Carcinog Risks Hum*, 82: 1–556.
- International Food Policy Research Institute (IFPRI) (2013). Aflatoxins: Finding Solutions for Improved Food Safety. Laurian Unnevehr and Delia Grace (Eds).
- Jackson, L. S. and Al-Taher, F. (2008). Factors Affecting Mycotoxin Production in Fruits. In: Barkai-Golan R, Paster N. (Ed), *Mycotoxins in Fruits and Vegetables*, Academic Press is an imprint of Elsevier, 75-104.
- Joint Expert Committee on Food Additives (JECFA) (2001). WHO Food additives series 47: Safety Evaluation of certain mycotoxins in foods. Joint FAO/WHO expert committee on food additives. 701pp.



- Juan, C., Rittieni, A., Font, G. and Mañes, J. (2010). Determination of mycotoxins produced by *Fusarium*, *Aspergillus* and *Penicillium* fungi species in bread. Proceedings of the International Conference on Food Innovation, 25-29 October, 2010.
- Kabak, B. and Var, I. (2005). Control of irradiation effects of mould and mycotoxin development. *Food* 30(1):197-201.
- Kensler, T.W., Roebuck, B.D., Wogan, G.N. and Groopman, J.D. (2011). Aflatoxin: A 50-Year odyssey of mechanistic and translational toxicology. *Toxicological Sci.* 120(1): 28-48.
- Krska, R., Welzig, E. and Boudra, H. (2007). Analysis of *Fusarium* toxins in feed. *Anim. Feed Sci. Technol.* 137: 241-264.
- Kumar V., Basu, M.S. and Rajendran, T.P. (2008). Mycotoxin research and mycoflora in some commercially important agricultural commodities. *Crop Protection*. 27 (6):891-905.
- Lanyasunya, T.P., Wamae, L.W., Musa, H.H., Olowofeso, O. and Lokwaleput, I.K. (2005). The risk of mycotoxins contamination of dairy feed and milk on smallholder dairy farms in Kenya. *Pak. J. Nutr.* 4:162-169.
- Lewis, L., Onsongo, M., Njapau, H., Schurz-Rogers, H., Lubber, G., Kieszak, S., Nyamongo, J., Backer, L., Dahiye, A.M., Misore, A., Decoet, K. and Rubin, C. (2005). Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in Eastern and Central Kenya. *Environ Health Persp.* 113:1763-1767.
- Liddell, C.M. (2003). Systematics of *Fusarium* species and allies associated with fusarium head blight. In *Fusarium Head Blight of Wheat and Barley*; Leonard, K.J., Bushnell, W.R., Eds.; The American Phytopathological Society: St. Paul, Minnesota, USA, 2003; pp. 35-43.
- Makun, H.A., Adeniran, A.L., Mailafiya, S.C., Ayanda, I.S., Mudashiru, A.T., Ojukwu, U.J., Jagaba, A.S., Usman, Z. and Salihu, D.A. (2013). Natural occurrence of ochratoxin A in some marketed Nigerian foods. *Food Contr.* 31(2): 566-571.
- Makun, H. A., Anjorin, S. T., Moronfoye, B., Adejo, F. O., Afolabi, O. A., Fagbayibo, G., Balogun, B. O. and Surajudeen, A. A. (2010). Fungal and aflatoxin contamination of some human food commodities in Nigeria. *Afr. J. Food Sci.* 4: 127-135.
- Makun, H.A., Dutton, F.M., Njobeh, P.B., Mwanza, M. and Kabiru, A.Y. (2011). Natural multi-occurrence of mycotoxins in rice from Niger State, Nigeria. *Mycotoxin Res.* 27(2):97-104.
- Marasas, W.F.O., Riley, R.T., Hendricks, K.A., Stevens, V.L., Sadler, T.W., Gelineau-van Waes, J., Missmer, S.A., Cabrera, J., Torres, O., Gelderblom, W.C.A., Allegood, J., Martinez, C., Maddox, J., Miller, J.D., Starr, L., Sullards, M.C., Roman, A., Voss, K.A., Wang, E. and Merrill, A.H. (2004). Fumonisin disrupt sphingolipid metabolism, folate transport, and neural tube development in embryo culture and in vivo: a potential risk factor for human neural tube defects among populations consuming fumonisin contaminated maize. *J. Nutr.* 134, 711-716.
- Marin, S., Ramos, A.J., Cano-Sancho, G. and Sanchis, V. (2013). Mycotoxins: Occurrence, toxicology, and exposure assessment. *Food Chem. Toxicol.* 60:218-237.
- Mazaheri, M. (2009). Determination of aflatoxins in imported rice to Iran. *Food Chem. Toxicol.* 47: 2064-2066.
- McDonald, D. (1964). Progress reports on research into aflatoxin production in groundnuts in Northern Nigeria. *Samaru Miscellaneous Papers* 2: 13p.
- Missmer, S. A., Suarez, L., Felkner, M., Wang, E., Merrill, A. H. and Rothman, K. J. (2006). Exposure to fumonisins and the occurrence of neural tube defects along the Texas-Mexico border. *Environ. Health Persp.* 114(2):237-241.
- Munimbazi, C. and Bullerman, L.B. (1998). Inhibition of aflatoxin production of *Aspergillus parasiticus* NRRL 2999 by *Bacillus pumilus*. *Mycopathologia* 140:163-169.
- Munkvold, G.P. (2003). Cultural and genetic approaches to managing mycotoxins in maize. *Ann. Rev. Phytopathol.* 41, 99-116.
- Nagwa, J. A., Alarousy, R.M. and Zaki, M.S. (2013). Mycotoxicosis in animals and fish. *Life Sci. J.* 10(1).
- National Agency for Food and Drug Administration and Control (NAFDAC) (2004). Consumer Safety Bulletin: NAFDAC Regulated Products. 6(1): 9.
- Nierman, W. C., Cleveland, T. E., Payne, G. A., Keller, N. P., Campbell, B. C., Bennett, J. W., Guo, B., Yu, J. and Robens, J. F. (2008). Mycotoxin Production and Prevention of Aflatoxin Contamination in Food and Feed. In: Goldman GH, Osmani SA. *The Aspergilli: Genomics, Medical Aspects, Biotechnology, and Research Methods*. Boca Raton: CRC Press, 457-472.
- Odoemelam, S.A. and Osu, C.I. (2009). Aflatoxin B1 contamination of some edible grains marketed in Nigeria. *E-J. Chem.* 6(2): 308-314.
- Okeke, K. S., Abdullahi, I. O., Makun, H. A. and Mailafiya, S. C. (2012). A preliminary survey of aflatoxin M<sub>1</sub> in dairy cattle products in Bida, Niger State, Nigeria. *Afr. J. Food Sci. Technol.* 3(10):273-276.



- Oladejo, D.A. and Adebayo-Tayo, B.C. (2011). Moulds, proximate mineral composition and mycotoxin contamination of banda ('kundi'/ 'tinko') sold in Ibadan, Oyo State, Nigeria. *U. J. Technol.* 15(1): 32-40.
- Olayiwola, I. O., Oganah, B. C., Oguntona, C. R. B., Popoola, A. R., Sanni, S. A. and Wobo, S.O. (2013). Status of aflatoxin and anti-nutritional contents of standardized maize-based dishes/snacks consumed in Nigeria. *Dis. J. Agric. Food Sci.* 1(5):93-96.
- Olorunfemi, M. F., Oyeboji, A. O., Awoite T. M., Agboola, A. A., Oyelakin, M. O., Alimi, J. P., Ikotun, I. O., Olagbaju, R. A. and Oyedele, A. O. (2014). Quality assessment of palm oil on sale in major markets of Ibadan, Nigeria. *Int. J. Food Res.* 1:8-15.
- Omar, H.E. (2013). Mycotoxins-Induced Oxidative Stress and Disease I n Mycotoxin and Food Safety in Developing Countries. In: H.A. Makun (Ed) *Mycotoxin and Food Safety in Developing Countries*. InTech, Rijeka, Croatia, pp63-92.
- Onyemelukwe, G.C., Ogoina, D., Ibiyam, G.E. and Ogbadu, G.H. (2012). Aflatoxins in body fluids and food of Nigerian children with Protein- Energy Malnutrition (PEM). *A. J. Food, Agric., Nutri. Dev.* 12(5):6553-6566.
- Oyelami, O.A., Maxwell, S., Adelusola, K.A., Aladekoma, T.A. and Oyelese, A.O. (1998). Aflatoxins in autopsy kidney specimens from children in Nigeria. *J Toxicol Environ Health* 55(5): 317-323.
- Oyenuga, V.A. (1967). Agriculture in Nigeria. Food and Agriculture Organization of the United Nations. FAO, Rome, Italy. 308 pp.
- Peraica, M., Radic, B., Lucic, A. and Pavlovic, M. (1999). Toxic effects of mycotoxins in humans. *Bulletin of the World Health Organization*, 77 (9).
- Phillips, T.D. (1997). Detection and Decontamination of aflatoxincontaminated food products. In: J.H. Williams, D.G. Cummins, G. Hutto, and A. King (eds). *Impacts and Scientific Advances through Collaborative Research on Peanut CRSP*, Griffin, GA. Pp. 117-130.
- Probst, C., Bandyopadhyay, R., Price, L.E. and Cotty, P.J. (2011). "Identification of Atoxigenic *Aspergillus flavus* Isolates to Reduce Aflatoxin Contamination of Maize in Kenya." *Plant Disease* 95: 212–218.
- Probst, C., Njapau, H. and Cotty, P.J. (2007). Outbreak of an acute aflatoxicosis in Kenya in 2004: identification of the causal agent. *Appl. Environ. Microbiol.* 73, 2762–2764.
- Qi, X., Yang, X, Chen, S., He, X., Dweep, H., Guo, M., Cheng, W., Xu, W., Luo, Y., Gretz, N., Dai, Q. and Huang, K. (2014). Ochratoxin A induced early hepatotoxicity: new mechanistic insights from microRNA, mRNA and proteomic profiling studies. *Scientific Rep.* 4:5163.
- Rachaputi, N.R., Wright, G.C., Kroschi, S., 2002. Management practices to minimise pre-harvest aflatoxin contamination in Australian groundnuts. *Austr. J. Exp. Agric.* 42, 595–605.
- Rai, K.M., Bonde, S.R., Ingle, A.P. and Aniket K.G. (2012). Mycotoxin: rapid detection, differentiation and safety *J Pharm Educ Res* 3(1): 22-34.
- Reddy, L. and Bhoola, K. (2010). Ochratoxins - food contaminants: Impact on human health *Toxins* 2:771-779.
- Refai, M. K. (1988). Aflatoxins and aflatoxicosis. *J. Egypt. Vet. Med. Ass* 48:1-19.
- Richard, J.L. (2007). Some major mycotoxins and their mycotoxicoses- an overview. *Int. J. Food Microbiol.* 119: 3-10.
- Rodrigues, I., Handl, J. and Binder, E.M. (2011). Mycotoxin occurrence in commodities, feeds and feed ingredients sourced in the Middle East and Africa. *Food Add. Cont.* B 4(3): 168–179.
- Salifu, A. (1981). Mycotoxins in short season sorghums in northern Nigeria. *Samaru J. Agric Res.* 1: 83–88.
- Sedmikova, M., Reisnerora, H., Dufkova, Z., Burta, I. and Jilek, F. (2001). Potential hazard of simultaneous occurrence of aflatoxin B1 and ochratoxin A. *Vet. Med.* 46: 169-174.
- Seifert, K.A., Aoki, T., Baayen, R.P., Brayford, D., Burgess, L.W., Chulze, S., Gams, W., Geiser, D., de Gruyter, J., Leslie, J.F., Logrieco, A., Marasas, W.F.O., Nirenberg, H.I., O'Donnell, K., Rheeder, J.P., Samuels, G.J., Summerell, B.A., Thrane, U. and Waalwijk, C. (2003). The name *Fusarium moniliforme* should no longer be used. *Mycological Res.* 107, 643-644.
- Senyuva, H.Z., Gilbert, J. and Stroka, J. (2010). Determination of fumonisins B1 and B2 in corn by LC/MS with immunoaffinity column cleanup: interlaboratory study *J. AOAC Int.* 93(2):611-621.
- Sherif, O.S., Salama, E.S. and Abdel-Wahhab, M.A. (2009). Mycotoxins and child health: The need for health risk assessment. *Int. Hyg. Environ. Health* 212, 347–368.
- Shima, J. S., Takase, S., Takahashi, Y., Iwai, Y., Fujimoto, H., Yamazaki, M. and Ochi, K. (1997). Novel detoxification of the trichothecene mycotoxin deoxynivalenol by a soil bacterium isolated by enrichment culture. *Appl. Environ. Microbiol.* 63:3825-3830.
- Smith, E.E., Kubena, L.F., Braithwaite, R.B., Harvey, R.B., Phillips, T.D. and Reine, A.H. (1992). Toxicological evaluation of aflatoxin and cyclopiazonic acid in broiler chickens. *Poult. Sci.* 71, 1136–1144.
- Somorin, Y.M., Bertuzzi, T., Battilani, P. and Pietri, A. (2012). Aflatoxin and fumonisin contamination of yam flour from markets in Nigeria. *Food Control* 25:53-58.

- Splensley, P.C. (1963). Aflatoxin, the active principle in Turkey X diseases. *Endeavour* 22: 75-79.
- Steyn, P.S. (1995). Mycotoxins, general view, chemistry and structure. *Toxicol. Let.* 82/83: 843-851.
- Sudakin, D.L. (2003). Trichothecenes in the environment: Relevance to human health. *Toxicol. Let.* 143: 97-107.
- Sun, G., Wang, S., Hu, X., Su, J., Huang, T. and Yu, J. (2007). Fumonisin B1 contamination on home-grown corn in high-risk areas for esophageal and liver cancer in China. *Food Additives and Cont.* 24(2):181-185.
- Sweeny, M.J. and Dobson, A.D.W. (1998). Mycotoxin production by *Aspergillum*, *Fusarium* and *Penicillium* species. *Int. J. Food Microbiol.* 43: 141-158.
- Sydenham, E. W., Thiel, P. G., Marasas, W. F. O., Shephard, G. S., Van Schalkwyk, D. J. and Koch, K. R. (1990). Natural occurrence of some *Fusarium* mycotoxins in corn from low and high esophageal cancer prevalence areas of the Transkei, Southern Africa. *J. Agric. Food Chem.* 38:1900-1903.
- Temur, C. and Tiryaki, O. (2013). Irradiation alone or combined with other alternative treatments to control postharvest diseases. *A. J. Agric. Res.* 8(5):421-434.
- Thomas, M.D. and Buddenhagen, I.W. (1980). Incidence and persistence of *Fusarium moniliforme* in symptomless maize kernels and seedlings in Nigeria. *Mycologia* 72: 882-887.
- Thrane, U. (2001). Developments in the taxonomy of *Fusarium* species based on secondary metabolites. In: Summerbell, B.A., Leslie, J.F., Backhouse, D., Bryden, W.L., Burgess, L.W. (Eds.), *Fusarium*. Paul E. Nelson Memorial Symposium. *APS Press*, St. Paul, Minnesota, pp. 29-49.
- Turner, P., Sylla, A., Gong, Y., Diallo, M., Sutcliffe, A., Hall, A. and Wild, C. (2005). Reduction of exposure to carcinogenic aflatoxins by postharvest intervention measures in West Africa: a communitybased intervention study. *Lancet* 365:1950-1959.
- Ueno, Y. (1983). *Trichothecenes - Chemical, Biological and Toxicological Aspects*. Elsevier, Amsterdam, The Netherlands.
- Uriah, N., Ibeh, I.N. and Oluwafemi, F. (2001). A study on the impact of aflatoxin on human reproduction. *A. J. Rep. Health* 5(1):106-110.
- Van der Merwe, K.J., Steyn, P.S. and Fourie, L. (1965). Mycotoxins. Part II. The constitution of ochratoxin A, B and C, metabolites of *Aspergillus ochraceus* Wilh. *J. Chem. Soc.* 5, 7083-7088.
- Van Egmond, H.P. (1989). Aflatoxin M1: occurrence, toxicity, regulation. In: Van Egmond, H.P. (Ed.), *Mycotoxins in Dairy Products*. Elsevier Applied Science, London, pp. 11-55.
- Voss, K.A., Riley, R.T., Norred, W.P., Bacon, C.W., Meredith, F.I., Howard, P.C., Plattner, R.D., Collins, T.F., Hansen, D.K. and Porter, J.K. (2001). An overview of rodent toxicities: liver and kidney effects of fumonisins and *Fusarium moniliforme*. *Environ Health Perspect.*, 109 (2):259-266.
- Whitlow, L.W. and Hagler, W.M. (2004). Mycotoxins in feeds. *Feedstuffs* 76 (38):66-76.
- Williams, J., Phillips, T.D., Jolly, P.E., Stiles, J.K., Jolly, C.M. and Aggarwal, D. (2004). Human aflatoxicosis in developing countries: a review of toxicology, exposure, potential health consequences, and interventions. *Am. J. Clin. Nutr.* 80, 1106-1122.
- World Health Organization (WHO) (2006). Mycotoxins in African foods: implications to food safety and health. AFRO Food Safety Newsletter. World Health Organization Food safety (FOS), [www.afro.who.int/des](http://www.afro.who.int/des).
- World Health Statistics (WHO) (2008). WHO Press, Geneva, [http://www.who.int/whosis/whostat/EN\\_WHS08\\_Full.pdf](http://www.who.int/whosis/whostat/EN_WHS08_Full.pdf).
- Wu, F. (2007). Measuring the economic impacts of *Fusarium* toxins in animal feeds. *Anim Feed Sci Technol.* 137(3-4): 363-374.
- Zhou, T., He, J. and Gong, J. (2008). Microbial transformation of trichothecene mycotoxins. *World Myc. J.* 1: 23-30.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:  
<http://www.iiste.org>

## CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

**Prospective authors of journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

## MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

## IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

