

Effect of Solvent on of 4- Phenyl-7-Hydroxy Coumarin - P – Chloro Anilide

DARSHAN SINGH (corresponding Author)

Department of Physics, Deenbandhu Chhotu Ram University of Science & Technology
 Murthal (Sonipat) Haryana, INDIA
 E-mail: darshansaini82@gmail.com

VINOD KUMAR

Department of Physics, Deenbandhu Chhotu Ram University of Science & Technology
 Murthal (Sonipat) Haryana, INDIA
 E-mail: vinodkumarrohj@gmail.com

YATEENDERA TIWARI

Department of Physics, Gurgaon college of Engineering, VPO – Pathreri, Bilaspur – Tauru Road,
 Gurgaon – 122413, Haryana, INDIA
 E-mail: yateendratiwari@rediffmail.com

S.K.SINGH

Department of Physics, Deenbandhu Chhotu Ram University of Science & Technology
 Murthal (Sonipat) Haryana, INDIA
 E-mail: sksingh2k6@gmail.com

Abstract

The absorption and fluorescence emission spectra of 4- phenyl-7-hydroxy Coumarin - p-chloroanilide have been reported in various polar and non polar solvents. The increase in solvent polarity shows the increase in absorption and emission wave length i.e $\pi \rightarrow \pi^*$ transition. This indicates the value of excited state dipole moments can't be calculated using the solvatochromic data. The results are explained in terms of bathochromic-shift, and possibility of hydrogen bonding.

1. Introduction

A large number of coumarin derivatives, with different substituent, occur as natural products and most of them are synthesized in laboratory¹. The photo- physical property of coumarin depends on the nature and position of substituent in the parent molecule and also change, due to change in surrounding media. The solvatochromic analysis using the Stoke's shift data is useful in calculating the excited state dipole moment (μ_e) of the molecule². The knowledge of excited state dipole moment and excited state charge distribution³ is important in understanding the photo – physical processes. A prior knowledge of the excited state is found helpful to assess the efficacies of quantum chemical derivation of wave function as well as electron correlation treatment. There has been a renewed interest in understanding excited state dipole moments, in connection with the design of organic molecules having large hyper – polarizabilities, to fabricate efficient nonlinear optical molecules⁴. A systematic study of solvent effect, in general, gives information about the first excited singlet states behavior of a molecule. Structural stability of the molecule can be inferred from the study of their polarization values in different organic solvents and also from the temperature dependent studies of fluorescence emission⁵⁻⁶. In the present work, the authors have studied the Stoke's shift data for 4- phenyl-7-hydroxy Coumarin - p-chloroanilide & and calculated values of function $F_1(D,n)$ & $F_2(D,n)$ for different solvents.

2. Experimental Details

The flavanone and its derivatives were synthesized and investigated, by adopting standard method published elsewhere⁷. The absorption spectra were obtained with a Shimadzu-260, UV-VIS recording spectro-photometer and the fluorescence spectra using an Aminco-Bowman spectrophoto-fluorometer. The recorded fluorescence emission wavelengths are accurate with in ± 2 nm and the absorption wavelengths within ± 1 nm.

3. Results and discussion

The structure of the various flavanones studied (I) are given in Fig. 1. The observed values of $\bar{\nu}_a$ (longest absorption wavelength maxima) and $\bar{\nu}_f$ (fluorescence band maxima) in various polar and non polar organic solvents along with the calculated values of the solvents parameters $F_1(D,n)$ $F_2(D,n)$ are parented in Table 1.

Solvent effect excited state dipole moment

Solvatochromic analysis is generally used for estimating the excited state dipole moment of the fluorescence

molecules^{5-6,9}. The following relations obtained from the theoretical consideration for the solvents effect, are useful on obtaining the excited state dipole moments².

$$(\bar{\nu}_a - \bar{\nu}_f) = S_1 F_1(D, n) + K_1 \dots \dots \dots (1)$$

And

$$\frac{1}{2}(\bar{\nu}_a - \bar{\nu}_f) = S_2 F_2(D, n) + K_2 \dots \dots \dots (2)$$

Here, $\bar{\nu}_a$ and $\bar{\nu}_f$ are absorption and fluorescence maxima, n is the refractive index and D the dielectric constant of the solvent. The expression for S_1 and S_2 are :

$$S_1 = \frac{2(\mu_e - \mu_g)^2}{hCa^3} \dots \dots \dots (3)$$

and

$$S_2 = \frac{-2(\mu_e^2 - \mu_g^2)^2}{hCa^3} \dots \dots \dots (4)$$

Where μ_e and μ_g are the excited and ground state dipole moment of the solute molecule, 'a' is the Onsager cavity radius, h is the Planck's constant and C is the velocity of light. Further $F_1(D, n)$ and $F_2(D, n)$ are given by the relation:

$$F_1(D, n) = \left[\frac{D-1}{D+2} - \frac{n^2-1}{n^2+1} \right] \left[\frac{2n^2+1}{n^2+2} \right] \dots \dots \dots (5)$$

And

$$F_2(D, n) = \frac{1}{2} F_1(D, n) + \frac{3}{2} \left[\frac{(n^4-1)}{(n^2+2)^2} \right] \dots \dots \dots (6)$$

If the included shifts are neglected in $\bar{\nu}_a$ and $\bar{\nu}_f$ due to complex formation, charge transfer and hydrogen bonding etc. and further assumed that, the angle between μ_e and μ_g is small and the cavity radius 'a' is same in both the ground and excited states, than from Eqs (1) and (2) the ratio of the dipole moments in the excited state and the ground state is given by:

$$\frac{\mu_e}{\mu_g} = \frac{|S_1 - S_2|}{|S_1 + S_2|} \dots \dots \dots (7)$$

Further the value of excited state dipole moment can be obtained if the Onsager cavity radius 'a' is known, using (3) and (4)

$$\mu_e = \frac{a}{2} (hCaS_1)^{1/2} \left[1 - \frac{S_1}{S_2} \right] \dots \dots \dots (8)$$

Also knowing the value of μ_e the value of μ_g can be obtained. The value of Onsager cavity radius has been calculated using the method of Vander Waal increments¹³ (Table2).

The plots of $(\bar{\nu}_a - \bar{\nu}_f)$ versus $F_1(D, n)$ and $\frac{1}{2}(\bar{\nu}_a + \bar{\nu}_f)$ versus $F_2(D, n)$ for compound (I) is drawn, Figs 1.1-1.2, using least square fit method. These plots are found not be linear, so excited state dipole moment can't be calculated.

4. Conclusions

The present investigation thus shows that:

1. The increase in solvent polarity causes abathochromic shift for the longest absorption band maxima as well as for the fluorescence peak for 4- phenyl-7-hydroxy Coumarin - p-chloroanilide. This shows $\pi \rightarrow \pi^*$ transition.
2. The value of excited state dipole moment (μ_e) of 4- phenyl-7-hydroxy Coumarin - p-chloroanilide can't calculate, due to possibility of hydrogen bonding.
3. Structure of 4- phenyl-7-hydroxy Coumarin - p-chloroanilide also shows the possibility of formation of hydrogen bonding or molecular complexes.

References

1. Murray, R. D. H., Mendez, J. & Brown, S. A. (1982). The Natural Coumarins. Chichester :Wiley Interscience, (Chapter 5).
2. Lakowicz, J. R. (2006). Principles of fluorescence spectroscopy. (3rd ed.). New York: Plenum Press.
3. Rohatgi, K.K. & Mukherjee (1986). Fundamentals of photochemistry. New Delhi: New Age (Chapter 4).
4. Chemla, D. S. & Zyss, J. (1987). Non-linear optical property of organic molecules and crystals. New Delhi Acadmic Press.
5. Gaplovsky, A., Donoyalova J., Magdolen, P., Toma, S. & Zahradnik, P. (2002). Spectroscopic study of 2-[2-(4-cyclaminophenyl)ethen-1-yl] benzothiazoles and their N-allylbenzothiazolium bromides. Solvent and substituent effects on their ultraviolet-visible and fluorescence spectra. Spectrochim Acta-A, 58(2),363- 371.
6. Kumar, S., Giri,R., Mishra, S. C. & Machwe, M. K. (1995). Photophysical characteristics of the laser dye 7-dimethylamino cyclopenta [c] coumarin. Spectrochim Acta-A, 51(9), 1459-1467.
7. Vogel, A. I. (2011). Elementary Practical Organic Chemistry: Small Scale Preparations Part 1. (2nd ed), India: Pearson Education, (Chapter 5).
8. Melhuish, W. H. (1962). Calibration of Spectrofluorimeters for Measuring Corrected Emission Spectra. J Opt Soc Am, 52(11),1256-1258.
9. Nikolai, V., Savintsev, I., Kovaler, A. & Wadkin, R. M. (2001). Solvatochromism of the Excitation and Emission Spectra of 7-Aminoactinomycin D: Implications for Drug Recognition of DNA Secondary Structures. J Phys Chem-B, 105(35), 8461-8467.
10. Edward, T. J. (1970). Molecular volumes and the Stokes-Einstein equation. J Chem Education B, 47(4), 261-269.

Table 1

| Sr.No | Solvents | λ_a (nm) | λ_f (nm) | $\bar{\nu}_a$ (cm^{-1}) | $\bar{\nu}_f$ (cm^{-1}) | $F_1(D,n)$ | $F_2(D,n)$ | $\bar{\nu}_a - \bar{\nu}_f$ (cm^{-1}) | $\frac{\bar{\nu}_a + \bar{\nu}_f}{2}$ (cm^{-1}) |
|-------|------------|---------------------|---------------------|--------------------------------|--------------------------------|------------|------------|--|--|
| 1 | Benzene | 279 | 305 | 35842 | 32786 | 0.005 | 0.34 | 3052 | 34314 |
| 2 | Dioxane | 288 | 328 | 34722 | 30487 | 0.04 | 0.31 | 4235 | 32604 |
| 3 | Methenol | 290 | 330 | 34482 | 30303 | 0.85 | 0.65 | 4179 | 32392 |
| 4 | Isobutanol | 393 | 342 | 34129 | 29239 | 0.75 | 0.65 | 4890 | 31684 |
| 5 | Ethanol | 292 | 343 | 34246 | 29154 | 0.81 | 0.65 | 5092 | 31700 |
| 6 | n -butanol | 292 | 342 | 34246 | 29239 | 0.77 | 0.65 | 5007 | 31742 |
| 7 | Aceton | 300 | *360 to 365 | 33333 | 27586 | 0.79 | 0.64 | 5747 | 30459 |
| 8 | Water | 304 | *360 to 370 | 32894 | 27397 | 0.91 | 0.68 | 5497 | 30145 |
| 9 | Formamide | 310 | *360 to 370 | 32258 | 27397 | 0.89 | 0.75 | 4861 | 29827 |

*Broad band, mean value taken.

Stoke's shift data for 4- phenyl-7-hydroxy Coumarin - p-chloroanilide & and calculated values of function $F_1(D,n)$ & $F_2(D,n)$ for different solvents.

Graph




Figure 1.1 : Plot of $F_1(D,n)$ versus Stoke's shifts according to Bakshiev's relation for 4- phenyl-7- hydroxy Coumarin - p-chloroanilide.




Figure 1.2 : Plot of $F_2(D,n)$ versus Stoke's shifts according to Chamma and Viallet's relation for 4- phenyl-7- hydroxy Coumarin - p-chloroanilide.