

VOLUME 6

ISSUE 2

JULY – DECEMBER 2018

Al-Kimia

Preparasi Material Sensor Kreatinin dengan Metode *Molecularly Imprinted* Menggunakan Prekursor MBAA dan AMPSA

Karmanto, Ahmad Amjad Muzani

Synthesis of N-Benzenesulfonyl-*p*-Coumaramide from *p*-Coumaric Acid

Nasriadi Dali, Arniah Dali

Penurunan Konsentrasi BOD₅, COD dan Padatan Tersuspensi pada Air Limbah dengan Menggunakan Teknologi Lahan Basah Buatan (*Constructed Wetland*)

Philiphi de Rozari, Sherly M.F. Ledoh

Uji Aktivitas Antibakteri Limbah Kulit Pisang Kepok (*Musa acuminata x balbisiana*), Kulit Pisang Uli (*Musa Paradisiaca Sapientum*), dan Kulit Pisang Nangka (*Musa sp L*)

Andi Nursanti, Irma Herawati Suparto, Tetty Kemala

Analisis Flavonoid Total Akar Tabar Kedayan (*Aristolochia foveolata* Merr)

Siti Jubaidah, Henny Nurhasnawati

Analisis Komposisi Asam Lemak Dari Mikroalga Laut *Navicula salinicola*

Liska Ramdanawati, Dewi Kurnia, Vita Aji Kusumaning Tyas, Zeily Nurachman

Deteksi Bakteri Patogen *Salmonella typhi* pada Sayuran Mentah Menggunakan Metode *nested Polymerase Chain Reaction*

Idar, Shinta Kusumawardhani, Mia Tria Novianti

Uric Acid Biosensor Based on Biofilm of *L. plantarum* using *Screen-Printed Carbon Electrode* Modified by Magnetite

Dian Siska RF, Deden Saprudin, Dyah Iswantini, Novik Nurhidayat

Kadar Fenolat dan Flavonoid Total serta Kapasitas Antioksidan Ekstrak Etanol dan Fraksi Jahe Merah (*Zingiber officinale var. Rubrum*)

Dwi Koko Pratoko, Firdha Aprillia Wardhani, Nia Kristiningrum, Fifteen Aprilia Fajrin, Dian Agung Pangaribowo

Sintesis dan Karakterisasi Hidroksiapatit Dari Tulang Ikan Tuna (*Thunnus Albacores*) Dengan Xrf, Ftir, Dan Xrd


Sitti Chadijah, Hardiyanti, Sappewali

Penentuan Sifat Fisikokimia Madu Hutan (*Apis dorsata*) Sulawesi Selatan

Sjamsiah*, Rismawati Sikanna, Azmalaeni Rifkah.A, Asri

Jurusan Kimia UIN Alauddin Makassar

p-ISSN: 2302-2736



Al-Kimia

EDITOR IN CHIEF

Sjamsiah

MANAGING EDITOR

Aisyah

REVIEWER

Ambara Rahmat Pradipta

Sarifah Fauziah

Suminar Setiati Achmadi

Muharram

Safri Ishmayana

Desi harneti Putri Huspa

Ajuk Sapar

Muhammad Qaddafi

St .Chadijah

Asri Saleh

Asriyani Ilyas

SECTION EDITOR

Rani Maharani

Ummi Zahra

Firnanelty Rasyid

A.Nurfitriani Abubakar

Chusnul Chatimah Asmad

Satriani

PUBLISHER

Department of Chemistry

Faculty of Science and Technology

Universitas Islam Negeri Alauddin Makassar

Jl. H. M. Yasin Limpo No. 36 Gowa South Sulawesi Indonesia

E -mail: al-kimia@uin-alauddin.ac.id

Al-Kimia

TABLE OF CONTENT

Preparasi Material Sensor Kreatinin dengan Metode <i>Molecularly Imprinted</i> Menggunakan Prekursor MBAA dan AMPSA Karmanto, Ahmad Amjad Muzani	97-112
Synthesis of N-Benzenesulfonyl- <i>p</i> -Coumaramide from <i>p</i> -Coumaric Acid Nasriadi Dali, Arniah Dali	113-119
Penurunan Konsentrasi BOD ₅ , COD dan Padatan Tersuspensi pada Air Limbah dengan Menggunakan Teknologi Lahan Basah Buatan (<i>Constructed Wetland</i>) Philiphi de Rozari, Sherly M.F. Ledoh	120-128
Uji Aktivitas Antibakteri Limbah Kulit Pisang Kepok (<i>Musa acuminata x balbisiana</i>), Kulit Pisang Uli (<i>Musa Paradisiaca Sapientum</i>), dan Kulit Pisang Nangka (<i>Musa sp L</i>) Andi Nursanti, Irma Herawati Suparto, Tetty Kemala	129-134
Analisis Flavonoid Total Akar Tabar Kedayan (<i>Aristolochia foveolata</i> Merr) Siti Jubaidah, Henny Nurhasnawati	135-140
Analisis Komposisi Asam Lemak Dari Mikroalga Laut <i>Navicula salinicola</i> Liska Ramdanawati, Dewi Kurnia, Vita Aji Kusumaning Tyas, Zeily Nurachman	141-149
Deteksi Bakteri Patogen <i>Salmonella typhi</i> pada Sayuran Mentah Menggunakan Metode <i>nested Polymerase Chain Reaction</i> Idar, Shinta Kusumawardhani, Mia Tria Novianti	150-159
Uric Acid Biosensor Based on Biofilm of <i>L. plantarum</i> using <i>Screen-Printed Carbon Electrode</i> Modified by Magnetite Dian Siska RF, Deden Saprudin, Dyah Iswantini, Novik Nurhidayat	160-170
Kadar Fenolat dan Flavonoid Total serta Kapasitas Antioksidan Ekstrak Etanol dan Fraksi Jahe Merah (<i>Zingiber officinale var. Rubrum</i>) Dwi Koko Pratoko, Firdha Aprillia Wardhani, Nia Kristiningrum, Fifteen Aprilia Fajrin, Dian Agung Pangaribowo	171-183
Sintesis dan Karakterisasi Hidroksiapatit Dari Tulang Ikan Tuna (<i>Thunnus Albacores</i>) Dengan Xrf, Ftir, Dan Xrd Sitti Chadijah, Hardiyanti, Sappewali	184-190
Penentuan Sifat Fisikokimia Madu Hutan (<i>Apis dorsata</i>) Sulawesi Selatan Sjamsiah, Rismawati Sikanna, Azmalaeni Rifkah.A, Asri Saleh	191-199

Synthesis of *N*-Benzenesulfonyl-*p*-Coumaramide from *p*-Coumaric Acid

Nasriadi Dali^{1*} and Arniah Dali²

¹Chemistry Department, Faculty of Mathematics and Natural Sciences,
University of Halu Oleo, Kendari.

²Department of Chemistry Education, Faculty of Teacher Training and Education,
University of Halu Oleo, Kendari.

*Email: nasriadidali@gmail.com

Received:Septemb,9,2018/Accepted:December,13,2018

doi:10.24252/al-kimia.v6i2.5891

Abstract: *N*-Benzenesulfonyl-*p*-Coumaramide has been synthesized from *p*-qumarid acid. This research aimed to synthesized *N*-Benzenesulfonyl-*p*-Coumaramide from *p*-qumarid acid. Targeted compound obtained from two steps reaction. Which were amidation and asilation. Synthesized product was identified and characterized by melting point, thin Layer Chromatography analysis, FTIR Spectrophotometer, ¹H-NMR, dan ¹³C-NMR. The result showed the obtained *N*-Benzenesulfonyl-*p*-Coumaramide was white solid (yield 78.58 %) with melting point was 145-147 °C and KLT (SiO₂, *n*-hexane : cloroform = 6 : 4 v/v, R_f = 0.45).

Keywords: asilation, amidation, *N*-Benzenesulfonyl-*p*-Coumaramide, *p*-qumarid acid.

1. INTRODUCTION

The *p*-coumaramide compound (Fig. 1) has been isolated from the root bark of *Paliassa Kleinhovia hospita* Linn and has a biological activity against shrimp larvae *Artemia salina* Leach with LC₅₀ = 180.53 µg/mL (Ilyas, 2008). The results of this research indicate that *p*-coumaramide compound have the potential to be developed as anticancer and antibacterial drugs.




Figure 1. Structure of *p*-coumaramide compound (Ilyas, 2008)

Based on the results of this study, Firdaus et al (2009) has synthesized *p*-coumaramide compound from *p*-coumaric acid and tested its biological activity against leukemia cancer cell P-388. The result of *p*-coumaramide compound test against leukemia cancer cell P-388 showed that *p*-coumaramide compound has biological activity with IC₅₀ = 44 µg/mL. According to Anderson et al (1990), the compound has strong anticancer activity if the IC₅₀ value < 20 µg/mL. Therefore, *p*-coumaramide compounds deserve to be the basic framework for obtaining more active compounds.

The biological activity of *p*-coumaramide compounds can be increased by modifying the molecular structure (Tang, 2005). For this reason, Firdaus et al (2011) has synthesized and tested the biological activity of P-388 leukemia cancer cell of 3 *p*-coumaramide derived compounds, *N,N*-diethyl-*p*-coumaramide (IC₅₀ = 23.50 µg/mL), *N*-propyl-*p*-coumaramide (IC₅₀ = 53.56 µg/mL), and piperidiny-*p*-coumaramide (IC₅₀ = 5.34 µg/mL). Rasyid (2014) has also synthesized and tested the biological activity of P-388 tumor cells from 2 *p*-coumaramide derivatives, namely methyl β-(*p*-hydroxyphenyl) acrylate (IC₅₀ = 16.51 µg/mL) and methyl β-(*p*-methoxyphenyl) acrylate (IC₅₀ = 21.18 µg/mL).

Similarly, Dali and Dali (2017) have also succeeded in synthesizing *N-p*-methylbenzyl-*p*-coumaramide (Fig. 2) from *p*-coumaric acid.




Figure 2. Structure of *N-p*-methylbenzyl-*p*-coumaramide compound (Dali and Dali, 2017)

The search for other *p*-coumaramide derivated compounds can be performed by choosing a secondary or primary amine group that is less polar, making it easier for these compounds to pass through cell membranes containing lipid compounds (Shargel et al., 2004). Besides the group polarity factor, the efficacy factor of the group is also used as the basis for consideration in determining the selection of group on N atom that bound to the basic framework. For example, amide compounds derived from sulfonic acid (RSO₃H) are efficacious as antibiotic or antibacterial drugs (Matta et al., 1996). Both of these reasons underlie the selection of target synthesis compounds in this research.

2. MATERIALS AND METHOD

Tools

Spectrometer FTIR Prestige-21 (Shimadzu), ¹H-NMR and ¹³C-NMR (FTNMR JEOL ECX500), melting point meter (Electrothermal 9100), analytical balance (Ohaus), Celsius thermometer, magnetic stirrer of 1 cm, heating mantle, three neck flask of 100 mL, spherical cooler, desiccator, refrigerator, oven, and glassware.

Materials

The *p*-coumaric acid, benzene, thionyl chloride, pyridine, commercial nitrogen, tetrahydrofuran, benzenesulfonamine, triethylamine, dichloromethane, *n*-hexane, chloroform, aquabidest (Onelab Waterone), anhydrous sodium sulfate, sodium iodide, potassium carbonate, hydrogen chloride, sodium chloride, and TLC plate. These chemicals are ordered from Merck and or Sigma Aldrich.

Procedures

Synthesis of p-coumaroyl chloride from p-coumaric acid

The *p*-coumaric acid compound of 0.44 g (0.5 mmol) is introduced into a round bottom three neck flask of 100 mL equipped with a spherical cooler. Subsequently 25 mL of dry benzene, 5 mL SOCl₂, and 3 drops of pyridine were added to a round bottom three neck flask. The mixture is stirred and refluxed for 8 hours at room temperature while flowed with nitrogen gas. Every 2 hours the mixture was tested by TLC method. The excess of thionyl chloride is separated by distillation (bp 71°C). The liquid formed is the *p*-coumaroyl chloride. The *p*-coumaroyl chloride solution is unstable, so this compound is used directly without further purification (Dali, 2016).

Synthesis of *N*-benzenesulfonyl-*p*-coumaramide from *p*-coumaroyl chloride

Chloride acid solution (0.3 g, 0.4147 mmol) in dry tetrahydrofuran (5 mL) was added dropwise under nitrogenous atmosphere conditions into a benzenesulfonamine solution (0.21 mL, 1.51 mmol) and triethylamine (0.21 mL, 1.51 mmol) in dry tetrahydrofuran (10 mL). The mixture is stirred with a magnetic stirrer for 24 hours at room temperature while flowing nitrogen gas. Every 8 hours the mixture is tested on TLC. Next the mixture is filtered and the filtrate is concentrated. The residue was dissolved in dichloromethane and the solution was washed with cold water (-5 °C), dried with Na₂SO₄ anhydrous. The product solution is filtered and the solvent evaporated. The formed solids are recrystallized with methanol-dichloromethane. The white solid formed is the target compound (Dali, 2016). The target compound is then dried in the desiccator and then characterized by KLT test, melting point, FTIR, ¹H-NMR, and ¹³C-NMR.

3. RESULTS AND DISCUSSION

Results

The *N*-benzenesulfonyl-*p*-coumaramide target compound is a white solid (yield 78.57%) with a melting point of 145-147 °C and TLC (SiO₂, *n*-hexane : chloroform = 6 : 4 v/v, R_f = 0.45). This target compound is obtained through two stages of the synthesis reaction. The first stage is an acylation reaction between *p*-coumaric acid as a starting material with thionyl chloride in a dry benzene solvent and produce *p*-coumaroyl chloride. In this acylation reaction stage there is a conversion of carboxylic acid group (RCO₂H) to acyl chloride group (RCOCl) (Fig. 3). This conversion reaction is supported by changes in R_f values of the reactant (0.75) to product (0.37). This is consistent with the expected product in which the product have a higher polarity than the reactant. The *p*-coumaroyl chloride is not spectroscopically analyzed because it is highly reactive to water vapor.




Figure 3. Synthesis reaction of *p*-coumaroyl chloride from *p*-coumaric acid

The second stage is an amidation reaction between *p*-coumaroyl chloride and benzene sulfonamine in a dry tetrahydrofuran solvent and we obtained the targeted molecule, namely a *N*-benzenesulfonyl-*p*-coumaramide target compound. At this stage of the amidation reaction there is conversion of the acyl chloride group (RCOCl) to the amide group (R'CONHR) (Fig. 4). This conversion reaction is supported by changes in R_f values of the reactant (0.37) to product (0.45). This is as expected, namely the product has a lower polarity compared to the reactant.




Figure 4. Synthesis reaction of *N*-benzenesulfonyl-*p*-coumaramide from *p*-coumaroyl chloride

Discussion

The FTIR spectrum of the target compound shows that two very strong absorption bands at 3377.36 cm^{-1} and 1687.71 cm^{-1} derived from the stretching of OH and C=O groups of *p*-coumaric acid as a starting materials are no longer visible. This indicates that the acylation and amidation reactions have occurred. This data is also reinforced by the appearance of three strong absorption bands at 3437.50 , 1651.07 , and 1510.26 cm^{-1} respectively derived from a stretching of N-H group of secondary amides, C=O group of secondary amides (amide I band), and C-N group of secondary amides (amide II band). Other supporting data are the appearance of three strong absorption bands at 1150.21 , 1170.79 , and 551.64 cm^{-1} respectively derived from the stretching of SO₂ group, SO₂ symmetry, and SO₂ scissor. Thus, from the FTIR data the target compound shows that the conversion reaction of the carboxylic acid group (RCO₂H) in *p*-coumaric acid as a starting material has changed to the amide group (R'CONHR) in *N*-benzenesulfonyl-*p*-coumaramide.

The success of the *N*-benzenesulfonyl-*p*-coumaramide synthesis reaction of *p*-coumaric acid is also supported by ¹H-NMR spectra data (Fig. 5).




Figure 5. Interpretation of ¹H-NMR (500 MHz, CDCl₃) spectrum data of *p*-coumaric acid and *N*-benzenesulfonyl-*p*-coumaramide synthesis product

The ¹H-NMR spectrum of the target compound did not show any signal at $\delta_{\text{H}} 10.3157$ ppm (1H, s, 1) derived from a proton of the *p*-coumaric acid (OH-1). This data is also reinforced by the appearance of a signal at $\delta_{\text{H}} 5.9762$ ppm (1H, s) derived from proton amide (NH).

Other supporting data are the appearance of three signals at δ_{H} 7.4301 ppm (1H, *d*, 2* and 6*), 7.4180 ppm (1H, *t*, 3* and 5*), and 7.2598 ppm (1H, *t*, 4*) derived from aryl protons (ArH) in *ortho*, *meta*, and *para* positions. Thus, the signals appearing in the $^1\text{H-NMR}$ spectrum of the target compound are in accordance with the proton character of the *N*-benzenesulfonyl-*p*-coumaramide compound (Fig. 5).

The interpretation of $^{13}\text{C-NMR}$ spectrum data (500 MHz, CDCl_3) of the target compound (Fig. 6) also strengthens the results of FTIR and $^1\text{H-NMR}$ spectrum analysis above.




Figure 6. Interpretation of $^{13}\text{C-NMR}$ spectrum data (500 MHz, CDCl_3) *p*-coumaric acid and *N*-benzenesulfonyl-*p*-coumaramide synthesis product

The $^{13}\text{C-NMR}$ spectrum data of the target compound shows the presence of the aryl carbon signals (C-aryl) not found in the spectrum of *p*-coumaric acid as a starting material. The signals of the aryl carbon atoms (C-aryl) are dispersed into four δ_{C} values, ie δ_{C} 138.7135 ppm [(C-terminal aryl) (C-SO₂)], δ_{C} 124.7254 ppm (C-*o* aryl), δ_{C} 126.9137 ppm (C-*m* aryl), and δ_{C} 126.2358 ppm (C-*p* aryl). While the signals of other carbon atoms that emerge from the target compound are in accordance with the signals of carbon atoms from *p*-coumaric acid (Fig. 6). Therefore, the signals appearing in the $^{13}\text{C-NMR}$ spectrum of the target compound are in accordance with the carbon skeleton of the *N*-benzenesulfonyl-*p*-coumaramide (Fig. 7).




Figure 7. The carbon framework of the *N*-benzenesulfonyl-*p*-coumaramide

4. CONCLUSION

The *N*-benzenesulfonyl-*p*-coumaramide has been successfully synthesized from *p*-coumaric acid through acylation and amidation reaction steps. The *N*-benzenesulfonyl-*p*-coumaramide is a white solid (yield 78.57%) with a melting point of 145-147 °C and TLC (SiO₂, *n*-hexane : chloroform = 6 : 4 v/v, R_f = 0.45).

Suggestion

Biological activity of *N*-benzenesulfonyl-*p*-coumaramide needs to be tested further so that these compounds can be used as raw materials for anticancer and antibacterial drugs.

Acknowledgements

Thanks to the Head of Organic Chemistry Laboratory and Head of Integrated Chemistry Laboratory of Chemistry Department FMIPA Unhas Makassar and Head of Laboratory of LIPI Chemical Research Center of PUSPIPTEK Area Serpong South Tangerang.

REFERENCES

- Anderson, J. E., Goetz, C. M., and McLaughlin, J. L. (1990). Blind Comparison of Simple Bench-top Bioassays and Human Tumour Cell Cytotoxicities as Antitumour Prescreen. *J. Phytochemical Analysis*. 6 : 107-111.
- Dali, N. (2016). *Synthesis, Characterization, and Application of p-tert-Butylcalix[4]-and-[6]arene Derivative Compounds as Ionophores in Ion Selective Electrode (ISE) Coated Wire Type for Analysis of Hg Heavy Metal Content Hg(II) and Au(I)*. Dissertation Not Published. Graduate School of Unhas, Makassar.
- Dali, N., and Dali, A. (2017). Synthesis of *N-p*-Methylbenzyl-*p*-Coumaramide from *p*-Coumaric Acid. *Al-Chemy*,. 5(2) : 154-160.
- Firdaus, Soekamto, N. H., and Karim, A. (2009). Synthesis of *p*-Hydroxyamide Compounds from *p*-Hydroxycinnamic Acid Through the Reaction of Esterification and Ammonolysis. *J. Chemica Acta*,. 2 : 2-8.
- Firdaus, Soekamto, N. H., Umar, U., Dali, S., Makmun, and Agustningsih, A. (2011). *Synthesis of p-Coumaramida Derivative Compounds and their Bioactivity Test against Cancer Cell Leukemia P-388*. Research Report. FMIPA Unhas, Makassar.
- Ilyas, A. (2008). *Isolation and Identification of Secondary Metabolites from Ethyl Acetate Extracts of Root Skin of Kleinhovia hospita Linn (Paliasa) and Toxicity Test Against Artemia salina Leach*. Thesis Not Published. Unhas Postgraduate Program, Makassar.
- Matta, M. S., Wilbraham, A. C., and Staley, D. D. (1996). *Introduction to Organic and Biological Chemistry*. D. C. Heath and Company, Lexington.
- Rasyid, H. (2014). *Synthesis of Methyl β-(p-hydroxyphenyl)acrylate and Methyl β-(p-Methoxyphenyl)acrylate and Bioactivity Test against P-388 Tumor Cells*. Thesis Not Published. FMIPA Unhas, Makassar.
- Shargel, L., Yu, A., and Wu-Pong, S. (2004). *Applied Biopharmaceutics and Pharmacokinetics*. 5th Edition. McGraw Hill Professional, New York.

Tang, P. (2005). Boroc Acid Catalyzed Amide Formation from Carboxylic Acid and Amines *N*-Benzyl-4-Phenylbutyramide. *J. Organic Syntheses*, 81 : 262-267.