



OIST

OKINAWA INSTITUTE OF SCIENCE AND TECHNOLOGY GRADUATE UNIVERSITY  
沖縄科学技術大学院大学


# Testing ontogenetic patterns of sexual size dimorphism against expectations of the expensive tissue hypothesis, an intraspecific example using oyster toadfish (*Opsanus tau*)

Author	Alex Dornburg, Dan L. Warren, Katerina L. Zapfe, Richard Morris, Teresa L. Iglesias, April Lamb, Gabriela Hogue, Laura Lukas, Richard Wong
journal or publication title	Ecology and Evolution
volume	8
number	7
page range	3609-3616
year	2018-03-02
Publisher	John Wiley & Sons Ltd.
Rights	(C) 2018 The Authors.
Author's flag	publisher
URL	<a href="http://id.nii.ac.jp/1394/00000608/">http://id.nii.ac.jp/1394/00000608/</a>

doi: info:doi/10.1002/ece3.3835

## ORIGINAL RESEARCH

# Testing ontogenetic patterns of sexual size dimorphism against expectations of the expensive tissue hypothesis, an intraspecific example using oyster toadfish (*Opsanus tau*)

Alex Dornburg<sup>1</sup>  | Dan L. Warren<sup>2</sup> | Katerina L. Zapfe<sup>1</sup> | Richard Morris<sup>1</sup> |  
Teresa L. Iglesias<sup>3</sup> | April Lamb<sup>1,4</sup> | Gabriela Hogue<sup>1</sup> | Laura Lukas<sup>1</sup> | Richard Wong<sup>5</sup>

<sup>1</sup>North Carolina Museum of Natural Sciences, Raleigh, NC, USA

<sup>2</sup>Senckenberg Institute for Biodiversity and Climate, Frankfurt am Main, Germany

<sup>3</sup>Physics and Biology Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan

<sup>4</sup>Department of Applied Ecology, North Carolina State University, Raleigh, NC, USA

<sup>5</sup>Delaware Division of Fish and Wildlife, Dover, DE, USA

## Correspondence

Alex Dornburg, North Carolina Museum of Natural Sciences, Raleigh, NC, USA.  
Email: alex.dornburg@naturalsciences.org

## Funding information

Trawl collections were funded by the U.S. Fish and Wildlife Service through the Federal Aid in Fisheries Restoration Act.

## Abstract

Trade-offs associated with sexual size dimorphism (SSD) are well documented across the Tree of Life. However, studies of SSD often do not consider potential investment trade-offs between metabolically expensive structures under sexual selection and other morphological modules. Based on the expectations of the expensive tissue hypothesis, investment in one metabolically expensive structure should come at the direct cost of investment in another. Here, we examine allometric trends in the ontogeny of oyster toadfish (*Opsanus tau*) to test whether investment in structures known to have been influenced by strong sexual selection conform to these expectations. Despite recovering clear changes in the ontogeny of a sexually selected trait between males and females, we find no evidence for predicted ontogenetic trade-offs with metabolically expensive organs. Our results are part of a growing body of work demonstrating that increased investment in one structure does not necessarily drive a wholesale loss of mass in one or more organs.

## KEYWORDS

evolutionary ecology, fishes, life history trade-offs, phenotypic evolution, reproductive physiology, swim bladder

## 1 | INTRODUCTION

Pronounced differences in ecology, life history, or morphology between males and females of the same species are common features of the vertebrate Tree of Life (Barrett & Hough, 2012; Herler, Kerschbaumer, Mitteroecker, Postl, & Sturmbauer, 2010; Karp et al., 2017; Lamb et al., 2017; Nottebohm & Arnold, 1976). Sexual size dimorphism (SSD), the variation between sexes in aspects of size, is a particularly striking pattern that has commanded the attention of researchers since Darwin (Clutton-Brock, Harvey, & Rudder, 1977; Darwin, 1871; Rohner, Blanckenhorn, & Puniamorthy, 2016; Scudder, 1876; Shine, 1978). The past several decades have yielded

remarkable insights into the eco-evolutionary dynamics (Legrand & Morse, 2000; Maan & Seehausen, 2011; Price, 1984; Sonerud et al., 2012), evolutionary trade-offs (Dunn et al., 2015; Gustafsson, Qvarnström, & Sheldon, 1995; Simmons & Emlen, 2006), and ontogeny underlying the rise of traits that exhibit SSD (German, 2004; Glassman, Coelho, Carey, & Bramblett, 1984; Hassell, Meyers, Billman, Rasmussen, & Belk, 2012; Holton, Alsamawi, Yokley, & Froehle, 2016). However, potential energetic trade-offs between selection for SSD and investment in metabolically expensive organs are rarely considered, precluding a broader understanding of how SSD shapes fundamental aspects of phenotypic evolution in vertebrates.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

The evolution of SSD requires selection to promote changes in some aspect of allometric growth (Bonduriansky, 2007). However, how modular these changes are remain unclear. Do such ontogenetic changes reflect trade-offs with other components of a given species' bauplan? This question is particularly relevant for SSD in metabolically or developmentally costly organs, as organisms are faced with a finite energy budget that they can invest into different structures in order to accumulate biomass. This raises the question of not only how organisms have evolved the sometimes extreme differences in organ size observed today, but also whether there are hidden costs to SSD. An often invoked answer to the generalized question of how organisms are able to change biomass investment in metabolically expensive organs was first conceptualized by Aiello and Wheeler (1995) in the form of the expensive tissue hypothesis (ETH). This hypothesis specifically posited that investment in a major metabolically expensive organ, the brain, should come at a cost to one or more other organ systems. As costly traits characterized by SSD (such as gonads or ornaments) become expressed, expectations of the ETH suggest that energy budgets will be differentially balanced between sexes, thereby driving reduced investment in the brain or other structures for the sex under selection. While the ubiquity of trade-offs in life history evolution provides intuitive appeal for the ETH, evidence supporting the expectations of this hypothesis has not been overwhelming.

Interspecific studies of metabolic trade-offs between organ systems have yielded mixed results for the ETH that include positive (Liao, Lou, Zeng, & Kotschal, 2016; Sukhum, Freiler, Wang, & Carlson, 2016; Tsuboi et al., 2015), negative (Bordes, Morand, & Krasnov, 2011; Jones & MacLarnon, 2004), or a lack of support (Isler & van Schaik, 2006; Lemaître, Ramm, Barton, & Stockley, 2009; Navarrete, van Schaik, & Isler, 2011; Schillaci, 2006) for the ETH. Likewise, intraspecific studies have yielded a mixture of positive support (Kotschal, Corral-Lopez, Szidat, & Kolm, 2015; Kotschal, Kolm, & Penn, 2016; Kotschal et al., 2013) and inconclusive/negative evidence (Warren & Iglesias, 2012). It is important to consider that the ETH was initially formulated with the intent of understanding size variation in the vertebrate brain (Aiello & Wheeler, 1995), and therefore, work investigating trade-offs has been almost entirely focused on the potential costs associated with increased brain size. Whether the ETH provides a predictive framework for understanding the impact of SSD in gonads or other costly organs in development remain unclear. Does SSD limit investment in the brain or other organs consistent with the expectations of the ETH?

Oyster toadfish (*Opsanus tau*; Figure 1) represent an exemplary species in which to investigate the impact of SSD on the ontogeny of metabolically costly traits. The physiology and life history of this species has been consistently studied for over a century (Clapp, 1891; Fine, 1975; Fine, McKnight, & Blem, 1995; Fine & Waybright, 2015; Gray & Winn, 1961; Schwartz & Dutcher, 1963; Tracy, 1926), and SSD has been well documented in one unusual metabolically expensive trait: the swim bladder. The swim bladders of oyster toadfish and their close relatives (Batrachoididae) are not primarily used for buoyancy, but instead serve as highly derived sound production organs that are unusual among teleost fishes (Fine, 1975). In oyster toadfish, both males and females use these organs for grunt-based



**FIGURE 1** Illustration of an oyster toadfish (*Opsanus tau*). Illustration by KLZ

communication (Fine & Waybright, 2015), but male swim bladders emit a specific "boat whistle"-like call to attract mates and exhibit a nearly twofold increase in swim bladder size as a result (Fine, 1975; Fine, Burns, & Harris, 1990). The demands of this call have driven a nearly 50% increase in the size and number of fibers in sonic muscles that surround the bladder, giving rise to one of the fastest twitching muscles found in vertebrates (Fine et al., 1990). Although there is a clear metabolic cost associated with the development of these traits, the allocation of energy by male toadfish to tissue investment and the potential cost to the development of their brains or other organs remains unclear.

Here, we analyze body, brain, liver, swim bladder, heart, and gonad mass collected from a population of oyster toadfish to test whether SSD drives ontogenetic trade-offs that support expectations of the ETH. We first quantify the allometric relationships of all organs to validate previous observations of SSD and test for differences in allometry between sexes. We then assess whether significant ontogenetic increases of organ masses with SSD are negatively correlated with the mass of other organs as anticipated by the predictions of the ETH. Results of our study provide a much needed ontogenetic investigation of SSD within the conceptual framework of the ETH and provide a critical perspective on the need for further development of theoretical expectations concerning the evolutionary relationship between sexual selection and organismal energy budgets.

## 2 | MATERIALS AND METHODS

Specimens for this study were collected by research trawl across several sites in the western Delaware Bay between August and November 2016 (Greco, 2017; Figure S1). For each fish, standard and total length were recorded in millimeters (mm), and wet body mass, liver mass, heart mass, swim bladder mass, brain mass, and eviscerated body mass were recorded in milligrams (mg). Voucher photographs and otoliths were taken from each specimen and deposited in the Fish Collections of North Carolina Museum of Natural Sciences (NCSM FO 17-28; 30; 35-40; 43; 45-55; 64-75; 80-89; 97-98; 100-106; 109-116). Tests for investment trade-offs expected under the ETH have repeatedly focused on the heart, brain, and liver as candidate organs (e.g., Aiello, 1997; Navarrete et al., 2011; Warren & Iglesias, 2012), as

these are metabolically expensive and likely subjected to evolutionary trade-offs (Konarzewski & Diamond, 1995). Since the inception of the ETH, several studies have also expanded the candidate pool of potential metabolically expensive organs subjected to trade-offs to include gonads (e.g., Bordes et al., 2011; Liu, Zhou, & Liao, 2014; Tsuboi, Shoji, Sogabe, Ahnesjö, & Kolm, 2016). As such our study captures tissues generally considered to be among the most likely targets for metabolic trade-offs. Thirty-six males ranging from 91 to 262 mm and 19 females ranging from 120 to 237 mm in total length were examined (Table S1).


To quantify allometric relationships between eviscerated body mass and each organ mass for males and females, we used analysis of covariance (ANCOVA). Prior to analysis, organ and eviscerated body masses were  $\log_{10}$  transformed. Although the use of a principle components analysis for extracting size as an independent measure has been used in allometric studies, this practice can drive spurious inferences of trait relationships in regression analyses (see Berner, 2011). As a result, we used eviscerated body mass as a representative of mass. Eviscerated mass was chosen over wet body mass as the stomach contents of individual fishes varied considerably, from no prey to individuals containing copious amounts of shellfish or conspecifics almost a quarter of their body size. A linear model was fit with tissue mass as the dependent variable and eviscerated body mass, sex, and  $\text{sex} \times \text{eviscerated body mass}$  as independent variables. Body mass was treated as a covariate, and sex difference was assessed by determining whether separate regression lines fit the data better than a single regression line. Sex difference was tested using an  $F$  test with two degrees of freedom ( $df$ ) in the numerator, one  $df$  for a sex difference in slope, and the other  $df$  for a sex difference in intercept. An alpha level of 0.05 was used to infer a sex difference in body mass allometry

for each organ. When an allometric difference between sexes was inferred, the result for a test for difference in slope was reported.


To evaluate ETH for male swim bladder as a hypothetically expensive tissue, we estimated partial correlation coefficients between swim bladder and each of four candidate tissues (gonad, heart, brain, and liver). As applied here, partial correlation measures linear association between swim bladder and a candidate tissue while controlling for the influence of body mass (see Supporting Information). When controlling for a single variable, in this case body mass, the partial correlation coefficient can be obtained from three correlation coefficients, swim bladder–body mass, target organ–body mass, and swim bladder–target organ (see Supporting Information). Again, all tissue masses were  $\log_{10}$  transformed prior to computing correlations, and 95% confidence intervals for partial correlation coefficients were based on a  $t$  distribution with  $n - 3$  degrees of freedom. A negative partial correlation coefficient for a target organ supports an ETH interpretation for swim bladder. Results obtained through partial correlation were additionally compared to multivariate and univariate regression (see Supporting Information). Additionally, to account for a potential confounding effect of sonic muscles in the fish trunk following evisceration, similar correlation analyses were repeated using standard length (mm) as a measure of body size (see Supporting Information).

### 3 | RESULTS

We found significant allometric differences between males and females for swim bladder mass ( $F_{2,56} = 13.38, p < .0001$ ), gonad mass ( $F_{2,55} = 42.02, p < .0001$ ), and liver mass ( $F_{2,56} = 12.13, p < .0001$ ). No significant difference between sexes was found for heart mass



**FIGURE 2** Patterns of sexual size dimorphism (SSD) in oyster toadfish for (a) swim bladder, (b) gonad, (c) liver, (d) heart, and (e) brain masses. Females are depicted in blue, and males are depicted in orange. Light shading of the plot indicates significant evidence for SSD based on an ANCOVA



**FIGURE 3** Partial regression coefficient estimates for linear relationships between mass of swim bladder and other putative metabolically expensive tissues. A 95% confidence interval for a negative correlation coefficient that excludes zero would support the expectation of the expensive tissue hypothesis (ETH) (light shading), while a confidence interval for a positive coefficient that excludes zero would provide contrary evidence (dark shading). Partial correlation intervals that include 0 provide no evidence for or against the expectations of the ETH

( $F_{2,56} = 2.902$ ,  $p = .063$ ) or brain mass ( $F_{2,56} = 1.924$ ,  $p = .155$ ). ANCOVA results supported a sex difference in slope for the swim bladder and body mass relationship ( $F_{1,56} = 20.06$ ,  $p < .001$ ), providing strong evidence for a difference in allometric growth of the swim bladder between males and females. Visualizations of the allometric trajectory for swim bladder mass show large males possess larger swim bladders than females of comparable body mass (Figure 2). Females appear to diverge from males in the slope of the regression of gonad on body mass (Figure 2); however, the one  $df$  test for a sex difference in slope did not achieve statistical significance ( $F_{1,56} = 3.762$ ,  $p = .058$ ). Likewise, no strong support was found for an allometric difference between sexes in liver mass ( $F_{1,56} = 0.978$ ,  $p = .327$ ; Figure 2).

Point estimates and associated 95% confidence intervals for partial correlation coefficients between swim bladder and four organs (gonad, heart, brain, and liver) are plotted in Figure 3. Point estimates for three partial correlation coefficients are negative and one is positive. However, all confidence intervals include zero (Figure 3), thereby providing no strong evidence for a negative linear association between swim bladder and any of the four internal organs individually examined. The sum of heart, brain, and liver mass was also evaluated for a negative association with swim bladder and exhibited the same result as individual organs. Collectively, all of our results provide no support for a negative correlation between swim bladder mass and the mass

of any other candidate organ and thereby do not support the expectations of the ETH.

## 4 | DISCUSSION

Swim bladder SSD has been well documented in oyster toadfish (Fine, 1975; Fine et al., 1990). Fine (1975) proposed that the size differences in the toadfish swim bladder are the result of different ontogenetic growth trajectories between males and females. Given little variation between the ontogenies of individual males and females, we would expect similar ontogenetic (individual) and static (population) allometric slopes. Our findings support this hypothesis, showing that an increase in static allometric slope in males relative to females underlies the sex-related difference in swim bladder size (see Figure 2). For species characterized by SSD, including those with highly exaggerated traits such as weapons and ornaments, positive changes in static allometric slope, where larger individuals have disproportionately larger traits, are far less common than negative changes or changes in intercept (isometry; reviewed in Bonduriansky, 2007). The lack of evidence for a tight coupling between positive change in allometry and SSD is consistent with the hypothesis that allometric slopes are not very evolvable below the species level (Voje, Hansen, Egset, Bolstad, & Pélabon, 2014). In species where changes in ontogenetic or static allometric



slope have been found, two explanations are commonly invoked. First, changes in slope driven by shifts in environment or resource utilization can promote rapid phenotypic divergence (e.g., Collins, Dornburg, Flores, Dombrowski, & Lewbart, 2016; Ruehl, Shervette, & Dewitt, 2011; Wund, Valena, Wood, & Baker, 2012). This explanation seems unlikely for oyster toadfish as swim bladders of both sexes are not used for buoyancy, but rather communication, and males and females overlap in habitat and diet (Wilson 1982). Alternatively, sexual selection theory predicts that strong selection on trait exaggeration with a low overall cost to the organism can promote allometric slope changes (Bonduriansky & Day, 2003). This hypothesis is appealing; however, our study does not offer a quantification of energetic costs associated with the swim bladder. Further, we do not believe this selection hypothesis offers a complete explanation. Instead, we propose that the changes in static allometric slope we report here may have deep evolutionary origins.

Other species of Batrachoidiformes, such as members of the genus *Porichthys* (Midshipman), have been found to be sexually dimorphic in swim bladder size (Mohr et al., 2017). Given that *Porichthys* and *Opsanus* share common ancestry at least 30 million years ago (Near et al., 2013), it is likely that the changes in ontogenetic slope found in our study may represent an axis of Batrachoidiform diversification that vastly predates the origin of *Opsanus tau*. Given these deep origins, the energetic costs associated with this organ may have long been integrated into the ontogeny of these organisms. Consequently, our study may represent a small component of a macroevolutionary shift in evolutionary allometry. Future work placing the evolutionary allometry of Batrachoidiform swim bladders into a phylogenetic framework and assessing the costs and developmental mechanisms of ontogenetic change between sexes of multiple species represents exciting frontiers that are necessary to evaluate the origin of this unusual organ and its related musculature.

Despite finding clear evidence of a significant change in the allometric slope of swim bladder growth between toadfish sexes, our analyses did not recover any evidence of a correlated trade-off in the mass of another organ during ontogeny (Figure 2). Male toadfish swim bladder mass has previously been found to be highly correlated with both sonic muscle size and the number of fibers (Fine, 1975), making swim bladder mass a good proxy for the heavy energetic cost associated with the development of the male toadfish acoustic repertoire. It is certainly possible that a trade-off between a tissue or life history trait and swim bladder mass may exist and was simply not examined here. Given that we find no significant negative change in brain, liver, or heart allometric trajectories, our study adds to the growing number of studies that have failed to recover support for the ETH in traits where an expectation of positive evidence is reasonable (reviewed in Warren & Iglesias, 2012).

The decoupling of ontogenetic changes between the swim bladder and other organs suggested by our results raises the possibility that the ontogeny of the swim bladder and related musculature may represent a morphological module. Modularity, the degree of separation of one axis of phenotype from other organismal parts, is a fundamental principle of biological organization (Esteve-Altava, 2017). However,

the expectations of the ETH suggest that the modular organization of biological forms can impact the development of the brain as a consequence of resource allocation constraints. We found no clear relationship between brain mass and the mass of any other organ. This raises the question: To what extent should we expect increased investment of one module to directly negatively impact the ontogenetic trajectory of another?

Studies of resource allocation trade-offs have suggested tissue proximity to be a potential predictor of changes in investment (Emlen, 2001). Although organisms must use a finite energy budget to accumulate body mass, the ubiquity of modularity in organismal systems ranging from mammals to fishes (Esteve-Altava, 2017; Larouche, Cloutier, & Zelditch, 2015) without obvious trade-offs between adjacent tissues (reviewed in Warren & Iglesias, 2012) suggests that simple economic predictions between morphological modules may not have much explanatory power for understanding the evolution of most ontogenetic pathways without a more detailed perspective of lineage-specific energy budgets and life history. While investment trade-offs between morphological modules have provided evidence for the expectations of the ETH in a few animal lineages (Emlen, 2001; Liao et al., 2016; Moczek & Nijhout, 2004), the large number of studies that have failed to recover support in other lineages suggests that the broad expectations of the ETH are far from a universal rule (reviewed in Warren & Iglesias, 2012).

While increases in the energetic cost of one morphological module do not often lead to negatively correlated changes in another, this does not preclude the possibility that some modules are in fact faced with a possible deficit in energy as sexual selection emphasizes trait investment (Moczek & Nijhout, 2004). However, without a detailed understanding of the ecology and energy requirements of a species, it is not clear to what extent organisms can offset deficits through changes in behavior or feeding ecology. Such subtle changes may in part explain the strong evidence for the ETH in experimental laboratory settings with tight controls (Kotrschal et al., 2013, 2015, 2016), despite limited support from wild populations such as the toadfishes in this study. Further, an organism is comprised of a suite of morphological modules that collectively use an energy budget to invest mass into their respective structures. Numerous subtle increases or decreases to energy consumption across any number of modules can therefore offset the cost of strong sexual selection to a specific module. Such a readjustment of energy budgets is reminiscent of many-to-one mapping of form to function, where numerous phenotypic solutions give rise to the same functional properties of a trait (Wainwright, Alfaro, Bolnick, & Hulsey, 2005). As such, the repeated lack of evidence for the ETH may not result from a lack of trade-offs, but rather from subtle and complex adjustments of ontogenetic investment between numerous morphological modules across an entire organism that cannot be detected through approaches based on tissue mass alone. Future studies assessing the behavior, feeding ecology, and energy costs of different morphological modules will be needed to determine whether any of these hypotheses explain the lack of trade-offs between the male toadfish swim bladder and other organs.

## 5 | CONCLUSION

Determining the impact of energetic trade-offs in the ontogenetic pathways that give rise to the diversity of phenotypes we observe today is fundamental to evolutionary biology. While the ETH holds promise of a general evolutionary principle, evidence for direct trade-offs between the brain and other metabolically expensive organs has been limited to few clades. In contrast, numerous studies, including this study, have reported negative evidence within and among a broad spectrum of clades. While a one-to-one mapping of brain investment increase to trait reduction does appear to exist in some species (Kotrschal et al., 2013; Tsuboi et al., 2015, 2016), these examples are few in number. Refinement of the ETH as well as the formulation of new metabolic investment hypotheses is warranted and needed to broaden our perspective on energetic trade-offs. Such hypotheses are critical if we are to develop new insights into the role of sexual selection in shaping other aspects of organismal form.

### ACKNOWLEDGMENTS

We thank Michael Greco, Joseph Mathews, Christian Pruneda, Cameron Day, and Captain Mike Garvilla for collecting specimens during the Delaware Division of Fish and Wildlife trawl surveys, and S. Guthrie, W. Brooks, L. Roupe, and Z. Creech for their help in processing fish used in this study. We additionally thank two anonymous reviewers for providing valuable feedback on an earlier version of this manuscript.

### CONFLICT OF INTEREST

None declared.

### AUTHOR CONTRIBUTIONS

A.D. and R.W. conceived of the study. R.W. collected samples. A.D., K.L.Z., A.L., G.H., and L.L. collected data. R.M., A.D., and D.W. performed analyses. A.D., D.W., K.L.Z., R.M., T.L.I., A.L., and R.W. wrote the initial manuscript. All other authors contributed to the subsequent writing and development of the manuscript. All data have been archived on Zenodo: (DOI pending formal acceptance).

### ORCID

Alex Dornburg  <http://orcid.org/0000-0003-0863-2283>

### REFERENCES

- Aiello, L. C. (1997). Brains and guts in human evolution: The expensive tissue hypothesis. *Brazilian Journal of Genetics*, 20, 141–148. <https://doi.org/10.1590/S0100-84551997000100023>
- Aiello, L. C., & Wheeler, P. (1995). The expensive-tissue hypothesis: The brain and the digestive system in human and primate evolution. *Current Anthropology*, 36, 199–221. <https://doi.org/10.1086/204350>
- Barrett, S. C. H., & Hough, J. (2012). Sexual dimorphism in flowering plants. *Journal of Experimental Botany*, 64, 67–82.
- Berner, D. (2011). Size correction in biology: How reliable are approaches based on (common) principal component analysis? *Oecologia*, 166(4), 961–971. <https://doi.org/10.1007/s00442-011-1934-z>
- Bonduriansky, R. (2007). Sexual selection and allometry: A critical reappraisal of the evidence and ideas. *Evolution*, 61, 838–849. <https://doi.org/10.1111/j.1558-5646.2007.00081.x>
- Bonduriansky, R., & Day, T. (2003). The evolution of static allometry in sexually selected traits. *Evolution*, 57, 2450–2458. <https://doi.org/10.1111/j.0014-3820.2003.tb01490.x>
- Bordes, F., Morand, S., & Krasnov, B. R. (2011). Does investment into “expensive” tissue compromise anti-parasitic defence? Testes size, brain size and parasite diversity in rodent hosts. *Oecologia*, 165, 7–16. <https://doi.org/10.1007/s00442-010-1743-9>
- Charles A., Wilson, John Mark, Dean, & Richard, Radtke (1982). Age, growth rate and feeding habits of the oyster toadfish, *Opsanus tau* (Linnaeus) in South Carolina. *Journal of Experimental Marine Biology and Ecology*, 62, 251–259. ISSN 0022-0981, [https://doi.org/10.1016/0022-0981\(82\)90205-2](https://doi.org/10.1016/0022-0981(82)90205-2). (<http://www.sciencedirect.com/science/article/pii/0022098182902052>)
- Clapp, C. M. (1891). Some points in the development of the Toad-fish (*Batrachus tau*). *Journal of Morphology*, 5, 494–501. [https://doi.org/10.1002/\(ISSN\)1097-4687](https://doi.org/10.1002/(ISSN)1097-4687)
- Clutton-Brock, T. H., Harvey, P. H., & Rudder, B. (1977). Sexual dimorphism, socionomic sex ratio and body weight in primates. *Nature*, 269, 797–800. <https://doi.org/10.1038/269797a0>
- Collins, S., Dornburg, A., Flores, J., Dombrowski, D., & Lewbart, G. (2016). A comparison of blood gases, biochemistry, and hematology to ecomorphology in a health assessment of pinfish (*Lagodon rhomboides*). *PeerJ*, 4, e2262. <https://doi.org/10.7717/peerj.2262>
- Darwin, C. R. (1871). *The descent of man, and selection in relation to sex* (Vol. 1, 1st edn). London: John Murray.
- Dunn, J. C., Halenar, L. B., Davies, T. G., Cristobal-Azkarate, J., Reby, D., Sykes, D., ... Knapp, L. A. (2015). Evolutionary trade-off between vocal tract and testes dimensions in howler monkeys. *Current Biology*, 25, 2839–2844. <https://doi.org/10.1016/j.cub.2015.09.029>
- Emlen, D. J. (2001). Costs and the diversification of exaggerated animal structures. *Science*, 291, 1534–1536. <https://doi.org/10.1126/science.1056607>
- Esteve-Altava, B. (2017). In search of morphological modules: A systematic review. *Biological Reviews of the Cambridge Philosophical Society*, 92, 1332–1347. <https://doi.org/10.1111/brv.12284>
- Fine, M. L. (1975). Sexual dimorphism of the growth rate of the swimbladder of the toadfish *Opsanus tau*. *Copeia*, 1975, 483. <https://doi.org/10.2307/1443646>
- Fine, M. L., Burns, N. M., & Harris, T. M. (1990). Ontogeny and sexual dimorphism of sonic muscle in the oyster toadfish. *Canadian Journal of Zoology*, 68, 1374–1381. <https://doi.org/10.1139/z90-205>
- Fine, M. L., McKnight, J. W., & Blem, C. R. (1995). Effect of size and sex on buoyancy in the oyster toadfish. *Marine Biology*, 123, 401–409. <https://doi.org/10.1007/BF00349218>
- Fine, M. L., & Waybright, T. D. (2015). Grunt variation in the oyster toadfish *Opsanus tau*: Effect of size and sex. *PeerJ*, 3, e1330. <https://doi.org/10.7717/peerj.1330>
- German, R. Z. (2004). The ontogeny of sexual dimorphism: The implications of longitudinal vs. cross-sectional data for studying heterochrony in mammals. In: F. Anapol et al. (Eds.), *Shaping primate evolution* (pp. 11–23). Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/CBO9780511542336>
- Glassman, D. M., Coelho, A. M. Jr, Carey, K. D., & Bramblett, C. A. (1984). Weight growth in savannah baboons: A longitudinal study from birth to adulthood. *Growth*, 48, 425–433.
- Gray, G.-A., & Winn, H. E. (1961). Reproductive ecology and sound production of the toadfish, *Opsanus Tau*. *Ecology*, 42, 274–282. <https://doi.org/10.2307/1932079>

- Greco, M. (2017). *Coastal finfish assessment survey. Final report, Project No. F16AF00047 (F-42-R-28)*. Dover, DE: Delaware Division of Fish and Wildlife.
- Gustafsson, L., Qvarnström, A., & Sheldon, B. C. (1995). Trade-offs between life-history traits and a secondary sexual character in male collared flycatchers. *Nature*, 375, 311–313. <https://doi.org/10.1038/375311a0>
- Hassell, E. M. A., Meyers, P. J., Billman, E. J., Rasmussen, J. E., & Belk, M. C. (2012). Ontogeny and sex alter the effect of predation on body shape in a livebearing fish: Sexual dimorphism, parallelism, and costs of reproduction. *Ecology and Evolution*, 2, 1738–1746. <https://doi.org/10.1002/ece3.278>
- Herler, J., Kerschbaumer, M., Mitteroecker, P., Postl, L., & Sturmbauer, C. (2010). Sexual dimorphism and population divergence in the Lake Tanganyika cichlid fish genus *Tropheus*. *Frontiers in Zoology*, 7, 4. <https://doi.org/10.1186/1742-9994-7-4>
- Holton, N. E., Alsamawi, A., Yokley, T. R., & Froehle, A. W. (2016). The ontogeny of nasal shape: An analysis of sexual dimorphism in a longitudinal sample. *American Journal of Physical Anthropology*, 160, 52–61. <https://doi.org/10.1002/ajpa.22941>
- Isler, K., & van Schaik, C. (2006). Costs of encephalization: The energy trade-off hypothesis tested on birds. *Journal of Human Evolution*, 51, 228–243. <https://doi.org/10.1016/j.jhevol.2006.03.006>
- Jones, K. E., & MacLarnon, A. M. (2004). Affording larger brains: Testing hypotheses of mammalian brain evolution on bats. *American Naturalist*, 164, E20–E31. <https://doi.org/10.1086/421334>
- Karp, N. A., Mason, J., Beaudet, A. L., Benjamini, Y., Bower, L., Braun, R. E., ... Fuchs, H. (2017). Prevalence of sexual dimorphism in mammalian phenotypic traits. *Nature Communications*, 8, 15475. <https://doi.org/10.1038/ncomms15475>
- Konarzewski, M., & Diamond, J. (1995). Evolution of basal metabolic rate and organ masses in laboratory mice. *Evolution*, 49, 1239–1248. <https://doi.org/10.1111/j.1558-5646.1995.tb04450.x>
- Kotrschal, A., Corral-Lopez, A., Szidat, S., & Kolm, N. (2015). The effect of brain size evolution on feeding propensity, digestive efficiency, and juvenile growth. *Evolution*, 69, 3013–3020. <https://doi.org/10.1111/evo.12784>
- Kotrschal, A., Kolm, N., & Penn, D. J. (2016). Selection for brain size impairs innate, but not adaptive immune responses. *Proceedings of the Royal Society B: Biological Sciences*, 283, 20152857. <https://doi.org/10.1098/rspb.2015.2857>
- Kotrschal, A., Rogell, B., Bundsen, A., Svensson, B., Zajitschek, S., Brännström, I., ... Kolm, N. (2013). Artificial selection on relative brain size in the guppy reveals costs and benefits of evolving a larger brain. *Current Biology*, 23, 168–171. <https://doi.org/10.1016/j.cub.2012.11.058>
- Lamb, A. D., Watkins-Colwell, G. J., Moore, J. A., Warren, D. L., Iglesias, T. L., Brandley, M. C., & Dornburg, A. (2017). Endolymphatic sac use and reproductive activity in the Lesser Antilles endemic gecko *Gonatodes antillensis* (Gekkota: Sphaerodactylidae). *Bulletin of the Peabody Museum of Natural History*, 58, 17–29. <https://doi.org/10.3374/014.058.0103>
- Larouche, O., Cloutier, R., & Zelditch, M. L. (2015). Head, body and fins: Patterns of morphological integration and modularity in fishes. *Evolutionary Biology*, 42, 296–311. <https://doi.org/10.1007/s11692-015-9324-9>
- Legrand, R. S., & Morse, D. H. (2000). Factors driving extreme sexual size dimorphism of a sit-and-wait predator under low density. *Biological Journal of the Linnean Society*, 71, 643–664. <https://doi.org/10.1111/j.1095-8312.2000.tb01283.x>
- Lemaître, J.-F., Ramm, S. A., Barton, R. A., & Stockley, P. (2009). Sperm competition and brain size evolution in mammals. *Journal of Evolutionary Biology*, 22, 2215–2221. <https://doi.org/10.1111/j.1420-9101.2009.01837.x>
- Liao, W. B., Lou, S. L., Zeng, Y., & Kotrschal, A. (2016). Large brains, small guts: The expensive tissue hypothesis supported within anurans. *American Naturalist*, 188, 693–700. <https://doi.org/10.1086/688894>
- Liu, J., Zhou, C. Q., & Liao, W. B. (2014). Evidence for neither the compensation hypothesis nor the expensive-tissue hypothesis in *Carassius auratus*. *Animal Biology*, 64(2), 177–187. <https://doi.org/10.1163/15707563-00002437>
- Moczek, A. P., & Nijhout, H. F. (2004). Trade-offs during the development of primary and secondary sexual traits in a horned beetle. *American Naturalist*, 163, 184–191. <https://doi.org/10.1086/381741>
- Mohr, R. A., Whitchurch, E. A., Anderson, R. D., Forlano, P. M., Fay, R. R., Ketten, D. R., ... Sisneros, J. A. (2017). Intra- and intersexual swim bladder dimorphisms in the plainfin midshipman fish (*Porichthys notatus*): Implications of swim bladder proximity to the inner ear for sound pressure detection. *Journal of Morphology*, 278, 1458–1468. <https://doi.org/10.1002/jmor.20724>
- Navarrete, A., van Schaik, C. P., & Isler, K. (2011). Energetics and the evolution of human brain size. *Nature*, 480, 91–93. <https://doi.org/10.1038/nature10629>
- Near, T. J., Dornburg, A., Eytan, R. I., Keck, B. P., Smith, W. L., Kuhn, K. L., ... Wainwright, P. C. (2013). Phylogeny and tempo of diversification in the superradiation of spiny-rayed fishes. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 12738–12743. <https://doi.org/10.1073/pnas.1304466110>
- Nottebohm, F., & Arnold, A. P. (1976). Sexual dimorphism in vocal control areas of the songbird brain. *Science*, 194, 211–213. <https://doi.org/10.1126/science.959852>
- Price, T. D. (1984). Sexual selection on body size, territory and plumage variables in a population of Darwin's finches. *Evolution*, 38, 327. <https://doi.org/10.1111/j.1558-5646.1984.tb00291.x>
- Rohner, P. T., Blanckenhorn, W. U., & Puniamoorthy, N. (2016). Sexual selection on male size drives the evolution of male-biased sexual size dimorphism via the prolongation of male development. *Evolution*, 70, 1189–1199. <https://doi.org/10.1111/evo.12944>
- Ruehl, C. B., Shervette, V., & Dewitt, T. J. (2011). Replicated shape variation between simple and complex habitats in two estuarine fishes. *Biological Journal of the Linnean Society*, 103, 147–158. <https://doi.org/10.1111/j.1095-8312.2011.01626.x>
- Schillaci, M. A. (2006). Sexual selection and the evolution of brain size in primates. *PLoS ONE*, 1, e62. <https://doi.org/10.1371/journal.pone.0000062>
- Schwartz, F. J., & Dutcher, B. W. (1963). Age, growth, and food of the oyster toadfish near Solomons, Maryland. *Transactions of the American Fisheries Society*, 92, 170–173. [https://doi.org/10.1577/1548-8659\(1963\)92\[170:AGAFOT\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1963)92[170:AGAFOT]2.0.CO;2)
- Scudder, S. H. (1876). Antigeny, or sexual dimorphism in butterflies. *Proceedings of the American Academy of Arts and Sciences*, 12, 150. <https://doi.org/10.2307/25138444>
- Shine, R. (1978). Sexual size dimorphism and male combat in snakes. *Oecologia*, 33, 269–277. <https://doi.org/10.1007/BF00348113>
- Simmons, L. W., & Emlen, D. J. (2006). Evolutionary trade-off between weapons and testes. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 16346–16351. <https://doi.org/10.1073/pnas.0603474103>
- Sonerud, G. A., Steen, R., Løw, L. M., Røed, L. T., Skar, K., Selås, V., ... Slagsvold, T. (2012). Size-biased allocation of prey from male to offspring via female: Family conflicts, prey selection, and evolution of sexual size dimorphism in raptors. *Oecologia*, 172, 93–107.
- Sukhum, K. V., Freiler, M. K., Wang, R., & Carlson, B. A. (2016). The costs of a big brain: Extreme encephalization results in higher energetic demand and reduced hypoxia tolerance in weakly electric African fishes. *Proceedings of the Royal Society B: Biological Sciences*, 283, 20162157. <https://doi.org/10.1098/rspb.2016.2157>
- Tracy, H. C. (1926). The development of motility and behavior reactions in the toadfish (*Opsanus tau*). *Journal of Comparative Neurology*, 40, 253–369. [https://doi.org/10.1002/\(ISSN\)1096-9861](https://doi.org/10.1002/(ISSN)1096-9861)
- Tsuboi, M., Husby, A., Kotrschal, A., Hayward, A., Buechel, S. D., Zidar, J., ... Kolm, N. (2015). Comparative support for the expensive tissue hypothesis: Big brains are correlated with smaller gut and greater parental investment in Lake Tanganyika cichlids. *Evolution*, 69, 190–200. <https://doi.org/10.1111/evo.12556>



- Tsuboi, M., Shoji, J., Sogabe, A., Ahnesjö, I., & Kolm, N. (2016). Within species support for the expensive tissue hypothesis: A negative association between brain size and visceral fat storage in females of the Pacific seaweed pipefish. *Ecology and Evolution*, 6, 647–655. <https://doi.org/10.1002/ece3.1873>
- Voje, K. L., Hansen, T. F., Egset, C. K., Bolstad, G. H., & Pélabon, C. (2014). Allometric constraints and the evolution of allometry. *Evolution*, 68, 866–885. <https://doi.org/10.1111/evo.12312>
- Wainwright, P. C., Alfaro, M. E., Bolnick, D. I., & Hulsey, C. D. (2005). Many-to-one mapping of form to function: A general principle in organismal design? *Integrative and Comparative Biology*, 45, 256–262. <https://doi.org/10.1093/icb/45.2.256>
- Warren, D. L., & Iglesias, T. L. (2012). No evidence for the “expensive-tissue hypothesis” from an intraspecific study in a highly variable species. *Journal of Evolutionary Biology*, 25, 1226–1231. <https://doi.org/10.1111/j.1420-9101.2012.02503.x>
- Wund, M. A., Valena, S., Wood, S., & Baker, J. A. (2012). Ancestral plasticity and allometry in threespine stickleback reveal

phenotypes associated with derived, freshwater ecotypes. *Biological Journal of the Linnean Society*, 105, 573–583. <https://doi.org/10.1111/j.1095-8312.2011.01815.x>

#### SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

**How to cite this article:** Dornburg A, Warren DL, Zapfe KL, et al. Testing ontogenetic patterns of sexual size dimorphism against expectations of the expensive tissue hypothesis, an intraspecific example using oyster toadfish (*Opsanus tau*). *Ecol Evol*. 2018;8:3609–3616. <https://doi.org/10.1002/ece3.3835>