

**OIST**OKINAWA INSTITUTE OF SCIENCE AND TECHNOLOGY GRADUATE UNIVERSITY
沖縄科学技術大学院大学

Elucidating the multiple genetic lineages and population genetic structure of the brooding coral *Seriatopora* (Scleractinia: Pocilloporidae) in the Ryukyu Archipelago

Author	Yuichi Nakajima, Akira Nishikawa, Akira Iguchi, Tomofumi Nagata, Daisuke Uyeno, Kazuhiko Sakai, Satoshi Mitarai
journal or publication title	Coral Reefs
volume	36
number	2
page range	415-426
year	2017-03-14
Publisher	Springer Berlin Heidelberg
Rights	(C) 2017 Springer-Verlag Berlin Heidelberg This is a post-peer-review, pre-copyedit version of an article published in Coral Reefs. The final authenticated version is available online at: http://dx.doi.org/10.1007/s00338-017-1557-x
Author's flag	author
URL	http://id.nii.ac.jp/1394/00000821/

doi: [info:doi/10.1007/s00338-017-1557-x](https://doi.org/10.1007/s00338-017-1557-x)

**Elucidating the multiple genetic lineages and population genetic structure
of the brooding coral *Seriatopora* (Scleractinia: Pocilloporidae) in the
Ryukyu Archipelago**

Yuichi Nakajima^{1,*}, Akira Nishikawa², Akira Iguchi³, Tomofumi Nagata⁴, Daisuke Uyeno⁵,
Kazuhiko Sakai², Satoshi Mitarai¹

¹Okinawa Institute of Science and Technology Graduate University, Tancha, Onna, Okinawa
904-0495, Japan

²Sesoko Station, Tropical Biosphere Research Center, University of the Ryukyus, Sesoko,
Motobu, Okinawa 905-0227, Japan

³National Institute of Technology, Okinawa College, Henoko, Nago, Okinawa 905-2192,
Japan

⁴Okinawa Prefecture Environment Science Center, Kyo-zuka, Urasoe, Okinawa 901-2111,
Japan

⁵Graduate School of Science and Engineering, Kagoshima University, Korimoto, Kagoshima,
Kagoshima 890-0065, Japan

*Correspondence: yuichi.nakajima@oist.jp; yuichi.nakajima@outlook.com

Keywords: Genetic differentiation; Genetic diversity; Microsatellite; Mitochondrial DNA;

Species diversity

Abstract

The elucidation of species diversity and connectivity is essential for conserving coral reef communities and for understanding the characteristics of coral populations. To assess the species diversity, intraspecific genetic diversity, and genetic differentiation among populations of the brooding coral *Seriatopora* spp., we conducted phylogenetic and population genetic analyses using a mitochondrial DNA control region and microsatellites at ten sites in the Ryukyu Archipelago, Japan. At least three genetic lineages of *Seriatopora* (*Seriatopora*-A, -B, and -C) were detected in our specimens. We collected colonies morphologically similar to *Seriatopora hystrix*, but these may have included multiple, genetically distinct species. Although sexual reproduction maintains the populations of all the genetic lineages, *Seriatopora*-A and -C had lower genetic diversity than *Seriatopora*-B. We detected significant genetic differentiation in *Seriatopora*-B among the three populations as follows: pairwise $F_{ST} = 0.064\text{--}0.116$ (all $P = 0.001$), pairwise $G_{ST} = 0.107\text{--}0.209$ (all $P = 0.001$). Additionally, only one migrant from an unsampled population was genetically identified within *Seriatopora*-B. Because the peak of the settlement of *Seriatopora* larvae is within 1 d and almost all larvae are settled within 5 d of spawning, our observations may be related to a low dispersal ability. Populations of *Seriatopora* in the Ryukyu Archipelago will

probably not recover unless there is substantial new recruitment from distant populations.

Introduction

Reef-building corals are important marine organisms in tropical and subtropical reef areas because they constitute the framework of the ecosystem and enhance the productivity and diversity of the coral-reef community. In addition, coral species diversity helps to maintain the functionality and resilience of coral-reef communities to environmental stresses (e.g., Hughes et al. 2003; Richards and Hobbs 2014). However, stony corals are threatened by climate change and anthropogenic disturbances at both local and global scales (e.g., Hoegh-Guldberg 1999; Hoegh-Guldberg et al. 2007). Accurate identification of coral species is essential to monitor, conserve, and manage coral populations (e.g., Bickford et al. 2007). However, in some cases, species definitions that are based solely on colony morphological traits have been misleading, and morphological homoplasy conceals the cryptic evolutionary relationships (Richards et al. 2013, 2016 and references therein). Because traditional morphological species definitions have been problematic in corals, genetic tools have more recently proven to be useful in discriminating between morphologically similar species (e.g., Nakajima et al. 2012, 2016; Pinzón et al. 2013; Warner et al. 2015; Suzuki et al. 2016).

Population genetic approaches provide potential for distinguishing inter- and intraspecific relationships more clearly even when the relationships are impacted by recent speciation and/or hybridization. This is because changes in allele frequencies within a population are expected to occur at an ecological timescale, which can be properly assessed by genetic markers (e.g., Nakajima et al. 2012).

Mitochondrial loci and polymorphic nuclear microsatellite loci provide fundamental information about species boundaries and intraspecific genetic diversity and differentiation. Mitochondrial DNA generally shows a higher mutation rate than nuclear DNA and is suitable as a genetic marker for phylogeny and species identification in animals (e.g., Avise et al. 1987; Hebert et al. 2003). However, low mutation rates in coral species make this analysis inappropriate (van Oppen et al. 1999, 2001; Fukami et al. 2000). Recent studies have elucidated species boundaries in the family Pocilloporidae. Species within the genus *Pocillopora* were delineated using mitochondrial loci, an open reading frame with an unknown function and a control region, and a nuclear internal transcribed spacer 2 (ITS2) region. These loci suggested a mismatch between genetic clusters and species definitions based on colony morphological traits (e.g., Pinzón and LaJeunesse 2011; Pinzón et al. 2013; Schmidt-Roach et al. 2013, 2014). Furthermore, species within the genus *Stylophora* were

divided into three clusters using mitochondrial loci or ITS2, and into four clusters using ITS1, and these were morphologically more variable than previously thought (Flot et al. 2011). *Seriatopora* collected from Okinawa, New Caledonia, and the Philippines consisted of four genetic clusters on a phylogenetic tree constructed based on the sequences of mitochondrial loci, D-loop, a control region between *atp8* and *cox1*, along with the control region between *atp6* and *nad4* (Flot et al. 2008). Furthermore, *Seriatopora hystrix* was classified into four genetic lineages, including cryptic species, on the Great Barrier Reef (GBR) (Warner et al. 2015). Therefore, it can be difficult to estimate the species diversity within the family Pocilloporidae based on traditional morphological descriptions. Clarification of the genetic structure of sympatric populations using nuclear multilocus microsatellites is a robust method for estimating species diversity (*Pocillopora*: Pinzón et al. 2013; *Seriatopora*: Warner et al. 2015).

If genetic tools detect genetic homogeneity that is maintained within a species through genetic exchanges, populations could be genetically differentiated due to geographical and oceanographic barriers and population fluctuations, including genetic drift and bottlenecks that are caused by historical events (e.g., Bay et al. 2008). Microsatellites can be used to describe the genetic connectivity among populations and species. Genetic connectivity

among populations within species is maintained by larval transport via oceanographic systems such as sea currents (e.g., Pineda et al. 2007). The Kuroshio Current is a strong oceanographic current that flows southwest to northeast along the Ryukyu Archipelago of Japan. This current is considered a major factor in the expansion and maintenance of coral reefs and reef-dwelling organisms around the islands (Nishihira and Veron 1995). The Ryukyu Archipelago is a subtropical area in the northwestern Pacific and includes the northern limit for most tropical coral species (Nishihira and Veron 1995). It is composed of multiple stepping-stone, isolated islands and a well-developed reef system. In the reef system, distances between islands range from tens to several hundreds of kilometers in most cases, and the area of suitable coral reef habitat is small compared to regions such as the GBR. Thus, we predict that larval recruitment among the isolated reefs is rare, especially in brooding corals, but occurs occasionally over long periods owing to the Kuroshio Current and its branches.

Seriatopora is mainly distributed on tropical and sub-tropical reefs in the western Indo-Pacific region (Veron 2000). Nishihira and Veron (1995) described two species of *Seriatopora* (*S. hystrix* and *S. caliendrum*) that inhabit the Ryukyu Archipelago. Veron (2000) and the Corals of the World website (<http://coral.aims.gov.au/>) suggest that four

morphological species of *Seriatopora* are probably distributed in the Ryukyu Archipelago: *S. hystrix*, *S. caliendrum*, *S. guttata*, and *S. stellata*. The Japanese Coral Reef Society and Ministry of the Environment (2004) reported that three species of *Seriatopora* (*S. hystrix*, *S. caliendrum*, and *S. stellata*) inhabit the area, but *S. stellata* is only in the Miyako and Yaeyama regions of the Ryukyu Archipelago.

The extent of coral recovery after a mass bleaching event is highly variable among species. The population recovery of *S. hystrix* is very slow in the Ryukyu Archipelago and it is considered a long-term loser species that does not recover after a mass bleaching event (e.g., Loya et al. 2001; van Woesik et al. 2011). We assess multiple genetic lineages of *Seriatopora* and the genetic structure and connectivity among sites using microsatellite markers to understand the historical interspecific delimitation, intraspecific population dynamics including reproductive strategies, and recent migration patterns. In addition, we discuss the potential for recovery of *Seriatopora* corals along the Ryukyu Archipelago.

Materials and methods

Sampling and genomic DNA extraction

We collected *Seriatopora* samples from ten sampling sites in four regions of the

Ryukyu Archipelago (Fig. 1, ESM Table S1). While at least three morphological species of *Seriatopora* appear to be distributed in the Ryukyu Archipelago, we collected colonies with a *S. hystrix* morphology and avoided colonies with a different morphology, which were rare in the sampling areas. A small piece of branch was preserved in 99.5% ethanol and transferred to the laboratory. Total DNA was extracted from the ethanol-preserved tissue pellets using the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) or the AquaPure Genomic DNA Isolation Kit (BioLad, Hercules, CA, USA) according to the manufacturer's protocols.

Sequencing of the mitochondrial putative control region

As the most comparable sequence with other publications, the mitochondrial putative control region (mtCR) between *atp6* and *nad4* was amplified by polymerase chain reaction (PCR) using Ex Taq DNA polymerase (Takara, Kusatsu, Shiga, Japan) with the primers SerCtl-F1 (5'-GTCTGCTCACATTAATTTAT-3') and SerCtl-R1 (5'-AGAGATCGAACTAAGAGTCG-3') (Bongaerts et al. 2010). These primers were designed from the mitochondrial genome of *S. hystrix* (Bongaerts et al. 2010; and see Chen et al. 2008; GenBank EF633600). The mtCR was defined as a control region in Chen et al. (2008), but almost all the sequences can be translated into an open reading frame (ORF) of a hypothetical protein-encoding gene (Flot et al. 2008). The location corresponds to a novel

mitochondrial ORF that is next to *atp6* in the case of *Pocillopora*, as shown by Flot and Tillier (2007). The PCR conditions were as follows: 40 cycles of 30 s at 94°C, 1 min at 56°C, 1 min at 72°C, followed by a final extension for 5 min at 72°C, carried out in a total reaction volume of 10 μ L. For direct sequencing, each PCR product was treated with ExoSAP-IT (Affymetrix, Santa Clara, CA, USA) following the manufacturer's protocol. The primers used for the sequencing reactions were the same as those for the PCR amplification. The products were sequenced using an automated capillary-based DNA sequencer, the ABI 3130xl Genetic Analyzer (Thermo Fisher Scientific, Waltham, MA, USA).

Phylogenetic analyses for mtCR haplotypes

Tandem repeat sections in the intergenic spacer were identified using Tandem Repeat Finder ver. 4.09 (Benson 1999); the alignment parameters for mismatches and indels were three and five, respectively. If two or more patterns were detected in a haplotype, the pattern with the larger score was adopted. We used maximum likelihood (ML) and Bayesian inference (BI) analyses to construct a phylogenetic tree of the generated mtCR haplotypes, with the 23 *Seriatopora* sequences classified according to Chen et al. (2008), Flot et al. (2008), Bongaerts et al. (2010), and Warner et al. (2015), and one *Pocillopora* sequence identified from Flot and Tillier (2007) as the outgroup. These sequences were aligned using

MUSCLE on MEGA, ver. 6.0 (Tamura et al. 2013). The consensus sequences, including repeat motifs, ranged from 513 bp to 741 bp of the mtCR. Prior to analysis, we searched for the best-fit model of evolution using MEGA, and the HKY (Hasegawa–Kishino–Yano) +G (Gamma distributed) model was selected. The robustness of each ML tree was tested by bootstrapping (1,000 replicates) under the condition of complete deletion of gaps. Furthermore, this model was applied for the BI analysis using MrBayes, ver. 3.2.5 (Ronquist and Huelsenbeck 2003; Ronquist et al. 2012). Two isolated runs were conducted under the following conditions: four Markov chain Monte Carlo (MCMC) iterations, 50,000,000 generations, a print frequency of 10,000, a sample frequency of 100 generations, and a 0.25 burn-in fraction.

Scoring of microsatellite genotypes

We attempted to use nine microsatellite loci with amplification in our specimens of *Seriatopora*. Seven loci characterized by Underwood et al. (2006) were effective for our genetic analysis (ESM Table S2). Of the two remaining loci, Sh3-008 (Underwood et al. 2006) showed no amplification for specimens from some sites, and Sh4.28 (Maier et al. 2001) contained very few heterozygotes. The reaction mixture (5 μ L) contained template DNA (<100 ng), AmpliTaq Gold 360 Master Mix (Thermo Fisher Scientific), and three

primers for each locus. The primers were a non-tailed forward primer (0.5 μ M), a reverse primer with a U19 sequence tail (0.5 μ M), and a U19 (5'-GGTTTTCCCAGTCACGACG-3') primer (0.5 μ M) fluorescently labeled with FAM, VIC, or NED. Amplifications for all microsatellite loci were carried out under the following conditions: 95°C for 9 min, followed by 35 cycles at 95°C for 30 s, 54°C for 30 s, 72°C for 1 min, and a final extension at 72°C for 5 min. PCR products amplified with an internal size standard, GeneScan 600 LIZ, (Thermo Fisher Scientific) were analyzed using an ABI 3130xl Genetic Analyzer and GeneMapper ver. 3.7 (Thermo Fisher Scientific).

Estimating the genetic groupings using microsatellites

We inferred the genetic structure of *Seriatopora* through Bayesian clustering of microsatellites using STRUCTURE, ver. 2.3.4 (Pritchard et al. 2000). First, we analyzed the genetic structure with the sampling location information. A burn-in period of 100,000 iterations followed by 1,000,000 MCMC replications was used for population clustering without LOCPRIOR model under the admixture model and correlated allele frequencies were assumed (Falush et al. 2003). Individuals were assumed to be drawn exclusively from the genetic K clusters and were allowed to have mixed ancestry in the admixture model (Pritchard et al. 2000; Falush et al. 2003). Simulations included 10 iterations, and the number

of assumed subpopulations was 1 to 11. After calculation of the mean log probability, $\ln P(D)$, determination of the number of K clusters that best fit the data was conducted based on the highest ΔK value using the method of Evanno et al. (2005) as implemented in STRUCTURE HARVESTER (Earl and vonHoldt 2012). To obtain adequate output panels to unify the colors, merged run data were implemented by CLUMPAK (Kopelman et al. 2015). Second, we analyzed the genetic structure and inferred the number of genetic lineages with the mtCR haplotype information using STRUCTURE. The run conditions of STRUCTURE were the same as the first step analysis except for the sampling location information. Furthermore, discriminant analysis of principal components (DAPC) was conducted in R ver. 3.0.2 (<http://www.r-project.org>) using the package adegenet, ver. 1.3–9.2 (Jombart et al. 2010), to represent genetic clusters formed by microsatellite genotypes for each mtCR haplotype. This clustering method does not assume Hardy–Weinberg equilibrium (HWE) or linkage equilibrium. All multilocus genotypic data of microsatellites were plotted both with and without six clusters defined a priori based on mtCR haplotypes. A total of 40 principal components were retained, accounting for 97.9 % of the genetic variability.

For multilocus genotypes (including replicated genotypes), we also calculated the probability of a given multilocus genotype occurring n times for each population within a

genetic lineage, repeated as a consequence of different recombination events (P_{SEX} , calculated taking into account the F_{IS} estimates in the data set) using GenClone, ver. 2.0 (Arnaud-Haond and Belkhir 2007). We retained the replicated multilocus genotype if two or more multilocus genotypes were the same but occurred by chance because of sexual reproduction (the threshold of the P_{SEX} value is 0.01). Based on the microsatellite genotypes for each genetic lineage and site, clonal diversity was estimated with the following index: $R = (N_{MLG} - 1)/(N - 1)$, where N_{MLG} is the number of retained multilocus genotypes and N is the number of colonies analyzed. We removed duplicated genotypes of clonemates within a population to analyze the genetic differentiation and structure, genetic diversity, and migrants within genetic lineage. Genetic differentiation based on microsatellites was estimated using a hierarchical analysis of molecular variance (AMOVA) (Excoffier et al. 1992), partitioning among lineages, among populations within lineages, and within populations, by GenAlEx, ver. 6.501 (Peakall and Smouse 2006).

Population genetic analyses between sites within a genetic lineage

After removal of the duplicate genotypes of clonemates, the genetic structure within each genetic lineage was also inferred using STRUCTURE. The run conditions were the same as previously, except for the number of assumed subpopulations ($K = 1$ to 6 for each

genetic lineage). Additionally, we ran a LOCPRIOR model because the detailed genetic structure as shown in all lineages was not detected without the LOCPRIOR model (see Results). Only populations with 20 or more retained multilocus microsatellite genotypes within each lineage were used in further analyses. For each suitable population, the number of alleles and values of observed and expected heterozygosity (H_O and H_E , respectively) for each site were evaluated with GenAlEx. We calculated the fixation index (F_{IS}) and the allelic richness for genetic diversity at each site using FSTAT ver. 2.9.3.2 (Goudet 1995). Additionally, the significance of the F_{IS} was tested by randomization using FSTAT. The genetic differentiation index between sites was calculated using GenAlEx. The significance of each pairwise F_{ST} value was tested with 999 permutations. Furthermore, the G-statistic method was used to calculate the pairwise G_{ST} values to account for the small number of populations (Meirmans and Hedrick 2011), and the significance of each value was tested with 999 permutations. Migration patterns over the recent ecological timescale were estimated using GeneClass2 (Piry et al. 2004) and the Bayesian method developed by Rannala and Mountain (1997). We assigned each multilocus genotype to a source site using a 10,000 resampling algorithm. A multilocus genotype with a >99% ($P < 0.01$) confidence interval was excluded from the sampled population, and we then assigned the colony as an immigrant to

the site of highest probability when the confidence of assignment was $\geq 10\%$ ($P \geq 0.1$). If a given multilocus genotype had low probability with $< 10\%$ ($P < 0.1$) confidence of assignment for all sites, the origin of this multilocus genotype was considered an immigrant from an unsampled site.

Results

Genetic differentiation and structure among the genetic lineages

Of the 195 colonies of *Seriatopora* collected from ten sites (Fig. 1; ESM Table S1), we succeeded in obtaining mtCR haplotypes (6 haplotypes; Ser-1 to -6) and multilocus genotypes with seven microsatellites in 182 colonies (93.3%). The community composition of individuals from different haplotypes differed among sites (Fig. 1; ESM Table S1). While some sites were dominated by a single haplotype (e.g., sites O-NKJ and K-AKA; see Fig. 1 for site names), others had population mixtures with up to five different haplotypes (e.g., Y-TKT). The haplotype sequences are available in the GenBank database (accession numbers: LC107881–LC107886). The composition of each haplotype group differed greatly among sites (Fig. 1; ESM Table S1). Thirty-five replicated multilocus microsatellite genotypes were detected, and 32 replicated multilocus microsatellite genotypes were detected

considering the mtCR haplotypes. Namely, three multilocus microsatellite genotypes were shared between the Ser-1 and Ser-2 mtCR haplotypes in O-OUR (one multilocus microsatellite genotype was shared between the two Ser-1 and two Ser-2 colonies, one other multilocus genotype was in the five Ser-1 and 1 Ser-2 colonies, and the last multilocus genotype was also in the five Ser-1 and one Ser-2 colonies). Multilocus microsatellite genotypes were not shared between the sites.

For the Bayesian clustering, the Evanno method indicated that the most probable number of genetic populations was three: best $K = 3$ ($\Delta K = 2712.52$ when each site was designated as the population, and $\Delta K = 4632.23$ when each mtCR haplotype was designated as the population). The STRUCTURE analyses suggested that there was no obvious agreement with geographical sites (Fig. 2). For example, multiple clusters were mixed in O-ONA and Y-TKT, and a common cluster was shared between populations from different regions (e.g., O-ONN and K-AKA, K-TKS and Y-IRO). STRUCTURE found at least three distinct genetic lineages among the sampled individuals, and the lineages include different mtCR haplotypes (Seriatorpora-A: Ser-1 and Ser-2; Seriatorpora-B: Ser-3 and Ser-4; Seriatorpora-C: Ser-5 and Ser-6), although the lineages of five colonies in Ser-5 and Ser-6 were not defined (Fig. 2; ESM Table S1). DAPC detected three main clusters when each

haplotype was assigned to each multilocus genotype (ESM Fig. S1). Of the five colonies with lineage undefined by STRUCTURE, four were assigned to *Seriatopora*-B without prior information of mtCR haplotype and to *Seriatopora*-C with prior information. A hierarchical AMOVA quantified the extent of the genetic differentiation among the genetic lineages based on STRUCTURE after removal of the five colonies with undefined lineage (total $F_{RT} = 0.349$, $P = 0.001$; Table 1). Thus, we concluded there were at least three genetic lineages of *Seriatopora* in the Ryukyu Archipelago.

On the ML (Fig. 3) and BI (ESM Fig. S2) phylogenetic trees, four haplotypes (Ser-1, Ser-2, Ser-3, and Ser-4) were assigned to the clusters including some haplotypes of *S. caliendrum*, and two (Ser-5 and Ser-6) were assigned to the group including several *S. hystrix* lineages. Within the mtCR analyzed, Ser-2 was identical to the haplotype previously obtained from Okinawa and Taiwan, and had one nucleotide gap compared with the haplotype from the Philippines. Ser-5 is identical to *S. hystrix* HostU from the GBR, which is the same haplotype from Okinawa and Taiwan. In contrast, the other four haplotypes have not been reported before. However, the relationship of the phylogenetic trees defined according to mtCR haplotype was not in agreement with the relationship identified through the genetic structure from microsatellites. All Ser-5 and one Ser-6 were the same clusters by

microsatellites, but were separated on the phylogenetic tree using mtCR, although there were only two samples in Ser-6. In addition, tandem repeats (composed of 51 bp) were observed in the mtCR, and these were separated into three groups based on the type of tandem repeat [Ser-1: 8.5 times; Ser-2, Ser-3, and Ser-4: 4.4 times; Ser-5 and Ser-6: 5.2 times (ESM Fig. S3)], but this pattern was also not completely related to the genetic structure obtained using microsatellites.

Clonal diversity and the population genetic index within the genetic lineages

Although 35 multilocus microsatellite genotypes (32 when considering mtCR haplotypes) were replicated the P_{SEX} value revealed that the number of clonal replicates was low, with only six of the 182 colonies appearing to be derived from asexual reproduction (Table 2). The clonal diversity for each genetic lineage and site was relatively high, especially considering the P_{SEX} value ($R = 0.75-1.00$) (Table 2).

We retained 176 multilocus genotypes after removing the six duplicate genotypes of clonemates. We calculated the allelic richness (A_R), standardized against 21 multilocus genotypes (i.e., $N_{MLG} = 21$; lowest number of multilocus genotypes). We also calculated the expected heterozygosity (H_E) at each site and for each genetic lineage to estimate the genetic diversity. The results for Seriatopora-A colonies at O-OUR and Seriatopora-C colonies at

Y-IRO indicated that these populations were maintained under low genetic diversity (Table 3). The genetic differentiation indexes, the pairwise F_{ST} and G_{ST} , were 0.064–0.116 (all $P = 0.001$) and 0.107–0.209 (all $P = 0.001$), respectively, among the three sites (with 20 or more retained multilocus microsatellite genotypes) within *Seriatopora*-B (ESM Fig. S4). The genetic structure was established for each genetic lineage without the LOCPRIOR model (*Seriatopora*-A: best $K = 2$, $\Delta K = 188.38$; *Seriatopora*-B: best $K = 2$, $\Delta K = 35.80$; *Seriatopora*-C: best $K = 2$, $\Delta K = 790.50$) and with the LOCPRIOR model (*Seriatopora*-A: best $K = 2$, $\Delta K = 284.83$; *Seriatopora*-B: best $K = 2$, $\Delta K = 39.54$; *Seriatopora*-C: best $K = 2$, $\Delta K = 67.90$) (ESM Fig. S4). The migration estimation using GeneClass2 indicated that migration from other sites was rare within the analyzed genetic lineages. Only one migrant was found in *Seriatopora*-B (to M-IKM from an unsampled site); therefore, the migration direction was not estimated.

Discussion

Population genetic analyses using microsatellite markers revealed that there are at least three distinct genetic lineages and population genetic structure of *Seriatopora* corals in the Ryukyu Archipelago. However, the phylogenetic trees that were based on mtCR

haplotypes did not always correspond to the genetic structure that was derived using microsatellites. The community composition of genetic lineages was variable among the sites, and only Y-TKT included all three genetic lineages in our data set. Considering the P_{SEX} value, the clonal diversity was high in all three genetic lineages, and clonal replicates were found in only six of our 182 genotyped colonies. Therefore, our data suggest that sexual reproduction is the main contributor to population maintenance of *Seriatopora* in the Ryukyu Archipelago. Nevertheless, there was large genetic differentiation among the populations within *Seriatopora*-B. This suggests that larval recruitment from other populations rarely occurred over the long-term history in the lineage.

Genetic lineages of *Seriatopora* in the Ryukyu Archipelago

We identified and collected all specimens of *Seriatopora* spp. in the Ryukyu Archipelago, but previous studies by Flot et al. (2008) noted that morphological traits do not align with genetic clusters of *Seriatopora*. Warner et al. (2015) suggested the existence of cryptic species within *S. hystrix*; four genetic types of *S. hystrix* were detected on the GBR. Two microsatellites showed no amplification for specimens from some sites, which may also reflect the difference of species. *Seriatopora* has a very limited population size and is endangered in the Ryukyu Archipelago (in our field observation, see also van Woesik et al.

2011). Therefore, it may take considerable time or not be possible to identify more genetic lineages due to the difficulty of collecting samples from only a limited number of populations or colonies even if more minor lineages are distributed in this region.

On the phylogenetic tree, the Ser-1 haplotype of the mtCR, found only at the O-OUR site, appeared to be the most recent common ancestor type for Ser-2, Ser-3, and Ser-4. Tandem repeats in this locus were about twice as long in Ser-1 (8.5 times) as in haplotypes Ser-2, Ser-3, and Ser-4 (4.4 times). However, these mitochondrial groupings revealed by the phylogenetic tree and tandem repeats were not necessarily supported by levels of genetic differentiation and Bayesian clustering using microsatellite genotypes, which suggests that *Seriatopora-A* is one genetic lineage, comprised of haplotypes Ser-1 and Ser-2. Although Chen et al. (2008) and Flot et al. (2008) investigated and registered the D-loop, a control region between *atp8* and *cox1*, sequences on GenBank, the number of sequences is fewer than that of the mtCR and therefore it is not possible to use the D-loop to compare haplotypes from multiple regions on the phylogenetic tree. By increasing the number of registered sequences and using a greater number of genetic loci and amino acid sequences, including other nuclear loci, it may be possible to resolve the evolutionary processes and relationships among *Seriatopora* species.

Habitat selection related to depth and geographical location

Although we could not perform statistical analysis along depth gradients due to limited sample sizes, we found that *Seriatopora*-A appeared to occur mostly at relatively deep sites (>13 m) compared to the distributions of the other two lineages. In particular, one of the two associated haplotypes (Ser-1) was only found at ≥ 30 m (O- OUR). Our data set indicated that all three genetic lineages of *Seriatopora* inhabit both shallow (2–10 m) and deep (30–32 m) habitats. Flot et al. (2008) also confirmed that all genetically different clusters of *Seriatopora* were distributed in various water depths (Cluster 1: 6.7–40.3 m; Cluster 2: 0.6–34.0 m; Cluster 3: 0.5–30.2 m; Cluster 4: 1.0–34.0 m). Recent studies, however, have suggested that there is strong genetic structuring with depth in *Seriatopora* populations (Bongaerts et al. 2010; van Oppen et al. 2011). Larval habitat selection for settlement among depths is an important factor in determining coral distribution patterns (Baird et al. 2003). Further sampling efforts including expanding or changing the focal area may clarify any potential depth structure among lineages.

Our data did not indicate habitat differentiation between sheltered vs. exposed sites, or among latitudes. However, the community composition of genetic lineages was clearly variable among the sites sampled, although genetic lineages of *Seriatopora* were separated

between sheltered and exposed sites on the GBR, and habitat selection been found to be related to the symbionts hosted (Warner et al. 2015). The geographical structure of the Ryukyu Archipelago consists of a series of isolated reefs that may restrict the genetic connectivity among populations and the homogeneity of community composition of genetic lineages in each population. In addition, there are populations with only one cryptic species in the large habitat of the GBR (Warner et al. 2015). The relationship between species distribution and geographical distance is also somewhat complicated in the Ryukyu Archipelago. For example, while K-TKS and K-AKA are geographically isolated from each other by approximately 6 km, the community composition of the genetic lineage is completely different. Conversely, one haplotype (Ser-2) in *Seriatopora-A* was shared between O-OUR and Y-TKT, regions that are isolated by approximately 470 km. Historical reef events and other environmental factors likely determine the distributions of the different lineages. For example, Y-TKT is located on Sekisei Reef, which constitutes the largest reef area and is also the only site we sampled that contained corals from all three lineages. Non-random sampling, standardization of site size and sample density, and further standardized demographic, ecological and biological information for each population will be helpful to solve the outstanding questions about habitat selection and population dynamics of

Seriatopora.

Population connectivity and the reproductive system of *Seriatopora*

We found large genetic differentiation among sites in *Seriatopora*-B. Previous studies have also found a high degree of genetic subdivision among the populations of *S. hystrix* (Ayre and Hughes 2000, 2004; Underwood et al. 2007, 2009; van Oppen et al. 2008; Noreen et al. 2009), which corresponds to the findings of this study. Some previous studies showed complex distribution patterns of multiple genetic clusters unrelated to geographical location (e.g., Noreen et al. 2009), but may not have divided the genetic lineage and may have overestimated the genetic differentiation among sites. Nevertheless, large genetic differentiation might be related to the low dispersal ability of the brooding coral *Seriatopora*. The low dispersal ability is because the larvae settle soon after release from the natal colony, the peak of settlement is within 1 d, and almost all larvae are settled within 5 d of spawning (Atoda 1951; Ayre and Hughes 2000). The range of almost all dispersal in *Seriatopora* will occur at the local habitat scale.

Mean larval duration of *Seriatopora* is much shorter than for broadcast-spawning species such as *Acropora*. The settlement peak of *Acropora* is 5 to 8 d after spawning (Suzuki et al. 2011), and the maximum rearing durations of two *Acropora* species have been found to

be more than 50 d under artificial rearing conditions (Nishikawa and Sakai 2005). However, some brooding corals have been found to have the potential for long distance dispersal with successful settlement competency suggested by both experiment and model predictions (e.g., approximately 100 d in *Pocillopora damicornis*; Richmond 1987). The genetic differentiation of *Acropora* in this region has often been shown to be significant, but lower overall among islands (Nishikawa et al. 2003; Nishikawa and Sakai 2005; Nakajima et al. 2010; Zayasu et al. 2016). Estimates of migration in the present study did not reveal the source population of recruitment. Migration analysis also suggested that larval dispersal of *Seriatopora* occurs mostly within the local site. Nevertheless, gene flow among sites within a genetic lineage might contribute to long-term population persistence because few long-distance migrants are likely between populations of *S. hystrix* (van Oppen et al. 2008, Noreen et al. 2009). Inbreeding by self-recruiting in a restricted geographical range may enhance the fixation of a small number of genotypes because most larval recruitment occurs within 100 m of the natal habitat (Underwood et al. 2007), but oceanographic conditions and high reproductive output may facilitate the occasional dispersal of brooded larvae over 10 km for *S. hystrix* (Underwood et al. 2009).

Clonal diversity in the Ryukyu Archipelago suggests that all three genetic lineages

have been maintained by sexual reproduction. Recruitment by migration of *S. hystrix* on the GBR was mainly by sexually produced larvae (van Oppen et al. 2008). However, some multilocus genotypes were the same among colonies, indicating that populations are maintained with low genetic diversity (*Seriatopora*-A in O-OUR and *Seriatopora*-C in Y-IRO). While there is no obvious relationship between genetic diversity by neutral loci and rarity in corals (Richards and van Oppen 2012), populations including these colonies may be endangered due to low genotypic variation because genetic diversity is an important index for evolutionary responses to rapid climate change (Ayre and Hughes 2004). More populations of *Seriatopora* should be surveyed for further estimation of genetic diversity and gene flow within each lineage, especially from the southern region near the center of the geographical distribution including the Coral Triangle. Different reef systems should be examined to understand overall genetic diversity and differentiation and the possibility for migration

When the population size of *Seriatopora* is unusually large and the colonies are also large, large numbers of larvae are produced and larvae that reach other reefs probably help to establish and maintain populations (van Oppen et al. 2008). Reef areas in the Ryukyu Archipelago are fewer and smaller than those in other regions such as the Coral Triangle and GBR, and therefore the population size of *Seriatopora* will not be large compared with

populations on large reefs. This difference in the reef system is likely to dictate the number of migrants between sites. Using individual-based ecological modeling with ecological parameters for a *Seriatopora* population in the Ryukyu Archipelago, constant recruitment or occasional abundant recruitment from other populations was found to be necessary to recover the population after disturbances (Muko et al. 2014). Our population genetic analysis found evidence of only a single migrant in three populations of *Seriatopora*-B. Without continuous recruitment from other populations, *Seriatopora* in the Ryukyu Archipelago is in danger of extinction locally if the local habitat suffers from disturbances.

Acknowledgments

This study was supported by a grant from the Japan Society for the Promotion of Science (JSPS) KAKENHI (25840152 to Y.N.) and a Pro Natura Foundation grant for natural conservation of the Nansei Islands to Y.N. We express our gratitude for financial support from the Okinawa Institute of Science and Technology Graduate University. We are grateful to anonymous reviewers for their helpful comments on the manuscript. All coral specimens were collected in strict accordance with good animal practice as defined by the relevant national and/or local animal welfare bodies, and all sampling was conducted with the

appropriate permissions within the Okinawa prefecture in Japan.

References

- Arnaud-Haond S, Belkhir K (2007) GENCLONE: A computer program to analyse genotypic data, test for clonality and describe spatial clonal organization. *Mol Ecol Notes* 7:15–17
- Atoda K (1951) The larva and postlarval development of the reef-building corals. V. *Seriatopora hystrix* Dana. The Science Reports of the Tohoku University, Series 4, pp 33–39
- Avise JC, Arnold J, Ball RM, Bermingham E, Lamb T, Neigel JE, Reeb CA, Saunders NC (1987) Intraspecific phylogeography: the mitochondrial DNA bridge between population genetics and systematics. *Ann Rev Ecol Syst* 18:489–522
- Ayre DJ, Hughes TP (2000) Genotypic diversity and gene flow in brooding and spawning corals along the Great Barrier Reef, Australia. *Evolution* 54:1590–1605
- Ayre DJ, Hughes TP (2004) Climate change, genotypic diversity and gene flow in reef-building corals. *Ecol Lett* 7:273–278
- Baird AH, Babcock RC, Mundy CP (2003) Habitat selection by larvae influences the depth distribution of six common coral species. *Mar Ecol Prog Ser* 252:289–293
- Bay LK, Caley MJ, Crozier RH (2008) Meta-population structure in a coral reef fish demonstrated by genetic data on patterns of migration, extinction and re-colonisation. *BMC Evol Biol* 8:248
- Bickford D, Lohman DJ, Sodhi NS, Ng PKL, Meier R, Winker K, Ingram KK, Das I (2007) Cryptic species as a window on diversity and conservation. *Trends Ecol Evol* 22:148–155
- Benson G (1999) Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res* 27:573–580
- Bongaerts P, Riginos C, Ridgway T, Sampayo EM, van Oppen MJH, Englebert N, Vermeulen F, Hoegh-Guldberg O (2010) Genetic divergence across habitats in the widespread coral *Seriatopora hystrix* and its associated *Symbiodinium*. *PLoS One* 5:e10871
- Chen C, Dai C-F, Plathong S, Chiou C-Y, Chen CA (2008) The complete mitochondrial genomes of needle corals, *Seriatopora* spp. (Scleractinia: Pocilloporidae): an idiosyncratic *atp8*, duplicated *trnW* gene, and hypervariable regions used to determine

species phylogenies and recently diverged populations. *Mol Phylogenet Evol* 46:19–33

- Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet Resour* 4:359–361
- Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software Structure: a simulation study. *Mol Ecol* 14:2611–2620
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131:479–491
- Falush D, Stephens M, Pritchard JK (2003) Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* 164:1567–1587
- Flot J-F, Tillier S (2007) The mitochondrial genome of *Pocillopora* (Cnidaria: Scleractinia) contains two variable regions: the putative D-loop and a novel ORF of unknown function. *Gene* 401:80–87
- Flot J-F, Licuanan WY, Nakano Y, Payri C, Cruaud C, Tillier S (2008) Mitochondrial sequences of *Seriatopora* corals show little agreement with morphology and reveal the duplication of a tRNA gene near the control region. *Coral Reefs* 27:789–794
- Flot J-F, Blanchot J, Charpy L, Cruaud C, Licuanan W, Nakano Y, Payri C, Tillier S (2011) Incongruence between morphotypes and genetically delimited species in the coral genus *Stylophora*: phenotypic plasticity, morphological convergence, morphological stasis or interspecific hybridization? *BMC Ecol* 11:22
- Fukami H, Omori M, Hatta M (2000) Phylogenetic relationships in the coral family Acroporidae, reassessed by inference from mitochondrial genes. *Zool Sci* 17:689–696
- Goudet J (1995) FSTAT (version 1.2): a computer program to calculate F-statistics. *J Hered* 86:485–486
- Hebert PDN, Cywinska A, Ball SL (2003) Biological identifications through DNA barcodes. *Proc R Soc Lond B Biol Sci* 270:313–321
- Hoegh-Guldberg O (1999) Coral bleaching, climate change, and the future of the world's coral reefs. *Marine and Freshwater Research* 50:839–866
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, Hatziolos ME (2007) Coral reefs under rapid climate change and ocean acidification. *Science* 318:1737–1742
- Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R,

- Hoegh-Guldberg O, Jackson JBC, Kleypas J, Lough JM, Marshall P, Nyström M, Palumbi SR, Pandolfi JM, Rosen B, Roughgarden J (2003) Climate change, human impacts, and the resilience of coral reefs. *Science* 301:929–933
- Jombart T, Devillard S, Balloux F (2010) Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genet* 11:94
- Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I (2015) CLUMPAK: a program for identifying clustering modes and packaging population structure inferences across *K*. *Mol Ecol Resour* 15:1179–1191
- Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H, van Woesik R (2001) Coral bleaching: the winners and the losers. *Ecol Lett* 4:122–131
- Maier E, Tollrian R, Nürnberger B (2001) Development of species-specific markers in an organism with endosymbionts: microsatellites in the scleractinian coral *Seriatopora hystrix*. *Mol Ecol Notes* 1:157–159
- Meirmans PG, Hedrick PW (2011) Assessing population structure: F_{ST} and related measures. *Mol Ecol Resour* 11:5–18
- Muko S, Arakaki S, Tamai R, Sakai K (2014) An individual-based model for population viability analysis of the brooding coral *Seriatopora hystrix*. *Ecol Model* 277:68–76
- Nakajima Y, Nishikawa A, Iguchi A, Sakai K (2010) Gene flow and genetic diversity of a broadcast-spawning coral in northern peripheral populations. *PLoS One* 5:e11149
- Nakajima Y, Nishikawa A, Iguchi A, Sakai K (2012) The population genetic approach delineates the species boundary of reproductively isolated corymbose acroporid corals. *Mol Phylogenet Evol* 63:527–531
- Nakajima Y, Zayasu Y, Shinzato C, Satoh N, Mitarai S (2016) Genetic differentiation and connectivity of morphological types of the broadcast-spawning coral *Galaxea fascicularis* in the Nansei Islands, Japan. *Ecol Evol* 6:1457–1469
- Nishihira M, Veron JEN (1995) Hermatypic corals of Japan. Kaiyusha, Tokyo, Japan (in Japanese)
- Nishikawa A, Sakai K (2005) Settlement-competency period of planulae and genetic differentiation of the scleractinian coral *Acropora digitifera*. *Zool Sci* 22:391–399
- Nishikawa A, Katoh M, Sakai K (2003) Larval settlement rates and gene flow of broadcast-spawning (*Acropora tenuis*) and planula-brooding (*Stylophora pistillata*) corals. *Mar Ecol Prog Ser* 256:87–97
- Noreen AME, Harrison PL, van Oppen MJH (2009) Genetic diversity and connectivity in a brooding reef coral at the limit of its distribution. *Proc R Soc Lond B Biol Sci* 276:3927–3935

- Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes* 6:288–295
- Pineda J, Hare JA, Sponaugle S (2007) Larval transport and dispersal in the coastal ocean and consequences for population connectivity. *Oceanography* 20:22–39
- Pinzón JH, LaJeunesse TC (2011) Species delimitation of common reef corals in the genus *Pocillopora* using nucleotide sequence phylogenies, population genetics and symbiosis ecology. *Mol Ecol* 20:311–325
- Pinzón JH, Sampayo E, Cox E, Chauka LJ, Chen CA, Voolstra CR, LaJeunesse TC (2013) Blind to morphology: genetics identifies several widespread ecologically common species and few endemics among Indo-Pacific cauliflower corals (*Pocillopora*, Scleractinia). *J Biogeogr* 40:1595–1608
- Piry S, Alapetite A, Cornuet J-M, Paetkau D, Baudouin L, Estoup A (2004) GENECLASS2: a software for genetic assignment and first-generation migrant detection. *J Hered* 95:536–539
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155:945–959
- Rannala B, Mountain JL (1997) Detecting immigration by using multilocus genotypes. *Proc Natl Acad Sci U S A* 94:9197–9201
- Richards ZT, Hobbs JPA (2014) Predicting coral species richness: the effect of input variables, diversity and scale. *PLoS One* 9:e83965
- Richards ZT, van Oppen MJH (2012) Rarity and genetic diversity in Indo-Pacific *Acropora* corals. *Ecol Evol* 2:1867–1888
- Richards ZT, Miller DJ, Wallace CC (2013) Molecular phylogenetics of geographically restricted *Acropora* species: implications for threatened species conservation. *Mol Phylogenet Evol* 69:837–851
- Richards ZT, Berry O, van Oppen MJH (2016) Cryptic genetic divergence within threatened species of *Acropora* coral from the Indian and Pacific Oceans. *Conserv Genet* 17:577–591
- Richmond RH (1987) Energetics, competency, and long-distance dispersal of planula larvae of the coral *Pocillopora damicornis*. *Mar Biol* 93:527–533
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* 61:539–542

- Schmidt-Roach S, Miller KJ, Lundgren P, Andreakis N (2014) With eyes wide open: a revision of species within and closely related to the *Pocillopora damicornis* species complex (Scleractinia; Pocilloporidae) using morphology and genetics. *Zool J Linn Soc* 170:1–33
- Schmidt-Roach S, Lundgren P, Miller KJ, Gerlach G, Noreen AME, Andreakis N (2013) Assessing hidden species diversity in the coral *Pocillopora damicornis* from eastern Australia. *Coral Reefs* 32:161–172
- Suzuki G, Arakaki S, Hayashibara T (2011) Rapid in situ settlement following spawning by *Acropora* corals at Ishigaki, southern Japan. *Mar Ecol Prog Ser* 421:131–138
- Suzuki G, Keshavmurthy S, Hayashibara T, Wallace CC, Shirayama Y, Chen CA, Fukami H (2016) Genetic evidence of peripheral isolation and low diversity in marginal populations of the *Acropora hyacinthus* complex. *Coral Reefs* 35:1419–1432
- Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729
- The Japanese Coral Reef Society, Ministry of the Environment (2004) Coral reefs of Japan. Ministry of the Environment, Tokyo, Japan
- Underwood JN, Smith LD, van Oppen MJH, Gilmour J (2007) Multiple scales of genetic connectivity in a brooding coral on isolated reefs following catastrophic bleaching. *Mol Ecol* 16:771–784
- Underwood JN, Smith LD, van Oppen MJH, Gilmour JP (2009) Ecologically relevant dispersal of corals on isolated reefs: implications for managing resilience. *Ecol Appl* 19:18–29
- Underwood JN, Souter PB, Ballment ER, Lutz AH, van Oppen MJH (2006) Development of 10 polymorphic microsatellite markers from herbicide-bleached tissues of the brooding pocilloporid coral *Seriatopora hystrix*. *Mol Ecol Notes* 6:176–178
- van Oppen MJH, Willis BL, Miller DJ (1999) Atypically low rate of cytochrome b evolution in the scleractinian coral genus *Acropora*. *Proc R Soc Lond B Biol Sci* 266:179–183
- van Oppen MJH, McDonald BJ, Willis BL, Miller DJ (2001) The evolutionary history of the coral genus *Acropora* (Scleractinia, Cnidaria) based on a mitochondrial and a nuclear marker: reticulation, incomplete lineage sorting, or morphological convergence? *Mol Biol Evol* 18:1315–1329
- van Oppen MJH, Lutz A, De'ath G, Peplow L, Kininmonth S (2008) Genetic traces of recent long-distance dispersal in a predominantly self-recruiting coral. *PLoS One* 3:e3401
- van Oppen MJH, Bongaerts P, Underwood JN, Peplow LM, Cooper TF (2011) The role of deep reefs in shallow reef recovery: an assessment of vertical connectivity in a

- brooding coral from west and east Australia. *Mol Ecol* 20:1647–1660
- van Woesik R, Sakai K, Ganase A, Loya Y (2011) Revisiting the winners and the losers a decade after coral bleaching. *Mar Ecol Prog Ser* 434:67–76
- Veron JEN (2000) *Corals of the world*. Australian Institute of Marine Science, Townsville, Australia
- Warner PA, van Oppen MJH, Willis BL (2015) Unexpected cryptic species diversity in the widespread coral *Seriatopora hystrix* masks spatial-genetic patterns of connectivity. *Mol Ecol* 24:2993–3008
- Zayasu Y, Nakajima Y, Sakai K, Suzuki G, Satoh N, Shinzato C (2016) Unexpectedly complex gradation of coral population structure in the Nansei Islands, Japan. *Ecol Evol* 6:5491–5505

Figure legends

Fig. 1 Map of the sampling sites, depths, numbers of samples, and number of haplotypes of the mitochondrial control region (mtCR) of *Seriatopora* in the Ryukyu Archipelago. Location data, sample size, and the number of colonies of each genetic lineage are shown in ESM Table S1

Fig. 2 STRUCTURE analysis for 182 microsatellite multilocus genotypes. Analyses for each genetic lineage were conducted with 176 genotypes after removing the duplicate genotypes of clonemates. Probable cluster number, shown as K , for each site (**a**) was assumed as three ($K = 3$) based on the method described by Evanno et al. (2005). Probable cluster number K for each genetic lineage (**b**) was assumed as $K = 3$ to 6. The arrow shows the undefined lineage detected by this analysis

Fig. 3 Maximum likelihood phylogenetic tree composed of 30 sequences using the mtCR of *Seriatopora* and one sequence of *Pocillopora damicornis* as an outgroup. Values on the branches indicate bootstrap values (%). Twenty-three haplotypes shown by their GenBank accession numbers were cited from Chen et al. (2008), Flot et al. (2008), Bongaerts et al. (2010), and Warner et al. (2015). Sequences from other references were selected to avoid duplication of sequence, region, or publication






Fig. 2





Fig. 3



Hierarchical group	Source	d.f.	SS	Var. (%)	<i>F</i> -statistics	<i>P</i> value
Lineages-haplotypes ($N_{MLG} = 170$)	Among lineages	2	197.778	0.794 (35%)	$F_{RT} = 0.352$	0.001
	Among haplotypes within lineages	2	22.291	0.158 (7%)	$F_{SR} = 0.108$	0.001
	Within haplotypes	335	436.502	1.303 (58%)	$F_{ST} = 0.422$	0.001
	Total	339	656.571	2.255 (100%)		
Lineages-sites ($N_{MLG} = 171$)	Among lineages	2	206.244	0.788 (35%)	$F_{RT} = 0.349$	0.001
	Among sites within lineages	10	78.473	0.297 (13%)	$F_{SR} = 0.202$	0.001
	Within sites	341	399.218	1.171 (52%)	$F_{ST} = 0.481$	0.001
	Total	353	683.935	2.256 (100%)		

Table 1

Code	Seriatorpora-A				Seriatorpora-B				Seriatorpora-C				Undefined			
	<i>N</i>	<i>G</i>	<i>N</i> _{MLG}	<i>R</i>	<i>N</i>	<i>G</i>	<i>N</i> _{MLG}	<i>R</i>	<i>N</i>	<i>G</i>	<i>N</i> _{MLG}	<i>R</i>	<i>N</i>	<i>G</i>	<i>N</i> _{MLG}	<i>R</i>
O-OUR	37	14	35	0.94	-	-	-	-	-	-	-	-	-	-	-	-
O-NKJ	7	7	7	1.00	-	-	-	-	-	-	-	-	-	-	-	-
O-ONN	5	4	4	0.75	-	-	-	-	2	2	2	1.00	-	-	-	-
O-GNW	-	-	-	-	23	22	22	0.95	-	-	-	-	-	-	-	-
K-TKS	-	-	-	-	-	-	-	-	23	21	23	1.00	-	-	-	-
K-AKA	-	-	-	-	23	23	23	1.00	-	-	-	-	-	-	-	-
M-IKM	-	-	-	-	23	21	21	0.91	-	-	-	-	-	-	-	-
Y-NGR	-	-	-	-	2	2	2	1.00	-	-	-	-	3	3	3	1.00
Y-TKT	7	7	7	1.00	10	10	10	1.00	3	3	3	1.00	2	2	2	1.00
Y-IRO	-	-	-	-	-	-	-	-	12	6	12	1.00	-	-	-	-
Total	56	32	53	0.95	81	78	78	0.96	40	32	40	1.00	5	5	5	1.00

Table 2

Genetic lineage	Site	N_{MLG}	A_R	H_O	H_E	F_{IS}
Seriatopora-A	O-OUR	35	1.66	0.151	0.178	0.165*
Seriatopora-B	O-GNW	22	4.81	0.390	0.413	0.079
	K-AKA	23	4.87	0.348	0.421	0.195**
	M-IKM	21	5.00	0.442	0.474	0.091
Seriatopora-C	K-TKS	23	2.92	0.193	0.258	0.275**

Table 3