

ORIGINAL ARTICLE

Antimicrobial susceptibility patterns of the gram-negative bacteria isolated from septicemia in Children's Medical Center, Tehran, Iran

N. RABIRAD¹, M. MOHAMMADPOOR¹, A. RASTEGAR LARI^{2,3}, A. SHOJAIE⁴, R. BAYAT³, M. ALEBOUYEH⁵

¹Paediatric Medical Center, Emam Hospital, Tehran University of Medical Sciences, Tehran, Iran; ²Razidrug Resaerch Center, Iran University of Medical Sciences, Tehran, Iran; ³Department of Microbiology, Iran University of Medical Sciences, Tehran, Iran; ⁴Tehran Heart center, Tehran University of Medical Sciences, Tehran, Iran; ⁵Department of Anesthesia, Iran University of Medical Sciences, Tehran, Iran

Key words

Antimicrobial resistance • Antimicrobial resistant profile • Bloodstream infection

Summary

Introduction. The choice of antimicrobial treatment for septicemia is often empirical and based on the knowledge of local antimicrobial activity patterns of the most common bacteria causing such bloodstream infections. The current study aimed to study the prevalence of bacterial pathogens causing septicemia and their antimicrobial resistant profiles in hospital admitted patients.

Methods. This cross sectional study done at Children's Medical Center, Tehran, Iran. We examined 168 bacterial strains isolated from 186 clinically diagnosed septicemia cases refereed at Children's Medical Center, Tehran, Iran Over a period of twelve months from July 2010 to 2011 July. 11446 blood samples from patients of clinically suggestive septicemia were evaluated.

Results. Bacterial strains were isolated from 910 (7.95%) of blood cultures. Gram-negative bacteria identified were *Pseudomonas* species (20.5%), *Pseudomonas aeruginosa* (1.86%), *Salmonella* spp (1.09%), *Acinetobacter naumannii* (8.13%), *Escherichia coli*

(4.06%), *Klebsiella* spp (5.16%). Gram-negative pathogens were more than gram positive in bloodstream infections. Antimicrobial susceptibility testing was done according to Clinical and Laboratory Standards Institute (CLSI, USA) guidelines against: amikacin, ampicillin, amoxicillin, amoxiclav, cefuroxime, cefotaxime, ceftazidime, cefoperazone, tetracycline, chloramphenicol, ciprofloxacin, gentamicin. Resistanc to different antibiotics in the most important isolated bacteria were: 32.1 %, 10.8%, 87.8%, 96%, 39.1%, 35.2, 49.4%, 69%, 80.02%, 22%, 59%, 30.1% respectively, for *Pseudomonas* spp, 32%, 3.7%, 84.2 %, 83.2%, 80.1%, 75.4%, 44.8%, 45.2%, 33.3%, 19%, 34.1, 11.5% respectively for *Acinetobacter* species.

Discussion. Resistant to majority of the antimicrobial agents for several pathogens implicated in bloodstream infections, particularly in Gram-negative bacteria, can make complication in treatment of infection cause by them.

Introduction

The increasing bloodstream infections mainly in developing countries is the one of most important health care systems concerne. Although the bloodstream infections mortality rate has reduced in latest two decades. Many studies worldwide have determined rising antibiotics resistance among causative bacteria for septicemia [1-3]. It can lead to make a major challenge for the physicians for treatment of them, because of they face to increase of antibiotics resistance in causative bacteria for septicemia [3-5].

Both Gram-negative and Gram-positive bacteria can cause bloodstream infections and that can be differs from different locality and/or different time [1, 2, 8]. Rapid and accurate diagnosis and select the appropriate antimicrobial treatment for patients with septicemia can help to decrease mortality and morbidity [7, 9]. Actually, for selection of empirical antibiotic therapy for bloodstream infections consider local antibiotic resistance patens in frequent pathogens associated with septicemia is necessary and can be one of important point.

So, this study was performed to find the frequency of Gram-negative bacteria that causing septicemia in Children's Medical Center hospital admitted patients and determined antimicrobial resistance profile till clinicians can select the best choice antibiotic therapy.

Methods

This study was performed at a Children's Medical Center, Tehran, Iran, over a period of twelve months from July 2010 to 2011 July. During the current study, 803 blood samples from patients of clinically suggestive bloodstream infection were surveyed, out of 4162 patients admitted. An informed consent was achieved from the patients. Blood samples from neonates and pediatrics were collected by pediatricians. Blood samples were gathered after careful cleaning of the venous with 70% alcohol and consequently followed by povidone iodine. Under the aseptic conditions 5 ml of blood was drawn by venipuncture and transmitted into two culture bottles each containing 50 ml of 0.5% bile-broth and 50 ml of 0.5% glucose-broth. Both the bottles were incubated

at 37°C for 10 days aerobically. Routine subculturing was performed on MacConkey agar and 5% sheep blood agar after 24 hours, 48 hours, 5th day and 10th day. In between these time points, subculturing was performed until there was observable turbidity. Microorganisms that isolated from blood culture were recognized by cultural characters, morphology and variety of standard biochemical tests such as oxidase, TSI, SIM, citrate, OF, Lysine decarboxylase, gelatinase, MR/VP, etc. [10-15]

Antimicrobial resistant patterns of the bacterial isolates to several antimicrobials was detected by using the Kirby Bauer disc diffusion method by Clinical and Laboratory Standards Institute (CLSI) guidelines [11]. The antibiotic concentration for each disc was as follows: gentamicin (10µg), amikacin (30µg), ampicillin (10µg), amoxicillin (30µg), amoxiclav (20:10µg), cefoxitin (30µg), cefotaxime (30µg), cefuroxime (30µg), ceftazidime (30µg), cefoperazone (75µg), ciprofloxacin (5µg), tetracycline (30µg) and chloramphenicol (30µg) for all organisms. All the antibiotic discs were purchased from MAST company (UK).

Results

Demographics

Over a period of twelve months from July 2010 to 2011 July. 11446 blood samples from patients of clinically suggestive septicemia were evaluated. The causative organism could be identified in 910 (7.95%) bacterial strains. Gram-negative bacteria identified were *Pseudomonas* species 187 (20.5%), *Acinetobacter baumannii* 74 (8.13%), *Klebsiella* spp 47 (5.16%), *Escherichia coli* 37 (4.06%), *Salmonella* spp 10 (1.09%), *Pseudomonas aeruginosa* 17 (1.86%) (Tab. I).

The rates of resistance in *Pseudomonas* spp to various antimicrobials are given below: gentamicin (30.1%), amikacin (10.8%), ampicillin (87.8%), amoxicillin (96%), amoxiclav (80.02%), cefuroxime (69%), cefotaxime (49.4%), cefazidime (35.2%), cefoperazone (39.1%), ciprofloxacin (22%), chloramphenicol (59%) and tetracycline (32.1 %) (Tab. II).

The rates of resistance in *Acinetobacter* species to various antimicrobials are given below: gentamicin (11.5%), amikacin (3.7%), ampicillin (84.2 %), amoxicillin (83.2%), amoxiclav (80.1%), cefuroxime (75.4%), cefotaxime (44.8%), cefazidime (45.2%), cefoperazone (33.3%), ciprofloxacin (19%), chloramphenicol (34.1%) and tetracycline (32%) (table. 2).

Tab. I. Number and percentage of isolated Species of Bacterial.

Species of isolated Bacterial	Number of isolated bacteria	Percentage of isolated bacteria
<i>Pseudomonas</i> species	187	20.5%
<i>Acinetobacter baumannii</i>	74	8.13
<i>Klebsiella</i> spp	47	5.16%
<i>Escherichia coli</i>	37	4.06%
<i>Pseudomonas aeruginosa</i>	17	1.86%
<i>Salmonella</i> spp	10	1.09%

amoxicillin (83.2%), amoxiclav (80.1%), cefuroxime (75.4%), cefotaxime (44.8%), ceftazidime (45.2%), cefoperazone (33.3%), ciprofloxacin (19%), chloramphenicol (34.1%) and tetracycline (32%) (table. 2).

The rates of resistance in *Salmonella* spp to various antimicrobials are given below: cefuroxime (21%), cefotaxime (39%), ampicillin (46.4%), amoxicillin (27.3%), amoxiclav (15.4%), cefoperazone (32.5%), ciprofloxacin (6.3%), chloramphenicol (9.1%) and tetracycline (50%) (Tab. II).

The rates of resistance in *Escherichia coli* to various antimicrobials are given below: gentamicin (25%), amikacin (18%), ampicillin (92%), amoxicillin (90.9%), amoxiclav (73.9%), cefuroxime (54.5%), cefotaxime (52.9%), cefoperazone (36.8%), ciprofloxacin (31.6%), chloramphenicol (39%) and tetracycline (33.7%) (Tab. II).

The rates of resistance in *Pseudomonas aeruginosa* to various antimicrobials are given below: gentamicin (42.2%), amikacin (30.8%), ampicillin (96.1%), amoxicillin (92.9%), amoxiclav (95.8%), cefuroxime (81%), cefotaxime (48.4%), ceftazidime (34.2%), cefoperazone (47.3%), ciprofloxacin (22.4%), chloramphenicol (32%) and tetracycline (42.6 %) (table. 2).

The rates of resistance in *Klebsiella* spp to various antimicrobials are given below: gentamicin (54%), chloramphenicol (49%) cefuroxime (77.4%), cefotaxime (52%), ampicillin (98.7%), amoxicillin (71.4%), amoxiclav (52%), cefoperazone (32%) and tetracycline (62%). *Klebsiella pneumoniae* did not show resistance to amikacin and ciprofloxacin (Tab. II).

Tab. II. Frequency of antibiotics resistance.

	gentamicin	amikacin	ampicillin	amoxicillin	amoxiclav	cefuroxime	cefotaxime	cefazidime	cefoperazone	ciprofloxacin	chloramphenicol	tetracycline
<i>Pseudomonas</i> spp.	30.1%	10.8%	87.8%	96%	80.02%	69%	49.4%	35.2%	39.1%	22%	59%	32.1%
<i>Acinetobacter</i> species	11.5%	3.7%	84.2 %	83.2%	80.1%	75.4%	44.8%	45.2%	33.3%	19%	34.1%	32%
<i>Salmonella</i> spp.	21%	39%	46.4%	27.3%	15.4%	-	-	-	32.5%	6.3%	9.1%	50%
<i>Escherichia coli</i>	25%	18%	92%	90.9%	73.9%	54.5%	52.9%	-	36.8%	31.6%	39%	39%
<i>Pseudomonas aeruginosa</i>	42.2%	30.8%	96.1%	92.9%	95.8%	81%	48.4%	34.2%	47.3%	22.4%	32%	42.6%
<i>Klebsiella</i> spp.	54%	0%	98.7%	52%	52%	77.4%	52%	-	32%	0%	49%	62%

Discussion

Bacterial antibiotics resistance is remaining as an alarming problem in the therapy of bloodstream infections [16, 17]. Bacterial bloodstream infection mostly caused by strains that are resistant to a wide range of antimicrobial agents [18]. The current study investigated the antimicrobial resistance profile of 372 Gram-negative bacteria isolated from bloodstream infections. The data demonstrated the frequency of antimicrobial resistance among bacterial pathogens isolated from bloodstream infections.

Gram-negative bacteria is the most cause of bloodstream infections in many countries [1, 2, 7, 18]. Also, this type of bacteria have been the most common contributing pathogens of bloodstream infections in the present study. This is considerable that, different etiological agents of bloodstream infections can be related to the varying demography of bloodstream infections in developing countries because of different geographical area.

In some studies demonstrated that *Acinetobacter* species, *Salmonella typhi* and *Escherichia coli* were the most common Gram-negative bacteria that involve in that bloodstream infections [19-21]. These results are approximately similar to our results in current study.

Gram-negative bacteria is more resistance to antibiotics than Gram-positive. *Acinetobacter* species demonstrated very high levels of resistance to beta-lactam antibiotics in some reports [22, 23], that it is similar to the results of current study. *Acinetobacter* species showed the lower levels of resistance to aminoglycoside and low level resistance rates to quinolones. Similarly, *Escherichia coli* demonstrated distinctly higher levels of resistance to beta-lactam antibiotics than other classes of antimicrobials. The organism demonstrated maximum resistance to ampicillin and minimum to amikacin [22, 23]. These results can support the finding of this study, because of our results of antibiotic susceptibility testing are similar to them.

Conclusion

Extensive frequency of antimicrobial resistance levels were distinguished in our study, is similar to another studies [17, 18]. The high frequency of antibiotic resistance rates in Children's Medical Center might be because of indifferently and excess of use of treatment in our country because of their easy availability [19]. The other cause might be frequency of ESBL production among Gram-negative isolates from neonatal bloodstream infection [20]. Also, other cause could be the altering patterns of antibiotic utilize and varies in lifestyle. In the light of over findings there continues an increasing requirement for new agents, though a recent study reported polymyxin are being restored to as potential choices for the therapy [21]. Proper antimicrobial treatment for bloodstream infections is important in declining morbidity and mortality among patients with bloodstream infections caused by bacteria. Therefore, accurate

microbiological diagnosis and their antimicrobial resistance profile can be very important for the rapid initiation of sufficient treatment for bloodstream infections [23].

References

- [1] Pittet D, Li N, Woolson RF, et al. *Microbiological factors influencing the outcome of nosocomial bloodstream infections: a 6-year validated, population-based model*. Clin Infect Dis 1997;24:1068-78.
- [2] Mylotte JM, Tayara A. *Blood cultures: clinical aspects and controversies*. Eur J Clin Microbiol Infect Dis 2000;19:157-63.
- [3] Reimer LG, Wilson ML, Weinstein MP. *Update on detection of bacteremia and fungemia*. Clin Microbiol Rev 1997;10:444-65.
- [4] Cleven BE, Palka-Santini M, Gielen J, et al. *Identification and characterization of bacterial pathogens causing bloodstream infections by DNA microarray*. J Clin Microbiol 2006;44:2389-97.
- [5] Mylotte JM, Tayara A. *Blood cultures: clinical aspects and controversies*. Eur J Microbiol Infect. Dis 2001;9:157-63.
- [6] Haug JB, Harthug S, Kalager T, et al. *Bloodstream infections at a Norwegian university hospital, 1974-1979 and 1988-1989: changing etiology, clinical features, and outcome*. Clin Infect Dis 1994;19:246-56.
- [7] Hsueh PR, Teng LJ, Chen CY, et al. *Pandrug-resistant Acinetobacter baumannii causing nosocomial infections in a university hospital, Taiwan*. Emerg Infect Dis 2002;8:827-32.
- [8] Castagnola E, Caviglia I, Pistorio A, et al. *Blood stream infections and invasive mycoses in children undergoing acute leukaemia treatment*. Eur J Cancer 2005;41:1439-45.
- [9] Pfaller MA, Jones RN, Doern GV, et al. *Bacterial pathogens isolated from patients with bloodstream infection: frequencies of occurrence and antimicrobial susceptibility patterns from the SENTRY Antimicrobial Surveillance Program (United States and Canada)*. Antimicrob Agents Chemother 1998;42:1762-70.
- [10] Vahdani M, Azimi L, Asghari B, et al. *Phenotypic screening of extended-spectrum β -lactamase and metallo- β -lactamase in multidrug-resistant pseudomonas aeruginosa from infected burns*. Ann Burns Fire Disasters 2012;7:78-81.
- [11] Azimi L, Rastegar Lari A, Alaghebandan R, et al. *KPC-Producer Gram Negative Bacteria Among Burned Infants In Motahari Hospital, Tehran: First Report From Iran*. Ann Burns Fire Disasters 2012;7:2.74-77.
- [12] Owlia P, Azimi L, Gholami A, et al. *ESBL- and MBL-mediated resistance in Acinetobacter baumannii: a global threat to burn patients*. Le Infezioni in Medicina 2012;3:182-7.
- [13] Arbabi L, Rahbar M, Jabbari M, et al. *Extended-spectrum β -lactamase-producing E.coli and Klebsiella pneumoniae isolated from urinary tract infections in Milad Hospital, Tehran, Iran*. HealthMED 2012;6:2818-22.
- [14] Rastegar Lari A, Validi N, Ghaffarzadeh K, et al. *Vitro activity of cefixime versus ceftizoxime against Salmonella typhi*. Pathol Biol (Paris) 1997;45:415-9.
- [15] MacFadin JF. *Biochemical Test for Identification of Medical Bacteria*. 2nd ed. USA: Waverly Press, Inc. 2000.
- [16] *Nosocomial infection rates for interhospital comparison: limitations and possible solutions. A report from the National Nosocomial Infections Surveillance (NNIS) System*. Infect Control Hosp Epidemiol 1991;12:609-21.
- [17] Amita J, Indranil R, Mahendra KG, et al. *Prevalence of extended-spectrum β -lactamase-producing gram-negative bacteria in bacteremic neonates in a tertiary care hospital*. J Med Microbiol 2003;52:421-5.
- [18] Ni-Chung L, Shu-Jen C, Ren-Bin T, et al. *Neonatal bacteremia in a Neonatal Intensive Care Unit: analysis of causative organisms and antimicrobial susceptibility*. J Chin Med Assoc 2004;67:15-20.

- [19] Karlowsky JA, Jones ME, Draghi DC, et al. *Prevalence and antimicrobial susceptibilities of bacteria isolated from blood cultures of hospitalized patients in the United States in 2002*. Ann Clin Microbiol Antimicrob 2004;3:7.
- [20] Hanberger H, Diekema D, Fluit A, et al. *Surveillance of antibiotic resistance in European ICUs*. J Hosp Infect 2001;48:161-76.
- [21] Neuhauser MM, Weinstein RA, Rydman R, et al. *Antibiotic resistance among Gram-negative bacilli in US intensive care units: implications for fluoroquinolone use*. JAMA 2003;289:885-8.
- [22] Ni-Chung L, Shu-Jen C, Ren-Bin T, et al. *Neonatal bacteremia in a Neonatal Intensive Care Unit: analysis of causative organisms and antimicrobial susceptibility*. J Chin Med Assoc 2004;67:15-20.
- [23] Weinstein M, Reller L, Murray J, et al. *The clinical significance of positive blood cultures: a comprehensive analysis of 500 episodes of bacteremia and fungemia in adults. Laboratory and epidemiologic observations*. Rev Infect Dis 1983;5:35-53.

■ Received on January 12, 2014. Accepted on January 21, 2014.

■ Correspondence: Razagh Bayat, Iran University of Medical Sciences, Tehran, Iran - Tel. +98 21 86703181 - Fax +98 21 86703183 - P.O. Box: 14515-717, Tehran, Iran - Email: azizlari@gmail.com; razaghbayatmd@gmail.com