



POLITECNICO DI TORINO  
Repository ISTITUZIONALE

Spatial investigation of wall turbulence via complex networks

*Original*

Spatial investigation of wall turbulence via complex networks / Iacobello, Giovanni; Scarsoglio, Stefania; Kuerten, J. G. M.; Ridolfi, Luca. - (2018), pp. 731-731. ((Intervento presentato al convegno The 12th European Fluid Mechanics Conference tenutosi a Vienna nel 09/09/2018 - 13/09/2018.

*Availability:*

This version is available at: 11583/2713083 since: 2018-09-17T08:53:37Z

*Publisher:*

Pierre-Emmanuel des Boscqs, Stefan Braun

*Published*

DOI:

*Terms of use:*

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# Spatial investigation of wall turbulence via complex networks

G. Iacobello\*, S. Scarsoglio\*, J. G. M. Kuerten† and L. Ridolfi‡

A turbulent channel flow numerically solved at  $Re_\tau = 180$  is studied through a complex network-based approach. By combining graph theory and statistical physics, complex networks theory recently emerged as a powerful tool to analyze complex systems, in particular to spatially characterize turbulent flows<sup>1,2</sup>. In this work, portions of volume of the physical domain are associated to the nodes of the network. Links between pairs of nodes,  $(i, j)$ , are active if their correlation coefficient,  $C_{i,j}$ , based on the streamwise velocity is (in modulus) above a suitable threshold. By doing so, the strongest kinematic inter-relations are highlighted and the spatial information of the correlation is preserved. The importance of nodes in the domain is evaluated through the *volume-weighted connectivity*,  $VWC$ , which quantifies the fraction of volume connected to a node. Accordingly,  $VWC(i)$  represents the fraction of volume strongly correlated with the domain location associated to  $i$ . Nodes with high values of  $VWC$  are the *hubs* of the network and they tend to cluster into streamwise-elongated regions of hubs, RoHs (see Fig. 1). By focusing on high- $VWC$  nodes, the network analysis reveals the presence of a high recurrence of *teleconnections*, that is long-range links between distant locations. Therefore, teleconnections represent distant parts of domain with similar kinematic information. Specifically, teleconnections mainly appear between near-wall regions and they are associated with the temporal persistence of coherent patterns, namely high- and low-speed streaks.

The proposed network-based approach provides a versatile and powerful framework to study turbulent flows with different levels of detail, ranging from a global to a local scale. Based on the observed findings, the current approach can pave the way for an enhanced spatial interpretation of the turbulence dynamics and for a systematic network-based investigation of turbulence.




Figure 1: 3D view of the regions of hubs, RoHs. Colors indicate the fraction of volume occupied by each RoH.

\*Dep. Mechanical and Aerospace Engineering, Politecnico di Torino, Turin, Italy

†Dep. Mechanical Engineering, Eindhoven University of Technology, Eindhoven, The Netherlands

‡Dep. Environmental, Land and Infrastructure Engineering, Politecnico di Torino, Turin, Italy

<sup>1</sup>Scarsoglio et al., *Int. J. Bifurcat. Chaos* **26**,1650223 (2016).

<sup>2</sup>Taira et al., *J. Fluid Mech.* **795**,R2 (2016)