

Antibacterial Properties of an Aqueous Extract of *Solanum torvum* (Solanaceae) on a Few Multidrug Resistant Bacterial Strains to Common Antibiotics

N'guessan Koffi Joseph^{a*}, Toty Abalé Anatole^b, Irié Bi Jean Sévérin^c, Konan Kouadio Fernique^d, Guessennd Nathalie^e, Abo Kouakou Jean-Claude^f

^{a,c,f}Laboratory of animal physiology, faculty of Biosciences, University of Félix Houphouët-Boigny (Abidjan, Côte d'Ivoire)

^{b,d,e}Institut Pasteur of Côte d'Ivoire (Abidjan)

^aEmail: nguessanphy@gmail.com

Abstract

This study aimed to evaluate the antibacterial activity of the aqueous extract of *Solanum torvum* leaves (EAS_t) on multidrug-resistant bacterial strains to common antibiotics. This plant is used in traditional medicine against diarrhoea. Solid agar diffusion and liquid dilution methods were respectively used to assess the sensitivity of bacterial strains to EAS_t and to determine the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC). Tests performed on ten (10) bacterial strains, isolated from various biological products, showed that *Staphylococcus aureus* 234 UB/17 and *Escherichia coli* 135 UB/17, were significantly sensitive to EAS_t, with a higher sensitivity for the latter strain. This sensitivity was shown by an inhibition of bacterial growth in solid agar and liquid medium, by a decrease in turbidity caused by EAS_t sensitive germs growth as the concentration of this plant extract increased (25 to 100 mg/ml). This demonstrate that EAS_t is an antibacterial substance. Thus, treated with EAS_t at a dose of 100 mg/ml, the MICs of *S. aureus* 234 UB/17 and *E. coli* 135 UB/17 were 6.25 and 12.5 mg/ml respectively, and the MBCs were 25 and 50 mg/ml respectively. The ratio of these bactericidal parameters (MBC/MIC) equaled to four (4) for these two bacterial strains, indicating that EAS_t is bactericidal for these bacterial strains. The antibacterial and bactericidal properties of EAS_t on multiple antibiotic resistant germs could justify the use of this plant in traditional medicine for the treatment of some bacterial infections causing diarrhoea.

Keywords: *Solanum torvum*; antibacterial; bactericidal; multidrug-resistant bacteria.

* Corresponding author.

1. Introduction

Since their discovery a century ago, antibiotics played a key role in medical therapeutic [2]. Nowadays, the upsurge and widespread of infections caused by antibiotics resistant in human populations is due to therapeutic failures. Indeed, antibiotic resistance has become a public health issue and a real concern for scientific communities, the challenge is to find new natural antibacterial agents from medicinal plants used for the treatment of infectious diseases [6].

It is the case of *Solanum torvum* (Solanaceae), used in traditional medicine against diarrhoea [4]. Previous studies showed that the leaf extract of this medicinal plant exhibited an antibacterial and antiviral activities [9] and can be used to relieve anite and shingles [3].

The objective of this work was to evaluate the antibacterial properties of *Solanum torvum* (Solanaceae). For this purpose, the effects of the aqueous leaf extract of this plant were evaluated on the *in vitro* growth of bacterial strains resistant to usual antibiotics.

2. Material and methods

2.1. Material

2.1.1. Plant material

Fresh leaves of *Solanum torvum* (Solanaceae) were collected from the district of Cocody (Abidjan, Ivory Coast). Plant material was identified and authenticated by the National Floristic Center of the University of Felix Houphouet-Boigny (Abidjan, Ivory Coast). The voucher specimen number was 8377.

2.1.2. Bacterial strains

For this study, seven (7) strains of enterobacteriaceae and three (3) strains of multidrug-resistant *Staphylococcus aureus*, provided by ASSURMI (Institut Pasteur of Côte d'Ivoire), were used (**Table1**).

[1])

ESBL : *Extended-spectrum beta-lactamases (ESBL)*

KT : Kanamycin-Tobromycin

KTG : Kanamycin-Tobromycin-Gentamycin

Meti-R : Meticillin-resistant

KGNet : Kanamycin-Gentamycin-Netilmicin

Table 1: Profiles of tested bacteria

Code	Bacterial strain	Phenotypic profiles	Origin
076 CO/17	<i>Shigella sp</i>	-Fluoroquinolone-resistant strain	Faeces
		-Betalactamine-resistant strain	
		-Phenotype KTG	
150 PI/17	<i>Salmonella para typhi B</i>	-Fluoroquinolone resistant strain	Faeces
		-Betalactamine-resistant strain	
214 CA/17	<i>Shigella flexneri</i>	-Aminioside-resistantstrain	Faeces
		-Bêtalactamine-resistantstrain	
118 UB/17	<i>Escherichia coli</i>	- <i>ESBL</i>	Blood
		- Fluoroquinolone-resistantstrain	
135 UB/17	<i>Escherichia coli</i>	<i>ESBL</i>	Blood
146 PI/17	<i>Escherichia coli</i>	<i>ESBL</i>	Blood
		- <i>ESBL</i>	
238 UB/17	<i>Escherichia coli</i>	- Fluoroquinolone-resistant strain	Blood
		- Betalactamase	
129 UB/17	<i>Staphylococcus aureus</i>	- KT Phenotype	Blood
		- Phenotype KTG	
234 UB/17	<i>Staphylococcus aureus</i>	- Meti-R Phenotype	Blood
		- KTG Phenotype	
241 UB/17	<i>Staphylococcus aureus</i>	- Meti-R Phenotype	Blood

3. Methods

3.1. Preparation of plant extract

The preparation of the aqueous extract of *Solanum torvum* (EAS_t) was carried out according to the method described by [7]. Two hundred fifty (250) g of fresh leaves of this plant were sliced into small pieces and boiled for 15 minutes in 1.5 L of distilled water. The hot decoction was filtered twice on hydrophilic cotton, then once on Wattman filter paper No. 1. Filtrate was dried in an oven at 50 °C for 72 hours. After drying, the aqueous dry

extract of *S. torvum* (EAS_t) was obtained.

3.2. Determination of the extraction yield of *Solanum torvum* (Solanaceae)

The dry matter yield was determined by calculating the ratio between the weight of the dry matter and the weight of plant material used for extraction (fresh *Solanum torvum* leaves), according to the following formula :

$$\text{Yield \%} = \frac{m_0}{m_1} \times 100$$

m_0 = Weight (g) of evaporated crude extract of *S. torvum*

m_1 = Weight (g) of fresh leaves of *S. torvum*

3.3. Evaluation of the antibacterial activity of plant extract

3.3.1. Preparation of concentration range of the aqueous extract of *Solanum torvum*

The concentration range of the aqueous extract of *Solanum torvum*(EAS_t) was prepared in six test tubes numbered from 1 to 6 and one growth control tube (CT) by twofold dilution method, according to the method described by [8]. The prepared concentrations were 100; 50; 25; 25; 12.5; 6.25 and 3.12 mg/ml for EAS_t. However, for CT, the concentration was 0 mg/ml.

3.3.2. Inoculum preparation

For the inoculum preparation, two young colonies were collected from bacterial culture using a platinum loop calibrated at 2 μ l and homogenized in 10 ml of Mueller Hinton broth, incubated for 3 hours at 37 °C, to obtain a pre-culture. Then, 0.1 ml of the pre-culture broth was collected for enterobacteriaceae and 0.3 ml for *Staphylococci* and introduced into a tube containing 10 ml of Mueller Hinton broth. This bacterial suspension was evaluated at about 10⁶ cells/ml, representing the pure inoculum (inoculum 10⁰) [15].

3.3.3. Inoculum count

The bacterial inoculum was diluted from 1/10 to 10⁻⁴ dilution. Four (4) successive dilutions, from 10⁻¹ to 10⁻⁴, were also obtained. The initial bacterial inoculum and the four successive dilutions are inoculated with a 2 μ l calibrated loop on Mueller Hinton agars with 5 cm long streaks, constituting box A [11]

3.3.4. Sensitivity test

Strains sensitivity to EAS_t was tested by the solid agar diffusion technique. Mueller Hinton agar was swabbed

as recommended by the Antibiogram Committee of the French Society of Microbiology [1]. Wells of approximately 6 mm in diameter were made in the agar using a sterile punch and each well received 80 μ l of EASt at concentrations of 100; 50; 25; 12.5; 6.25 and 3.12 mg/ml. A cup, taken as a control, received 80 μ l of sterile distilled water. After 30 minutes of pre-diffusion of the EASt at laboratory temperature (26 °C), the petri dishes were incubated at 37 °C for 24 hours. The presence or absence of an inhibition zone was observed after 24 hours of incubation [11]. The interpretation was made according to the indications of [13,5].

3.3.5. Inoculation

In a series of six (6) hemolysis tubes numbered T1 to T6, and in the growth control tube (CT), 1 ml of pure inoculum was introduced. Then 1 ml of EASt was added to the 6 test tubes according to the prepared concentration range (100; 50; 25; 25; 12.5; 6.25 and 3.12 mg/ml). The control tube (CT) corresponding to the control received, instead of plant extract, 1 ml of sterile Mueller Hinton broth.

3.3.6. Determination of the minimum inhibitory concentration (MIC)

The MIC is the lowest concentration of the substance for which there is no growth visible to the naked eye after an incubation time of 24 h (Moroh and his colleagues 2008). Its determination is made by observation of the turbidity induced by the growth of organisms studied in each tube (T1 to T6 and CT).

3.3.7. Determination of the minimum bactericidal concentration (MBC)

MBC is the lowest concentration of substance that leaves at most 0.01 % of surviving germs [12]. Using a loop calibrated at 2 μ l, the contents of the tubes in which no turbidity was observed, were collected and sown on a Mueller Hinton agar starting with the tube containing the MIC, as well as the growth control. Seeding were done by parallel striations 5 cm long on the surface of the agar (Box B).

After 24 hours of incubation in an oven at 37 °C, the number of colonies on the striae was compared to those in the inoculum count box (Box A).

4. Statistical analysis

Data analyses and graphical representations were carried out using Graph Pad InStat and Graph Pad Prism 4 softwares (Microsoft, San Diego, California, USA).

Values were expressed as mean \pm standard of deviation. Data analysis was carried out using one way analysis of variance (ANOVA ONE WAY), followed by Tukey-kramer multi-comparison test and significance of difference was observed at $p < 0.05$.

5. Results

5.1. Yield of dry extract of *Solanum torvum* (Solanaceae)

19.66 g of dry aqueous extract (EAS_t) were obtained from 250 g of fresh leaves of *S. torvum*, with an extraction yield of 7.86%.

5.2. Sensitivity of bacterial strains to aqueous extract of *Solanum torvum* (EAS_t)

The sensitivity tests carried out showed that EAS_t, at concentrations ranging from 3.12 to 100 mg/ml, has a good growth inhibitory activity on *Escherichia coli* 135 UB/17 and *Staphylococcus aureus* 234 UB/17. Indeed, compared to control, the inhibition diameters induced by EAS_t on *E. coli* 135 UB/17 at concentrations of 50 and 100 mg/ml were 14.00 ± 0.00 mm ($P < 0.01$) and 18.00 ± 1.16 mm ($P < 0.001$) respectively and, on *S. aureus* 234 UB/17, the inhibition diameter was 13.33 ± 0.33 mm ($P < 0.01$) at a concentration of 100 mg/ml. However, a low sensitivity to EAS_t was observed at some other concentrations for these two bacterial strains; the growth inhibition diameters were 6 ± 0.00 mm ($P > 0.05$). Similarly, for the eight (8) other bacterial strains tested (*Shigella sp* 076 CO/17, *S. para typhi* B 150 PI/17, *S. flexneri* 214 CA/17, *E. coli* 118 UB/17, *E. coli* 146 PI/17, *E. coli* 238 UB/17, *S. aureus* 129 UB/17 and *S. aureus* 241 UB/17), the growth inhibition diameters were all 6 ± 0.00 mm ($P > 0.05$) at concentrations ranging from 3.12 to 100 mg/ml (Table 2). Figure 1 shows the growth inhibition diameters of different bacterial strains at a concentration of 100 mg/ml.

Table 2: Growth inhibition of bacterial strains by EAS_t

	Strains and codes	Origin	Extract of <i>Solanum torvum</i>		
			From 3,12 to 25 mg/ml	50 mg/ml	100 mg/ml
Diameter of bacterial growth inhibition zones (mm)	<i>Shigella. sp</i> 076 CO/17	Faeces	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$
	<i>S. para typhi</i> B 150 PI/17	Faeces	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$
	<i>S. flexneri</i> 214 CA/17	Faeces	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$
	<i>E. coli</i> 118 UB/17	Blood	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$
	<i>E. coli</i> 135 UB/17	Blood	$6 \pm 0,00$	$14,00 \pm 0,00^{**}$	$18,00 \pm 1,16^{***}$
	<i>E. coli</i> 146 PI/17	Blood	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$
	<i>E. coli</i> 238 UB/17	Blood	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$
	<i>S. aureus</i> 129 UB/17	Blood	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$
	<i>S. aureus</i> 234 UB/17	Blood	$6 \pm 0,00$	$6 \pm 0,00$	$13,33 \pm 0,33^{**}$
	<i>S. aureus</i> 241 UB/17	Blood	$6 \pm 0,00$	$6 \pm 0,00$	$6 \pm 0,00$

$n = 3$

** $P < 0,01$

*** $P < 0,001$ compared to control




Figure 1: Sensitivity of bacterial strains to EASSt at concentration of 100 mg/ml

$n = 3$

** $P < 0,01$

*** $P < 0,001$ compared to control

5.3. Determination of antibacterial parameters (MBC and MIC) of the aqueous extract of *Solanum torvum* (EASSt)

For susceptible strains (*E. coli* 135 UB/17 and *S. aureus* 234 UB/17), as the concentration of the plant extract increased, a gradual decrease in the intensity of turbidity induced by bacterial growth was observed. *S. aureus* 234 UB/17 gave the lowest values of MIC and MBC. Indeed, at a concentration of 100 mg/ml *S. aureus* 234 UB/17 and *E. coli* 135 UB/17 respectively exhibited MICs of 6.25 and 12.5 mg/ml and CMBs of 25 and 50 mg/ml. Thus, the MBC/MIC ratios for these 2 bacterial strains sensitive to EASSt were 4 (Table 3).

Table 3: Antibacterial parameters and bactericidal activity of EASSt

Bacterial strain	MIC (mg/ml)	MBC (mg/ml)	MBC/MIC	Activity of EASSt
<i>Escherichia coli</i> 135 UB/17	12,5	50	4	Bactericidal
<i>Staphylococcus aureus</i> 234 UB/17	6,25	25	4	Bactericidal

6. Discussion

The choice of *Solanum torvum* (fresh leaves) and the extraction method (decoction) was based on testimonies collected from the population and traditional therapists in Abidjan (Côte d'Ivoire). The extraction yield of *S. torvum* is similar to that of *Justicia secunda* (Acanthaceae) [7]. According to the results, this extraction method is satisfactory for the detection of the active ingredients of *Solanum torvum* and justifies the traditional use of the plant.

This study focuses on the antibacterial activity of the aqueous extract of *Solanum torvum* (EAS_t) on the *in vitro* growth of multi-resistant bacterial strains to common antibiotics. The sensitivity test of bacterial strains showed that *Escherichia coli* 135 UB/17 and *Staphylococcus aureus* 234 UB/17 were significantly sensitive to EAS_t. These two bacterial strains growth were significantly inhibited by EAS_t, their inhibition diameter zones were greater than 10 mm. On the one hand, the growth of eight (8) other bacterial strains studied has an insignificant inhibition diameter zone of 6 mm when tested by EAS_t, so these 8 bacteria were resistant to EAS_t. Indeed, according to [13, 5], an extract has an antibacterial activity if its inhibition diameter is greater than 10 mm. On the other hand, when the inhibition diameter is less than 10 mm, the bacterium is said to be resistant. In addition, for the two susceptible bacterial strains (*E. coli* 135 UB/17 and *S. aureus* 234 UB/17), compared to the growth control, as the concentration of EAS_t increased, a decrease in turbidity caused by the growth of EAS_t susceptible organisms was observed. Inhibition of bacterial growth justifies the antibacterial activity of the aqueous extract of *S. torvum*. This helped to determine the different antibacterial parameters for susceptible strains, such as MIC and MBC. At a concentration of 100 mg/ml, the lowest values of antibacterial parameters were obtained with the methicillin-resistant (Meti-R) strain of *S. aureus* 234UB/17, with a MIC of 6.25 mg/ml and a MBC of 25 mg/ml. The highest values were obtained with the *extended-spectrum beta-lactamases* (ESBL) producing *E. coli* 135 UB/17 strain, with a MIC of 12.5 mg/ml and a MBC of 50 mg/ml. Thus, for EAS_t, *S. aureus* strain was the most sensitive, while *E. coli* strain was the least sensitive. Comparison of antibacterial properties of this extract with other anti-infective plants showed that the aqueous extract of *S. torvum* was twice as active on *E. coli* as the aqueous extract of *Harungana madagascariensis* (Hypericaceae) [8]. and the aqueous extract of *Chromolaena odorata* (Asteraceae) [10]. According to these authors, both extracts inhibited the growth of *E. coli in vitro* with CMBs at concentrations of 100 mg/ml and 117.50 mg/ml respectively, while the CMB for EAS_t was at a concentration of 50 mg/ml. The CMB/CMI ratios for both bacterial strains susceptible to EAS_t were four (4). According to [15], when the CMB/CMI ratio value is less than or equal to 4, the activity of the extract studied is qualified as bactericidal. Thus, EAS_t is a bactericidal substance for *S. aureus* 234 UB/17 and *E. coli* 135 UB/17.

The antibacterial and bactericidal properties of the aqueous extract of *Solanum torvum* on multidrug-resistant bacterial strains to common antibiotics justify the use of this plant in traditional medicine for the treatment of some bacterial infections and specifically against diarrhea of bacterial origin.

Acknowledgments

We are grateful to the Ivorian National Floristic Center (University of Félix Houphouët-Boigny Abidjan, Côte

d'Ivoire), for plant authentication, and to Institut Pasteur of Côte d'Ivoire, for providing equipment and technical support for this study.

References

- [1] CASFM/EUCAST, 2017. Comité de l'Antibiogramme de la Société Française de Microbiologie (Recommandation 2017), Edition Mars 2017, Paris (France), 128 p.
- [2] Coustès T., 2016. Loi d'avenir agricole, réglementation du médicament vétérinaire et lutte contre l'antibiorésistance. Thèse de Doctorat de Vétérinaire, Ecole Nationale Vétérinaire d'Alfort, Faculté de Médecine de Créteil (France), 106 p.
- [3] Nga E. N., Pouka C. K., Ngo B. G. P. C., Dibong S. D., Mpondo M. E., 2016. Inventaire et caractérisation des plantes médicinales utilisées en thérapeutique dans le département de la Sanaga Maritime : Ndom, Ngambe et Pouma. *Journal of Applied Biosciences*, 106: 10333-10352.
- [4] Ambé A. S. A., Ouattara D., Tiebre M-S., Vroh Bi T. A., Zirihi G., N'guessan K. E., 2015. Diversité des plantes médicinales utilisées dans le traitement traditionnel de la diarrhée sur les marchés d'Abidjan (Côte d'Ivoire). *Journal of Animal & Plant Sciences*, 26(2) : 4081-4096.
- [5] Bentabet L. N., 2015. Etude phytochimique et évaluation des activités biologiques de deux plantes *Fredolia aretioides* et *Echium vulgare* de l'ouest Algérien. Thèse de Doctorat, Université Aboubekr Belkaïd Tlemcen (Algérie), 113 p.
- [6] Saffidine K., 2015. Etude analytique et biologique des flavonoïdes extraits de *Carthamus caeruleus* L. et de *Plantago major* L. Thèse de Doctorat en Sciences, Filière : Biologie, Université Ferhat Abas-Sétif (Algérie), 132 p.
- [7] Abo K. J. C., 2013. De la plante à la molécule : toxicité, effets pharmacologiques et mécanisme d'action de *Justicia secunda* (Acanthaceae), plante antihypertensive, sur le système cardio-vasculaire de Mammifères. Thèse de Doctorat d'Etat ès Sciences Naturelles, Université Félix Houphouët-Boigny (Abidjan, Côte d'Ivoire), n° 752/2013, 351 p.
- [8] Toty A., Guessennnd N., Bahi C., Kra A. M., Otokore D. A., Dosso M., 2013. Évaluation in-vitro de l'activité antibactérienne de l'extrait aqueux de l'écorce de tronc de *Harungana madagascariensis* sur la croissance de souches multi-résistantes. *Bulletin de la Société Royale des Sciences de Liège* (Belgique), 82 : 12-21.
- [9] Zubaida Y., Ying W., Elias B., 2013. Phytochemistry and pharmacological studies on *Solanum torvum* Swartz. *Journal of Applied Pharmaceutical Science*, 3(04) : 152-160.
- [10] Agnem E. C., Medanga S. A., Abba A., Nyonbourg E., 2011. Evaluation in vitro de l'activité

antibactérienne de cinq plantes de la pharmacopée traditionnelle de l'Adamaoua (Cameron). *Journal of Experimental Biology*, 7 : 22-27.

- [11] Bssaibis F., Gmira N., Meziane M., 2009. Activité antibactérienne de *Dittrichia viscosa* (L.) W. Greuter. *Rev Microbiol Ind San et Environn*, 3(1) :44-45.
- [12] Moroh J. L. A., Dje K., Loukou Y. G., Guédé-G., 2008. Etude de l'activité antibactérienne de l'extrait acétatique (EAC) de *Morinda morindoides* (Baker) Milneredheat (Rubiaceae) sur la croissance in vitro des souches d'*Escherichia coli*. *Bulletin de la Société Royale des Sciences de Liège (Belgique)*,77(4) : 44-61.
- [13] Ponce A. G., Fritz R., Del Valle C., Roura S. I., 2003. Antibacterial activity of essential oils on the native microflora of organic Swiss chard. *Society of Food Science and Technology Elsevier*, 36 : 679-684.
- [14] Savard P. Y., 2003. Caractérisation structurale et dynamique de la bêta-lactamase TEM-1 de la bactérie *Escherichia coli* par RMN liquide. *Philosophiae*. Thèse de Doctorat de Biochimie et de Microbiologie, Faculté des Sciences et de Génie, Université Laval (Québec),224 p.
- [15] Dosso M., Faye-Kette H., 1995. Documents Techniques Antibiotiques. Université Nationale de Côte d'Ivoire, Faculté de Médecine, Département de Microbiologie, Laboratoire de Bactériologie-Virologie, (Abidjan, Côte d'Ivoire), 178 p.