



1-2017

Sharp-Tailed Grouse Nest Survival and Nest Predator Habitat Use in North Dakota's Bakken Oil Field

Paul C. Burr

Aaron C. Robinson

Randy T. Larsen

Robert A. Newman

Susan N. Ellis-Felege

University of North Dakota, susan.felege@und.edu

Follow this and additional works at: <https://commons.und.edu/bio-fac>

 Part of the [Biology Commons](#)

Recommended Citation

Burr, Paul C.; Robinson, Aaron C.; Larsen, Randy T.; Newman, Robert A.; and Ellis-Felege, Susan N., "Sharp-Tailed Grouse Nest Survival and Nest Predator Habitat Use in North Dakota's Bakken Oil Field" (2017). *Biology Faculty Publications*. 3.
<https://commons.und.edu/bio-fac/3>

This Article is brought to you for free and open access by the Department of Biology at UND Scholarly Commons. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of UND Scholarly Commons. For more information, please contact zeinebyousif@library.und.edu.

RESEARCH ARTICLE

Sharp-Tailed Grouse Nest Survival and Nest Predator Habitat Use in North Dakota's Bakken Oil Field


Paul C. Burr^{1☯✉*}, Aaron C. Robinson^{2☯}, Randy T. Larsen^{3,4☯}, Robert A. Newman^{1☯}, Susan N. Ellis-Felege^{1☯}

1 Department of Biology, University of North Dakota, Grand Forks, ND, United States of America, **2** North Dakota Game and Fish Department, Dickinson, North Dakota, United States of America, **3** Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, United States of America, **4** Monte L. Bean Life Sciences Museum, Brigham Young University, Provo, UT, United States of America

☯ These authors contributed equally to this work.

✉ Current address: Department of Wildlife, Fisheries, and Aquaculture, Mississippi State University, Mississippi State, MS, United States of America

* pcb124@msstate.edu



OPEN ACCESS

Citation: Burr PC, Robinson AC, Larsen RT, Newman RA, Ellis-Felege SN (2017) Sharp-Tailed Grouse Nest Survival and Nest Predator Habitat Use in North Dakota's Bakken Oil Field. PLoS ONE 12(1): e0170177. doi:10.1371/journal.pone.0170177

Editor: Mark S. Boyce, University of Alberta, CANADA

Received: May 12, 2016

Accepted: December 30, 2016

Published: January 12, 2017

Copyright: © 2017 Burr et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: This work was supported by Wildlife Restoration Funds, U.S. Fish and Wildlife Service (<http://wsfrprograms.fws.gov/Subpages/GrantPrograms/WR/WR.htm>) and North Dakota Game and Fish (<http://gf.nd.gov/>), Grant Number: CFDA 15-6111, SNE, RAL. It also received funding from Energy Initiative Seed Funding for Non Engineering or Geology Disciplines, University of

Abstract

Recent advancements in extraction technologies have resulted in rapid increases of gas and oil development across the United States and specifically in western North Dakota. This expansion of energy development has unknown influences on local wildlife populations and the ecological interactions within and among species. Our objectives for this study were to evaluate nest success and nest predator dynamics of sharp-tailed grouse (*Tympanuchus phasianellus*) in two study sites that represented areas of high and low energy development intensities in North Dakota. During the summers of 2012 and 2013, we monitored 163 grouse nests using radio telemetry. Of these, 90 nests also were monitored using miniature cameras to accurately determine nest fates and identify nest predators. We simultaneously conducted predator surveys using camera scent stations and occupancy modeling to estimate nest predator occurrence at each site. American badgers (*Taxidea taxus*) and striped skunks (*Mephitis mephitis*) were the primary nest predators, accounting for 56.7% of all video recorded nest depredations. Nests in our high intensity gas and oil area were 1.95 times more likely to succeed compared to our minimal intensity area. Camera monitored nests were 2.03 times more likely to succeed than non-camera monitored nests. Occupancy of mammalian nest predators was 6.9 times more likely in our study area of minimal gas and oil intensity compared to the high intensity area. Although only a correlative study, our results suggest energy development may alter the predator community, thereby increasing nest success for sharp-tailed grouse in areas of intense development, while adjacent areas may have increased predator occurrence and reduced nest success. Our study illustrates the potential influences of energy development on the nest predator—prey dynamics of sharp-tailed grouse in western North Dakota and the complexity of evaluating such impacts on wildlife.

North Dakota's Institute for Energy Studies, <https://engineering.und.edu/institute-for-energy-studies/>, to SNE; SportDog Conservation Fund, SportDOG Brand, <http://www.sportdog.com/>, to SNE; Pheasants Forever Scholarship, Pheasants Forever, <http://www.pheasantsforever.org/>, to PCB; National Science Foundation, <http://www.nsf.gov/>, Grant Number: 13197000, to SNE; University of North Dakota's Faculty Seed Grant Program, UND's Research Development and Compliance, <https://und.edu/research/resources/>, to SNE; University of North Dakota Collaborative Seed Grant, North Dakota EPSCoR (<https://www.ndepscor.ndus.edu/>) and UND's Research Development and Compliance, <https://und.edu/research/resources/>, to SNE. Co-Author Aaron Robinson from funder ND Game and Fish played a role in study design, data collection, and manuscript edits.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

The United States has been gradually reducing its reliance on imported petroleum products by significantly increasing its own production [1]. This was primarily made possible through the advent of hydraulic fracturing in conjunction with horizontal drilling [2]. These technological developments, amongst others, have increased the amount of recoverable oil within formations and has made commercial-scale energy production possible throughout the nation [2,3]. North Dakota is currently one of the leading producers of oil in the U.S. [4], with more than 9,600 active oil wells at the end of 2013, approximately double that of eight years prior, with most development occurring in the Northwest [5].

The Bakken and Three Forks formations, which span the grasslands of western North Dakota into eastern Montana and southern Saskatchewan, contain most of the oil produced in the state [6,7]. The portion of the Bakken formation contained in North Dakota alone may sustain more than an estimated 38,000 oil wells and have the potential to impact more than one seventh of the state's total land area [3]. This large-scale energy development results in economic growth and employment opportunities for the state, but also brings challenges in understanding and managing potential environmental impacts [8].

Various efforts have been made to understand how wildlife are affected while energy development continues to expand across the country [9]. Frequently this research has focused on species that pique public interest. For example, large mammals [10–14], sage-grouse [15], and various gamebirds and songbirds [16–19] all show varying signs of avoidance to structures related to energy development. In addition, lekking locations for certain grouse species appear to be sensitive to this activity [20]. However, little is currently known about the ecology of sharp-tailed grouse (*Tympanuchus phasianellus*) in the presence of energy development [8,16,20].

Sharp-tailed grouse (hereafter; sharp-tail) are a popular game bird throughout their range, and are recognized as an indicator species of grassland ecosystems health [8,21]. As such, this species is of particular concern for the U.S. Forest Service and North Dakota Game and Fish when making future prairie management decisions and understanding how landscape changes may influence grassland birds [21]. Sharp-tails are considered common throughout North Dakota, but immediate potential threats to their habitat include disturbances related to gas and oil development [8,16]. These disturbances have the potential to impact multiple aspects of sharp-tail ecology, both directly and indirectly. Common disturbances associated with gas and oil development include noise and light pollution, dust, traffic, road and housing development, and fragmentation of the landscape [3,9,16,18,22–24]. Possible influences from these disturbances on the nest survival of sharp-tails is an area of particular concern as nest survival is one important factor influencing reproductive success [25].

Similar to other prairie grouse species, sharp-tail nest failure is primarily caused by nest depredation [22,26–29]. Therefore, potential influences of energy development on the nest predator community are of equal concern when examining sharp-tail nest survival. Medium sized mammals, or meso-mammals, are common nest predators of ground nesting bird in western North Dakota [30], which primarily include coyotes (*Canis latrans*), striped skunks (*Mephitis mephitis*), American badgers (*Taxidea taxus*), raccoons (*Procyon lotor*), and red fox (*Vulpes vulpes*) [30,31]. In addition, these species often perform central roles in ecosystems as predators to a variety of other prey species [28,32–34]. However, research is lacking on how meso-mammals may be influenced by energy development. Any potential influences on the meso-mammal community may have indirect consequences for sharp-tails, as well as other prey species.

Our objectives for this study were to evaluate sharp-tail nest success and meso-mammal occupancy in northwestern North Dakota at two study areas varying in energy development.

During the summers of 2012 and 2013 we used radio-marked hens and nest cameras to estimate daily nest survival of sharp-tails and camera scent stations to estimate meso-mammal occupancy rates at each site. We hypothesized that areas of increased energy development may have less favorable nesting habitat and experience increased disturbances which has been shown to elicit avoidance behaviors, lek abandonment, and reduced adult survival for grouse species (see [16]). We predicted nests within these areas may therefore have reduced nest success compared to its less developed counterpart. Similarly, we predicted the possibly more fragmented landscape of developed areas would facilitate predator travel and increase nest discovery efficiency thereby decreasing nest success as well [35,36].

Study Areas

We selected two study areas, Belden and Blaisdell, with the goal of gathering data from areas with similar land use but differing levels of oil and gas intensity based on relative oil well densities (Fig 1). Study boundaries were constructed using 95% minimum convex polygons around sharp-tail nesting locations from 2010–2011 (A. C. Robinson, North Dakota Game and Fish Department, unpublished data). Study areas were located in Mountrail County, North Dakota and were approximately 15 kilometers apart from each other.

Belden covered 147.2 km² (centroid: N 48.107922, W -102.393517), and was classified as intense oil activity with numerous active oil wells present within and around its boundary (Fig 1). Based on data from the North Dakota Industrial Commission [5], oil well density in Belden was 0.767 wells/km² in August of 2012 and 0.950 wells/km² in August of 2013. Belden was characterized as 61% grassland/hay land, 31% cropland, 6% wetland, and 2% trees/shrubs [37].

Blaisdell represented an area of minimal oil development and covered 38.7 km² in 2012 (centroid: N 48.300744, W -102.130655), but was expanded to 158.3 km² in 2013 (centroid: N 48.262096, W -102.077418) in order to create a more equal-sized study area and increase sample sizes. We expanded Blaisdell's 95% minimum convex polygon by including sharp-tail nesting locations recorded during our first field season in 2012. We also included two additional lek locations discovered in 2012 to the polygon, and added a 3.22 km (2 mile) buffer around each of these leks to encompass potential nesting locations for sharp-tails using these leks ([38]; Fig 1). No active oil wells were within the 2012 boundary, but one oil well was within the extended 2013 boundary resulting in a density of 0.006 wells/km² [5]. Although no active drilling of oil wells occurred within Blaisdell during this study, there was activity present around the study area, primarily to the west (Fig 1). Since it was still susceptible to some disturbances associated with oil development, we considered it an area of minimal development rather than no activity. The larger, 2013 Blaisdell polygon, was characterized as 44% grassland/hay land, 45% cropland, 11% wetland, and 0% trees/shrubs [37].

All study area land cover estimations, along with lands cover metrics used in subsequent analysis, were calculated using the U.S. Fish and Wildlife Service land cover layer [37]. We estimated classification accuracy of this data by randomly generating 200 points in each study area, proportional to the amount of each land cover (S1 Table). We compared these points to high resolution imagery (sub-meter) taken and mosaicked by the National Agriculture Imagery Program in 2012. Classification accuracy at Belden was 87.0%, and Blaisdell was 85.5%, giving a combined accuracy of 86.3%.

Methods

Field Methods

Nesting ecology of sharp-tailed grouse. We captured sharp-tail hens beginning in late April of 2012 and 2013 using walk-in funnel traps at leks [39]. Five leks were trapped at each




Fig 1. Two study areas established in Mountrail County of western North Dakota used to monitor sharp-tailed grouse nests and their nest predators in 2012 and 2013. Belden served as an area of intense oil development, whereas Blaisdell served as an area of minimal oil development for comparison. The dashed line within Blaisdell represents its boundary in 2012, and points represent active oil wells during our study years.

doi:10.1371/journal.pone.0170177.g001

study area, with the exception of Blaisdell in 2013 when we included two additional leks to expand its size (see study area). We checked traps four times a day, twice in the morning and twice in the evening. We removed sharp-tails from traps using a handheld net and placed them into a small fabric kennel before data collection. All grouse were individually aged (juvenile or adult), weighed, and banded. We fit hens with a VHF necklace style radio collar (10.7 or 16.0 grams; Advanced Telemetry Systems) and released all birds at the capture site immediately after data collection was over. We tracked hens weekly from May to July by radio telemetry using hand held, vehicle mounted, and fixed winged aircraft mounted units and recorded all locations on GPS units. Once hens were found incubating a nest they were flushed to record the nest location and number of eggs. We confirmed nests remained active every 4–5 days using telemetry. If a hen was not found at her nest we then examined the nest bowl to determine if a depredation or hatching event occurred. A nest was considered successful if at least one egg hatched.

We monitored a subset of nests using 24-hour video surveillance nest cameras to accurately determine nest fates and identify nest predators. We opportunistically placed cameras at nests as they were found at the beginning of the nesting season to maximize the total number of nests video monitored. We were unable to monitor all nests due to our limited supply of cameras. Nests that received a camera later in the nesting season were systematically selected based on their general location within each study area. We placed cameras in this manner to ensure representative coverage throughout each study area, and all nests were subject to video monitoring, regardless of accessibility. Cameras were only placed on a nest once we believed incubation began (i.e. ≥ 11 eggs present; [38]). During camera installation, field technicians wore latex gloves to avoid leaving human scent. Cameras were clamped to a steel bar that was inserted into the ground approximately 0.5 m from the nest. A cord at ground level connected the camera to a digital video recorder (DVR) placed inside a waterproof box at least 30 m from the nest. The DVR and camera were powered by a 12 volt, 35 amp battery. Footage was recorded by the DVR unit and saved onto a 32-GB SD card. All camera equipment was camouflaged using paint and surrounding vegetation.

After installation of the camera was complete, we used telemetry to confirm the hen returned to its nest the following day. If the hen did not return we assumed camera presence influenced regular nesting and removed the camera. Batteries and memory cards were changed every 4–5 days to insure continuous recording. If the hen was absent from the nest at this time, we visually inspected the nest to determine if the nest was still actively being incubated. After a nest had been depredated or successfully hatched, all camera equipment was removed and placed at another active nest. All video footage was later reviewed to assess nest fates accurately, and all nest predators were identified to species if possible.

Nest Predator Surveys

We conducted predator surveys using camera-scent stations during the sharp-tail nesting season (May–July; [38]) in 2012 and 2013. Each station consisted of a PC900 Hyperfire™ Reconyx passive infrared field camera mounted approximately 1 m above the ground and placed approximately 5 m in front of a scent lure. During camera installation field technicians wore latex gloves to conceal human scent. Vegetation between the camera and scent lure was removed or reduced to create a clear line of sight for the camera. Each camera was set to take 3 consecutive photographs, 3 seconds apart. After the third photograph was taken the camera could not be triggered again for 5 minutes.

We deployed stations across each study area using a two-stage sampling design. First, we created a grid system with a cell size of 1 km by 1 km on each study area using ArcGIS (ESRI

2012, v10.1). A primary and secondary random point was generated for each grid cell along a habitat edge, as meso-mammals are thought to use such edges while traveling and foraging [35,36]. Edges were identified using the U.S. Fish and Wildlife Service [37] land cover layer in ArcGIS and were characterized as areas where water, grassland, agriculture, or trees/shrubs intersected. Each grid cell and its associated random point served as a potential location for one predator survey. Grid cells to be sampled were systematically chosen to ensure broad and even sampling across the two study areas. In some instances, selected grid cells or the primary point within the cell were not able to be sampled due to access limitations on private land. In such cases, we sampled using the secondary random point or the next closest grid cell not included in the sample.

We sampled over three separate periods spanning from 21 May to 30 July to increase coverage of the area. In 2012 each sample period lasted approximately 14 days. The three periods began on 21 May, 4 June, and 18 June, respectively. After discovering our detection probabilities were lower than expected in 2012, we increased the sample period length in 2013 [40]. Each sample period in 2013 lasted approximately 22 days and began on 24 May, 16 June, and 8 July, respectively.

Within the first half of each survey, we used a fatty acid scented predator disk (Pocatello Supply Depot, Pocatello, Idaho) to lure mammalian predators to the camera. To avoid predator acclimation to the scent of the predator disks, we replaced them half way through the survey with Caven's "Violator 7" predator lure (Minnesota Trapline Products, Inc., Pennock, Minnesota) with the goal of maintaining predator interest and increasing detection probabilities. This second scent lure was placed inside of a hollowed golf ball that was mounted on a wooden dowel and staked into the ground. The golf ball served as a visual stimulus thought to elicit explorative behavior in meso-mammals such as coyotes when in unfamiliar environments [41,42]. In the event of major precipitation, we replaced scent lures to avoid scent being washed out. We identified animals in photos to species and recorded them per sampling occasion. We classified one sampling occasion as a 72 hour period (three calendar days). We omitted detections observed during the day of installation or termination of the camera-scent station as these did not encompass a full day.

All sampling procedures, including sharp-tail capturing, handling, and camera monitoring as well as predator scent station protocols were approved by the Institutional Animal Care and Use Committee of the University of North Dakota and strictly followed their procedures (#A3917-01, Protocol 1201-2). This work was done in collaboration with the North Dakota Game and Fish department, who served as a regulatory agency for our research. All efforts were made to minimize the amount of handling time and reduce the amount of stress on sharp-tails during the study. Permission to conduct research activities on private land was obtained directly from all land owners within our study areas. We also obtained permission from the North Dakota Game and Fish department for use of all state owned land, and from the U.S. Fish and Wildlife Service for federally owned land. This field study did not involve endangered or protect species throughout its duration.

Data Analysis

Nesting ecology of Sharp-tailed grouse. We used video footage to identify nest predators, and modeled daily nest survival rates using Program MARK [43–46]. We included all nests monitored in both years, regardless of individual hen re-nests within or between years. There were 18 total re-nests (11% of all nests, 8 from Belden and 10 from Blaisdell) within a given nesting season used in the analysis. However, program MARK does not allow for random effects, so we first ran our analysis without including re-nest events, and then ran the same

analysis while including re-nests for comparison. We chose to treat each of the 18 re-nests as individual nesting events if the results were similar, primarily to maintain a high sample size. We did not include any nests that appeared to fail due to abandonment caused by researcher disturbance or camera presence.

We explored the influence of multiple covariates in our daily nest survival models (S2 Table). Study area was included as a grouping variable, and year and nest camera presence were included as binary covariates. We hypothesized oil wells and roads may influence nest survival by potentially impacting local nest predator activities. However, due to model convergence issues, we could not include these variables as continuous data. This was likely the result of low variability associated with distance to nearest road, and the fact all Blaisdell nests were a considerable distance from the closest oil well. Therefore, we included Euclidean distance to the nearest active oil well and nearest road as categorical covariates. Nearest active oil well was classified as less than 450 m, 450–1,000 m, or greater than 1,000 m from the nest. Nearest road was classified as either less than 450 m or greater than 450 m from the nest. We selected these incremental distance categories relative to the approximate 450 m average movement distance of sharp-tail hens while laying and incubating eggs [47].

We included habitat composition around nest locations using multiple spatial scales in model construction. To classify composition, we used the U.S. Fish and Wildlife Service 30 m resolution land cover layer [37] and lumped land cover categories as either water, grassland, agriculture, or trees/shrubs in ArcGIS. We calculated percent composition for each category within a 50 m, 200 m, and 450 m buffer centered at the nest, again focusing on the 450 m buffer of habitat use by hens while incubating [47]. The 50 m buffer captured differences between nests at the microsite level, and the 200 m buffer was used as an intermediate measure. Edge density (m/km^2) was also calculated at the 450 m extent with edges characterized as roads and where land cover types changed. We hypothesized this edge metric may influence survival as numerous mammalian nest predators exploit habitat edges when traveling and foraging for prey items [35,36,48,49]. We did not mix spatial extents when including habitat composition within models.

We based model selection on Akaike's Information Criterion scores corrected for small sample sizes (AIC_c) to determine which models had the most support [50]. We estimated daily nest survival rates (S), as well as individual covariate beta estimates (β) using model averaging of all models constructed [50]. We back-transformed beta estimates to their respective odds ratio (OR) for interpretation. Odds ratio confidence intervals including 1.0 are not considered statistically significant, but may be biologically important if estimates are trending away from 1.0. For this study, we refer to these potentially important biological results as trending.

Nest predator surveys. We constructed single season occupancy models in program MARK to estimate predator occupancy [46]. Because our study included only two years of data, and not all survey sites were sampled in both years, we chose to include year as a covariate in our analysis rather than using a robust model design. Moreover, our goal was to determine if differences in species occurrence existed between study areas, rather than directly modeling changes in occupancy between years.

We developed a candidate set of biologically relevant models for all species detections pooled together. Specifically, a survey location was considered occupied if any target species was photographed at least once during a sampling occasion. We did not explore individual species occupancy models due to low detections. We used combinations of multiple covariates for occupancy and detection and compared these models using AIC_c (S3 Table; [50]). We first identified the best covariates that explained the detection parameter while keeping occupancy constant in our model construction. We explored the effects of year and sampling period (time of summer) as covariates (S3 Table). Year was a binary variable with 2012 serving as the baseline for comparison. We used dummy variables to model sampling period, where period one and

period two were compared to the baseline of period three. We also evaluated if detection rates differed between individual occasions or between scent lures.

After determining which covariates best explained the detection parameter, we modeled covariates for the occupancy parameter. We examined the covariates of study area (Blaisdell as our baseline), year, Euclidean distance to nearest oil well (m), Euclidean distance to nearest road (m), and oil well density and habitat composition within a 500 m radius of the survey location (S3 Table). A 500 m buffer was chosen to limit the amount of overlap between neighboring survey locations while maintaining independence between sampling sites. We lumped habitat composition into similar land cover categories and classified them as water, grassland, agriculture, or trees/shrubs. We then calculated the percentage of area covered by each of these categories within the 500 m buffer around the survey locations in ArcGIS.

We model averaged estimates of occupancy (ψ), detection (p), and individual covariate beta's (β ; [50]). Beta estimates were back-transformed to their respective odds ratios (OR) for interpretation as previously described for nest survival. We used model deviance to assess fit of models using the most general model with the distribution of deviance values obtained from 1,000 parametric bootstrapping replicates from program MARK [51]. Program MARK is unable to perform the bootstrap procedure with the incorporation of individual covariates [52] so we selected a general model without covariates for this step.

We tested for multicollinearity among all continuous variables by calculating Pearson's correlation coefficients for both the nest survival and nest predator analyses. We took a conservative approach and did not use both covariates in the analysis if $r^2 > 0.3$ [53]. We also tested for spatial autocorrelation to verify we did not violate the assumption of spatial independence by calculating Moran's I in program Spatial Analysis in Macroecology (SAM Version 4.0, <http://www.ecoevol.ufg.br/sam/>, accessed 12 September 2013). To calculate this we used nest success (successful [0] vs. failed [1]) for the nest survival analysis and species detection (present [1] vs. not-present [0]) for the predator occupancy analysis. We assessed spatial autocorrelation by calculating Moran's I values and associated p-values for each study area and year.

Results

Nesting Ecology of Sharp-Tailed Grouse

We monitored 163 sharp-tail nests across both study areas and years (S1 Dataset). Apparent nest success at Belden (i.e. intense gas and oil development area) was 62% based on 79 nest events across years, and 44% at Blaisdell (i.e. minimum gas and oil development area) based on 84 nest events across years (Table 1). Ninety of these nests were also monitored using nest cameras, with 42 deployed at Belden and 48 at Blaisdell, across years. Overall apparent nest success was 58.9% for nests monitored with cameras, 45.2% for those not monitored with cameras (Table 1). Eleven total nest abandonments occurred across both study areas and years after the hen was initially flushed or after nest camera installation. These were not included in any subsequent analysis.

Seventy-seven nests (47.2%) failed across both study areas and years. Depredation was the leading cause of nest failures, accounting for 81% ($n = 62$) of all failed nests (Table 1). Belden had fewer depredations ($n = 19$, with 6 captured on camera) compared to Blaisdell ($n = 44$, with 24 captured on camera). Hen mortality accounted for 9% of all nest failures, followed by cattle trampling (6%), and farm machinery (4%; Table 1). Twenty-nine of the 37 nest failures captured on camera were depredation events. American badgers and skunks were the primary nest predators accounting for 30.0% ($n = 9$) and 26.7% ($n = 8$) of all recorded depredations, respectively. Raccoons were responsible for the third most depredations (16.7%, $n = 5$), all of which occurred at Blaisdell. Coyotes accounted for 10.0% of depredations ($n = 3$), followed by

Table 1. Summary of sharp-tailed grouse nests monitored in 2012 and 2013 in western North Dakota. Nests are delineated by study area, monitoring method, and categories of nest failures. Belden study area represents intense gas and oil development, whereas Blaisdell represents minimal development.

	Total Nests Monitored	Depredated	Hen Mortality	Cattle Trampling	Farm Machinery	Apparent Nest Success
All Nests	163	62	7	5	3	52.8%
By Study Area						
Blaisdell	84	43	3	0	1	44.0%
Belden	79	19	4	5	2	62.0%
By Monitoring Method						
Nest Camera	90	29	4	3	1	58.9%
Telemetry Only	73	33	3	2	2	45.2%
Combined						
Blaisdell						
Nest Camera	48	23	2	0	0	47.9%
Telemetry Only	36	20	1	0	1	38.9%
Belden						
Nest Camera	42	6	2	3	1	71.4%
Telemetry Only	37	13	2	2	1	51.4%

doi:10.1371/journal.pone.0170177.t001

red fox (6.7%, n = 2). We observed two raptor depredations of eggs, one from a northern harrier (*Circus cyaneus*) and one species unidentifiable because of vegetation in front of camera.

We found evidence of correlation among percent grass and percent agriculture at each spatial extent ($r^2 = 0.95$ for all extents). We hypothesized percent grass cover around the nest would have a larger influence on nest survival and therefore excluded percent agriculture from our analysis. In addition, we also excluded percent trees as there was extremely low variation among nest locations. In fact, 78.5% (128 out of 163) of the nest locations had 0% trees within the largest spatial extent of 450m, and average percent trees for all nests was lower than 1.4% at each spatial extent. Moran's I correlograms revealed no evidence of spatial autocorrelation ($p > 0.05$) among nest fates (i.e., success or failure).

We constructed 59 models to evaluate sharp-tail nest survival. Models sets with, and without re-nests included produced extremely similar results. We are therefore only reporting models with re-nests included here. Study area and camera presence were in the top ranked model as the best predictors describing daily nest survival rates (Table 2). These two covariates

Table 2. Models within two ΔAIC_c scores from the highest ranked daily nest survival (S) model constructed for sharp-tailed grouse in western North Dakota, 2012–2013. Relative model weight (w), likelihood estimate (L), parameter count (K), and deviance are also displayed. Null deviance for nest survival = 531.17.

Model	AIC_c	ΔAIC_c	w	L	K	Deviance
S(Study Area + Nest Cam)	520.29	0.00	0.16	1.00	3	514.28
S(Study Area + Nest Cam + 50m Grass)	520.77	0.48	0.13	0.79	4	512.75
S(Study Area + Nest Cam + 200m Grass)	521.67	1.38	0.08	0.50	4	513.65
S(Study Area + Nest Cam + 50m Water)	521.80	1.51	0.08	0.47	4	513.78
S(Study Area + Nest Cam + 450m Grass)	521.89	1.60	0.07	0.45	4	513.88
S(Study Area + Nest Cam + Year)	521.99	1.70	0.07	0.43	4	513.98
S(Study Area + Nest Cam + DistRoad)	522.06	1.77	0.07	0.41	4	514.04
S(Study Area + Nest Cam + 200m Water)	522.17	1.88	0.06	0.39	4	514.15
S(Study Area + Nest Cam + 450m Water)	522.24	1.95	0.06	0.38	4	514.23

Covariate description: Study Area, study area as a grouping variable; Nest Cam, nest camera presence on the nest; DistRoad, distance to nearest road from the nest; Year, year as a categorical variable; Also shown are specific habitat compositions within the displayed distance from the nest.

doi:10.1371/journal.pone.0170177.t002

Table 3. Model-averaged beta (β), standard error (SE), and 95% confidence interval (CI) estimates from Program MARK for all covariates included in the sharp-tailed grouse daily nest survival analysis from North Dakota, 2012–2013. Associated odds ratios (OR) and 95% confidence intervals are also calculated for interpretation of results.

Model covariate	β Estimate	β SE	β 95% LCI	β 95% UCI	Odds Ratio (OR)	OR 95% LCI	OR 95% UCI
Intercept	2.566	0.350	1.879	3.253			
Study Area*	0.669	0.267	0.147	1.191	1.952	1.158	3.292
Camera*	0.708	0.237	0.244	1.172	2.029	1.276	3.227
Year	-0.128	0.241	-0.601	0.346	0.880	0.548	1.413
Distance to Road (m)	-0.109	0.249	-0.596	0.379	0.897	0.551	1.460
Distance to Well (0-450m)	-0.091	0.616	-1.298	1.117	0.913	0.273	3.056
Distance to Well (451-1000m)	0.043	0.556	-1.047	1.134	1.044	0.351	3.107
50m Grass %	0.005	0.004	-0.003	0.013	1.005	0.997	1.013
50m Water %	-0.023	0.031	-0.084	0.037	0.977	0.919	1.038
200m Grass %	0.004	0.005	-0.005	0.013	1.004	0.995	1.013
200m Water %	-0.014	0.040	-0.092	0.064	0.986	0.912	1.067
450m Grass %	0.004	0.006	-0.007	0.015	1.004	0.993	1.015
450m Water %	0.009	0.037	-0.064	0.082	1.009	0.938	1.085

* Indicates statistically significant terms (i.e., odds ratio did not overlap 1.0)

doi:10.1371/journal.pone.0170177.t003

were also included together in the next top 11 models, containing 89% of all AIC_c weight. Additionally, study area and camera presence were included in combination or alone with a combination of other covariates in models containing over 99% of all AIC_c weight. Belden nests were 1.95 times more likely to succeed than nests at Blaisdell (Table 3). Model-averaged daily nest survival was 0.975 (95% CI = 0.963–0.984) in Belden, and 0.955 (95% CI = 0.937–0.967) in Blaisdell. Nest survival over the average 23 day incubation period of sharp-tails was 55.9% at Belden and 34.7% at Blaisdell. Camera monitored nests were 2.03 times more likely to succeed than non-camera monitored nests (Table 3).

Other covariates contained in candidate models within 2 Δ AIC_c scores from the top model included habitat composition metrics from each spatial extent, year, and distance to roads (Table 2). However, model-averaged beta estimates and associated odds ratios revealed these covariates had little influence on daily nest survival (Table 3). All models containing the covariate of edge density within 450 m failed to converge and were not reported.

Nest Predator Surveys

We deployed 62 scent-stations in 2012, and increased this to 101 in 2013 to expand study area coverage, yielding 940 separate trap occasions over both areas and years (S1 Dataset). Fifty of the original 62 survey locations were resampled in 2013. Nest predator detections are summarized in Table 4.

We found no evidence of lack of fit for any of the general models that successfully converged. Percent agriculture and percent grassland were the only covariates exhibiting multicollinearity ($r^2 = 0.912$). We used grassland habitat to examine composition on predator occurrence. We also excluded percent trees as there was extremely low variation among the scent stations similar to our nest survival analysis. We found no evidence of spatial autocorrelation for any of the individual species detections or when species were lumped together ($p > 0.05$). Due to model convergence issues, we had to omit the covariates of distance to nearest active oil well and nearest road.

Table 4. Mammalian nest predator detections recorded from camera scent-stations deployed in 2012 and 2013 at two study areas in western North Dakota. Numbers listed for each species represent the number of stations a species was detected at least one time.

	2012		2013		Total
	Belden (Intense Development)	Blaisdell (Minimal Development)	Belden (Intense Development)	Blaisdell (Minimal Development)	
Number of Stations	33	29	51	50	163
Sampling Occasions (72 hour period)	132	116	349	343	940
Coyote	5	8	22	26	61
American badger	0	6	7	13	26
Raccoon	2	3	5	16	26
Striped skunk	1	4	5	9	19
Red fox	0	1	0	2	3

doi:10.1371/journal.pone.0170177.t004

Sampling period and year were the best explanatory variables for detection while holding occupancy constant. We constructed 24 models exploring various relationships between covariates and occupancy. Model averaged estimates for daily detection probabilities for the predator community were 0.286 (95% CI = 0.231–0.347). Detections were estimated to be 1.2 times greater during the first sampling period compared with sampling period three, but the difference was not statistically significant. We found sampling period two was 1.9 times less likely to detect a meso-mammal compared to period three, and year did not show a strong influence on detection (Table 5).

Study area and year appeared in the top eight models, which contained 95.5% of the total AICc weight (Table 6). Based on model-averaged estimates, we found Belden was 6.9 times less likely to be occupied by meso-mammals than Blaisdell and occupancy was 6.5 times greater in 2013 compared to 2012 (Table 5). Occupancy for the predator community as a whole was 0.896 (95% CI = 0.647–0.976) for Blaisdell, and 0.574 (95% CI = 0.356–0.767) for Belden. No other single covariates used in the analysis showed a significant trend on occupancy (Table 5).

Table 5. Model-averaged beta (β), standard error (SE), and 95% confidence interval (CI) estimates for all covariates included within the occupancy and detection parameter based on all models constructed. Associated odds ratios (OR) and 95% confidence intervals were also calculated for result interpretation. Models include detections from coyotes, American badger, raccoons, skunks, and red fox.

Model parameter	β Estimate	β SE	β 95% LCI	β 95% UCI	Odds Ratio (OR)	OR 95% LCI	OR 95% UCI
Occupancy							
Intercept	0.770	0.958	-1.108	2.649			
Study Area*	-1.942	0.819	-3.547	-0.338	0.143	0.029	0.713
Year*	1.872	0.593	0.709	3.035	6.503	2.032	20.808
Percent Grass	-0.007	0.011	-0.029	0.015	0.993	0.972	1.015
Percent Water	0.161	0.096	-0.027	0.348	1.174	0.973	1.417
Well Density	0.039	0.206	-0.365	0.442	1.040	0.694	1.557
Detection							
Intercept	-0.748	0.390	-1.513	0.017			
Period 1	0.198	0.233	-0.257	0.654	1.219	0.773	1.924
Period 2*	-0.633	0.261	-1.144	-0.122	0.531	0.319	0.886
Year	-0.057	0.338	-0.718	0.605	0.945	0.488	1.831

* Indicates statistically significant terms (i.e., odds ratio did not overlap 1.0)

doi:10.1371/journal.pone.0170177.t005

Table 6. Candidate models within two AIC_c scores from the highest ranked model constructed for occupancy analysis in Program MARK. Relative model weight (*w*), likelihood estimate (L), parameter count (K), and deviance are also displayed. Species used in analysis include coyotes, American badger, raccoons, skunks, and red fox detected in western North Dakota, 2012–2013. Null deviance for nest predator occupancy = 928.46.

Model	AIC _c	ΔAIC _c	<i>w</i>	L	K	Deviance
p(P1 + P2 + Year) Ψ(Study Area + Year)	906.86	0.00	0.30	1.00	7	892.13
p(P1 + P2 + Year) Ψ(Study Area + PerWater + Year)	907.56	0.71	0.21	0.70	8	890.63
p(P1 + P2 + Year) Ψ(Study Area + PerGrass + Year)	908.32	1.46	0.14	0.48	8	891.38

Occupancy (Ψ) and detection (p) covariate description: Study Area, study area as a grouping variable; P1, survey period from 20 May to 18 June; P2, survey period from 19 June to 8 July; Year, year as a categorical variable; PerWater, percent water habitat within 500 meters; PerGrass, percent grass habitat within 500 meters.

doi:10.1371/journal.pone.0170177.t006

Discussion

This study is one of the first to simultaneously examine differences in both predator and prey relative to high and low levels of energy development. Although only correlative, we found a relationship between predator occurrence and nest survival relative to our study sites. Based on current literature we originally expected the direction of this relationship to be opposite of our findings. Instead, we observed higher meso-mammal occupancy and lower daily nest survival rates at our area of minimal gas and oil development (i.e., Blaisdell), and the opposite was observed at the adjacent area of intense gas and oil development (i.e., Belden). We hypothesize energy development in North Dakota is indirectly influencing sharp-tail nest success by altering local predator activity.

A number of studies have examined the effects of energy development on multiple ecological aspects of prairie grouse [54]. Most of these have reported overall negative effects such as potential reduced yearling and adult survival rates [20,55], behavioral avoidance of infrastructure [20,22,56–58], and reduced lek attendance [20,59–62]. However, sharp-tailed grouse nest success was found to be similar in areas with and without gas and oil development in the Little Missouri National Grasslands of North Dakota [63]. Similarly, no difference in nest success of greater sage-grouse was found between disturbed and undisturbed areas in Wyoming [56], and evidence of higher nest success for Greater prairie-chickens (*Tympanuchus cupido*) was found within more developed areas of Kansas [64]. In our study, apparent nest success at Belden was comparable to that reported elsewhere for sharp-tails, whereas Blaisdell was slightly lower [63,65–67].

Previous work has found a negative impact of human built infrastructure on mammalian abundance [68]. Gas and oil development introduces a variety of infrastructure such as access roads, buildings, camp sites, drill pads, and power lines [69]. Meso-mammals may be avoiding such structures and seeking out other habitat. In addition, areas under development experience large increases of vehicle traffic associated with the construction, drilling, and maintenance of oil wells [24]. This increase may reduce meso-mammal occurrence either by eliciting an avoidance behavior, or directly through mortality caused by vehicle collisions. One primary cause of mortality for many meso-mammal species is vehicle collisions [70–74]. Likewise, energy development produces high levels of chronic noise which can negatively impact a variety of different taxa [23,75]. Meso-mammals may be responding to some or all of these factors to different degrees, resulting in the reduced occurrence observed at our intense energy development study area.

Neither distances to active oil well or road were good predictors of sharp-tail daily nest survival, and were inestimable for meso-mammal occupancy. Generally, there was little variation in distance to roads among nests or predator scent-stations. This was most likely a result of the

grid system for roads that exists across our study areas with roads located approximately 1.6 km (1 mile) apart. Further, much of the well pad development occurred directly along these roads. Although not captured in our analysis, we did notice substantially more vehicle traffic across our entire Belden study area than the Blaisdell site. This may have underlying impacts on the predator community, nest site selection, and hen stress, all of which may affect nest success and other demographic rates. Distance to active oil wells was most likely correlated with study area, as all nests except one in Blaisdell were greater than 1,000 m from an oil well.

The process of gas and oil development can commonly be broken down into four general stages: exploration, drilling, production, and abandonment. During the two years this study took place, we worked in an area dominated by the drilling stage, which includes the active construction and drilling of oil wells. These activities are often blamed for the fragmentation of landscapes through the introduction of roads, well pads, buildings, power lines, and other infrastructure [3,69,76]. Eventually this area is expected to be saturated with wells and will then be in the production stage, at which point human presence is only required for regular maintenance and inspection. This progression will result in a landscape left altered to a certain degree, but likely experiencing less disturbance than occurred during our study. Such fragmentation and the associated habitat edges are often exploited by predators when foraging and can be linked to decreased nest survival of ground nesting birds [35,36,77]. If disturbances associated with gas and oil are currently displacing predators and reducing nest depredations, it is possible as energy development progresses that predators may ultimately return and reduce nest success below natural variation in the newly fragmented landscape. This idea illustrates the importance of future research continuing to assess the impacts on wildlife as the dynamic process of gas and oil development progresses through each stage.

Unlike other studies (i.e., [38,47,49,77]), habitat composition had little influence in our daily nest survival analysis. For example, the habitat composition of agriculture has been found to be particularly influential on sharp-tail nest survival at rather broad extents (i.e. 1,600 m) [47]. We did not measure such a large extent as the nests we monitored were spatially clumped together and larger buffers would have resulted in substantial overlap. Overall, composition within 450 m of nests differed little between study areas (S4 Table). Similarly, no spatial covariates measured at the 500m scale extent used in our meso-mammal analysis were strong predictors of occupancy. Our findings of the importance of study area as a covariate may be indicative of the importance of larger spatial scales for both sharp-tails and meso-mammals, or that study area as a factor better represented the composite effects of habitat composition and disturbances than any linear combination of individual factor included as independent covariates.

Similar to other prairie grouse species, we found depredation to be the leading cause of nest failure for sharp-tails [22,26–29], and like other ground-nesting species, most were depredated by meso-mammals [30]. We confirmed the identity of primary nest predators for sharp-tails in western North Dakota using nest cameras, avoiding any potential bias of predator identification from sign left at the nest [26,78,79]. American badgers and skunks were responsible for the most depredations captured on camera at either site. Unfortunately, due to the low sample size of depredated nests recorded at our Belden site, we could not confidently make inferences regarding differences in nest predator frequencies between study areas. We observed two instances of raptors depredating eggs (not just hens). We did not observe any other avian nest predators such as members of the Corvidae family, which have been reported widely for ground nesting, grassland bird species [26,30,47,80]. Unfortunately, we do not have corvid abundance data in our study area to explain this lack of depredations. However, anecdotally we did observe some crows (*Corvus brachyrhynchos*) and magpies (*Pica hudsonia*), but only in very small numbers. Given the prairie landscape and few trees, we would not expect a high

abundance of these species. This potential low abundance may explain the lack of Corvidae depredations, something that is commonly reported for grassland nesting bird species.

We observed a positive influence on daily nest survival with the presence of nest cameras. Similarly, a meta-analysis also found an overall positive effect of nest cameras on daily nest survival for a number of monitored avian species [81]. We believe two possible explanations, or a combination of each, could be driving this result. First, predators may be avoiding the novel structures of nest camera systems. If this is the case, then nest cameras may directly increase nest success by eliciting predator avoidance. Secondly, we generally deployed nest cameras later in incubation due to logistical restrictions, which may therefore be biasing our result as nests farther along in incubation are more likely to succeed [44]. Similar findings have been reported for the monitoring of greater sage-grouse nests [26,82].

Meso-mammal occupancy was consistently greater in the year 2013 compared to 2012. These findings agree with North Dakota's annual rural mail carrier survey of furbearer species [83]. These surveys encompass large geographical regions of the state and are primarily used to evaluate trends in species populations. Coyotes, striped skunks, red fox, and raccoons all showed increases in the number of observations per 1,000 miles between 2012 and 2013 in our study region, whereas badgers showed only a slight six percent drop [83]. These reports also generally do not indicate differences in predator communities existed between the two adjacent study areas historically.

Time of summer consistently influenced our ability to detect meso-mammals, and is most likely linked to species activity patterns relating to their reproductive ecology (see [31]). Detection probabilities were greatest during the beginning of summer. With the exception of badgers, the meso-mammals observed here generally begin to breed at this time, and activity levels typically increase for mate selection and foraging purposes. Detection probabilities were lowest during the mid-summer when male activity generally decreases once the breeding season concludes. Female's activity also decreases as the gestation period begins and families are typically restricted to dens as nursing takes place.

Ecological impacts of energy development have received considerable attention in recent years over the concern for the management and conservation of wildlife and their habitats. Here, we found a negative correlation with the meso-mammal community and a positive correlation with sharp-tail nest success and energy development at areas with similar habitat composition. However, the fast pace and large scale of gas and oil development occurring in western North Dakota makes before and after studies of the impacts on wildlife extremely difficult to conduct and causation problematic to establish. Further, given our lack of spatial replication at the study area level we were unable to evaluate the generality of the associations we observed. Similarly, additional factors such as chick and hen survival also have direct impacts on sharp-tail population dynamics and require further investigation.

Wildlife managers need to recognize that the process of gas and oil development is very dynamic, and each phase has differing implications for wildlife populations. Early stages of development include increased disturbances created from machinery, traffic, and human presence. Later stages of development have reduced human presence but may result in a landscape left altered. Such habitat changes may result in altered predator community dynamics that have potential impacts on local prey species found in areas of energy development. We estimated higher sharp-tail nest success at our high intensity study area and greater predation rates at the low intensity study area, demonstrating direct and indirect influences of energy development on wildlife populations. Development will continue to create significant habitat change which may influence wildlife as energy demands continue to increase. Continued research on this subject will ultimately help to understand these processes as well as mitigate impacts on local ecosystems.

Supporting Information

S1 Table. Classification accuracy estimated for the U.S. Fish and Wildlife Service land cover layer used in this study for two study areas established in Mountrail County, North Dakota. Accuracy was calculated by randomly generating 200 reference points in each study area, proportional to the amount of each land cover type, and comparing them to high resolution imagery.

(DOCX)

S2 Table. Explanatory covariates used for evaluating sharp-tailed grouse daily nest survival rates in western North Dakota, 2012–2013.

(DOCX)

S3 Table. Explanatory covariates used for analyzing occupancy and detection rates of the mammalian nest predator community in western North Dakota, 2012–2013.

(DOCX)

S4 Table. Mean habitat composition within 450 meters of sharp-tailed grouse nests monitored at Belden and Blaisdell.

(DOCX)

S1 Dataset. Sharp-tailed grouse nest data, nest predator occupancy data, and all accompanied covariate data used for analysis.

(XLSX)

Acknowledgments

We thank C. Felege, S. Fox, J. Johnson, R. Kludt, J. Kolar, A. Mattson, P. O’Neel, A. Pachl, and A. Veroline for their involvement and dedication with field work associated with the project. We appreciate the GIS consulting provided by M. Niedzielski. We also thank the U.S. Fish and Wildlife Service, particularly staff at Lostwood National Wildlife Refuge for assisting in logistics and housing during our work.

Author Contributions

Conceptualization: PCB ACR RTL SNE.

Data curation: PCB ACR SNE.

Formal analysis: PCB RAN SNE.

Funding acquisition: ACR RTL SNE.

Investigation: PCB ACR SNE.

Methodology: PCB ACR RAN SNE.

Project administration: PCB ACR SNE.

Resources: ACR RTL SNE.

Software: PCB SNE.

Supervision: PCB ACR SNE.

Validation: PCB ACR RTL SNE.

Visualization: PCB SNE.

Writing – original draft: PCB SNE.

Writing – review & editing: PCB ACR RTL RAN SNE.

References

1. U.S. Energy Information Administration. Annual Energy Outlook 2015 With Projections to 2040. U.S. Department of Energy, Washington, DC, USA. 2015.
2. Wiseman H. Untested Waters: The Rise of Hydraulic Fracturing in Oil and Gas Production and the Need to Revisit Regulation. *Fordham Environ Law Rev.* 2009; 20.
3. Mason J. Oil Production Potential of the North Dakota Bakken. *Oil Gas J.* 2012;
4. Ausick P, Sauter MB. The 10 Most Oil-Rich States. USA Today. 2013. Available: <http://www.usatoday.com/story/money/business/2013/08/03/the-most-oil-rich-states/2613497/>. Accessed 14 Jan 2014.
5. North Dakota Industrial Commission. Department of Mineral Resources, Oil and Gas Division. 2013. Available: <https://www.dmr.nd.gov/oilgas/>. Accessed 12 Feb 2012.
6. Gaswirth S, Marra K, Cook T, Charpentier RR, Gautier DL, Higley DK, et al. Assessment of Undiscovered Oil Resources in the Bakken and Three Forks Formations, Williston Basin Province, Montana, North Dakota, and South Dakota,. *US Geol Surv.* 2013; Available: <http://pubs.usgs.gov/fs/2013/3013/fs2013-3013.pdf>
7. Meissner FF. Petroleum Geology of the Bakken Formation, Williston Basin, North Dakota and Montana. Guidebook to Geology and Horizontal Drilling of the Bakken Formation. Montana Geological Society, Billings, Montana, USA. 1991.
8. Dyke S, Fryda D, Kleyer D, Williams J, Hosek B, Jensen W, et al. Potential Impacts of Oil and Gas Development on Select North Dakota Natural Resources: A Report to the Director. North Dakota Game and Fish Department. 2010.
9. Copeland HE, Doherty KE, Naugle DE, Pocewicz A, Kiesecker JM. Mapping Oil and Gas Development Potential in the US Intermountain West and Estimating Impacts to Species. *PLoS One.* 2009; 4: 1–8.
10. Nellemann C, Cameron RD. Cumulative Impacts of an Evolving Oil-field Complex on the Distribution of Calving Caribou. *Can J Zool.* 1998; 76: 1425–1430.
11. Wolfe SA, Griffith B, Wolfe CAG. Response of Reindeer and Caribou to Human Activities. *Polar Res.* 2000; 19: 63–73.
12. Sawyer H, Nielson RM, Lindzey F, McDonald LL. Winter Habitat Selection of Mule Deer Before and During Development of a Natural Gas Field Winter Habitat Selection of Mule Deer Before and During Development of a Natural Gas Field. *J Wildl Manage.* 2006; 70: 396–403.
13. Tietje WD, Ruff RL. Responses of Black Bears to Oil Development in Alberta. *Wildl Soc Bull.* 1983; 11: 99–112.
14. Van Dyke F, Klein WC. Response of Elk to Installation of Oil Wells. *J Mammal.* 1996; 77: 1028–1041.
15. Naugle DE, Doherty KE, Walker BL, Holloran MJ, Copeland HE. Energy Development and Greater Sage-grouse. *Stud Avian Biol.* 2011; 38: 505–516.
16. Beck JL. Impacts of Oil and Natural Gas on Prairie Grouse: Current Knowledge and Research Needs. National Meeting of the American Society of Mining and Reclamation. 2009. pp. 66–87.
17. Gilbert MM, Chalfoun AD. Energy Development Affects Populations of Sagebrush Songbirds in Wyoming. *J Wildl Manage.* 2011; 75: 816–824.
18. Lawson AL. Impacts of Oil and Gas Development on Wintering Grassland Birds at Padre Island National Seashore, Texas. *Southeast Nat.* 2011; 10: 303–320.
19. Thompson SJ, Johnson DH, Niemuth ND, Ribic CA. Avoidance of Unconventional Oil Wells and Roads Exacerbates Habitat Loss for Grassland Birds in the North American Great Plains. *Biol Conserv. Elsevier B.V.*; 2015; 192: 82–90.
20. Hovick TJ, Elmore RD, Dahlgren DK, Fuhlendorf SD, Engle DM. Evidence of Negative Effects of Anthropogenic Structures on Wildlife: A Review of Grouse Survival and Behaviour. *J Appl Ecol.* 2014; 51: 1680–1689.
21. U.S. Forest Service. Record of Decision for Dakota Prairie Grasslands: Final Environmental Impact Statement and Land Resource Management Plan. Forest Service, U.S. Department of Agriculture, Missoula, Montana, USA. 2002.
22. Pitman JC, Hagen C a, Robel RJ, Loughin TM, Applegate RD. Location and Success of Lesser prairie-Chicken Nests in Relation to Vegetation and Human Disturbance. *J Wildl Manage.* 2005; 69: 1259–1269.

23. Barber JR, Crooks KR, Fristrup KM. The Costs of Chronic Noise Exposure for Terrestrial Organisms. *Trends Ecol Evol.* 2010; 25: 180–189. doi: [10.1016/j.tree.2009.08.002](https://doi.org/10.1016/j.tree.2009.08.002) PMID: [19762112](https://pubmed.ncbi.nlm.nih.gov/19762112/)
24. Wilke PW, Harrell M. Assessment of Impact of Energy Development Projects on Local Roads. Conference of the Transportation Association, Edmonton, Alberta, Canada. 2011.
25. Akcakaya HR, Radeloff VCVC, Mladenoff DJ, He HS. Integrating Landscape and Metapopulation Modeling Approaches: Viability of the Sharp-Tailed Grouse in a Dynamic Landscape. *Conserv Biol.* 2004; 18: 526–537.
26. Coates PS, Connelly JW, Delehanty DJ. Predators of Greater Sage-Grouse Nests Identified by Video Monitoring. *J F Ornithol.* 2008; 79: 421–428.
27. Webb SL, Olson C V, Dzialak MR, Harju SM, Winstead JB, Lockman D. Landscape Features and Weather Influence Nest Survival of a Ground-Nesting Bird of Conservation Concern, the Greater Sage-grouse, in Human-altered Environments. *Ecol Process.* 2012; 1. Available: <http://www.ecologicalprocesses.com/content/1/1/4>
28. Ricklefs RE. An Analysis of Nesting Mortality in Birds. *Smithson Contrib to Zool.* 1969; 9: 1–48.
29. Pitman JC, Hagen CA, Jamison BE, Robel RJ, Loughin M, Applegate RD, et al. Nesting Ecology of Lesser Prairie-Chickens in Sand Sagebrush Prairie of Southwestern Kansas. *Wilson J Ornithol.* 2006; 118: 23–35.
30. Sargeant AB, Sovada MA, Greenwood RJ. Interpreting Evidence of Depredation of Duck Nests in the Prairie Pothole Region. U.S. Geological Survey, Northern Prairie Wildlife Research Center, Jamestown, ND and Ducks Unlimited, Inc., Memphis, Tennessee, USA. 1998.
31. Seabloom R. The Mammals of North Dakota. Institute for Regional Studies, North Dakota State University, Fargo, North Dakota, USA. 2011.
32. Korpimäki E, Norrdahi K. Experimental Reduction of Predators Reverses the Crash Phase of Small-Rodent Cycles. *Ecology.* 1998; 79: 2448–2455.
33. Crooks K, Soulé M. Mesopredator Release and Avifaunal Extinctions in a Fragmented System. *Nature.* 1999; 400: 563–566.
34. Palomares F, Gaona P, Ferreras P, Delibes M. Positive Effects on Game Species of Top Predators by Controlling Smaller Predator Populations: An Example with Lynx, Mongooses, and Rabbits. *Conserv Biol.* 1995; 9: 295–305.
35. Andren H. Effects of Landscape Composition on Predation Rates at Habitat Edges. In: Hansson L, Farig L, Marriam G, editors. *Mosaic Landscapes and Ecological Processes.* Springer Netherlands, London, United Kingdom; 1995. pp. 225–255.
36. Dijk WD, Thompson FR III. Landscape and Edge Effects on the Distribution of Mammalian Predators in Missouri. *J Wildl Dis.* 2000; 64: 209–216.
37. U.S. Fish and Wildlife Service. US Fish and Wildlife Service 2002 Landcover. Region 6. iso:19115:2003/19139. Available: <http://www.nd.gov/gis/>. Accessed 12 February 2012. 2002.
38. Connelly JW, Gratson MW, Resse KP. Sharp-tailed Grouse (*Tympanuchus phasianellus*). Ithica: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online. Accessed 29 May 2013. 1998.
39. Schroeder MA, Braun CE. Walk-in Traps for Capturing Greater Prairie-Chickens on Leks. *J F Ornithol.* 1991; 62: 378–385.
40. MacKenzie DI, Royle JA, Nichols JD, Pollock KH, Bailey LL, Hines JE. *Occupancy Estimation and Modeling: Inferring Patterns and Dynamics of Species Occurrence.* Academic, New York, New York, USA; 2006.
41. Windberg LA. Coyote Responses to Visual and Olfactory Stimuli Related to Familiarity with an Area. *Can J Zool.* 1996; 74: 2248–2253.
42. Harris CE, Knowlton FF. Differential Responses of Coyotes to Novel Stimuli in Familiar and Unfamiliar Settings. *Can J Zool.* 2001; 79: 2005–2013.
43. Rotella JJ, Dinsmore SJ, Shaffer TL. Modeling Nest-survival Data: A Comparison of Recently Developed Methods that can be Implemented in MARK and SAS. *Anim Biodivers Conserv.* 2004; 27: 187–205.
44. Mayfield HF. Suggestions for Calculating Nest Success. *Wilson Bull.* 1975; 87: 456–466.
45. Dinsmore SJ, White GC, Knopf FL. Advanced Techniques for Modeling Avian Nest Survival. *Ecology.* 2002; 83: 3476–3488.
46. White GC, Burnham KP. Program MARK: Survival Estimation from Populations of Marked Animals. *Bird Study.* 1999; 46: S120–S139.
47. Manzer DL, Hannon SJ. Relating Grouse Nest Success and Corvid Density To Habitat: a Multi-Scale Approach. *J Wildl Manage.* 2005; 69: 110–123.

48. Kuehl AK, Clark WR. Predator Activity Related to Landscape Features in Northern Iowa. *J Wildl Manage.* 2002; 66: 1224–1234.
49. Batary P, Baldi A. Evidence of an Edge Effect on Avian Nest Success. *Conserv Biol.* 2004; 18: 389–400.
50. Burnham KP, Anderson DR. *Model Selection and Multimodel Inference: a Practical Information-Theoretic Approach.* 2nd ed. Springer, New York, New York, USA; 2002.
51. White GC, Burnham KP, Anderson DR. Advanced Features of Program MARK. In: Field R, J. W. H. O, Sievert PR, editors. *Wildlife, Land, and People: Priorities for the 21st Century: Proceedings of the Second International Wildlife Management Congress* The Wildlife Society, Bethesda, Maryland, US. 2001. pp. 368–377.
52. Cooch E, White G. Program MARK: a Gentle Introduction. Thirteenth edition. Volume 7. Available: <http://www.phidot.org/software/mark/index.html>. Accessed 12 June 2013. 2006.
53. Moore DR, McCabe GP. *Introduction to the Practice of Statistics.* 2nd ed. W. H. Freeman and Company, New York, New York, USA; 1993.
54. Hagen C. Impacts of Energy Development on Prairie Grouse Ecology: a Research Synthesis. *Transactions of the North American Wildlife and Natural Resources Conference.* 2010. pp. 96–103.
55. Holloran MJ, Kaiser RC, Hubert W. Yearling Greater Sage-Grouse Response to Energy Development in Wyoming. *J Wildl Manage.* 2010; 74: 65–72.
56. Lyon AG, Anderson SH. Potential Gas Development Impacts on Sage Grouse Nest Initiation and Movement. *Wildl Soc Bull.* 2003; 31: 486–491.
57. Doherty KE, Naugle DE, Walker BL, Graham JM. Greater Sage-Grouse Winter Habitat Selection and Energy Development. *J Wildl Manage.* 2008; 72: 187–195.
58. Hagen C, Pitman JC, Loughin TM, Sandercock BK, Robel RJ, Applegate RD. Impacts of Anthropogenic Features on Habitat Use by Lesser Prairie-Chickens. *Stud Avian Biol No* 39. 2011; 63–76.
59. Walker BL, Naugle DE, Doherty KE. Greater Sage-Grouse Population Response to Energy Development and Habitat Loss. *J Wildl Manage.* 2007; 71: 2644–2654.
60. Blickley JL, Blackwood D, Patricelli GL. Experimental Evidence for the Effects of Chronic Anthropogenic Noise on Abundance of Greater Sage-Grouse at Leks. *Conserv Biol.* 2012; 26: 461–471. doi: [10.1111/j.1523-1739.2012.01840.x](https://doi.org/10.1111/j.1523-1739.2012.01840.x) PMID: [22594595](https://pubmed.ncbi.nlm.nih.gov/22594595/)
61. Hess JE, Beck JL. Disturbance Factors Influencing Greater Sage-grouse Lek Abandonment in North-central Wyoming. *J Wildl Manage.* 2012; 76: 1625–1634.
62. Taylor RL, Tack JD, Naugle DE, Mills LS. Combined Effects of Energy Development and Disease on Greater Sage-Grouse. *PLoS One.* 2013; 8.
63. Williamson RM. Impacts of Oil and Gas Development on Sharp-tailed Grouse on the Little Missouri National Grasslands, North Dakota. Thesis, South Dakota State University, Brookings, USA. 2009.
64. Mcnew LB, Gregory AJ, Wisely SM, Sandercock BK. Human-Mediated Selection on Life-History Traits of Greater Prairie-Chickens. In: Sandercock BK, Martin K, Segelbacher G, editors. *Ecology, Conservation, and Management of Grouse.* University of California Press, Berkeley, CA; 2011. pp. 255–266.
65. Kantrud HA, Higgins KF. Nest and Nest Site Characteristics of same Ground-nesting, Non-passerine Birds of Northern Grasslands. *Prairie Nat.* 1992; 24: 67–84.
66. Kirby DR, Grosz KL. Cattle Grazing and Sharp-Tailed Grouse Nesting Success. *Rangelands.* 1995; 17: 124–126.
67. Norton MA. Reproductive Success and Brood Habitat Use of Greater Prairie Chickens and Sharp-Tailed Grouse on the Fort Pierre National Grassland of Central South Dakota. Thesis, South Dakota State University, Brookings, USA. 2005.
68. Benítez-López A, Alkemade R, Verweij PA. The Impacts of Roads and Other Infrastructure on Mammal and Bird Populations: A Meta-analysis. *Biol Conserv.* 2010; 143: 1307–1316.
69. Weller C, Thomson J, Morton P, Aplet G. *Fragmenting Our Lands: The Ecological Footprint from Oil and Gas Development.* The Wilderness Society, Washington, DC, USA. 2002.
70. Ashley EP, Robinson JT. Road Mortality of Amphibians, Reptiles and Other Wildlife on the Long Point Causeway, Lake Erie, Ontario. *Can F Nat.* 1996; 110: 403–412.
71. Tigas LA, Van Vuren DH, Sauvajot RM. Behavioral Responses of Bobcats and Coyotes to Habitat Fragmentation and Corridors in an Urban Environment. *Biol Conserv.* 2002; 108: 299–306.
72. Gehrt SD. Seasonal Survival and Cause Specific Mortality of Urban Striped Skunks in the Absence of Rabies. *J Mammal.* 2005; 86: 1164–1170.

73. Gosselink TE, Deelen T, Warner RE, Mankin PC, Gosselink TE, Natural I, et al. Survival and Cause-Specific Mortality of Red Foxes in Agricultural and Urban Areas of Illinois. *J Wildl Manage.* 2007; 71: 1862–1873.
74. Kinley T, Newhouse N. Ecology and Translocation-Aided Recovery of an Endangered Badger Population. *J Wildl Manage.* 2008; 72: 113–122.
75. Francis CD, Ortega CP, Cruz A. Noise Pollution Changes Avian Communities and Species Interactions. *Curr Biol.* 2009; 19: 1415–1419. doi: [10.1016/j.cub.2009.06.052](https://doi.org/10.1016/j.cub.2009.06.052) PMID: [19631542](https://pubmed.ncbi.nlm.nih.gov/19631542/)
76. Copeland HE, Doherty KE, Naugle DE, Pocewicz A, Kiesecker JM. Mapping Oil and Gas Development Potential in the US Intermountain West and Estimating Impacts to Species. *PLoS One.* 2009; 4.
77. Paton PWC. The Edge Effect on Avian Nest Success: How Strong is the Evidence? *Conserv Biol.* 1994; 8: 17–26.
78. Marini M, Melo C. Predators of Quail Eggs, and the Evidence of the Remains: Implications for Nest Predation Studies. *Condor.* 1998; 100: 395–399.
79. Larivière S. Reasons Why Predators Cannot be Inferred From Nest Remains. *Condor.* 1999; 101: 718–721.
80. Dzialak MR, Olson C V., Harju SM, Webb SL, Mudd JP, Winstead JB, et al. Identifying and Prioritizing Greater Sage-grouse Nesting and Brood-rearing Habitat for Conservation in Human-modified Landscapes. *PLoS One.* 2011; 6.
81. Richardson TW, Gardali T, Jenkins SH. Review and Meta-analysis of Camera Effects on Avian Nest Success. *J Wildl Manage.* 2009; 73: 287–293.
82. Moynahan BJ, Lindberg MS, Thomas JW. Factors Contributing to Process Variance in Annual Survival of Female Greater Sage-Grouse in Montana. *Ecol Appl.* 2006; 16: 1529–1538. PMID: [16937816](https://pubmed.ncbi.nlm.nih.gov/16937816/)
83. Tucker S. Study No. E-11: Furbearer Harvest Regulations Study. North Dakota Game Fish Dep Bismarck, North Dakota, USA.