

In vitro antifilarial activity, antioxidant potential and phenolic constituents of *Quisqualis indica* L.

Subha Rastogi^{*1,+}, Madan Mohan Pandey^{1,^}, Ajay Kumar Singh Rawat¹, Vikas Kushwaha^{2,3,#} & P Kalpana Murthy^{2,4,@}

¹Pharmacognosy & Ethnopharmacology Division, CSIR-National Botanical Research Institute, Lucknow 226 001, Uttar Pradesh, India

²Division of Parasitology, CSIR-Central Drug Research Institute, New Campus, BS 10/1, Sector 10, Jankipuram Extension, Sitapur Road, Lucknow 226 031, Uttar Pradesh, India

³Zoology Department, Panjab University, Chandigarh, Sector 14, 160 014, India

⁴CSIR-Emeritus Scientist, Department of Zoology, University of Lucknow, University Road, Lucknow 226 007, Uttar Pradesh, India

E-mail: ⁺subharastogi1@rediffmail.com, mmp78@rediffmail.com; [#]kushwahavikas0@gmail.com;

[@]drpkmurthy@yahoo.com; drpkmurthy@gmail.com

Received 04 September 2018; revised 21 May 2019

Quisqualis indica L., commonly known as 'Rangoon-ki-bel' or 'Madhumalti', has been used by the traditional healers as it is active against some of the commonly occurring diseases like boils, fevers diarrhea and helminthiasis. However, no systematic and scientifically validated studies on antifilarial activity of *Q. indica* are available. In the present study, we report *in vitro* antifilarial activity of ethanolic and hydroethanolic extracts of the leaves (QILE and QILEW) and flowers (QIFE and QIFEW) of this plant on microfilariae (mf) and female adult worms of human lymphatic filariid *Brugia malayi* using motility and or 3-(4,5-dimethylthiazol-2-yl)-2,5 diphenyltetrazolium bromide (MTT)-reduction assays. The hydroalcoholic extract of flowers (QIFEW) was found effective as it killed adult female worms (LC₁₀₀: 62.5 µg/mL) and mf (LC₁₀₀: 125 µg/mL); IC₅₀ values for the respective parasite stages were 34.50 and 31.88 µg/mL. SI values recorded with respect to motility of female parasite and mf was more than 20. The active principle(s) responsible for antifilarial activity may thus be present in QIFEW. The antioxidant activity results also indicated QIFEW to possess better antioxidant potential than the other extracts studied. HPLC analysis showed that the 02 keyphenolics present in hydroalcoholic extract of the flowers (QIFEW) were gallic acid and ellagic acid. In the different extracts, the concentration of gallic acid was found to vary from 26.9 mg/g to 2.50 mg/g while ellagic acid ranged between 11.5 mg/g to 6.77 mg/g. It was also observed that the leaves were rich in flavonoids whereas the flowers were rich in phenolics. The findings indicate that active molecule (s) of hydroalcoholic extract from *Q. indica* flowers may help in providing new leads for developing antifilarial agents. We believe that this is the first systematically studied report on the *in vitro* antifilarial activity of the hydroalcoholic extract of *Q. indica* flowers.

Keywords: Antifilarial, Antioxidant, *Brugia malayi*, HPLC, *In vitro* assays, Phenolics, *Quisqualis indica*

IPC Code: Int. Cl.¹⁹: A61K 31/4709, A61P 17/18, A01N 43/56, A61K 38/00, C07K 14/02, C07C 39/04, A61K 31/19

Lymphatic filariasis (LF), commonly known as elephantiasis, is a neglected tropical disease. The

malayi and *B. timore*. It is a painful and profoundly disfiguring disease, causing the body parts to enlarge abnormally. It leads to physical disabilities and the person infected with this disease becomes a social outcast. Filariasis damages the lymphatic system. India, Indonesia, Nigeria and Bangladesh contribute about 70% of the infection worldwide. LF causes a variety of clinical manifestations, including lymphoedema, hydrocele and swelling of the scrotum. The vast majority of infected people from endemic

areas are asymptomatic microfilaremics and do not display any clinical manifestations even during their

are 52 countries with a population of 500 million that require preventive methods for stopping the spreading of this disease. According to WHO, in the year 2000 the number of people found infected was more than 120 million including nearly 40 million who were disfigured and incapacitated¹. Current treatment is based on combination of a broad spectrum anthelmintic albendazole (ALB) and diethylcarbamazine (DEC)/ivermectin (IVM), or DEC fortified with cooking salt. DEC and IVM principally act on microfilariae (mf; a larval stage of the parasite) and do not exert any appreciable effect on adult parasites².

*Corresponding author

Although treatment with mass drug administration (MDA) is considerably reducing the incidence of the infection and even eliminating it in some countries³, unfortunately there are reports of reemergence of infection in some areas especially in Sri Lanka⁴. Also, the existing drugs are known to produce adverse effects in treated subjects^{5,6}. Therefore, new antifilarials are required to act as macrofilaricidal agents as well as to replace current drugs threatened with development of IVM resistance, as reported in onchocerciasis patients⁷.

In recent years, a lot of emphasis is being given to the development of antifilarial agents from plant products. Antifilarials from plants have great potential to be used as medicine because they are generally safer and have lesser side effects than which are often associated with synthetic drugs⁸. India is rich in medicinal plants. In our drug development programme, CSIR Network project NWP-0037, continuous efforts have been made to discover anti-filarial agents through research on medicinal plants^{2,9,10}. *Quisqualis indica* Linn. Family: Combretaceae, was one of the plants listed in that project and was selected for detailed studies. *Q. indica* (syn. *Combretum indicum* (L.) De Filippis, popularly called Rangoon-ki-bel or Madhumalti, is a woody climber having vigorous growth that needs sturdy support and can grow up to 8 m tall. Its roots are not very strong nor do they go deep inside the soil. The leaves are arranged in opposite pairs along the stem and are elliptic to elliptic-oblong in shape and measure about 5-18 cm by 2.5-7 cm. The fragrant flowers are borne on a pendant raceme and are white in color, the upper surface of the flower turns pink then red over a period of time. The fruit is a red drupe, turning dark brown when mature. *Q. indica* is an ornamental plant found growing in several countries. Investigations have led to the isolation and identification of several phytoconstituents such as pelargonidin-3-glucoside, trigonelline, L-proline, L-asparagine, quisqualic acid, rutin and isoenzyme A and isoenzyme B and fixed oil. *Quisqualis* fruits are used for roundworm infection in Traditional Chinese Medicine (TCM)¹¹. Preparations from different parts of this plant are used for the treatment of diarrhea, nephritis, rheumatism, headache, boils, ulcers, inflammation, and as anthelmintic^{12,13}.

The aim of the present study was to investigate the *in vitro* antifilarial activity of flowers and leaves of *Q. indica*, which have been reported to have anthelmintic properties, and evaluate their antioxidant

activity in order to evaluate their utility in providing relief from the oxidative stress that is generated as a result of different physiological changes taking place in the body of the patient with lymphatic filariasis. Since it is well known that polyphenols are good antioxidants, different samples were also subjected to HPLC-PDA analysis in order to determine the different polyphenols present in them.

Materials and methods

Chemicals and reagents

Standards (Ellagic acid, Gallic acid, Protocatechuic acid and Rutin) as well as DPPH, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), dimethyl sulfoxide, Hanks Balanced Salt Solution (HBSS), Diethylcarbamazine-citrate and Ivermectin were obtained from Sigma-Aldrich. HPLC-grade solvents and other chemicals were procured from Merck Chemicals.

Plant material

The plant *Q. indica* L. (Specimen number NBR/PH/227365) was collected from Lucknow, Uttar Pradesh (India), authenticated and herbarium deposited in the division.

Extraction and sample preparation

The leaves (QIL) and flowers (QIF) (100 g each) of *Q. indica* were dried, coarsely grinded, separately, and both extracted with ethanol and 50% aqueous ethanol (200 mL x 3) at room temperature (25°C) for 24 h on a shaker (60 rpm). The combined extracts were concentrated in a rotatory evaporator (Buchi, Switzerland) under reduced pressure at 45°C and lyophilized to yield dried ethanolic (QILE and QIFE) and 50% hydro alcoholic (QILEW and QIFEW) extracts. Their percentage yields were QILE-3.12%, QILEW- 10.23%, QIFE- 7.30% and QIFEW-12.61%. These extracts were used for evaluation of *in vitro* antifilarial and antioxidant activities as well as for HPLC analysis and determination of total phenolic and total flavonoid contents.

In vitro evaluation of antifilarial activity

For *in vitro* evaluation of antifilarial activity, the maintenance of the animals and strains as well as the isolation of parasites was done according to the methods reported earlier^{10,14,15}. Motility and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction assays were carried out for assessing the antifilarial activity^{15,16}. The solutions of test samples as well as IVM were prepared in

dimethylsulphoxide (DMSO) while DEC-c was prepared in sterile distilled water. IC_{50} and CC_{50} values were also determined and computed^{10,17,18}.

Antioxidant assays

Total antioxidant capacity (TAC) by phosphomolybdate antioxidant assay- The phosphomolybdenum method¹⁹ was used to determine the total antioxidant capacity of all the different samples. In this method, the sample under analysis reduces Mo(VI) to Mo(V) with the formation of green compounds, the absorbance of which was measured spectrophotometrically at a wavelength of 695 nm. Ascorbic acid was used as the standard and the results were expressed as mmol ascorbic acid per gram (mmol AA/g) extract.

Ferric reducing antioxidant power (FRAP) assay- The method described by Oyaizu was used to determine the FRAP of all the different samples, but with slight modifications^{20,21}. In this method, the sample under analysis reduces potassium ferricyanide (Fe^{3+}) to potassium ferrocyanide (Fe^{2+}). This potassium ferrocyanide (Fe^{2+}) subsequently forms a Pearl's Prussian blue ferric-ferrous complex with ferric chloride. The absorbance is measured at 700 nm using a spectrophotometer which is an indicator of the amount of Fe^{3+} ions present.

Estimation of Total Phenolic content (TPC) and Flavonoid content (TFC)

Folin-Ciocalteu method²² was used for determining TPC of all the different samples. Different concentrations of gallic acid were used for preparing the standard curve. The total phenolic content of the samples studied was calculated and expressed as mg GAE g^{-1} of extract. Colorimetric method using aluminium chloride¹⁹ was used for quantification of TFC of the different samples. Quercetin was used at different concentrations for preparing the standard curve which was used to calculate the total flavonoid content of the samples studied, expressed as mg QE g^{-1} of extract.

HPLC analysis for Phenolics and Flavonoids

HPLC analysis was carried out on Waters HPLC system using the same conditions as described earlier by us²³. The column used for separation was Polaris 5 C18-A (250×4.6 mm i.d.; 5 μ m particle size) and PDA at 280 nm was used for detection. Samples of a concentration of 15 mg/mL were used and the chromatograms were visualized under 3 different wavelengths viz. 254, 270 and 350 nm considering

the different compounds that may be present. Comparison of the retention times and spectra of the peaks obtained with the known standards was used for identifying the peaks in the chromatograms. Gallic acid, protocatechuic acid, ellagic acid and rutin (0.1 mg/mL) were used as reference standards.

Results

In vitro antifilarial activity

The antifilarial activity of the different ethanolic and hydroethanolic samples was determined *in vitro*. 62.5 μ g/mL was set as the cutoff concentration for positive antifilarial activity 'hit' of natural products *in vitro* and 500 μ g/mL as the limit concentration for testing. None of the ethanolic extracts (QILE and QIFE) showed any adulticidal or microfilaricidal activity up to the acceptable concentration of 62.5 μ g/mL; they showed no activity even up to 500 μ g/mL. Results of *in vitro* activity of the hydroalcoholic extract of leaves (QILEW) and flowers (QIFEW) of the plant on adult worms and mf of *Brugia malayi*, evaluated by motility and MTT reduction assays, are shown in Table 1. The hydroalcoholic extract of flowers (QIFEW) killed the female adult worms (LC_{100} : 62.5 μ g/mL) and mf (LC_{100} : 125 μ g/mL). The extract produced remarkable inhibition (>90%) in MTT reduction potential of the adult worms. IC_{50} values for the respective parasite stages were found to be 34.50 and 31.88 μ g/mL. SI values of the extract recorded with respect to motility of female adult worm and mf were 22.89 and 24.78, respectively. QILEW required 250 μ g/mL or more than 250 μ g/mL concentration to kill the adult worms and failed to affect MTT-reduction potential of the female parasites. Thus, *in vitro* activity of QILEW was not encouraging and it was, therefore, not taken up for further studies.

DEC-c killed the adult worms and mf at 800 μ M (LC_{100}) and 500 μ M (LC_{100}), respectively: the IC_{50} values of the drug for respective parasite stages were 289.00 μ M and 354 μ M. IVM required much lower concentrations to kill both adult worms (LC_{100} : 5 μ M) and mf (LC_{100} : 2.5 μ M) and IC_{50} values were found to be 3.05 and 1.57 μ M, respectively. SI values of the reference drugs with respect to motility of the parasites were >25–159.

In summary, only the hydroalcoholic extract of flowers (QIFEW) of the plant remarkably affected viability of female adult worms and mf *in vitro*. SI values of the extract were 23 and 25 with respect to adult worm and mf, respectively.

Table 1 — *In vitro* efficacy of *Quisqualis indica* and the reference drugs ivermectin and DEC citrate on adult worms (AW) and microfilariae (Mf) of *Brugia malayi* evaluated by motility assay (MA) and MTT reduction assay.

Antifilarial agent (Extract)	Macrofilaricidal				Microfilaricidal			CC ₅₀
	LC ₁₀₀ ^a (µg/mL) for AW in MA	IC ₅₀ ^b (µg/mL) for AW in MA	Mean % inhibition in MTT reduction by AW	SI in MA	LC ₁₀₀ (µg/mL) for Mf in MA	IC ₅₀ (µg/mL) for Mf in MA	SI in MA	
	QIFEW	62.5	34.50	94.52	22.89	125	31.88	
QILEW	250	ND	NI	ND	>250	ND	ND	ND
DEC-c*	800	289	64	31.50	500	354	25.71	9103
IVM*	5	3.05	5.80	81.97	2.50	1.57	159.23	250

Control: Motility score of parasite+DMSO only wells= 4 (=parasites were motile with highly active); MTT absorbance values (OD_{510nm}) of parasites+DMSO only= 0.64; ^aLC₁₀₀= Lethal concentration of the test agent which causes 100% irreversible immobility (death) in *B. malayi* adult and mf. ^bIC₅₀= The concentration of the agent at which 50% inhibition in motility of the parasites is achieved; ^cCC₅₀= concentration at which 50% of cells are killed; SI= Selectivity Index (CC₅₀/IC₅₀); DEC-c= Diethylcarbamazine-citrate; IVM= Ivermectin; *µM; Experiments were repeated twice. NI= No inhibition; ND= Not done.

Table 2 — TPC, TFC and antioxidant activities of leaves and flowers of *Q. indica*

Sample	Total phenolic content* (mg GAE/g)	Total flavonoid content* (mg QE/g)	Total Antioxidant Capacity by Phosphomolybdate Antioxidant Assay (mmol AA/g extract)
QILE	11.49	5.21	0.539
QILEW	10.81	3.75	0.437
QIFE	11.08	2.63	0.715
QIFEW	16.86	1.02	0.732
AA	-	-	-

*Values are means of three determinations each

***In vitro* antioxidant activity**

Total antioxidant capacity by phosphomolybdate antioxidant assay– The total antioxidant capacity of the ethanolic and 50% hydroalcoholic extracts of QIL and QIF are given in Table 2 & Fig. 1. The order of total antioxidant capacity was QIFEW>QIFE>QILE>QILEW. It was observed that there was a direct correlation between the phosphomolybdenum complex formed and the amount of the extracts used.

Ferric reducing anti-oxidant power (FRAP) assay–The reducing power of the ethanolic and 50% hydroalcoholic extracts of QIL and QIF was measured using Fe³⁺ to Fe²⁺ reduction assay. The results obtained are shown in Fig. 2. All the samples exhibited reducing power which had a direct correlation with increasing concentration of the samples. However ascorbic acid exhibited a higher reducing power.

Total phenolic and flavonoid contents

TPC and TFC of the ethanolic and 50% hydroalcoholic extracts of QIL and QIF were determined (Table 2, Fig. 1). QIFEW showed the




Fig. 1 — TPC, TFC, and TAC (mmol AA/g extract) of methanolic (M) and 50% hydroalcoholic (MW) extracts of different parts of *Q. indica*. (QIL: *Q. indica* leaves; QIF: *Q. indica* flowers)




Fig. 2 — Ferric reducing anti-oxidant power of the methanolic (M) and 50% hydroalcoholic (MW) extracts of different parts of *Q. indica*. (QIL: *Q. indica* leaves; QIF: *Q. indica* flowers)

highest TPC with 16.86 mg GAE/g extract while QILE showing the highest TFC with 5.21 mg QE/g extract. The TPC and TFC of the samples were in the order QIFEW>QILE>QIFE>QILEW and QILE>QILEW>QIFE>QIFEW respectively.

HPLC analysis

The ethanol and the ethanol-water extracts obtained from leaves and flowers of *Q. indica* were subjected to HPLC analyses. Different phenolics and flavonoids that may be present were analyzed under the conditions described earlier. The HPLC chromatograms of the standard mixture comprising of gallic acid (Rt: 6.31 min), protocatechuic acid (Rt: 10.33 min), rutin (Rt: 19.73 min) and ellagic acid (Rt: 20.90 min), and the different extracts, monitored at 254 nm, are shown in Fig. 3 & Fig. 4 respectively. Their chemical structures as well as their respective UV spectra are given in Fig. 5. Confirmation of the identity of the individual compounds was done by comparing with the UV spectra and the retention time of the standards and expressed as mg/g of dry extract. The qualitative and quantitative results are summarized in Table 3 & Table 4 respectively. Results indicated that gallic acid and ellagic acid were the two major phenolics found in all the samples. The concentration of gallic acid varied from 26.9 mg/g (QIFE) to 2.50 mg/g (QILEW) while ellagic acid ranged between 11.5 mg/g (QIFEW) to 6.77 mg/g (QILE). Protocatechuic acid was present in all the extracts, although in very minor quantity (0.25-0.56 mg/g extract). Besides these compounds the chromatograms also showed the presence of other peaks, many of them being classified as either phenolics or flavonoids based on the analysis of their UV spectra. Based on the analysis of the spectra of the eluted peaks, QILEW contained other compounds with spectra similar to that of ellagic acid, thereby indicating the presence of ellagic acid related compounds. Although several flavonoids were present in the extracts, none of them corresponded with rutin. Rutin was, therefore, used as reference for determining the presence of other flavonoids. A peak eluting after ellagic acid, at about 21.95 min, was observed in all the samples. Although this compound,

tentatively designated as QI-1, was neither isolated nor identified, it appeared to be a flavonoid or its glycoside based on its UV spectrum. Its concentration was calculated with respect to rutin and was found highest in QILE (15.70 mg/g extract) and lowest in QIFEW (1.04 mg/g extract). It was also observed that




Fig. 4 — HPLC chromatograms of the methanolic and 50% hydroalcoholic extracts of different parts of *Q. indica*. (QIL: *Q. indica* leaves; QIF: *Q. indica* flowers)




Fig. 3 — HPLC chromatogram of standard mixture (GA-Gallic acid, PC-Protocatechuic acid, R-Rutin and EA-Ellagic acid)

Table 3 — Qualitative HPLC analysis of phenolics and flavonoids present in leaves and flowers of *Q. indica*

Sample	Phenolics and flavonoids					
	Gallic acid	Protocatechuic acid	Ellagic acid	Other phenolics*	Rutin	Other flavonoids*
QILE	+	+	+	+	-	++
QILEW	+	+	+	+	-	++
QIFE	+	+	+	++	-	+
QIFEW	+	+	+	++	-	+

*Unidentified

Table 4 — Quantitative HPLC analysis of phenolics and flavonoids present in leaves and flowers of *Q. indica*

Sample	Phenolics and flavonoids (mg/g ext)			
	Gallic acid	Protocatechuic acid	Ellagic acid	QI-1*
QILE	2.96	0.48	6.77	15.7
QILEW	2.50	0.56	7.62	9.43
QIFE	26.9	0.40	10.2	5.20
QIFEW	17.2	0.25	11.5	1.04

*Unidentified




Fig. 5 — Chemical structures and spectral profiles of Gallic acid, Protocatechuic acid, Rutin and Ellagic acid

the leaves were rich in flavonoids whereas the flowers were rich in phenolics.

Discussion

Filariasis damages the lymphatic system, causing the various body parts to enlarge abnormally. It leads to certain disabilities that result not only in physical impairment but also reduced economic productivity, discrimination and serious psychosocial consequences. Thus, a global programme was launched by WHO in 2000 to eliminate lymphatic filariasis. DEC and IVM are available for treatment of filariasis but they have

limitations. They are principally microfilaricidal and their effect on adult worms is doubtful². Their repetitive use may give rise to drug resistance. Considering the limitations of the existing drugs and the huge socio-economic requirements, there is need to search for new antifilarials.

Developing countries mainly use herbal medicines for the primary health care needs of their people. This is because they have been found to be safe, effective, with lesser side effects as well as due to their cultural acceptability. It is believed that the phytoconstituents present in them are more compatible with the human body²⁴. As indicated by the results obtained, the hydroalcoholic extract of *Q. indica* flowers possesses promising *in vitro* antifilarial as well as antioxidant activities. QIFEW also contains a good amount of phenolic compounds, some of which are known to exhibit antinematodal activity. HPLC analysis indicated that ellagic acid was the major phenolic acid present in the samples, the content of which was highest in QIFEW. Earlier, the effect of gallic acid, ellagic acid and genistic acid was studied on the bovine parasite *Onchocerca ochengi* and drug-resistant strains of the free living nematode *Caenorhabditis elegans*²⁵. The results indicated that ellagic acid binds differently in the worms and anthelmintic (levamisole, ALB and IVM)-resistant strains of *C. elegans* and could therefore have potential for the treatment even against those nematodal infections which are resistant to anthelmintic drugs. In case of *Q. indica* too, ellagic acid and its related compounds were the major phenolics present in QIFEW, indicating that they may be responsible for the good level of antifilarial activity exhibited by QIFEW in both motility and MTT reduction assays.

Conclusion

The present studies, therefore, indicate that the hydroalcoholic extract of the flowers of *Q. indica* has the potential to yield compounds that may be used not only for developing novel antifilarial candidates but also for providing relief from the oxidative stress that

is generated as a result of different physiological changes taking place in the body of the patient with lymphatic filariasis.

Acknowledgement

The authors thank the Directors, CSIR-NBRI and CSIR-CDRI, Lucknow, for their keen interest and encouragement, and for providing the necessary facilities to carry out this work. Part of this study was supported by the CSIR Network project NWP-0037. Thanks are due to CSIR, New Delhi for award of Emeritus Scientist Project [No. 21 (0963)/13/EMR-II; 29-10-2014] to PKM and to the Honorable Vice Chancellor, Lucknow University, Lucknow, for providing office/laboratory space and computational facilities to PKM for data analysis and manuscript preparation at the Department of Zoology.

References

- 1 <http://www.who.int/mediacentre/factsheets/fs102/en/>
- 2 Kalani K, Kushwaha V, Sharma P, Verma R, Srivastava M, Khan F, Murthy, PK & Srivastava SK, *In vitro*, *in silico* and *in vivo* studies of ursolic acid as an anti-filarial agent. *PLoS One*, 9 (2014) e111244.
- 3 WHO 2018. [<https://www.who.int/westernpacific/news/detail/08-10-2018-three-more-countries-eliminate-lymphatic-filariasis>]
- 4 Rao RU, Samarasekera SD, Nagodavithana KC, Dassanayaka TDM & Weil GJ, Reassessment of areas with persistent lymphatic filariasis nine years after cessation of mass drug administration in Sri Lanka, *PLoS Negl Trop Dis*, 11 (2017) e0006066.
- 5 Sashidhara KV, Kumar A, Rao KB, Kushwaha V, Saxena K & Murthy PK, *In vitro* and *in vivo* antifilarial activity evaluation of 3,6-epoxy [1,5] dioxocines: a new class of antifilarial agents. *BMCL*, 22 (2012) 1527-1532.
- 6 Sashidhara KV, Rao KB, Kushwaha V, Modukuri RK, Verma R & Murthy PK, Synthesis and antifilarial activity of chalcone-thiazole derivatives against a human lymphatic filarial parasite, *Brugia malayi*. *Eur J Med Chem*, 81 (2014) 473-480.
- 7 Atweneboana MYO, Eng JKL, Boakye DA, Gyapong JO & Prichard RK, Prevalence and intensity of *Onchocerca volvulus* infection and efficacy of ivermectin in endemic communities in Ghana: a two-phase epidemiological study, *Lancet*, 369 (2007) 2021-2029.
- 8 Narasimha Rao R, Bhavya B, Pavani K, Swapna A & Prasoona CH, Anthelmintic activity of *Symplocos racemosa*, *Int J Pharm Biol Sci*, 1 (2011) 198-230.
- 9 Chatterjee RK, Fatma N, Murthy PK, Sinha P, Kulshrestha DK & Dhawan BN, Macrofilaricidal activity of the stem bark of *Streblus asper* and its major active constituents, *Drug Dev Res*, 26 (1992) 67-68.
- 10 Kushwaha V, Saxena K, Verma R, Verma SK, Katoch D, Kumar N, Lal B, Murthy PK & Singh B, Antifilarial activity of diterpenoids from *Taxodium distichum*. *Parasit Vectors*, 9 (2016) 312.
- 11 http://www.tcm-basics.com/materiamedica/fructus_quisqualis.htm
- 12 <http://www.mpbd.info/plants/quisqualis-indica.php>
- 13 Singh S, Rai A, Maity S, Sarkar S, Maji S & Saha S. Effect of ethanolic extract of *Quisqualis indica* L. flower on experimental esophagitis in albino Wistar rats, *Indian J Exp Biol*, 2 (55) 122-126.
- 14 Murthy PK, Murthy PS, Tyagi K & Chatterjee RK, Fate of infective larvae of *Brugia malayi* in the peritoneal cavity of *Mastomys natalensis* and *Meriones unguiculatus*, *Folia Parasitol (Praha)*, 44 (1997) 302-304.
- 15 Murthy PK & Chatterjee RK, Evaluation of two *in vitro* test systems employing *Brugia malayi* parasite for screening of potential antifilarials, *Curr Sci*, 77 (1999) 1084-1089.
- 16 Lakshmi V, Joseph SK, Srivastava S, Verma SK, Sahoo MK, Dube V, Mishra SK & Murthy PK, Antifilarial activity *in vitro* and *in vivo* of some flavonoids tested against *Brugia malayi*, *Acta Trop*, 116 (2010) 127-133.
- 17 Murthy PK, Joseph SK & Murthy PS, Plant products in the treatment and control of filariasis and other helminth infections and assay systems for antifilarial/anthelmintic activity, *Planta Med*, 77 (2011) 647-661.
- 18 Huber W & Koella JC, A comparison of three methods of estimating EC₅₀ in studies of drug resistance of malaria parasites, *Acta Trop*, 55 (1993) 257-261.
- 19 Prieto P, Pineda M & Aguilar M, Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E, *Anal Biochem*, 269 (1999) 337-341.
- 20 Oyaizu M, Studies on products of browning reactions: antioxidative activities of products of browning reaction prepared from glucosamine, *Jpn J Nutr*, 44 (1986) 307-315.
- 21 Elmastas M, Gulcin D, Beydemir S, Kufrevioglu OD & Aboul-Enein HY, A study on the *in vitro* antioxidant activity of juniper (*Juniperus communis* L.) seeds extracts, *Anal Lett*, 39 (2006) 47-65.
- 22 Wolfe K, Wu X & Liu RH, Antioxidant activity of apple peels, *J Agric Food Chem*, 51 (2003) 609-614.
- 23 Pandey MM, Vijayakumar M, Rastogi S & Rawat AKS, Phenolic content and antioxidant properties of selected Indian spices of Apiaceae, *J Herbs Spices Med Plants*, 18 (2012) 246-256.
- 24 Kamboj VP, Herbal medicine, *Curr Sci*, 78 (2000) 35-39.
- 25 Ndjonga D, Abladam ED, Djafsa B, Ajonina-Ekoti I, Achukwi MD & Liebau E. Anthelmintic activity of phenolic acids from the axlewood tree *Anogeissus leiocarpus* on the filarial nematode *Onchocerca ochengi* and drug-resistant strains of the free-living nematode *Caenorhabditis elegans*, *J Helminthol*, 88 (2014) 481-488.