

3-2-2017

Hurwitz Number Fields

David P. Roberts

University of Minnesota - Morris, roberts@morris.umn.edu

Follow this and additional works at: <https://digitalcommons.morris.umn.edu/mathematics>



Part of the [Mathematics Commons](#)

Recommended Citation

David P. Roberts. Hurwitz number fields. *New York Journal of Mathematics* 23 (2017) 227-272.

This Article is brought to you for free and open access by the Faculty and Staff Scholarship at University of Minnesota Morris Digital Well. It has been accepted for inclusion in Mathematics Publications by an authorized administrator of University of Minnesota Morris Digital Well. For more information, please contact skulann@morris.umn.edu.

HURWITZ NUMBER FIELDS

DAVID P. ROBERTS

ABSTRACT. The canonical covering maps from Hurwitz varieties to configuration varieties are important in algebraic geometry. The scheme-theoretic fiber above a rational point is commonly connected, in which case it is the spectrum of a Hurwitz number field. We study many examples of such maps and their fibers, finding number fields whose existence contradicts standard mass heuristics.

CONTENTS

1. Introduction	1
2. A degree 25 introductory family	3
3. Background on Hurwitz covers	8
4. Specialization to Hurwitz number algebras	15
5. A degree 9 family: comparison with complete number field tables	19
6. A degree 52 family: tame ramification and exceptions to Principle B	21
7. A degree 60 family: non-full monodromy and a prime drop	26
8. A degree 96 family: a large degree dessin and Newton polygons	30
9. A degree 202 family: degenerations and generic specialization	33
10. A degree 1200 field: computations in large degree	37
References	38

1. INTRODUCTION

This paper is a sequel to *Hurwitz monodromy and full number fields* [19], joint with Venkatesh. It is self-contained and aimed more specifically at algebraic number theorists. Our central goal is to provide experimental evidence for a conjecture raised in [19]. More generally, our objective is to get a concrete and practical feel for a broad class of remarkable number fields arising in algebraic geometry, the Hurwitz number fields of our title.

1.1. Full fields, the mass heuristic, and a conjecture. Say that a degree m number field $K = \mathbb{Q}[x]/f(x)$ is *full* if the Galois group of $f(x)$ is either the alternating group A_m or the symmetric group S_m . For \mathcal{P} a finite set of primes, let $F_{\mathcal{P}}(m)$ be the number of isomorphism classes of full fields K of degree m unramified outside \mathcal{P} and ∞ . In the sequel, we suppress the words “of isomorphism classes” as it is understood that we are always counting fields up to isomorphism.

In [3, Eq. 10], Bhargava formulated a heuristic expectation $\mu_D(m)$ for the number $F_D(m)$ of degree m full number fields with absolute discriminant $D \in \mathbb{Z}_{\geq 1}$. The main theorems of [7], [2], and [4] respectively say that this heuristic is asymptotically correct for $m = 3, 4,$ and 5 . While Bhargava is clearly focused in [3] on this

“horizontal” direction of fixed m and increasing D , it also makes sense to apply the same mass heuristic in the “vertical” direction. In [16, Eq. 68], we summed over contributing D to obtain a heuristic expectation $\mu_{\mathcal{P}}(m)$ for the number $F_{\mathcal{P}}(m)$. It is a product of local contributions, one for each $p \in \mathcal{P}$. Figure 6 of [16] graphed the function $\mu_{\{2,3\}}$, while Figure 1.1 graphs the function $\mu_{\{2,3,5\}}$ which is more relevant for us here. All $\mu_{\mathcal{P}}$ share a common qualitative behavior: the numbers $\mu_{\mathcal{P}}(m)$ can be initially quite large, but by [16, Eq. 42 and Prop. 6.1] they ultimately decay super-exponentially to zero. From this decay, one might expect that for any fixed \mathcal{P} , the sequence $F_{\mathcal{P}}(m)$ would be eventually zero.




FIGURE 1.1. The heuristic approximation $\mu_{\{2,3,5\}}(m)$ to the number $F_{\{2,3,5\}}(m)$ of degree m full fields ramified within $\{2, 3, 5\}$. In contrast, §9 shows $F_{\{2,3,5\}}(202) \geq 2497$ and §10 shows $F_{\{2,3,5\}}(1200) \geq 1$.

The construction studied in [19] has origin in work of Hurwitz and involves an arbitrary finite nonabelian simple group T . Let \mathcal{P}_T be the set of primes dividing T . The construction gives a large class of separable algebras $K_{h,u}$ over \mathbb{Q} which we call *Hurwitz number algebras*. Infinitely many of these algebras have all their ramification with \mathcal{P}_T . Within the range of our computations here, these algebras are commonly number fields themselves; in all cases, they factor into number fields which we call *Hurwitz number fields*. The algebras come in families of arbitrary dimension $\rho \in \mathbb{Z}_{\geq 0}$, with the *Hurwitz parameter* h giving the family and the *specialization parameter* u giving the member of the family. Strengthening Conjecture 8.1 of [19] according to the discussion in §8.5 there, we expect that there are enough contributing $K_{h,u}$ to give the following statement.

Conjecture 1.1. *Suppose \mathcal{P} contains the set of primes dividing the order of a finite nonabelian simple group. Then the sequence $F_{\mathcal{P}}(m)$ is unbounded.*

From the point of view of the mass heuristic, the conjecture has both an unexpected hypothesis and a surprising conclusion.

1.2. Content of this paper. The parameter numbers $\rho = 0$ and 1 have special features connected to *dessins d'enfants*, and we present families with $\rho \in \{0, 1\}$ in [15]. To produce enough fields to prove Conjecture 1.1, it is essential to let ρ tend to infinity. Accordingly we concentrate here on the next case $\rho = 2$, with our last example being in the setting $\rho = 3$.

Section 2 serves as a quick introduction. Without setting up any general framework, it exhibits a degree 25 family. Specializing this family gives more than 10,000 number fields with Galois group S_{25} or A_{25} and discriminant of the form $\pm 2^a 3^b 5^c$.

Section 3 introduces Hurwitz parameters and describes how one passes from a parameter to a Hurwitz cover. Full details would require deep forays into moduli problems on the one hand and braid group techniques on the other. We present information at a level adequate to provide a framework for our examples to come. In particular, we use the Hurwitz parameter $h = (S_5, (2111, 5), (4, 1))$ corresponding to our introductory example to illustrate the generalities.

Section 4 focuses on specialization, meaning the passage from a Hurwitz cover to its fibers. In the alternative language that we have been using in this introduction, a Hurwitz cover gives a family of Hurwitz number algebras, and then specialization is passing from the entire family to one of its members. The section elaborates on the heuristic argument for Conjecture 1.1 given in [19]. It formulates Principles A, B, and C, all of which say that specialization behaves close to generically. Proofs of even weak forms of Principles A and B would suffice to prove Conjecture 1.1. Here again, the introductory example is used to illustrate the generalities.

The slightly shorter Sections 5-10 each report on a family and its specializations, degrees being 9, 52, 60, 96, 202, and 1200. Besides describing its family, each section also illustrates a general phenomenon.

Sections 5-10 together indicate that the strength with which Principles A, B, and C hold has a tendency to increase with the degree m , in strong support of Conjecture 1.1. In particular, our two largest degree examples clearly show that Hurwitz number fields are not governed by the mass heuristic as follows. In the degree 202 family, Principles A, B, and C hold without exception. One has $\mu_{\{2,3,5\}}(202) \approx 2 \cdot 10^{-17}$, but the family shows $F_{\{2,3,5\}}(202) \geq 2947$. Similarly, $\mu_{\{2,3,5\}}(1200) \approx 10^{-650}$ while the one specialization point we look at in the degree 1200 family shows $F_{\{2,3,5\}}(1200) \geq 1$.

There are hundreds of assertions in this paper, with proofs in most cases involving computer calculations, using *Mathematica* [23], *Pari* [21], and *Magma* [5]. We have aimed to provide an accessible exposition which should make all the assertions seem plausible to a casual reader. We have also included enough details so that a diligent reader could efficiently check any of these assertions. Both types of readers could make use of the large *Mathematica* file `HNF.mma` on the author's homepage. This file contains seven large polynomials defining the seven families considered here, and miscellaneous further information about their specialization to number fields.

1.3. Acknowledgements. This paper was started at the same time as [19]. It is a pleasure to thank Akshay Venkatesh whose careful reading of early drafts of this paper in the context of its relation with [19] improved it substantially. It is also a pleasure to thank the Simons foundation which partially supported this work through grant #209472. Finally, I thank the anonymous referee whose careful reading of the paper improved the exposition.

2. A DEGREE 25 INTRODUCTORY FAMILY

In this section, we begin by constructing a single full Hurwitz number field, of degree 25 and discriminant $2^{56} 3^{34} 5^{30}$. We then use this example to communicate the general nature of Hurwitz number fields and their explicit construction. We close by varying two parameters involved in the construction to get more than ten

thousand other degree twenty-five full Hurwitz number fields from the same family, all ramified within $\{2, 3, 5\}$.

2.1. The 25 quintics with critical values $-2, 0, 1$ and 2 . Consider polynomials in $\mathbb{C}[s]$ of the form

$$(2.1) \quad g(s) = s^5 + bs^3 + cs^2 + ds + x.$$

We will determine when the set of critical values of $g(s)$ is $\{-2, 0, 1, 2\}$.

The critical points of such a polynomial are of course given by the roots of its derivative $g'(s)$. The critical values are then given by the roots of the resultant

$$r(t) = \text{Res}_s(g(s) - t, g'(s)).$$

Explicitly, this resultant works out to $r(t) =$

$$\begin{aligned} & 3125t^4 \\ & + 1250(3bc - 10x)t^3 \\ & + (108b^5 - 900b^3d + 825b^2c^2 - 11250bcx + 2000bd^2 + 2250c^2d + 18750x^2)t^2 \\ & - 2(108b^5x - 36b^4cd + 8b^3c^3 - 900b^3dx + 825b^2c^2x + 280b^2cd^2 \\ & \quad - 315bc^3d - 5625bcx^2 + 2000bd^2x + 54c^5 + 2250c^2dx - 800cd^3 + 6250x^3)t \\ & + (108b^5x^2 - 72b^4cdx + 16b^4d^3 + 16b^3c^3x - 4b^3c^2d^2 - 900b^3dx^2 + 825b^2c^2x^2 \\ & \quad + 560b^2cd^2x - 128b^2d^4 - 630bc^3dx + 144bc^2d^3 - 3750bcx^3 \\ & \quad + 2000bd^2x^2 + 108c^5x - 27c^4d^2 + 2250c^2dx^2 - 1600cd^3x + 256d^5 + 3125x^4). \end{aligned}$$

This large expression conforms to the *a priori* known structure of $r(t)$: it is a quartic polynomial in the variable t depending on the four parameters b, c, d , and x . The computation required to obtain the expression is not at all intensive; for example, *Mathematica*'s `Resultant` does it nearly instantaneously.

Now consider in general the problem of classifying quintic polynomials (2.1) with prescribed critical values. Clearly, if the given values are the roots of a monic degree four polynomial $\tau(t)$, then we need to choose the b, c, d , and x so that $r(t)$ is identically equal to $3125\tau(t)$. Equating coefficients of t^i for $i = 0, 1, 2$, and 3 gives four equations in the four unknowns b, c, d , and x . If (b, c, d, x) is a solution then so is $(\omega^2b, \omega^3c, \omega^4d, \omega^5x)$ for any fifth root of unity ω . Thus the solutions come in packets of five, each packet having a common x .

In our explicit example, $\tau(t) = (t+2)t(t-1)(t-2)$. *Mathematica* determines in less than a second that there are 125 solutions (b, c, d, x) . The twenty-five possible x 's are the roots of a degree twenty-five polynomial,

$$(2.2) \quad f(x) = 2^{98}3^8x^{25} - 2^{96}3^85^2x^{24} + \dots + 4543326944239835953052526892234.$$

The algebra $\mathbb{Q}[x]/f(x)$ is our first explicit example of a Hurwitz number algebra. In this case, $f(x)$ is irreducible in $\mathbb{Q}[x]$, so that $\mathbb{Q}[x]/f(x)$ is in fact a Hurwitz number field.

2.2. Real and complex pictures. Before going on to arithmetic concerns, we draw two pictures corresponding to the Hurwitz number field $\mathbb{Q}[x]/f(x)$ we have just constructed. Any Hurwitz number algebra K would have analogous pictures. Our objective is to visually capture the fact that any Hurwitz number algebra K is involved in a very rich mathematical situation. Indeed if K has degree m , then one has m different geometric objects, with their arithmetic coordinated by K .




FIGURE 2.1. Graphs $t = g_x(s)$ of five quintic polynomials with critical points (s_i, t_i) ordered from bottom to top and marked by $i \in \{1, 2, 3, 4\}$.

Of the twenty-five solutions x to (2.2), five are real. Each of these x corresponds to exactly one real solution (b, c, d, x) . The corresponding polynomials $g_x(s)$ are plotted in the window $[-2.1, 2.1] \times [-2.4, 2.4]$ of the real s - t plane in Figure 2.1. The critical values t_i are indexed from bottom to top so that always $(t_1, t_2, t_3, t_4) = (-2, 0, 1, 2)$, with i printed at the corresponding turning point (s_i, t_i) . The labeling of each graph encodes the left-to-right ordering of the critical points s_i . For example, in the upper left rectangle the critical points are $(s_2, s_1, s_4, s_3) \approx (-1.5, -0.6, 0.7, 1.4)$ and the graph is accordingly labeled by $L = 2143$. The labeling is consistent with the labeling in Figure 2.4 below.

To get images for all twenty-five roots x , we consider the semicircular graph \circlearrowleft in the complex t -plane drawn in Figure 2.2. We then draw in Figure 2.4 its preimage $g_x^{-1}(\circlearrowleft)$ in the complex s -plane under twenty-five representatives g_x . Each of the four critical values $t_i \in \{-2, 0, 1, 2\}$ has a unique critical preimage $s_i \in \mathbb{C}$, and we print i at s_i in Figure 2.4. There are braid operations $\sigma_1, \sigma_2, \sigma_3$ corresponding to universal rules which permute the figures, given in this instance by Figure 2.3. Here the σ_i all have cycle type $3^5 2^5$ with σ_2 preserving the letter and incrementing the index modulo 3. The fact that this geometric action has image all of S_{25} suggests that the Galois group of (2.2) will be A_{25} or S_{25} as well.




FIGURE 2.2. A graph connecting the roots of the specialization polynomial $(t + 2)t(t - 1)(t - 2)$




FIGURE 2.3. Actions of σ_1 (vertical arrows), σ_2 (symbols), and σ_3 (horizontal arrows)

The twenty-five preimages are indeed topologically distinct. Thus for the twelve $\gamma_{abc} = \gamma_{cba}$, the critical points a , b , and c are connected by a triangle and the middle index b is connected also to the remaining critical point. Similarly the indexing for the twelve $\gamma_{abcd} = \gamma_{dcba}$ describes how the critical points are connected. The five graphs corresponding to the real x treated in Figure 2.1 are easily identified by the horizontal line present in Figure 2.4. We touch on the braid-theoretic infrastructure




FIGURE 2.4. The preimage of the semicircular graph of Figure 2.2 under twenty-five quintic polynomials

of Hurwitz number fields in this paper only very lightly. Our point in presenting Figures 2.2-2.4 is simply to give some idea of the topology behind the existence of Hurwitz number fields.

2.3. A better defining polynomial $\phi(x)$ and field invariants. We are not so much interested in the polynomial $f(x)$ from (2.2) itself, but rather in the field $\mathbb{Q}[x]/f(x)$ it defines. *Pari's* command `polredabs` converts $f(x)$ into a monic polynomial $\phi(x)$ which defines the same field and has minimal sum of the absolute squares of its roots. It returns

$$\begin{aligned} \phi(x) = & x^{25} - 5x^{24} + 15x^{23} - 5x^{22} - 380x^{21} + 1290x^{20} - 4500x^{19} - 28080x^{18} \\ & + 183510x^{17} + 74910x^{16} - 3033150x^{15} + 4181370x^{14} + 27399420x^{13} \\ & - 48219480x^{12} - 124127340x^{11} + 266321580x^{10} + 466602765x^9 \\ & - 592235505x^8 - 905951965x^7 + 1232529455x^6 + 2423285640x^5 \end{aligned}$$

$$+664599470x^4 - 814165000x^3 - 517891860x^2 - 58209720x + 2436924.$$

For fields of sufficiently small degree, one applies the reduction operation `polredabs` as a matter of course: the new smaller-height polynomials are more reflective of the complexity of the fields considered, isomorphic fields may be revealed, and any subsequent analysis of field invariants is sped up.

Pari's nfdisc calculates that the discriminant of $\mathbb{Q}[x]/\phi(x)$ is

$$D = 1119186718586212624367616000 = 2^{56}3^{34}5^{30}.$$

The fact that D factors into the form $2^a3^b5^c$ is known from the general theory presented in Sections 3 and 4, using that 2, 3, and 5 are the primes less than or equal to the degree 5, and the polynomial discriminant of $\tau(t) = (t+2)t(t-1)(t-2)$, namely $2304 = 2^83^2$, has this form too. Note that since all the exponents of the field discriminant are greater than the degree 25, the number field is wildly ramified at all the base primes, 2, 3, and 5.

To look more closely at $\mathbb{Q}[x]/\phi(x)$, we factorize the p -adic completion $\mathbb{Q}_p[x]/\phi(x)$ as a product of fields over \mathbb{Q}_p . We write the symbol e_c^f to indicate a factor of degree ef , ramification index e , and discriminant p^{fc} . One gets

$$\begin{aligned}
 \text{(2.3)} \quad \text{2-adically: } & \quad \mathbf{16}_{50} 3_2 3_2 3_2, \\
 \text{3-adically: } & \quad \mathbf{9}_{18} 4_3 4_3 \mathbf{3}_5 \mathbf{3}_5 1_0^2, \\
 \text{5-adically: } & \quad \mathbf{25}_{30},
 \end{aligned}$$

with wild factors printed in bold. Thus, the first line means that $\mathbb{Q}_2[x]/\phi(x)$ is a product $K_1 \times K_2 \times K_3 \times K_4$, where K_1 is a wild totally ramified degree sixteen extension of \mathbb{Q}_2 with discriminant 2^{50} , while K_2 , K_3 , and K_4 are tame cubic extensions of discriminant 2^2 . The behavior for the three primes is roughly typical, although, as we will see in Figure 4.2, a little less ramified than average.

Because the field discriminant is a square, the Galois group of $\phi(x)$ is in A_{25} . Many small collections of p -adic factorization patterns for small unramified p each suffice to prove that the Galois group is indeed all of A_{25} . Most easily, $\phi(x)$ factors in $\mathbb{Q}_{19}[x]$ into irreducible factors of degrees 17, 6, and 2, so that the Galois group contains an element of order 17. Jordan's criterion now applies: a transitive subgroup of S_m containing an element of prime order in $(m/2, m-2]$ is all of A_m or S_m . We will use this easy technique without further comment for all of our other determinations that Galois groups of number fields are full. One could also use information from ramified primes as above, but unramified primes give the easiest computational route.

2.4. A family of degree 25 number fields. We may ask, more generally, for the quintics with any fixed set of critical values. This amounts to repeating our previous computation, replacing the polynomial $\tau(t) = (t+2)t(t-1)(t-2)$ of the three previous subsections with other separable quartic polynomials

$$(2.4) \quad \tau(t) = t^4 + b_1t^3 + b_2t^2 + b_3t + b_4.$$

From each such τ , we obtain a degree 25 algebra over \mathbb{Q} , once again the algebra determined by the possible values of the variable x .

Changing τ via a rational affine transformation $t \rightarrow \alpha t + \beta$ does not change the degree twenty-five algebra constructed. Accordingly, one can restrict attention to specialization polynomials $\tau(t)$ with $b_1 = 0$, and consider only a set of representatives for the equivalence $(b_2, b_3, b_4) \sim (\alpha^2 b_2, \alpha^3 b_3, \alpha^4 b_4)$, where α is allowed to be

in \mathbb{Q}^\times . In particular, if b_2 and b_3 are nonzero, any such polynomial is equivalent to a unique polynomial of the form

$$(2.5) \quad \tau(u, v, t) = t^4 - 2t^2v - 8tv^2 - 4uv^2 + v^2.$$

Here the reason for the complicated form on the right is explained in the discussion around (3.4). We will treat in what follows only the main two-parameter family where b_2 and b_3 are both nonzero. Note, however, that two secondary one-parameter families are also interesting: if $b_3 = 0$, one gets degree 25 algebras with Galois group in $S_5 \times S_2 \wr S_{10}$, because of the symmetry induced from $t \mapsto -t$; the case $b_2 = 0$ gives rise to full degree 25 algebras, just like the main case.

One can repeat the computation of §2.1, now with the parameters u and v left free. The corresponding general degree twenty-five moduli polynomial $f_{25}(u, v, x)$ has 129 terms as an expanded polynomial in $\mathbb{Z}[u, v, x]$. After replacing x by $5x/4$ and clearing a constant, coefficients average about 16 digits. We will not write this large polynomial explicitly here, instead giving a simpler polynomial that applies only in the special case $u = 1/3$ at the end of §4.2.

2.5. Keeping ramification within $\{2, 3, 5\}$. Suppose $\tau(t)$ from (2.4) normalizes to $\tau(u, v, t)$ from (2.5). We write the corresponding Hurwitz number algebra as $K_{u,v}$. Inclusion (3.12) below says that if $\tau(t)$ is ramified within $\mathcal{P} = \{2, 3, 5\}$, then so is $K_{u,v}$. By a computer search we have found 11031 such (u, v) . From irreducible $f_{25}(u, v, x)$, we obtain $F_{\mathcal{P}}(25) \geq 10938$. The remaining $f_{25}(u, v, x)$ all have a single rational root and from these polynomials we obtain $F_{\mathcal{P}}(24) \geq 93$. The behavior of the 11031 different $K_{u,v}$ will be discussed in more detail in Section 4 below.

A point to note is that ramification is obscured by the passage to standardized coordinates. In the case of our first example $\tau(t) = (t+2)t(t-1)(t-2)$, the corresponding (u, v) is $(37/175, 9/1715)$. The standardized polynomial $\tau(37/175, 9/1715, t)$ after clearing denominators has a 7 in its discriminant.

3. BACKGROUND ON HURWITZ COVERS

In this section, we provide general background on Hurwitz covers. Most of our presentation is in the setting of algebraic geometry over the complex numbers. In the last subsection, we shift to the more arithmetic setting where Hurwitz number fields arise.

3.1. Hurwitz parameters. We use the definition in [19, §1B] of Hurwitz parameter: Let $r \in \mathbb{Z}_{\geq 3}$. An r -point Hurwitz parameter is a triple $h = (G, C, \nu)$ where

- G is a finite group;
- $C = (C_1, \dots, C_k)$ is a list of conjugacy classes whose union generates G ;
- $\nu = (\nu_1, \dots, \nu_k)$ is a list of positive integers summing to r such that $\prod [C_i]^{\nu_i} = 1$ in the abelianization G^{ab} . We henceforth always take the C_i distinct and not the identity, and normalize so that $\nu_i \geq \nu_{i+1}$. The number ν_i functions as a multiplicity for the class C_i .

Table 3.1 gives the Hurwitz parameters of the seven Hurwitz covers described in this paper. It also gives the associated degrees m and bad reduction sets \mathcal{P}_h , each to be discussed later in this section. In the case that G is a symmetric group S_n , we label a conjugacy class C_i by the partition λ_i of n giving the lengths of the cycles of any of its elements. We describe classes for general G in a similar way. Namely we choose a transitive embedding $G \subseteq S_n$. We then label classes C_i by

Section	G	C	ν	m	\mathcal{P}_h
§2	S_5	(2111, 5)	(4, 1)	25	{2, 3, 5}
§5	$S_3 \wr S_2$	(21111, 33, 222)	(3, 1, 1)	9	{2, 3}
§6	S_6	(21111, 222, 3 ₁ 111, 3 _∞ 2 ₀ 1)	(2, 1, 1, 1)	52	{2, 3, 5}
§7	$PSL_3(3)$	(2 ⁴ 1 ⁵ , 3 ³ 1 ⁴)	(3, 2)	2 · 60	{2, 3}
§8	$GL_3(2)$	(22111, 421)	(4, 1)	2 · 96	{2, 3, 7}
§9	S_6	(21111, 3 ₀ 21, 3 ₁ 111, 4 _∞ 11)	(2, 1, 1, 1)	202	{2, 3, 5}
§10	S_6	(21111, 321, 411)	(4, 1, 1)	1200	{2, 3, 5}

TABLE 3.1. Hurwitz parameters for the seven covers pursued in this paper, two of them with normalizations given via subscripts

their induced cycle partitions λ_i , removing any ambiguities which arise by further labeling. In none of our examples is further labeling necessary.

Our concept of Hurwitz parameter emphasizes multiplicities more than other similar concepts in the literature. For example, the first line of Table 3.1 says that our introductory example comes from the parameter $h = (S_5, (2111, 5), (4, 1))$. In e.g. [13], the indexing scheme would center on the class vector $(2111, 2111, 2111, 2111, 5)$.

3.2. Covers indexed by a parameter. An r -point parameter $h = (G, C, \nu)$ determines an unramified cover of r -dimensional complex algebraic varieties

$$(3.1) \quad \pi_h : \text{Hur}_h \rightarrow \text{Conf}_\nu.$$

The base is the variety whose points are tuples (D_1, \dots, D_k) of disjoint subsets D_i of the complex projective line \mathbb{P}^1 , with D_i consisting of ν_i points. Above a point $u = (D_1, \dots, D_k) \in \text{Conf}_\nu$, the fiber has one point for each solution of a moduli problem indexed by (h, u) .

Note that we are using a sans-serif font to indicate smooth complex algebraic varieties, to be thought of simply as complex manifolds in the classical topology. This fonting convention was introduced in [19, §3] and is followed also throughout [15]. As explained around (3.11) below, we switch to a different font when we need to descend to algebraic varieties over \mathbb{Q} . Most of our work takes place at the conceptually simpler complex level, despite the fact that our ultimate concern is the construction of number fields. As another notational convention, we sometimes subscript a projective line by the coordinate we are using; thus \mathbb{P}_t^1 has function field $\mathbb{C}(t)$.

The moduli problem described in [19, §2] involves degree $|G|$ Galois covers $\Sigma \rightarrow \mathbb{P}_t^1$, with Galois group identified with G . An equivalent version of this moduli problem makes reference to the embedding $G \subseteq S_n$ used to label conjugacy classes. When G is its own normalizer in S_n , which is the case for all our examples, the equivalent version is easy to formulate: above a point $u = (D_1, \dots, D_k) \in \text{Conf}_\nu$, the fiber $\pi_h^{-1}(u)$ consists of points x indexing isomorphism classes of degree n covers

$$(3.2) \quad S_x \rightarrow \mathbb{P}_t^1.$$

These covers are required to have global monodromy group G , local monodromy class C_i for all $t \in D_i$, and be otherwise unramified. In this equivalent version, the

ramification numbers of the preimages of $t \in D_i$ in S_x together form the partition λ_i .

We prefer the equivalent version for the purposes of this paper, since it directly guides our actual computations. For example, in our introductory example, the quintic polynomials prominent there can be understood as degree five rational maps $\mathbb{P}_s^1 \rightarrow \mathbb{P}_t^1$. Here \mathbb{P}_s^1 is a common coordinatized version of all the S_x . Also the preimage of ∞ consists of the single point ∞ , explaining why polynomials rather than more general rational functions are involved. At no point did degree 120 maps explicitly enter into the computations of Section 2.

3.3. Covering genus. Let $h = (G, C, \nu)$ be a Hurwitz parameter with $G \subseteq S_n$ a transitive permutation group. Let ℓ_i be the the number of parts of the partition λ_i induced by C_i , and let $d_i = n - \ell_i$ be the corresponding drop. Consider the Hurwitz covers $S_x \rightarrow \mathbb{P}_t^1$ parametrized by $x \in \text{Hur}_h$. By the Riemann-Hurwitz formula, the curves S_x all have genus $g = 1 - n + \frac{1}{2} \sum \nu_i d_i$.

Given G , let d be the minimal drop of a nonidentity element. If h is an r -point Hurwitz parameter based on G , then necessarily $g \geq 1 - n + dr/2$. To support Conjecture 1.1, one needs to draw fields from cases with arbitrarily large r and thus arbitrarily large g . However explicit computation of families rapidly becomes harder as g increases, and in this paper we only pursue cases with genus zero.

3.4. Normalization. The three-dimensional complex group PGL_2 acts by fractional linear transformations on Conf_ν . Since PGL_2 is connected, the action lifts uniquely to an action on Hur_h making π_h equivariant. To avoid redundancy, it is important for us to use this action to replace (3.1) by a cover of varieties of dimension $\rho = r - 3$. Rather than working with quotients in an abstract sense, we work with explicit codimension-three slices as follows.

We say that a Hurwitz parameter is *base normalizable* if $k \geq 3$ and $\nu_{k-2} = \nu_{k-1} = \nu_k = 1$. For a base normalizable Hurwitz parameter, we replace (3.1) by a map of ρ -dimensional varieties,

$$(3.3) \quad \pi_h : X_h \rightarrow U_\nu.$$

Here the target U_ν is the subvariety of Conf_ν with $(D_{k-2}, D_{k-1}, D_k) = (\{0\}, \{1\}, \{\infty\})$. The domain X_h is just the preimage of U_ν in Hur_h . This reduction in dimension is ideal for our purposes: each PGL_2 orbit on Conf_ν contains exactly one point in U_ν .

We say that a base normalizable genus zero Hurwitz parameter is *fully normalizable* if the partitions λ_{k-2} , λ_{k-1} , and λ_k have between them at least three singletons. A normalization is then obtained by labeling three of the singletons by 0, 1, and ∞ , as illustrated twice in Table 3.1. This labeling places a unique coordinate function s on each S_x . Accordingly, each point of X_h is then identified with an explicit rational map from $\mathbb{P}_s^1 \rightarrow \mathbb{P}_t^1$.

When the above normalization conventions do not apply, we modify the procedure, typically in a very slight way, so as to likewise replace the cover of r -dimensional varieties (3.1) by a cover of ρ -dimensional varieties (3.3). For example, two other multiplicity vectors ν figuring into some of our examples are $(4, 1)$ and $(3, 1, 1)$. For these cases, we define

$$\tau_4(t) = t^4 - 2t^2v - 8tv^2 - 4uv^2 + v^2, \quad \tau_3(t) = t^3 + t^2 + ut + v.$$

The form for $\tau_4(t)$ is chosen to make discriminants tightly related:

$$(3.4) \quad \text{disc}_t(\tau_4(t)) = -2^{12}v^6d, \quad \text{disc}_t(t\tau_3(t)) = vd,$$

with

$$(3.5) \quad d = 4u^3 - u^2 - 18uv + 27v^2 + 4v.$$

In the respective cases, we say that a divisor tuple is normalized if it has the form

$$(D_1, D_2) = ((\tau_4(t)), \{\infty\}), \quad (D_1, D_2, D_3) = ((\tau_3(t)), \{0\}, \{\infty\}).$$

These normalization conventions define subvarieties $U_{4,1} \subset \text{Conf}_{4,1}$ and $U_{3,1,1} \subset \text{Conf}_{3,1,1}$. As explained in the (4, 1) setting in §2.4, we are throwing away some perfectly interesting PGL_2 orbits on Conf_ν by our somewhat arbitrary normalization conventions. However all these orbits together have positive codimension in Conf_ν and what is left is adequate for our purposes of supporting Conjecture 1.1. Always, once we have $U_\nu \subset \text{Conf}_\nu$ we just take $X_h \subset \text{Hur}_h$ to be its preimage.

The two base varieties just described are identified by their common coordinates: $U_{4,1} = U_{3,1,1} = \text{Spec } \mathbb{C}[u, v, 1/vd]$. This exceptional identification has a conceptual source as follows. With (u, v) fixed, let $D_1 = (\tau_4(u, v, t))$ so that $(D_1, \{\infty\}) \in U_{4,1}$. Let V be the four-element subgroup of PGL_2 consisting of fractional transformations stabilizing the roots of $\tau_4(u, v, t)$. One then has a degree four map q from \mathbb{P}_t^1 to its quotient $\mathbb{P} := \mathbb{P}_t^1/V$. There are three natural divisors on \mathbb{P} : the divisor Δ_1 consisting of the three critical values, and the one-point divisors $\Delta_2 = \{q(D_1)\}$ and $\Delta_3 = \{q(\infty)\}$. Uniquely coordinatize \mathbb{P} so that $(\Delta_1, \Delta_2, \Delta_3) = ((\tau_3(u', v', t)), \{0\}, \{\infty\})$. Then $u' = u$ and $v' = v$.

3.5. The mass formula and braid representations. The degree m of a cover $X_h \rightarrow U_\nu$ can be calculated by group-theoretic techniques as follows. Define the mass \bar{m} of an r -point Hurwitz parameter $h = (G, C, \nu)$ via a sum over the irreducible characters of G :

$$(3.6) \quad \bar{m} = \frac{\prod_i |C_i|^{\nu_i}}{|G|^2} \sum_{\chi \in \hat{G}} \frac{\prod_i \chi(C_i)^{\nu_i}}{\chi(1)^{r-2}}.$$

Then $\bar{m} \geq m$ always, because $\bar{m} - m$ comes from covers with monodromy group strictly containing G , while m counts covers with the desired monodromy group. In particular, suppose that no proper subgroup $H \subset G$ contains elements from all the conjugacy classes C_i , as is the case in §§2, 5, 6. Then $\bar{m} = m$. When there exist such H , as in §§7, 8, 9, and 10, one can still get exact degrees by applying (3.6) to all such H and computing via inclusion-exclusion. Chapter 7 of [20] gives (3.6) as Theorem 7.2.1 and works out several examples in the setting $r = 3$.

As a one-parameter collection of examples, consider $h(j) = (S_5, (2111, 5), (j, 1))$ for $j \geq 4$ even. Since S_5 is generated by any 5-cycle and any transposition, one has $\bar{m} = m$ for $h(j)$. From 0's in the character table of S_5 , only the characters 1, ϵ , χ , and $\chi\epsilon$ contribute, with ϵ the sign character and $\chi + 1$ the given degree 5 permutation character. We can ignore ϵ and $\chi\epsilon$ by doubling the contribution of 1 and χ :

$$m = \frac{10^j 24}{120^2} \left(2 + 2 \frac{\chi(2111)^j \chi(5)}{\chi(1)^{r-1}} \right) = \frac{10^{j-2}}{6} \left(2 + 2 \cdot \frac{2^j(-1)}{4^{j-1}} \right) = \frac{1}{3} (10^{j-2} - 5^{j-2}).$$

For $j = 4$, one indeed has $m = 25$, as in the introductory example.

The monodromy group of a cover $X_h \rightarrow U_\nu$ can be calculated by group-theoretic techniques [19, §3]. These techniques center on braid groups and underlie the mass formula. The output of these calculations is a collection of permutations in S_m which generate the monodromy group, with $\langle \sigma_1, \sigma_2, \sigma_3 \rangle = S_{25}$ from Figure 2.3 being completely typical. Fullness of these representations is important for us: once we switch over to the arithmetic setting in §3.8, it implies fullness of generic specializations.

Theorem 5.1 of [19] proves a general if-and-only-if result about fullness. In one direction, the important fact for us here is that to systematically obtain fullness one needs for G to be very close to a nonabelian simple group T . Here “very close” includes subgroups of $\text{Aut}(T)$ of the form $T.2$, such as $G = S_n$ for $T = A_n$. This direction accounts for the hypothesis of Conjecture 1.1. In the other direction, fullness is the typical behavior for these G . This statement is the main theoretical reason we expect that the conclusion of Conjecture 1.1 follows from the hypothesis.

3.6. Accessible families. The groups A_n and S_n give rise to many computationally accessible families with $\rho \in \{0, 1, 2\}$. Table 3.2 presents families with $\rho = 2$ and $n \in \{5, 6\}$, omitting 1’s from partitions to save space. The table gives the complete list of h with covering genus $g = 0$ and degree $m \in \{1, \dots, 250\}$. We have verified by a braid group computation that the 58 families listed all have full monodromy group.

$\nu = (3,1,1)$						
n	C_1	C_1	C_1	C_2	C_3	m
6	3	3	3	2	32	216
6	2	2	2	32	5	150
6	2	2	2	32	42	120
6	3	3	3	2	4	96
6	2	2	2	4	5	75
6	2	2	2	4	42	72
6	22	22	22	2	222	60
6	2	2	2	22	6	$C54$
6	2	2	2	32	33	$C54$
5	2	2	2	3	4	48
5	2	2	2	22	4	48
5	2	2	2	3	32	45
6	3	3	3	2	222	44
6	2	2	2	3	6	36
6	2	2	2	4	33	$B36$
5	2	2	2	22	32	$B36$
6	2	2	2	222	5	A25

$\nu = (2,1,1,1)$						
n	C_1	C_1	C_2	C_3	C_4	m
6	22	22	2	3	4	240
6	2	2	3	4	32	202
6	3	3	2	22	4	168
6	2	2	3	22	5	125
6	2	2	3	22	42	100
6	2	2	22	32	222	60
6	22	22	2	3	222	57
6	2	2	3	32	222	52
6	2	2	3	22	33	48
6	3	3	2	22	222	42
6	2	2	22	4	222	40
6	2	2	3	4	222	36

$\nu = (2,2,1)$						
n	C_1	C_1	C_1	C_2	C_3	m
6	2	2	22	22	5	175
6	2	2	4	4	22	158
6	2	2	22	22	42	128
6	2	2	4	4	3	89
6	2	2	3	3	42	80
6	2	2	3	3	5	75
6	2	2	22	22	33	54
5	2	2	3	3	22	58
5	2	2	22	22	3	48
6	2	2	3	3	33	39

$\nu = (5)$						
n	C_1	C_1	C_1	C_1	C_1	m
6	3	3	3	3	3	96

$\nu = (4,1)$						
n	C_1	C_1	C_1	C_1	C_2	m
6	3	3	3	3	22	192
5	2	2	2	2	5	A25

$\nu = (3,2)$						
n	C_1	C_1	C_1	C_2	C_2	m
5	3	3	3	2	2	55
5	22	22	22	2	2	40

TABLE 3.2. Fifty-eight computationally accessible two-parameter families. One, eight, one, and forty-eight of these families respectively have $G = A_5, S_5, A_6,$ and S_6 .

Table 3.2 reveals that our introductory example has the lowest degree m in this context. It and the only other degree 25 family are highlighted in bold. Two of the six families we pursue in §5-10 are likewise put in bold. The remaining families from these sections are not on the table because three of them have group different from A_n and S_n and one has $\rho = 3$.

A remarkable phenomenon revealed by braid computations is what we call *cross-parameter agreement*. There are three instances on Table 3.2: covers given with the same label, be it A , B , or C , are isomorphic. Note that the first instance involves the exceptional isomorphism $U_{4,1} = U_{3,1,1}$ from §3.4, with the cover of $U_{4,1}$ being our introductory family. Many instances of cross-parameter agreement are given with defining polynomials in [15]. Völklein [22] explains some instances of cross-parameter agreement via the Katz middle convolution operator [10].

3.7. Computation and rational presentation. Our general method of passing from a Hurwitz parameter $h = (G, C, \nu)$ to an explicit Hurwitz cover is well illustrated by our introductory example. Very briefly, one writes down all covers $S \rightarrow \mathbb{P}_t^1$ conforming to h and satisfying the chosen normalization conditions. From this first step, one extracts a generator x of the function field of the variety X_h . For all ν we are considering, one has also coordinates u_1, \dots, u_ρ on the base variety U_ν . By computing critical values, one arrives at a degree m polynomial relation $f(u_1, \dots, u_\rho, x) = 0$ describing the degree m extension $\mathbb{C}(X_h)/\mathbb{C}(U_\nu)$. In all the examples of both Table 3.2 and §5-10, the covering variety X_h is connected and so $\mathbb{C}(X_h)$ is a field. In general, as illustrated many times in [15], the polynomial $f(u_1, \dots, u_\rho, x)$ may factor, making X_h disconnected and $\mathbb{C}(X_h)$ a product of fields.

When X_h is a connected rational variety, one can seek a more insightful presentation as follows. One finds not just the above single element x of the function field, but rather elements x_1, \dots, x_ρ which satisfy $\mathbb{C}(X_h) = \mathbb{C}(x_1, \dots, x_\rho)$. Then, working birationally, the map $\pi_h : X_h \rightarrow U_\nu$ is given by ρ rational functions,

$$(3.7) \quad u_i = \pi_{h,i}(x_1, \dots, x_\rho).$$

We call such a system a *rational presentation*.

As an example of a rational presentation, consider the Hurwitz parameter $\hat{h}_{25} = (S_6, (21111, 222, 51), (3, 1, 1))$, chosen because it relates to our introductory example h_{25} by cross-parameter agreement. We partially normalize via $5_\infty 1_0$. We complete our normalization by requiring the coefficient of s^2 in the cubic in the numerator of $g(s)$ be 1:

$$g(s) = \frac{(s^3 + s^2 + zs + y)^2}{as}, \quad \frac{g'(s)}{g(s)} = \frac{5s^3 + 3s^2 + zs - y}{s(x + s^3 + s^2 + sz)}.$$

In the logarithmic derivative of $g(s)$ to the right, let $\Delta(s)$ be its numerator. Writing $g(s) = g_0(s)/g_\infty(s)$, one requires that the resultant $\text{Res}_s(g_0(s) - g_\infty(s)t, \Delta(s))$ be proportional to $t^3 + t^2 + ut + v$. Working out this proportionality makes $a = 4(27 - 225z + 500z^2 + 375y - 5625yz)/3125$.

We have thus identified X_h birationally with the plane $\mathbb{C}_y \times \mathbb{C}_z$. But moreover, the proportionality gives

$$(3.8) \quad u = \frac{5^5 \left(\begin{array}{c} -2025y^3 + 2700y^2z^2 - 405y^2z - 12y^2 - 660yz^3 \\ + 301yz^2 - 36yz + 16z^5 - 8z^4 + z^3 \end{array} \right)}{(-5625yz + 375y + 500z^2 - 225z + 27)^2},$$

$$(3.9) \quad v = -\frac{5^{10}y(27y^2 - 18yz + 4y + 4z^3 - z^2)^2}{(-5625yz + 375y + 500z^2 - 225z + 27)^3}.$$

Equations 3.8 and 3.9 together form a rational presentation of the form (3.7). In general, one can always remove all but one of the x_i by resultants, thereby returning to a ρ -parameter univariate polynomial.

To see the cross-parameter agreement between h_{25} and \hat{h}_{25} explicitly, we proceed as in [17, (5.3) or (5.5)] to identify the root of $f_{25}(u, v, x)$ in the function field $\mathbb{C}(y, z)$. It turns out to be

$$(3.10) \quad x = \frac{3 \cdot 5^7 z (4y^3 - y^2 - 18yz + 27z^2 + 4z)}{2(500y^2 - 5625yz - 225y + 375z + 27)^2}.$$

Thus the natural function x in the first approach has only a rather complicated presentation in the second approach.

3.8. Rationality, descent, and bad reduction. We have been working over \mathbb{C} so far in this section to emphasize that large parts of our subject matter are a mixture of complex geometry and group theory. In the construction of Hurwitz number fields, arithmetic enters “for free” and only at the end. For example, the final equations (3.8) and (3.9) have coefficients in \mathbb{Q} , even though we were thinking only in terms of complex varieties when deriving them.

Following [19, §2D] we say that a Hurwitz parameter $h = (G, C, \nu)$ is *strongly rational* if all the conjugacy classes C_i are rational. This is the case in all our examples, as each C_i is distinguished from all the other classes in G by its partition λ_i . We henceforth work only with strongly rational Hurwitz parameters. In this case, the cover (3.1) canonically descends to a cover of varieties defined over \mathbb{Q} ,

$$(3.11) \quad \pi_h : \text{HUR}_h \rightarrow \text{CONF}_\nu.$$

A standard reference for Hurwitz varieties is [1]. This reference is written from a very different viewpoint from the present paper. For example in the development culminating in §6.2 there, the existence of HUR_h is proved by moduli techniques without reference to \mathbb{C} ; the associated complex variety is recovered as $\text{Hur}_h = \text{HUR}_h(\mathbb{C})$.

Similarly, since all our normalizations are chosen rationally, the corresponding reduced cover (3.3) descends to a cover of \mathbb{Q} -varieties, $\pi_h : X_h \rightarrow U_\nu$. Computations as in our introductory example or the previous subsection end at polynomials $f(u_1, \dots, u_\rho, x) \in \mathbb{Q}[u_1, \dots, u_\rho, x]$ whose vanishing corresponds to (3.11).

Note that in the previous paragraph we conformed to the notational conventions of [19] and [15] by changing fonts as we passed from complex spaces to \mathbb{Q} -varieties. As a further example of this font change, U_ν has appeared many times already as conveniently brief notation for $U_\nu(\mathbb{C})$. In the future we will also need the subsets $U_\nu(R)$ for various subrings R of \mathbb{C} . In subsequent sections we will continue this convention: when working primarily geometrically we emphasize complex spaces, and when specializing we emphasize varieties over \mathbb{Q} .

Let \mathcal{P}_h be the set of primes at which (3.11) has bad reduction. Let \mathcal{P}_G be the set of primes dividing the order of G . Then a fundamental fact is

$$(3.12) \quad \mathcal{P}_h \subseteq \mathcal{P}_G.$$

This fact is essential for our argument supporting Conjecture 1.1, and enters our considerations through (4.1). The inclusion (3.12) follows from the standard reference [1] because all the results there hold for any ground field with characteristic not dividing $|G|$. This good reduction statement is not emphasized throughout [1], but is indicated by the standing convention introduced in §2.1.1 there, that p can be any prime not dividing $|G|$. Table 3.1 gives \mathcal{P}_h for our covers.

4. SPECIALIZATION TO HURWITZ NUMBER ALGEBRAS

This section discusses specializing a given Hurwitz cover $X_h \rightarrow U_\nu$ to number fields, taking the introductory example of Section 2 further to illustrate general concepts. The goal is to extrapolate from the observed behavior of the 11031 algebras $K_{u,v} = K_{(S_5, (2111, 5), (4, 1)), (u, v)}$ to the expected behavior of specialization in general. We dedicate a subsection each to Principles A, B, and C. The extent to which they hold will be discussed in connection with all of our examples in the sequel.

4.1. Algebras corresponding to fibers. Let $X_h \rightarrow U_\nu$ be a Hurwitz cover, as in §3.8. Let $u \in U_\nu(\mathbb{Q})$. The scheme-theoretic fiber $\pi_h^{-1}(u)$ is the spectrum of a separable \mathbb{Q} -algebra $K_{h,u}$. We call $K_{h,u}$ a Hurwitz number algebra. The homomorphisms of $K_{h,u}$ into \mathbb{C} are indexed by points of the complex fiber $\pi_h^{-1}(u) \subset X_h$. Like all separable algebras, the $K_{h,u}$ are products of fields. These factor fields are the Hurwitz number fields of our title. Whenever the monodromy group of $X_h \rightarrow U_\nu$ is transitive, the algebras $K_{h,u}$ are themselves fields for all but a thin set of u , by the Hilbert irreducibility theorem [20, Chapter 3].

For many ν , certainly including all ν containing three 1's, U_ν can be identified with an open subvariety of affine space $\text{Spec } \mathbb{Q}[u_1, \dots, u_\rho]$ as in [18, §8]. Birationally at least, the cover is given by a polynomial equation $f(u_1, \dots, u_\rho, x) = 0$. The point u corresponds to a vector $(u_1, \dots, u_\rho) \in \mathbb{Q}^\rho$. The algebra $K_{h,u}$ is then $\mathbb{Q}[x]/f(u_1, \dots, u_\rho, x)$. The factorization of $K_{h,u}$ into fields corresponds to the factorization of $f(u_1, \dots, u_\rho, x)$ into algebras.

4.2. Real pictures and specialization sets $U_\nu(\mathbb{Z}[1/\mathcal{P}])$. Figure 4.1 draws a window on $U_{4,1}(\mathbb{R})$. With the choice of coordinates made in §3.4, it is the complement of the drawn discriminant locus in the real u - v plane. One should think of the line at infinity in the projectivized plane as also part of the discriminant locus. An analogous picture for $\nu = (2, 1, 1, 1)$ is drawn in Figure 6.1.

Let \mathcal{P} be a finite set of primes with product N . Let $\mathbb{Z}[1/\mathcal{P}] = \mathbb{Z}[1/N]$ be the ring obtained from \mathbb{Z} by inverting the primes in \mathcal{P} . When the last three entries of ν are all 1 then U_ν is naturally a scheme over \mathbb{Z} . Accordingly it makes sense to consider $U_\nu(R)$ for any commutative ring. The finite set of points $U_\nu(\mathbb{Z}[1/\mathcal{P}])$ is studied in detail in [18], including complete identifications for many (ν, \mathcal{P}) . For general ν , one similarly has a finite subset $U_\nu(\mathbb{Z}[1/\mathcal{P}])$ of $U_\nu(\mathbb{Q})$. Its key property for us is that

$$(4.1) \quad \text{for any Hurwitz cover } X_h \rightarrow U_\nu \text{ and any } u \in U_\nu(\mathbb{Z}[1/\mathcal{P}]), \text{ the algebra } K_{h,u} \text{ is ramified within } \mathcal{P}_h \cup \mathcal{P}.$$

In §6.5 we take \mathcal{P} strictly containing \mathcal{P}_h so as to provide examples of ramification known *a priori* to be tame. Otherwise, we are always taking $\mathcal{P} = \mathcal{P}_h$ in this paper. Figure 4.1 shows the 8461 of the known 11031 points of $U_{4,1}(\mathbb{Z}[1/30])$ which fit into the window.




FIGURE 4.1. A window on $U_{4,1}(\mathbb{R})$, which is the complement of the two discriminantal curves in the u - v plane. Points are part of the specialization set $U_{4,1}(\mathbb{Z}[1/30])$, which applies in §2. For §5, §7, and §8, specialization sets $U_{3,1,1}(\mathbb{Z}[1/6])$, $U_{3,1,1}(\mathbb{Z}[1/6])$, and $U_{4,1}(\mathbb{Z}[1/42])$ are respectively used, and the corresponding pictures would have the same discriminant locus but different specialization points.

In our Hurwitz parameter formalism, we emphasize the multiplicity vector ν because of the following important point. Fix r and a non-empty finite set of primes \mathcal{P} , and consider all multiplicity vectors ν with total r . Then $U_\nu(\mathbb{Z}[1/\mathcal{P}])$ tends to get larger as ν moves from (1^r) to (r) . This phenomenon is represented by the two cases considered for $\mathcal{P} = \{2, 3, 5\}$ in this paper: $|U_{2,1,1}(\mathbb{Z}[1/30])| = 2947$, from [18, §8.5], and $|U_{4,1}(\mathbb{Z}[1/30])| \geq 11031$. In fact, as r increases the cardinality $|U_{1^r}(\mathbb{Z}[1/\mathcal{P}])|$ eventually becomes zero [18, §2.4] while $|U_{r-3,1,1}(\mathbb{Z}[1/\mathcal{P}])|$ increases without bound [18, §7]. This increase is critical in supporting Conjecture 1.1.

In both Figure 4.1 and the similar Figure 6.1, one can see specialization points from $U_\nu(\mathbb{Z}[1/30])$ concentrating on certain lines. These lines, and other less visible curves, have the property that they intersect the discriminant locus in the projective plane exactly three times. While the polynomial $f_{25}(u, v, x)$ of §2.4 was too complicated to print, variants over any of these curves are much simpler. For example, the most prominent of the lines is $u = 1/3$. Parametrizing this line by $v = (j - 1)/27j$, one has the simple equation

$$f_{25}(j, x) = 2^2(x + 2) \cdot (729x^8 - 486x^7 - 702x^6 - 8x^5 + 105x^4 + 1118x^3 - 1557x^2 + 1296x - 576)^3 + 5^{15}j(x - 1)^4x^9.$$

The ramification partitions above 0, 1, and ∞ are respectively 3^81 , $2^{10}1^5$, and $(12, 9, 4)$. A systematic treatment of these special curves in the cases $\nu = (3, 1, 1)$

and $\nu = (3, 2)$ is given in [17, §7]. For general ν , they play an important role in [15]. In this paper the above line $v = 1/3$ will play a prominent role in §8, and analogous lines for $\nu = (2, 1, 1, 1)$ will enter in §6.2 and §9.3.

4.3. Pairwise distinctness. For each of the 11031 algebras $K_{u,v}$ of §2.5, and each prime $p \geq 7$, one has a Frobenius partition $\alpha_{u,v,p}$ giving the degrees of the factor fields of $K_{u,v} \otimes \mathbb{Q}_p$. For $p = 7, 11, 13, 17, 19,$ and 23 , the number of partitions of 25 arising is 71, 126, 157, 205, 243, and 302. Taking now $p = 7, 11, 13, 17, 19,$ and 23 as cutoffs, the number of tuples $(\alpha_{u,v,7}, \dots, \alpha_{u,v,p})$ arising is 71, 2992, 10252, 10981, 11027, and 11031. Thus all the known algebras $K_{u,v}$ are pairwise non-isomorphic. There are many other quick ways of seeing this pairwise distinctness. For example, one could use that 6772 different discriminants $D_{u,v}$ arise as a starting point.

Abstracting this simple observation to a general Hurwitz map $X_h \rightarrow U_\nu$ gives

Principle A. *For almost all pairs of distinct elements u_1, u_2 in $U_\nu(\mathbb{Z}[1/\mathcal{P}])$, the algebras K_{h,u_1} and K_{h,u_2} are non-isomorphic.*

So, at least when one restricts to the known elements of $U_{4,1}(\mathbb{Z}[1/30])$, Principle A holds without exception for our introductory family.

In general, the reader should understand our principles as being statements which one could refine in several inequivalent ways into precise conjectures. For example, let G be a finite non-abelian simple group and let \mathcal{P} be the set of primes dividing its order. Then one rigorous refinement is that there is a sequence of Hurwitz parameters $h = (G, C, \nu)$ with $|U_\nu(\mathbb{Z}[1/\mathcal{P}])|$ tending to ∞ , so that Principles A, B, and C all hold with the word “almost” removed. Given the behavior of our examples, we think that this very strong assertion is plausible. However, various much weaker rigorizations of just Principles A and B would also suffice for Conjecture 1.1. We find it best at the moment to not try to speculate on the strongest true rigorization of the three principles. Our repeated use of the phrase “almost all” lets us meaningfully speak about exceptions to these principles. To summarize: we expect exceptions to be very rare in a way that it is premature to quantify.

4.4. Minimal Galois group drop. The Galois group of $f_{25}(u, v, x)$ over $\mathbb{Q}(u, v)$ is S_{25} . Some of the 11031 specialized algebras $K_{u,v}$ have smaller Galois groups as follows. First, in 93 cases, there is a factorization of the form $K_{u,v} = K'_{u,v} \times \mathbb{Q}$, with $K'_{u,v}$ a field. Second, the discriminant of the specializing polynomial $\tau(u, v, t)$ and the discriminant of the degree twenty-five algebra $K_{u,v}$ agree modulo squares. Thus one knows the total number of times that a given discriminant class $d \in \mathbb{Q}^\times / \mathbb{Q}^{\times 2}$ occurs, even without inspecting the $K_{u,v}$ themselves. The number of degree m fields obtained with discriminant class d is as follows:

$m \setminus d$	-30	-15	-10	-6	-5	-3	-2	-1	1	2	3	5	6	10	15	30
25	1050	547	310	363	641	1702	1000	480	557	360	576	572	1026	787	897	70
24	14	3	2	4	5	15	8	4	2	4	10	6	3	1	12	0

Galois groups are as large as possible given the above considerations. Thus A_{25} and A_{24} occur respectively 557 times and twice, leaving S_{25} and S_{24} occurring respectively 10381 and 91 times.

Let Gal_h be the generic Galois group of the cover $X_h \rightarrow U_\nu$.

Principle B. *For almost all elements u in $U_\nu(\mathbb{Z}[1/\mathcal{P}])$, the specialized Galois group $\text{Gal}(K_{h,u})$ contains the derived group Gal'_h of the generic Galois group.*

The most important case of this principle for us is when $X_h \rightarrow U_\nu$ is full, i.e. all of A_m or S_m . Then the principle says that $K_{h,u}$ is full for almost all $u \in U_\nu(\mathbb{Z}[1/\mathcal{P}])$. In our example, 93 of the 11031 known points of $U_{4,1}(\mathbb{Z}[1/30])$, thus slightly less than 1%, are exceptions to the principle. However, in terms of supporting Conjecture 1.1, these exceptions are relatively minor, in that they produce contributors to $F_{\{2,3,5\}}(24)$ rather than $F_{\{2,3,5\}}(25)$.

Principle B is formulated so that it includes other cases of interest to Conjecture 1.1. For example, let $m = m_1 + m_2$ with $m_1, m_2 \geq 3$. Suppose Gal_h is one of the five intransitive groups containing $A_{m_1} \times A_{m_2}$. Then Principle B holds for u if and only if $K_{h,u}$ factors as a product of two full fields. This case is illustrated many times in [15], with splittings of the form $25 = 10 + 15$ and $70 = 30 + 40$ being presented in detail in §6.1 and §6.2 respectively.

4.5. Wild ramification. Consider the discriminants $\text{disc}(K_{u,v}) = \pm 2^a 3^b 5^c$ as (u, v) varies over the known elements of $U_{4,1}(\mathbb{Z}[1/30])$. The left part of Figure 4.2 gives the distribution of the exponents a , b and c . There is much less variation in the exponents than is allowed for field discriminants of degree twenty-five algebras in general. For general algebras, the minimum value for a , b , and c is of course 0 in each case. The maximum values occur for the algebras defined by $(x^{16} - 2)(x^8 - 2)x$, $(x^{18} - 3)(x^6 - 3)x$, and $x^{25} - 5$, and are respectively 110, 64, and 74. The average values in our family are $(\langle a \rangle, \langle b \rangle, \langle c \rangle) \approx (56, 43, 42)$.




FIGURE 4.2. Left: distribution of the discriminant exponents $\text{ord}_p(D)$ the algebras $K_{u,v}$; the variation of $\text{ord}_p(D)$ is much less than is allowed by general discriminant bounds. Right: distribution of the wildness degrees $m_{p\text{-wild}}$ relevant for Principle C.

There are many open questions to pursue with regard to wild ramification. One could ask for lower bounds valid for all u , upper bounds valid for all u , or even exact formulas for wild ramification as a function of u . Principle C is in the spirit of lower bounds. Here we say that a global algebra K is wildly ramified at a prime p if one of the factor fields of its completion K_p is wildly ramified over \mathbb{Q}_p .

Principle C. *For almost all $u \in U_\nu(\mathbb{Z}[1/\mathcal{P}])$, the specialized algebra $K_{h,u}$ is wildly ramified at all primes $p \in \mathcal{P}_h$.*

Certainly, if $\text{ord}_p(K_{h,u}) \geq m$ then Principle C holds for $K_{h,u}$ and p . The left part of Figure 4.2 shows that, for each p , most $K_{u,v}$ satisfy this sufficient criterion. In fact, for $p = 2, 3$, and 5 , there are only 374, 568, and 179 algebras $K_{u,v}$ which do not. However to conform to Principle C at p , an algebra $K_{h,u}$ needs only to satisfy a much weaker condition. Define the *wild degree* of a p -adic algebra K to be the sum of the degrees of its wildly ramified factor fields. Thus in (2.3) these degrees $m_{p\text{-wild}}$ for $p = 2, 3$, and 5 are 16, 15, and 25 respectively. Then conformity to Principle C at p means simply that the p -adic wild degree is positive.

The right part of Figure 4.2 gives the distribution of $m_{p\text{-wild}}$. For example, for $p = 5$, there are 179 exceptions to Principle C, including all the 93 factorizing algebras. Besides these exceptions, all algebras have $m_{5\text{-wild}}$ at its maximum possible value of 25.

4.6. Expectations. As discussed in [19, §8], the Hilbert irreducibility theorem, applied to $X_h \rightarrow U_\nu$ and $X_h \times X_h \rightarrow U_\nu \times U_\nu$ respectively, already points in the direction of Principles A and B. In a wide variety of contexts, analogs of these principles hold with great strength. For example, in [14, §9] several covers are discussed in the setting $\mathcal{P} = \{2, 3\}$ and for most of them both Principles A and B hold without exception. However the situation we consider here, with fixed \mathcal{P} and arbitrarily large degree m , is outside the realm of previous experience. Explicitly verifying the principles in degrees large enough to contradict the mass heuristic is important for being confident that these standard expectations do indeed hold in this new realm.

We are confident that for a given G and varying $h = (G, C, \nu)$, one has strict inclusion $\mathcal{P}_h \subset \mathcal{P}_G$ for only finitely many (C, ν) . This expectation, together with Principle C, suggests that there are only finitely many full fields $K_{h,u}$ ramified strictly within \mathcal{P}_G . One possibility is that full number fields coming from Hurwitz-like constructions are the main source of outliers to the mass heuristic. If one believes this, then one is led to the first of the two extreme possible complements to Conjecture 1.1 discussed at the end of [19]: *The sequence $F_{\mathcal{P}}(m)$ always has support on a density zero set, and it is eventually zero unless \mathcal{P} contains the set of primes divisors of the order of a nonabelian finite simple group.* Our verification that Principle C holds with great strength in our examples is supportive of this very speculative assertion.

5. A DEGREE 9 FAMILY: COMPARISON WITH COMPLETE NUMBER FIELD TABLES

This section begins our sequence of six sample families of increasing degree. To start in very low degree, we take G solvable. The number fields coming from this first example are not full and so not directly relevant to Conjecture 1.1. This family is nonetheless a good place to begin our presentation of examples, for two reasons. First, the low degree makes comparison with complete tables of number

fields possible. Second, there are many exceptions to Principles A, B, and C. These exceptions form the first data-point arguing for the expectation already formulated in the introduction: as the degree of the Hurwitz family increases, the frequency of exceptions decreases.

5.1. A Hurwitz parameter with solvable G . Let G be the wreath product $S_3 \wr S_2$ of order 72, considered as a subgroup of S_6 . The group G has unique conjugacy classes with cycle type 21111, 222, and 33. Take $h = (G, (21111, 222, 33), (3, 1, 1))$. Then $\bar{m}_h = m_h = 9$.

5.2. A two-parameter polynomial. In the present context of $\nu = (3, 1, 1)$, our normalized specialization polynomials take the form

$$\tau(u, v, t) = (t^3 + t^2 + ut + v)t.$$

The discriminant of the cubic factor is $d = 4u^3 - u^2 - 18uv + 27v^2 + 4v$ from (3.5). A nonic polynomial capturing the family and a resolvent octic are as follows:

$$\begin{aligned} f_9(u, v, x) &= x^9 - 3x^8 + 12ux^7 - 4(u + 12v)x^6 + 42vx^5 - 6(4u + 1)vx^4 \\ &\quad + 4v(2u + 3v)x^3 - 12v^2x^2 + 3(4u - 1)v^2x - v^2(4u - 8v - 1), \end{aligned}$$

$$\begin{aligned} f_8(u, v, x) &= x^8 + x^4(18v - 6u^2) + x^2(8u^3 - 36uv + 108v^2) \\ &\quad + (-3u^4 + 18u^2v - 27v^2). \end{aligned}$$

Here $f_9(u, v, x)$ and $f_8(u, v, x)$ respectively have Galois group $9T26 = \mathbb{F}_3^2.GL_2(\mathbb{F}_3)$ and $8T23 = GL_2(\mathbb{F}_3) = \tilde{S}_4$. Because of the complete lack of singletons in the partitions 222 and 33, our computation of $f_9(u, v, x)$ required substantial *ad hoc* deviations from the procedure sketched in §3.7.

The discriminants of the two polynomials are respectively

$$D_9(u, v) = -2^{24}3^9v^{10}d^4(27v - 1)^6, \quad D_8(u, v) = -2^{24}3^{19}v^8d^4(u^2 - 3v)^2.$$

In each case, the discriminant modulo squares is -3 . Because of this constancy, the Galois groups of $f_9(u, v, x)$ and $f_8(u, v, x)$ over $\mathbb{C}(u, v)$ are respectively the index two subgroups $9T23 = \mathbb{F}_3^2.SL_2(\mathbb{F}_3)$ and $8T12 = SL_2(\mathbb{F}_3) = \tilde{A}_4$. The last factor of the discriminant in each case is an artifact of our particular polynomials. Because of these factors, one knows that that if $v = 1/27$ or $v = u^2/3$, the algebra $K_{u,v}$ has to be in some way degenerate. However if $v \neq 1/27$ and $v \neq u^2/3$, then these factors do not contribute to field discriminants in specializations.

5.3. Comparison of specializations with complete tables of number fields.

We work with 507 pairs (u, v) in $U_{3,1,1}(\mathbb{Z}[1/6])$. Twenty-one of them have $v = 1/27$ and so $f_9(u, v, x)$ is not separable. For forty more, $f_9(u, v, x)$ also reduces, with the factorization partitions 63, 81, 6111, and 333 occurring respectively 9, 29, 1, and 1 times. The remaining 446 specialization points yield only 129 different fields, as for example $(-13/12, 2/9)$, $(11/12, 1/9)$, $(-5/12, -8/27)$, $(1/4, -1/27)$, $(1/4, 2/27)$, $(35/108, 8/243)$, $(1/4, 64/3375)$, and $(19/2028, 1/59319)$ all yield the field defined by $x^9 - 9x^7 + 27x^5 - 27x^3 - 4$. Moreover, a wide variety of subgroups of $9T26$

appear, as follows.

Group G :	9T4	9T8	9T12	9T13	9T16	9T18	9T19	9T26
Size $ G $:	18	36	54	54	72	108	144	432
Number of fields in family:	2	1	10	1	5	20	8	82
Total number of fields:	4	1	12	3	5	23	8	87

The last line compares with the relevant complete lists at the website associated to [9]. It gives the total number of number fields with the given Galois group and with discriminant of the form $-2^a 3^b$ with a even and b odd. One can get even a larger fraction of the total number of fields by specializing outside of $U_{3,1,1}(\mathbb{Z}[1/6])$, both by considering the curve at infinity and then by specializing also at the rare-but-existent points of say $U_{3,1,1}(\mathbb{Z}[1/6p])$, where the auxiliary prime p does not divide the discriminant of the field constructed. The fact that such a large fraction of all fields of the type considered come from a single Hurwitz family is suggestive that other Hurwitz families may be essentially the only source of number fields with certain invariants.

5.4. Exceptions to Principles A, B, and C. The current family presents many examples of phenomena that Principles A, B, and C say are rare in general. The drop from 446 specialization points giving nonic fields to only 129 isomorphism classes of fields constitutes many exceptions to Principle A. The further drop from 129 fields to just 82 fields with the generic Galois group includes many exceptions to Principle B. Some of the specializations are tamely ramified or even unramified at 2, and thus correspond to exceptions to Principle C.

6. A DEGREE 52 FAMILY: TAME RAMIFICATION AND EXCEPTIONS TO PRINCIPLE B

In our introductory family, the only exceptions to Principle B were algebras of the form $K_{h,u} = \mathbb{Q} \times K'_{h,u}$ with $K'_{h,u}$ full. In specializing many other full families, most of the exceptions to Principle B we have found have this very same form. In this section, we present a family which is remarkable because some of its specializations have a much more pronounced drop in fullness. However we do not regard this more serious failure of Principle B as anywhere near extreme enough to raise doubts about Conjecture 1.1.

6.1. A Hurwitz parameter yielding a rational X_h . We start from the normalized Hurwitz parameter

$$h = (S_6, (21111, 222, 3_1 111, 3_\infty 2_0 1), (2, 1, 1, 1)).$$

All rational functions with this normalized Hurwitz parameter have the form

$$g(s) = \frac{(s^3 + bs^2 + cs + x)^2}{as^2(s - y)}.$$

The ramification requirement on g at 1 is that $(g(1), g'(1), g''(1)) = (1, 0, 0)$. These three equations allow the elimination of a , b , and c via

$$\begin{aligned} a &= -64(x + 1)^2(y - 1)^3, \\ b &= 4xy - 3x + 4y - 6, \\ c &= -8xy^2 + 12xy - 6x - 8y^2 + 12y - 3. \end{aligned}$$

Using a resolvent as usual, we find that the critical values of $g(s)$ besides 0, 1, and ∞ are the roots of $Wt^2 + (V - U - W)t + U$ where

$$\begin{aligned} U &= (4xy - x + 3y) (64x^2y^4 - 160x^2y^3 + 180x^2y^2 - 108x^2y + 27x^2 + 256xy^4 \\ &\quad - 736xy^3 + 864xy^2 - 540xy + 162x + 192y^4 - 576y^3 + 576y^2 - 216y + 27)^2, \\ V &= 3^3(2xy - x + 1)^4 (64xy^3 - 144xy^2 + 108xy - 27x + 64y^3 - 144y^2 + 81y), \\ W &= 2^{12}3^3(x + 1)^4(y - 1)^6y^3. \end{aligned}$$

Comparing with the standard quadratic $t^2 + (v - u - 1)t + u$, one gets the rational presentation

$$(6.1) \quad u = \frac{U}{W}, \quad v = \frac{V}{W}.$$

Summarizing, birationally we have $X_h = \mathbb{C}_x \times \mathbb{C}_y$, $U_\nu = \mathbb{C}_u \times \mathbb{C}_v$, and the equations (6.1) give the map $X_h \rightarrow U_\nu$. Removing y by a resolvent gives the single equation $f_{52}(u, v, x) = 0$. Likewise removing x by a resolvent gives the single equation $\phi_{52}(u, v, y) = 0$. The left sides have 2781 and 829 terms respectively.

The discriminants of $f_{52}(u, v, x)$ and $\phi_{52}(u, v, y)$ are both -3 times a square in $\mathbb{Q}(u, v)$. The Galois groups of these polynomials over $\mathbb{Q}(u, v)$ are S_{52} , but over $\mathbb{C}(u, v)$ they reduce to A_{52} . This general phenomenon appeared already in the previous section. It is not of central importance to us, which is why we generally refer to full fields and only sometimes make the distinction between S_m and A_m fields.

6.2. Specialization to curves. Using homogeneous coordinates U, V , and W , related to our standard coordinates u and v via (6.1), we can view $U_{2,1,1,1}$ as completed by the projective plane. Its complement in this projective plane has four components,

- A: the vertical line $U = 0$,
- B: the horizontal line $V = 0$,
- C: the line at infinity $W = 0$, and
- D: the conic $U^2 + V^2 + W^2 - 2UV - 2UW - 2VW = 0$.

Figure 6.1 draws A, B, and D. Note that lines A, B, and C pass through points $\mathbf{a} = (0 : 1 : 1)$, $\mathbf{b} = (1 : 0 : 1)$, and $\mathbf{c} = (1 : 1 : 0)$ respectively, while the conic D goes around $\mathbf{d} = (1 : 1 : 1)$. Note also that while this completion to a projective plane has the virtue of introducing a convenient S_3 symmetry, it is not particularly natural from a moduli-theoretic viewpoint.

A general line in the projective plane intersects the discriminant locus in five points. However the lines that go through two of the points in $\{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}\}$ intersect the discriminant locus only three times. These six lines are parametrized in Table 6.1, so that the three points become 0, 1, and ∞ . Exactly as in Figure 4.1 earlier, the lines are clearly suggested by the drawn specialization points. Having used homogeneous coordinates for two paragraphs to make an S_3 symmetry clear, we now return to our standard practice of focusing on the affine u - v plane.

When restricted to any one of the six lines, the cover X_h remains full. This preserved fullness is in the spirit of Principle B. Table 6.1 gives the ramification partitions of these restricted covers. Note that all partitions are even, reflecting the fact that the monodromy group is only A_{52} . Before beginning any computations




FIGURE 6.1. A window on $U_{2,1,1,1}(\mathbb{R})$, which is the complement of the three thick discriminantal curves in the $u-v$ plane. Points are part of the specialization set $U_{2,1,1,1}(\mathbb{Z}[1/30])$ which is used in both §6 and §9. The thin parabola $(u - v)^2 = 4v$ plays a role only in §6.

with polynomials, we knew these partitions and the fullness of the six covers from a braid group computation.

Line	u	v	λ_0	λ_1	λ_∞	genus
ad	$4t$	1	$12^2 6 4^4 3 2 1$	$3^8 2^{12} 1^4$	$20 12 5 4^3 2 1$	6
bd	1	$4t$	$10^2 8 6^3 5 1$	$3^8 2^{12} 1^4$	$10^2 6^2 5 4^2 2^2 1^3$	5
cd	$t/4$	$t/4$	$4^9 2^7 1^2$	$3^8 2^{12} 1^4$	$6^4 4^6 2^2$	0
bc	t	$t - 1$	$2^{22} 1^8$	$10^2 8 6^3 5 1$	$6^4 4^6 2^2$	2
ac	$t - 1$	t	$4^9 2^3 1^{10}$	$12^2 6 4^4 3 2 1$	$6^4 4^6 2^2$	5
ab	t	$1 - t$	$12^2 6 4^4 3 2 1$	$10^2 8 6^3 5 1$	$5^5 4^2 3^4 1^7$	9

TABLE 6.1. Six lines in $U_{2,1,1,1}$ and topological information on their preimages in X_h .

To give an explicit degree 52 polynomial coming from the cover $X_h \rightarrow U_{2,1,1,1}$, we work over the line cd. The preimage of cd is a curve in the $x-y$ plane with equation having x -degree 3, y -degree 6, and twenty-two terms. A parametrization

is

$$x = -\frac{(s^2 - 2s - 2)(s^4 - 4s^3 + 4s + 2)}{2s^3}, \quad y = -\frac{3(s^2 - 2s - 2)}{2(s-2)(s^2 - 4s - 2)}.$$

Using the domain coordinate s and the target coordinate t , the restricted rational function takes the form

$$(6.2) \quad t = \frac{-A}{C} = \frac{B}{C} + 1.$$

Here A , B , and C sum to zero and are given explicitly by

$$\begin{aligned} A &= (s+1)^4 (s^8 - 10s^7 + 34s^6 - 40s^5 - 2s^4 + 8s^3 + 8s^2 + 16s + 8)^4 \\ &\quad (s-2)^2 (s^2 - 4s - 2)^2 (s^4 - 6s^3 + 9s^2 - 6)^2 (s-4)s, \\ B &= -(s^8 - 12s^7 + 52s^6 - 92s^5 + 30s^4 + 96s^3 - 72s^2 - 48s + 8)^3 \\ &\quad (s^{12} - 12s^{11} + 48s^{10} - 52s^9 - 87s^8 + 108s^7 + 264s^6 - 216s^5 \\ &\quad - 312s^4 + 48s^3 + 192s^2 + 96s + 16)^2 (s^4 - 4s^3 + 4s + 2), \\ C &= 2^2 (2s^3 - 9s^2 + 6s + 2)^6 (s^6 - 6s^5 + 6s^4 + 10s^3 - 6s^2 - 12s - 4)^4 \\ &\quad (s^2 - 2s - 2)^2. \end{aligned}$$

This explicit slice is intended to give a sense of the full cover for h_{52} , just as the slices in §4.2 and §8.2 indicate the covers for h_{25} and h_{96}^* respectively. Here we have explicitly presented information at all three of the cusps, not just at 0 and ∞ as in §4.2 and §8.2.

6.3. No exceptions to Principles A and C. Unlike all our previous examples, the current ν contains at least three ones. It thus fits into the framework of [18], where many $U_\nu(\mathbb{Z}[1/P])$ for such ν are completely identified. We therefore can be more definitive in reporting specialization results.

The set $U_{2,1,1,1}(\mathbb{Z}[1/30])$ contains exactly 2947 points [18, §8.5] and is drawn in Figure 6.1. The Hurwitz number algebras $K_{h,u}$ are all non-isomorphic, so that Principle A holds without exception. All 2947 algebras are wildly ramified at all three of 2, 3, and 5, so that Principle C also holds without exception; in fact $\text{ord}_p(D) \geq 52$ fails at $p = 2, 3, \text{ and } 5$ only 0, 60, and 481 times, so the verification of Principle C is particularly easy at $p = 2$.

6.4. Easily explained exceptions to Principle B. Twenty-five of the 2947 specialization points (u, v) give exceptions to Principle B. Three of these, namely $(3/8, 1/8)$, $(1/16, -375/16)$, $(16, -375)$ are exceptions of the sort we have seen earlier: $K_{h,u} = \mathbb{Q} \times K'_{h,u}$ with $K'_{h,u}$ full. Exceptions of this nature are not surprising whenever the Hurwitz cover is rational. In this case the three points in question come respectively from points $(-1/6, 3/8)$, $(-4/3, 3/4)$, and $(-3, 3/4)$ in $X_h(\mathbb{Q})$.

In fact, any point $(x, y) \in X_h(\mathbb{Q})$ causes such a factorization, because it is a rational point above its image $u = (u, v) \in U_\nu(\mathbb{Q})$. However even for very low height (x, y) , the algebra $K_{h,u}$ is typically ramified at extraneous primes. For example, take $s = 1$ in the equations after (6.2), making

$$(6.3) \quad t = \frac{111936400}{43923} = \frac{2^4 5^2 23^4}{3 \cdot 11^4} = 1 + \frac{37^3 47^2}{3 \cdot 11^4}.$$

The number field K' defined by the degree 51 factor of $f_{52}(t, s)$ has discriminant $-2^{102}3^{95}5^{48}11^{16}37^{12}47^{16}$. Whenever we discuss exceptions to Principle B, we always have in mind a fixed \mathcal{P} , here $\{2, 3, 5\}$, and do not consider fields like K' to be exceptions.

6.5. Ramification at tame primes. We are confident that that ramification at p in a Hurwitz number algebra $K_{h,u}$ can only be wild if $p \in \mathcal{P}_h$ or $p \leq \max_i \nu_i$. The field K' from the previous subsection presents a convenient opportunity to illustrate how ramification in $K_{h,u}$ at the remaining primes should be calculable in purely group-theoretic terms.

To describe the factorization of the local algebras K'_p , we represent the fields appearing by symbols e_c^f as in (2.3). We simplify by just writing e^f for tame fields, since tameness implies $c = e - 1$. The factorizations are

$$\begin{aligned} 2: & \mathbf{16}_{38} \mathbf{16}_{38} \mathbf{8}_{16} \mathbf{2}_3 \mathbf{2}_3 \mathbf{2}_2 \mathbf{2}_2 1^2 1, & 11: & 3^6 3^2 1^{16} 1^2 1^2 1^2 1^2 1 1 1 \rightarrow 3^8 1^{27}, \\ 3: & \mathbf{18}_{39} \mathbf{12}_{21}^2 \mathbf{6}_{11} \mathbf{3}_3, & 37: & 2^5 2^4 2^2 2 1^{18} 1^3 1 1 1 1 1 1 \rightarrow 2^{12} 1^{27}, \\ 5: & \mathbf{25}_{40} 2^4 2^4 1^4 1^2 1 1 1 1, & 47: & 3^6 3^2 1^4 1^4 1^4 1^4 1^2 1 \rightarrow 3^8 1^{27}. \end{aligned}$$

The wild primes behave in a complicated way as always, with p -wildness at $p = 2, 3$, and 5 being 48, 51, and 25. However the tame primes are much more simply behaved.

To work at an even simpler level, we factor over the maximal unramified extension of \mathbb{Q}_p , rather than \mathbb{Q}_p itself. For tame primes, this corresponds to regarding the printed exponents f simply as multiplicities, and collecting together symbols with a common base. The resulting tame ramification partitions are indicated to the right, after arrows.

Note that there are actually four primes greater than 5 involved in (6.3). With their naturally occurring exponents, 11^4 is associated to ∞ , 23^4 to 0, and 37^3 and 47^2 to 1. In general, tame ramification partitions can be computed from the placement of the specialization point in $U_\nu(\mathbb{Q}_p)$ and braid group considerations. In the setting of three-point covers, the general formula is simple, and uses the standard notion of the power of a partition. Namely, if p^m is associated to $\tau \in \{0, 1, \infty\}$ its tame ramification is the power λ_τ^m of the geometric ramification partition λ_τ . Applying the cd line of Table 6.1, the partitions $\lambda_\infty^4 = 3^8 1^{27}$, $\lambda_0^4 = 1^{51}$, $\lambda_1^3 = 2^{12} 1^{27}$, and $\lambda_1^2 = 3^8 1^{27}$ do indeed agree with the partitions found by direct factorization of the polynomial defining K' .

The mass heuristic reviewed in §1.1 is based on an equidistribution principle. In the horizontal direction, it translates to the following conjecture, proved for $m \leq 5$: when one considers full degree m fields ordered by their absolute discriminant outside of p , all tame ramification partitions are asymptotically equally likely. We regard the fact that Hurwitz number fields escape the mass heuristic as being directly related to their highly structured ramification. In the current instance, there are 239,943 partitions of the integer 51, and the two partitions $2^{12} 1^{27}$ and $3^8 1^{27}$ are far from typical.

6.6. A curve of more extreme exceptions to Principle B. The twenty-two exceptions not discussed in §6.4 all have a common geometric source: above the base curve B given by $(u - v)^2 = 4v$, the cover splits into a full degree 42 cover of genus five and a full degree 10 cover C of genus zero. While decompositions $52 = 51 + 1$ are governed by rational points on X_h itself, decompositions of the

form $52 = 42 + 10$ are governed by rational points on a resolvent variety of degree $\binom{52}{10}$ over $U_{2,1,1,1}$. As this degree is about 15 billion, the existence of an entire curve of rational points is remarkable.

To reveal the structure of the cover $C \rightarrow B$, we parametrize the base curve B via

$$(6.4) \quad u = \frac{4t}{(t-1)^2}, \quad v = \frac{4}{(t-1)^2}.$$

In the decompositions $K_{h,(u,v)} = K_t^{42} \times K_t$, the twenty-two K_t^{42} are all full degree forty-two fields, with pairwise distinct discriminants.

The genus zero curve C is given by $x(4y-3)^3 = -24y^2(2y-3)$, and so y is a parameter. The map from the y -line C to the t -line B is given by the vanishing of

$$f_{10}(t, y) = (4y-3)(8y-3)(32y^4 - 192y^3 + 360y^2 - 252y + 27)^2 \\ + t(4y-9)(96y^4 - 256y^3 + 216y^2 - 108y + 27)^2.$$

Thus one has two visible ramification partitions $\lambda_0 = \lambda_\infty = 222211$. The discriminant of $f_{10}(t, y)$ is $-2^{136}3^{57}5^{25}t^4(t-1)^5(t-9)^5$. At the other singular values, the ramification partitions are $\lambda_1 = \lambda_9 = 32221$. In fact, the decic algebras K_t and $K_{9/t}$ are isomorphic via the involution $y \mapsto (6y-9)/(8y-6)$.

At the level of the decic cover only, we have just indicated a failure of Principle A: rather than 22 distinct decic algebras, there are ten pairs switched by $t \leftrightarrow 9/t$ and then two algebras K_3 and K_{-3} arising once each. The ten algebras arising twice are all full fields and wildly ramified at all three of 2, 3, and 5. However K_3 and K_{-3} are not full, and not wildly ramified at 5, giving failures of Principle B and C at this decic cover level.

In terms of supporting Conjecture 1.1 for $\mathcal{P} = \{2, 3, 5\}$, the exceptional behavior above B is in a sense good. Instead of twenty-two contributions to $F_{\mathcal{P}}(52)$, one gets twenty-two contributions to $F_{\mathcal{P}}(42)$ and then ten more to $F_{\mathcal{P}}(10)$. But in another sense this exceptional behavior is bad. It explicitly illustrates phenomena which, if occurring ubiquitously in high degree, might make Conjecture 1.1 false. However our computations suggest that, far from becoming ubiquitous, the phenomena exhibited here become rarer as degrees increase.

7. A DEGREE 60 FAMILY: NON-FULL MONODROMY AND A PRIME DROP

The statement of Conjecture 1.1 involves all finite nonabelian simple groups equally. In this paper, however, we focus on the simple groups A_5 and A_6 because of the computational accessibility of the corresponding families $X_h \rightarrow U_\nu$. In this section and the next, we add some balance by presenting results on covers coming from simple groups not of the form A_n . The family presented here has the particular interest that it is non-generic in two ways.

7.1. A Hurwitz parameter with unexpectedly non-full monodromy. The simple group $G = PSL_3(\mathbb{F}_3)$ has order $5616 = 2^4 \cdot 3^3 \cdot 13$ and outer automorphism group of order two. It has two non-isomorphic degree 13 transitive permutation representations, coming from an action on a projective plane $\mathbb{P}^2(\mathbb{F}_3)$ and its dual $\widehat{\mathbb{P}^2}(\mathbb{F}_3)$. These actions are interchanged by the outer involution. The two smallest non-identity conjugacy classes in G consist of order 2 and order 3 elements. In each of the degree 13 permutation representations, these elements act with cycle structure $2^4 1^5$ and $3^3 1^4$ respectively.

Let $h = (PSL_3(\mathbb{F}_3), (2^4 1^5, 3^3 1^4), (3, 2))$. To conform to our main reference [11] for this section, we make a quadratic base change and work over $U_{3,1,1}$ rather than $U_{3,2}$. A braid group computation reveals that the degree 120 Hurwitz cover π_h factors as a composition of three covers as indicated:

$$(7.1) \quad X_h \xrightarrow{2} X_h^* \xrightarrow{15} \text{Quart} \xrightarrow{4} U_{3,1,1}.$$

The intermediate cover X_h^* is just the quotient of X_h by the natural action of $\text{Out}(G)$. This failure of fullness illustrates one of the general phenomena treated at length in [19].

However, very unusually in comparison with Table 3.2, the reduced Hurwitz cover $\pi_h^* : X_h^* \rightarrow U_{3,1,1}$ is also not full. It clearly fails to be primitive, because of the intermediate cover **Quart**. Moreover, the degree fifteen map is not even full, as its monodromy group is S_6 in a degree 15 transitive representation.

The degree 13 covers of the projective line parametrized by X_h have genus zero. Using this fact as a starting point, König [11, §7] succeeded in finding coordinates a, b on X_h , with corresponding covers $P_s^1 \rightarrow P_t^1$ being as follows. Define

$$\begin{aligned} f_0 &= \frac{abs}{3} + \frac{ab}{9} + as^2 - \frac{a}{3} + s^3, \\ f_1 &= \frac{s^2(ab^2 - 4ab + 12a - 3b^2 - 9)}{(b-3)^2} + \frac{s(ab^2 - 4ab + 12a - 9b - 9)}{3(b-3)} + s^3 - 1, \\ g_0 &= \frac{abs}{3} + \frac{ab}{9} + as^2 - \frac{a}{3} + s^3, \\ g_1 &= \frac{1}{9}s(4ab^2 - 6ab + 9a + 9b - 27) + \frac{1}{3}s^2(4ab - 3a + 9) + as^3 - a. \end{aligned}$$

Then the two-parameter family is given by $g(a, b, t, s) := f_0^3 f_1 s - t g_0^3 g_1 = 0$.

König's interest in this family is in producing number fields with Galois group G . For example $(a, b, t) = (-9, -6, -3)$ gives a totally real such field with discriminant $3^{12} 251^4 353^4$. To systematically study specializations, it is important to determine the discriminant of $g(a, b, t, s)$. Computation shows that it has the following form:

$$D(a, b, t) = \left(-\frac{4}{3}ab^3 + a^2b^2 + 6ab^2 - 3b^2 - 4a^2b - 18ab + 18b + 12a^2 - 27 \right)^{28} a^{12}(b-3)^{18}t^6 (C_0t^3 + C_1t^2 + C_2t + C_3)^4.$$

Here $C_0, C_1, C_2,$ and C_3 as expanded elements of $\mathbb{Q}[a, b]$ have 24, 45, 53, and 36 terms respectively. Because of the complicated nature of this discriminant, it is hard to get field discriminants to be as small as the one exhibited above.

For König's purposes of constructing degree thirteen fields with Galois group G , he does not need the map to configuration space at all. To move over into our context of constructing Hurwitz number fields, we do need this map. Replacing t in $(C_0t^3 + C_1t^2 + C_2t + C_3)$ with C_1t/C_0 and setting the resulting cubic proportional to $t^3 + t^2 + ut + v$ gives a degree 120 map π_h from the a - b plane X_h to the u - v plane $U_{3,1,1}$. Removing a from the pair of equations gives a degree 120 polynomial $f_{120}(u, v, b) \in \mathbb{Q}(u, v)[b]$ describing the covering map.

7.2. Reduction to degree 60. To reduce from the degree 120 cover X_h to the degree 60 cover X_h^* , we proceed as follows. For $(a_i, b_i) \in \mathbb{Q}^2$, one gets $(u_i, v_i) = \pi_h(a_i, b_i) \in \mathbb{Q}^2$. Then $f_{120}(u_i, v_i, b) \in \mathbb{Q}[b]$ factors. For almost all choices of (a_i, b_i) , the degrees of the irreducible factors are 90, 6, 6, 4, 4, 4, 4, 1, and 1. One of the

linear factors is $b - b_i$ and we write the other one as $b - b'_i$. Then typically just one rational number a'_i satisfies the two equations $\pi_h(a'_i, b'_i) = (u_i, v_i)$. From enough datapoints we interpolate to get the canonical involution on X_h . It is

$$(7.2) \quad a' = \frac{(b-3)(4ab+6a+9)}{ab^2-4ab+12a-18b+18}, \quad b' = \frac{3b}{b-3}.$$

This involution is useful even in König's context. For example, specializing at $(a', b', t) = (171/58, 2, -3)$ gives the dual totally real number field, also with discriminant $3^{12}251^4353^4$.

A quantity stabilized by the involution is $x = b^2/(b-3)$. The resolvent

$$\text{Res}_b(f_{120}(u, v, b), (b-3)x - b^2)$$

is proportional the square of a degree 60 polynomial $f_{60}(u, v, x)$. This polynomial captures the cover $X_h^* \rightarrow U_{3,1,1}$.

7.3. Low degree resolvents. From the braid group computation, we know that the monodromy group has quotients of type S_3 , S_4 , and $8T40 = 2^3.S_4$. Here the S_4 quotient corresponds to the cover **Quart**. Equations for these quotients and their discriminants are

$$\begin{aligned} f_3(u, v, x) &= x^3 + x^2 + xu + v, & D_3 &= d, \\ f_4(u, v, x) &= x^4 - 2x^2v - 8xv^2 - 4uv^2 + v^2, & D_4 &= -2^{12}dv^6, \\ f_8(u, v, x) &= x^8 + 8x^4duv - 72x^4dv^2 & D_8 &= -2^{60}d^{17}v^{14}(u^3 - v)^4 \\ &\quad + 64x^2d^2v^2 - 16d^3v^2, \end{aligned}$$

Here we have seen the cubic and quartic polynomials in §3.4, with d being given explicitly in (3.5).

7.4. Reduction to degree 24. The equation $f_4(u, v, m) = 0$ is linear in u . Solving it gives $u = (m^4 - 2m^2v + v^2 - 8mv^2)/(4v^2)$. Expressing $f_{60}(u, v, x)$ in terms of m, v , and x and factoring, one gets $g_{15}(m, v, x)g_{45}(m, v, x)$. Here $g_{15}(m, v, x)$ has Galois group S_6 over $\mathbb{Q}(m, v)$, in a degree 15 permutation representation.

Abbreviate $e = m^3 - mv - 2v^2$. Then the polynomial for the standard sextic representation works out to

$$g_6(m, v, x) = 2x^6v^2 - 3x^4e(m^2 - v) - 8x^3e^2 - 6x^2e^2m + 2e^3.$$

Returning to the original base, one gets a degree 24 polynomial,

$$f_{24}(u, v, x) = \text{Res}_m(f_4(u, v, m), g_6(m, v, x)).$$

Similarly, by means of the outer automorphism of S_6 , one has a twin polynomial $g_6^t(m, v, x)$ and its degree 24 polynomial $f_{24}^t(u, v, x)$. While $f_{60}(u, v, x)$, $f_{24}(u, v, x)$, and $f_{24}^t(u, v, x)$ all have the same splitting field, the latter two are much easier to work with because of their lower degree.

7.5. Specialization to number fields. We have specialized at the 507 points in $U_{3,1,1}(\mathbb{Z}[1/6])$ considered in §5, obtaining 507 algebras with discriminant of the form $\pm 2^a 3^b$. Replacing (u, v) by $(v/u^2, v^2/u^3)$, corresponding to the involution of $U_{3,1,1}$ with quotient $U_{3,2}$, gives an isomorphic algebra. We report on the fields involved in these algebras, since Galois groups are small enough so that future comparison with other sources of fields with these groups seems promising.

For simplicity, we exclude the twenty-three (u, v) where $u = 0$, so that the involution above is everywhere defined. We switch coordinates to the coordinates used in [17, §7.1] via $(p, q) = (3u, 3v/u^2)$ and $(u, v) = (p/3, p^2q/27)$. In the new coordinates, the involution is simply $(p, q) \mapsto (q, p)$, and we normalize by requiring $p \leq q$. We then have 232 algebras $K_{p,q}$ with $p < q$ and 20 algebras $K_{p,p}$. Besides these algebras, we have their twins $K_{p,q}^t$, and their common octic and quartic resolvents $\tilde{R}_{p,q}$ and $R_{p,q}$.

Despite the non-generic behavior of the family in general, Principle A has no exceptions in the current context: the 252 algebras $K_{p,q}$ and their 252 twins $K_{p,q}^t$ form 504 non-isomorphic algebras. Principle C also has no exceptions, as all algebras are wildly ramified at both 2 and 3.

There are many exceptions to Principle B. For example $K_{153/1849,129/289}$ factors as $6 + 6 + 12$ with the factors having Galois group $6T9$, $6T15 = A_6$, and $12T299 = S_6 \wr S_2$. Its twin factors as $3 + 3 + 6 + 12$ with factors having Galois groups S_3 , S_3 , A_6 , and $S_6 \wr S_2$. The two A_6 factors are given by the polynomials

$$(7.3) \quad f_6(x) = x^6 - 3x^5 + 3x^4 - 6x^2 + 6x - 2,$$

$$(7.4) \quad f_6^t(x) = x^6 - 3x^4 - 12x^3 - 9x^2 + 1.$$

These polynomials will be discussed further at the end of the next subsection.

For the rest of this section, we avoid Galois-theoretic complications like those of the last paragraph by requiring that $R_{p,q}$ either has an irreducible cubic factor or is irreducible itself. There are 39 (p, q) of the first type, and 178 (p, q) of the second. Failures of Principle B in this restricted setting are very mild, as A_6^4 is a subgroup of the Galois group of all these specializations. In the case of a cubic-times-linear quartic resolvent, we change notation by focusing on the larger degree part, so that $K_{p,q}$, $K_{p,q}^t$, $\tilde{R}_{p,q}$, and $R_{p,q}$ now have degrees 18, 18, 6, and 3.

7.6. Some number fields with small root discriminant. Table 7.1 summarizes the fields under consideration, with resolvent Galois groups indicated by Q and \tilde{Q} . In all cases, if $K_{p,q}$ has some Galois group mTj then its twin $K_{p,q}^t$ has the same Galois group mTj . For each Galois group, the table gives a corresponding field in our collection with smallest root discriminant. Thus (p, q) is chosen because one of $\delta = \text{rd}(K_{p,q})$ and $\delta^t = \text{rd}(K_{p,q}^t)$ is small; the other is sometimes substantially larger. Galois groups were computed by *Magma*, making use thereby of the algorithms of [8] and works classifying permutation groups.

For almost all groups in degree ≤ 19 , the database of Klueners and Malle presents at least one corresponding field. The database also highlights the field presented with smallest absolute discriminant. For the five degree eighteen groups appearing in Table 7.1, our fields are well under the previous minima, these being 643.84, 51.78, 66.63, 71.35, and 57.52 in the order listed. For the twelve degree twenty-four groups, we similarly do not know of other fields with smaller root discriminants.

The small root discriminants of these fields is often reflected in the smallness of coefficients in the standardized polynomials returned by *Pari's* `polredabs`. For example, the degree eighteen field in the table of smallest root discriminant is $K_{-3/125,1}^t$. It is defined by

$$\begin{aligned} f_{18}(x) = & \\ & x^{18} + 9x^{16} - 18x^{15} + 18x^{14} - 36x^{13} + 72x^{12} - 18x^{11} + 36x^{10} \\ & - 180x^9 + 18x^8 + 54x^7 + 48x^6 - 108x^5 + 18x^4 - 30x^3 + 9x^2 - 1. \end{aligned}$$

Q	$ \tilde{Q} $	\tilde{Q}	p	q	$\text{Gal}(K_{p,q})$	$\#$	D	δ	D^t	δ^t
S_3	6	$6T2$	1/12	24	$18T971$	1	$2^{22}3^{46}$	38.66	$2^{24}3^{44}$	36.95
S_3	12	$6T3$	-3/125	1	$18T972$	7	$2^{26}3^{41}$	33.24	$2^{26}3^{39}$	29.42
A_3	6	$6T6$	9/121	11/27	$18T974$	3	$-2^{18}3^{46}$	33.14	$-2^{24}3^{44}$	36.95
S_3	24	$6T7$	-3/49	7/3	$18T976$	1	$2^{30}3^{46}$	52.60	$2^{32}3^{44}$	50.29
S_3	48	$6T11$	1/9	9	$18T977$	27	$-2^{33}3^{41}$	43.52	$-2^{27}3^{39}$	30.57
C_4	16	$8T7$	18/25	5/6	$24T24946$	2	$2^{87}3^{24}$	37.01	$2^{97}3^{24}$	49.41
V	16	$8T9$	-72	3/16	$24T24948$	3	$2^{54}3^{40}$	29.68	$2^{70}3^{38}$	43.00
V	16	$8T11$	-27	-1/3	$24T24949$	3	$2^{48}3^{38}$	22.78	$2^{52}3^{42}$	30.70
D_4	16	$8T6$	-9	1/3	$24T24952$	8	$2^{81}3^{24}$	31.12	$2^{85}3^{24}$	34.94
D_4	16	$8T8$	1	9/8	$24T24953$	5	$-2^{52}3^{43}$	32.14	$-2^{60}3^{45}$	44.38
A_4	24	$8T13$	-2	1/4	$24T24956$	5	$2^{54}3^{46}$	39.07	$2^{56}3^{46}$	41.39
D_4	32	$8T17$	-8/3	9/16	$24T24961$	3	$2^{83}3^{24}$	32.97	$2^{81}3^{24}$	31.12
D_4	32	$8T15$	-27/25	5/9	$24T24962$	17	$2^{52}3^{43}$	32.14	$2^{48}3^{45}$	31.38
C_4	32	$8T16$	16/27	27/32	$24T24964$	1	$2^{79}3^{24}$	29.38	$2^{97}3^{24}$	49.41
S_4	48	$8T23$	-1/3	1	$24T24968$	44	$-2^{46}3^{59}$	56.22	$-2^{46}3^{51}$	38.98
D_4	64	$8T26$	-135/289	17/25	$24T24974$	26	$2^{58}3^{35}$	26.50	$2^{50}3^{39}$	25.26
S_4	192	$8T40$	-7/12	32/49	$24T24982$	61	$2^{55}3^{59}$	72.91	$2^{49}3^{51}$	42.51

TABLE 7.1. Fields $K_{p,q}$ and $K_{p,q}^t$ with given Galois group and small root discriminant

Similarly, the degree twenty-four field in the table of smallest root discriminant is $K_{-27,-1/3}$. It is defined by

$$\begin{aligned}
 f_{24}(x) = & x^{24} - 8x^{21} + 64x^{18} - 36x^{17} \\
 & - 9x^{16} - 56x^{15} + 276x^{14} - 72x^{13} + 237x^{12} - 24x^{11} + 486x^{10} - 88x^9 \\
 & + 513x^8 + 36x^7 + 256x^6 + 48x^5 + 18x^4 + 20x^3 - 6x^2 + 1.
 \end{aligned}$$

Another particularly interesting case comes from the second to last line of Table 7.1, where both δ and δ^t are small.

For speculating where the fields of this section may fit into complete lists, it is insightful to compare with the polynomials from (7.3) and (7.4). The fields $\mathbb{Q}[x]/f_6(x)$ and $\mathbb{Q}[x]/f_6^t(x)$ have root discriminants $\delta = (2^8 3^8)^{1/6} \approx 10.90$ and $\delta^t = (2^{10} 3^8)^{1/6} \approx 13.74$. These root discriminants are 12th and 44th on the complete sextic A_6 list, substantially behind the first entry $(2^6 67^2)^{1/6} \approx 8.12$ [9]. On the other hand the common splitting field of $f_6(x)$ and $f_6^t(x)$ has root discriminant $2^{13/6} 3^{16/9} \approx 31.66$. This is the smallest root discriminant of a Galois A_6 field, substantially ahead of the second smallest $2^{7/6} 3^{25/18} 13^{1/2} \approx 37.23$ [9]. We expect that the degree 18 and 24 fields discussed in this subsection behave similarly to these sextic fields: their root discriminants should appear early on complete lists, and their Galois root discriminants should appear even earlier.

8. A DEGREE 96 FAMILY: A LARGE DEGREE DESSIN AND NEWTON POLYGONS

Almost all the full number fields presented so far in this paper have been ramified exactly at the set $\{2, 3, 5\}$. Conjecture 1.1 on the other hand envisions inexhaustible

supplies of full number fields with other ramification sets \mathcal{P} . This section presents examples with $\mathcal{P} = \{2, 3, 7\}$.

8.1. One of two similar Hurwitz parameters. Theorems 4.1 and 4.2 of Malle's paper [12] each give a two-parameter family of septic covers of the projective line with monodromy group $SL_3(\mathbb{F}_2) \subset S_7$ of order $168 = 2^3 \cdot 3 \cdot 7$. We focus on the family of Theorem 4.2 which is indexed by the Hurwitz parameter

$$h = (SL_3(\mathbb{F}_2), (22111, 421), (4, 1)).$$

The corresponding degree is $m = 192$.

The situation has much in common with König's situation from §7.1 and we can proceed similarly. Thus, by equating a discriminantal factor with a standard quartic, we realize X_h as a degree 192 cover of $U_{4,1}$. The outer involution of $SL_3(\mathbb{F}_2)$ coming from projective duality gives an explicit involution analogous to (7.2). Quotienting by this involution yields the degree 96 cover $X_h^* \rightarrow U_{4,1}$. Unlike the cover of the previous section, this cover is full.

The family from Theorem 4.1 of [12] is very similar: the partition 421 is replaced by 331, and the degree $192/2 = 96$ is replaced by $216/2 = 108$. We are working with the degree 96 family because the curve given by $f_{96}(j, x) = 0$ below has genus zero, while its analog for the degree 108 family has genus one.

8.2. A dessin. The reduced configuration space $U_{4,1}$ is the same as that for our introductory family and has been described in §3.4. However the specialization set is now $U_{4,1}(\mathbb{Z}[1/42])$ rather than the $U_{4,1}(\mathbb{Z}[1/30])$ drawn in Figure 4.1. We present here only a polynomial for the degree 96 cover of the vertical line $(u, v) = (1/3, (j-1)/27j)$ evident in Figure 4.1:

$$\begin{aligned} f_{96}(j, x) = & (7411887x^{32} - 316240512x^{31} + 5718682592x^{30} - 57608479936x^{29} \\ & + 345466405984x^{28} - 1143902168192x^{27} + 500924971008x^{26} + 20121596404224x^{25} \\ & - 178485128485440x^{24} + 1076315934382080x^{23} - 4902849972088320x^{22} \\ & + 16964516971136000x^{21} - 45252388465854976x^{20} + 95197078307043328x^{19} \\ & - 161987009378324480x^{18} + 229049096903122944x^{17} - 277106243726667264x^{16} \\ & + 295558502345637888x^{15} - 284898502452436992x^{14} + 250987121290100736x^{13} \\ & - 200876992270295040x^{12} + 143474999551229952x^{11} - 89556680876359680x^{10} \\ & + 47950288840949760x^9 - 21681369027919872x^8 + 8162827596988416x^7 \\ & - 2520589064601600x^6 + 626540088655872x^5 - 122178152300544x^4 \\ & + 17986994307072x^3 - 1878160048128x^2 + 123834728448x - 3869835264)^3 \\ & - 2^{20} j x^6 (3x - 2)^2 (x^2 + 2x - 2)^6 (7x^2 - 14x + 6)^{21} (2x^3 - 15x^2 + 18x - 6)^9. \end{aligned}$$

The printed degree thirty-two polynomial capturing behavior at $j = 0$ has Galois group A_{32} and field discriminant only $2^{64} 3^{36} 7^{18}$.

Figure 8.1 draws the dessin of $f_{96}(j, x)$, not in the copy of \mathbb{C} with coordinate x , but rather the copy of \mathbb{C} with coordinate $x' = 1/(1-x)$, for better geometric appearance. By definition, the figure consists of all x' corresponding to x satisfying $f_{96}(j, x) = 0$ with $j \in [0, 1]$. This figure has the natural structure of a graph with 96 edges, the preimages of $(0, 1)$. All vertices have degree ≤ 3 : there are thirty-two triple points, the preimages of 0, and forty double points and sixteen endpoints,




FIGURE 8.1. The dessin corresponding to $f_{96}(j, x)$. Besides the nine regions with indicated sizes there is a tenth region of size 1 immediately to the left of the centrally printed 21. This small region is adjacent to two triple points and near an endpoint. Also to the immediate right of each of the two left-printed nines, there is one triple point and two endpoints.

the preimages of 1. The forty double points are not readily visible in the figure, as they lie in the middle of forty double edges, but most of the triple points and endpoints are. There are also ten regions, of varying size, defined as half the number of bounding edges. The few aspects of all this structure which are not visible are described in the caption of Figure 8.1. The topological structure could also be deduced from a braid computation, rather than from the defining equation.

The polynomial $f_{96}(j, x)$ and Figure 8.1 illustrate the nature and complexity of the objects we are considering. Note that the existence of this cover shows that the Hurwitz number algebra indexed by $(A_{96}, (3^{32}, 2^{40} 1^{16}, 21^2 9^3 7 6^3 2), (1, 1, 1))$ has at least one factor of \mathbb{Q} . The entire Hurwitz algebra is way out of computational range, because the two main terms in the mass formula (3.6) give 3×10^{15} as an approximation for its degree.

A common feature of $f_{25}(j, x)$ from §4.2 and $f_{96}(j, x)$ is not accidental. In the braid group description of their monodromy, calculable purely group-theoretically, local monodromy operators about 0 and 1 are the images of braid group elements of order 3 and 2 respectively. Thus the preimage of $u = 1/3$ in $X_h \rightarrow U_{4,1}$ for any h with multiplicity vector $(4, 1)$ likewise has this property.

8.3. Specialization and Newton polygons. For greater explicitness, we report only on specializing $f_{96}(j, x)$ to $j \in U_{3,1}(\mathbb{Z}[1/42])$. From complete tables of elliptic curves [6], this specialization set has size 413. Supporting Principle A, all 413 algebras are non-isomorphic. Supporting Principle B, these algebras all have Galois group A_{96} . Investigating Principle C is more subtle. In lieu of completely factoring $f_{96}(j, x)$ over \mathbb{Q}_p and taking field discriminants of the factors, we use Newton polygons. To illustrate this computationally much simpler method, we take $j = 1/3$ as

a representative example, and work with

$$g(x) = 3f(1/3, x) = 3^7 7^{21} x^{96} - 2^7 3^7 7^{21} x^{95} + \dots + 2^{53} 3^{32} x - 2^{48} 3^{31}.$$

Factoring modulo 2, 3, and 7 gives

$$g(x) \stackrel{2}{\equiv} x^{96}, \quad g(x) \stackrel{3}{\equiv} x^{54}(x-2)^{33}, \quad g(x) \stackrel{7}{\equiv} h_2(x)h_{20}(x)h_{25}(x),$$

with $h_k(x)$ irreducible of degree k . The 2-adic Newton polygon of $g(x)$ has all slopes $1/2$, showing that all 96 roots $\alpha \in \overline{\mathbb{Q}}_2$ have $\text{ord}_2(\alpha) = 1/2$. Since the denominator is divisible by 2, one has that the 2-adic wild degree as in §4.5 is $m_{2\text{-wild}} = 96$. From a more complicated calculation with 3-adic Newton polygons, we get that the 96 roots $\alpha \in \overline{\mathbb{Q}}_3$ are distributed as follows:

9 roots with $\text{ord}_3(\alpha) = 4/9$,	22 roots with $\text{ord}_3(\alpha - 1) = 13/21$,
27 roots with $\text{ord}_3(\alpha) = 1/3$,	3 roots with $\text{ord}_3(3\alpha - (1 - i)) = 2/3$,
3 roots with $\text{ord}_3(\alpha) = -1/3$,	3 roots with $\text{ord}_3(3\alpha - (1 + i)) = 2/3$,
9 roots with $\text{ord}_3(\alpha/3 - i) = 5/9$,	
9 roots with $\text{ord}_3(\alpha/3 + i) = 5/9$,	12 roots with $\text{ord}_3(\alpha - 1) = 1/2$.

Only the last twelve α could possibly not contribute to the 3-adic wild degree, giving already $m_{3\text{-wild}} \geq 84$. But in fact these α satisfy $\text{ord}_3(\alpha^2/3 - 2) = 5/12$ so one has $m_{3\text{-wild}} = 96$. Finally, the 7-adic Newton polygon of $g(x)$ has slopes 0 and $-3/7$ with multiplicities 47 and 49 respectively. The slope of 0 corresponds to the isolated roots modulo 7 and the slope of $-3/7$ then gives $m_{7\text{-wild}} = 49$.

The Newton polygon process can be easily automated. It says that all 413 algebras are wildly ramified at both 2 and 3. It says also that all algebras are wildly ramified at 7 except for those coming from the specialization points $-3^{15} 5^3 7^3 / 2^8$, $-7^3 / 2^1 3^2$, $7^3 / 2^9$, $7^3 / 3^5$, $7^3 / 2^1 3^3$, $5^3 7^3 / 3^5$, $2^2 7^3 / 3$ and $7^4 / 2^6 3$, $-7^4 / 2^7 3^4$, 7^4 , $-7^5 / 2^1 3^8$. The first seven all have tame ramification at 7 corresponding to the partition $19^3 1^{39}$ while the last four have tame ramification at 7 corresponding to the partition $57^1 3^{13}$. This behavior comes from the fact that these specialization points are all 7-adically close to $j = 0$ and the degree 32 polynomial above has tame ramification at 7 given by the partition $19^1 1^{13}$.

9. A DEGREE 202 FAMILY: DEGENERATIONS AND GENERIC SPECIALIZATION

Continuing to increase degrees as we go through the last six sections, we now describe a family having degree 202. Our description emphasizes its degenerations, a relevant topic because how a family degenerates has substantial influence on how ramification behaves in the Hurwitz number fields within the family. We conclude by observing that specialization is generic, both in one of the degenerations of the family and in the family itself.

9.1. Some plane curves. To streamline the subsequent subsections, we first present some polynomials defining affine curves in the x - y plane. The next two subsections will place a natural function on each curve, and we index the polynomials by the degree of this function.

Eleven relatively simple polynomials are

$A_{10} = x,$	$B_4 = x - 1,$
$A_{13} = y,$	$B_8 = y - 1,$
$A_{14} = x - y,$	$B_{32} = x^2 y - 4x^2 - 8xy + 20x + 10y - 20,$

$$\begin{aligned}
A_{16} &= 3xy - 6x - 6y + 10, & C_{22} &= 3xy^2 - 12xy + 8x - 15y^2 + 40y - 24, \\
A_{20} &= x^2y - 3x^2 - 6xy & C_{25} &= 3x^2y - 6x^2 - 12xy + 20x + 10y - 15, \\
&+ 12x + 6y - 10, & D_{10} &= 3x^2 - 12x + 10.
\end{aligned}$$

For all but one of these polynomials P , the curve \mathbf{P} given by its vanishing is obviously rational, as at least one of the variables appears to degree one in P . In contrast, the curve \mathbf{D}_{10} consists of two genus zero components, neither one of which is defined over \mathbb{Q} .

B_{52}	1	x	x^2	x^3	x^4	D_{32}	1	x	x^2	x^3	x^4
1	1080	-2160	1296	-176	-24	1			160	-192	48
y	-1080	2052	-1164	156	12	y		-320	192	120	-48
y^2	135	-276	180	-36		y^2	250	12	-213	12	12
y^3	50	-84	36			y^3	-300	288	-18	-12	
y^4	15	-12				y^4	90	-108	27		

D_{48}	1	x	x^2	x^3	x^4	x^5	x^6
1				1600	-2880	1632	-288
y			2400	-8160	9048	-3960	576
y^2		1200	-6480	11448	-8712	2880	-324
y^3	-2500	6300	-4620	-108	1395	-513	54
y^4	1500	-3900	3780	-1692	351	-27	

TABLE 9.1. Three polynomials $\sum c_{ij}x^i y^j$, presented by listing their coefficients c_{ij} .

Three more complicated polynomials are given in matrix form in Table 9.1. The corresponding curves \mathbf{B}_{52} , \mathbf{D}_{32} , and \mathbf{D}_{48} have genus 1, 2, and 5 respectively. Each genus is much smaller than the upper bound allowed by the support in $\mathbb{Z}_{\geq 0}^2$ of the coefficients; this bound, being the number of ‘‘interior’’ coefficients, is 6, 6, and 12 respectively. In each case, there are several singularities causing this genus reduction, one of which is at $(1, 1)$.

9.2. Calculation of a rational presentation. This subsection is very similar to §6.1, illustrating that in favorable cases computation of Hurwitz covers following the outline of §3.7 is quite mechanical. As normalized Hurwitz parameter we take

$$h = (S_6, (21111, 3_021, 3_1111, 4_\infty 11), (2, 1, 1, 1)).$$

Any function governed by h is of the form

$$g(s) = \frac{s^3(s-x)^2(s-y)}{a(s^2 + s(d-e-1) + e)}.$$

The ramification requirement on g at 1 is that $(g(1), g'(1), g''(1)) = (1, 0, 0)$. These three equations let us express a , d , and e in terms of x and y . Namely

$$a = -C_{25}, \quad d = \frac{B_4^2 B_8}{C_{25}}, \quad e = \frac{A_{20}}{C_{25}}.$$

Using a resolvent as usual, we find that the critical values of $g(s)$ besides 0, 1, and ∞ are the roots of $Wt^2 + (V - U - W)t + Ut^2$ with

$$(9.1) \quad U = -2^2 3^3 A_{10}^5 A_{13}^4 A_{14}^3 A_{16} A_{20}^2,$$

$$(9.2) \quad V = -3^3 B_4 B_8^2 B_{32}^4 B_{52},$$

$$(9.3) \quad W = 2^8 C_{25}^5 C_{22}.$$

Comparing with the standard quadratic $t^2 + (v - u - 1)t + u$, one gets the rational presentation

$$(9.4) \quad u = \frac{U}{W}, \quad v = \frac{V}{W}.$$

So, appealing to (9.1)-(9.3) and the explicit polynomials in §9.1, Equations (9.4) express u and v as explicit functions of x and y .

9.3. A view of X_h . Recall from §6.2 and Figure 6.1 that the complement of $U_{2,1,1,1}$ in the projective u - v -plane consists of three lines A, B, C and a conic D. In the map from the affine x - y plane to the projective u - v plane, we can consider the preimages of these discriminantal curves.

Figure 9.1 draws the real points of these four preimages. Using as before a similar notation for an equation and its curve, inspection of our equations gives

$$\begin{aligned} \pi_h^{-1}(\text{A}) &= \text{A}_{10} \cup \text{A}_{14} \cup \text{A}_{13} \cup \text{A}_{16} \cup \text{A}_{20}, \\ \pi_h^{-1}(\text{B}) &= \text{B}_4 \cup \text{B}_8 \cup \text{B}_{32} \cup \text{B}_{52}, \\ \pi_h^{-1}(\text{C}) &= \text{C}_{25} \cup \text{C}_{22}, \\ \pi_h^{-1}(\text{D}) &= \text{D}_{10} \cup \text{D}_{32} \cup \text{D}_{48}. \end{aligned}$$

The figure is intended to indicate the rich geometry present in any Hurwitz surface. Other interesting curves present whenever $\nu = (2, 1, 1, 1)$ are the preimages of the lines ad, bd, cd, bc, ac, and ab introduced in §6. For the current h , all of them have a complicated real locus. Their genera are respectively 25, 18, 23, 35, 31, and 23. The curves ad, bd, and cd intersect at the preimage of the point d, and Figure 9.1 also draws the ten real points of this preimage.

9.4. Degree 202 polynomials and their degenerate factorizations. Removing y and x respectively from (9.4) by resultants gives degree 202 polynomials $f(u, v, x)$ and $\phi(u, v, y)$. Completely expanded, they have 10484 and 15555 terms respectively.

The structures studied in the previous two subsections appear when one factors specializations corresponding to the four discriminantal components:

$$\begin{aligned} f(0, v, x) &= x^{50}(x^2 - 4x + 6)a_{13}(v, x)^4 a_{14}(v, x)^3 a_{16}(v, x)a_{20}(v, x)^2, \\ f(u, 0, x) &= -(x - 1)^6 b_8(u, x)^2 b_{32}(u, x)^4 b_{52}(u, x), \\ \lim_{u \rightarrow \infty} \frac{f(u, wu, x)}{u^{10}} &= -2^{10}(x - 1)^7 c_{10}(w, x)^3 c_{22}(w, x)c_{25}(w, x)^5, \\ f(r^2, (1 - r)^2, x) &= -(3x^2 - 12x + 10)^5 d_{32}(r, x)^3 d_{48}(r, x)^2, \end{aligned}$$

and

$$\begin{aligned} \phi(0, v, y) &= y^{52}(3y^2 - 8y + 8)\alpha_{10}(v, y)^5 \alpha_{14}(v, y)^3 \alpha_{16}(v, y)\alpha_{20}(v, y)^2, \\ \phi(u, 0, y) &= (y - 1)^{16}(y - 4)^2 \beta_4(u, y)\beta_{32}(u, y)^4 \beta_{52}(u, y), \end{aligned}$$




FIGURE 9.1. $X_h(\mathbb{R})$ is the complement of the drawn curves in the real x - y plane. The drawn points are the ten real preimages of $(u, v) = (1, 1)$.

$$\begin{aligned} \lim_{u \rightarrow \infty} \frac{\phi(u, wu, y)}{u^{13}} &= -2^{36}(y-2)^3(y-1)^7\gamma_6(w, y)^2\gamma_{22}(w, y)\gamma_{25}(w, y)^5, \\ \phi(r^2, (1-r)^2, y) &= \delta_{10}(r, y)\delta_{32}(r, y)^3\delta_{48}(r, y)^2. \end{aligned}$$

Our notation coordinates the different viewpoints: for example, the equations $D_{48} = 0$, $d_{48}(r, x) = 0$, and $\delta_{48}(r, y) = 0$ all describe the genus five curve D_{48} .

As a sample degeneration, chosen because it makes an interesting comparison the degree 25 polynomials from our introductory example,

$$\begin{aligned} c_{25}(w, x) = & \\ & -(2x-5)(3x-5)^2(6x^4-40x^3+105x^2-120x+50)^4 \\ & (12x^6-60x^5-40x^4+760x^3-1800x^2+1750x-625) \\ & +4wx^5(x^2-5x+5)^3(3x^2-10x+10)^2(6x^2-20x+15)^4(6x^2-15x+10). \end{aligned}$$

Here the x -line is identified with X_h for $h = (S_6, (321, 3111, 51, 21111), (1, 1, 1, 1))$, and the w -line with $U_{1,1,1,1}$. All the other degenerations have a similar four-point description. The discriminant of $c_{25}(w, x)$ is $2^{212}3^{66}5^{285}w^{13}(w-1)^{19}$. All the other degenerations are likewise three-point covers, all full except for A_{14} , A_{16} , A_{20} , C_6 , and D_{10} . In every case, the target variable, be it v , u , w , or r , is chosen such that the singular values are 0, 1, and ∞ .

To be noted is that we are not expending any extra effort here to introduce a conceptually defined completion of X_h . Indeed the curves that consist of horizontal lines, namely A_{13} and B_8 , are seen clearly by f but only as vestigial factors by ϕ . In reverse, the curves that consist of vertical lines, namely A_{10} , B_4 , and D_{10} , are seen completely by ϕ but only partially by f . Finally, to see preimages corresponding to the factors $c_{10}(w, x)$ and $\gamma_6(w, x)$, one would have to go beyond the x - y plane as a partial completion of X_h .

A braid group computation gives the partition of 202 which captures how local sheets of X_h are interchanged as one goes around one of the four discriminantal components in the completion of $U_{2,1,1,1}$. These partitions are

$$\beta_A = 5^{10}4^{13}3^{14}2^{20}1^{16+2}, \quad \beta_B = 4^{32}2^{8+1}1^{52+4}, \quad \beta_C = 5^{25}3^{10+2}2^{6+2}1^{22+3},$$

and $\beta_D = 3^{32}2^{48}1^{10}$. The boldface exponents correspond to components not seen by our simple calculations. Thus we are missing only 2, 2, 13, and 0 of the 202 sheets near the preimages of A, B, C, and D respectively.

9.5. Specialization. The degenerations can be specialized, and the computations support Principles A, B, and C. For example, consider $c_{25}(w, x)$ specialized to w in the known set $U_{1,1,1,1}(\mathbb{Z}[1/30])$. The 99 algebras are all distinct, they are all full, and they are all wildly ramified at each of 2, 3, and 5.

Specialization of the full family at the 2947 points of $U_{2,1,1,1}(\mathbb{Z}[1/30])$ can also be satisfactorily studied, despite the large degree. The 2947 algebras are all distinct and they all have Galois group A_{202} or S_{202} . From Newton polygons, we know they are all wildly ramified at 2, 3, and 5. Thus, in this family, Principles A, B, and C hold without exception.

10. A DEGREE 1200 FIELD: COMPUTATIONS IN LARGE DEGREE

Conjecture 1.1 says that for certain finite sets of primes \mathcal{P} , there exist full number fields of arbitrarily large degree with ramification set in \mathcal{P} . A natural computational challenge for a given \mathcal{P} is then to produce an explicit full Hurwitz number field $K_{h,u}$ with degree m as large as possible. In this short final section, we take $\mathcal{P} = \{2, 3, 5\}$ and produce such a field for degree $m = 1200$.

Taking $h = (S_6, (21111, 3_02_11, 4_\infty11), (4, 1, 1))$ and normalizing as indicated, the functions to consider are

$$g(s) = \frac{as^3(s-1)^2(s-x)}{s^2+bs+c}.$$

As specialization point, we take $u = ((t^4 - 4t - 6), \{0\}, \{\infty\})$. This specialization point indeed keeps ramification within $\{2, 3, 5\}$ as the discriminant of $t^4 - 4t - 6$ is -2^83^5 .

The condition that the critical values besides 0 and ∞ are the roots of $t^4 - 4t - 6$ gives four equations in the four unknowns x, a, b, c . Of the unknowns, we focus on

x because its special values 0, 1, and ∞ are all meaningful, corresponding to degenerations. Eliminating a and then c is easy. Eliminating b then has a ten-minute run-time on *Magma* to get a degree 3700 polynomial. Factorizing this polynomial to find the relevant factor has a one-minute run-time. The resulting monic polynomial $f_{1200}(x) \in \mathbb{Z}[1/30][x]$ defining $K_{h,u}$ satisfies $f_{1200}(0) = 2^{880}/5^{500}$ and $f_{1200}(1) = 3^{684}/2^{256}5^{500}$. After removing all factors of 2, 3, and 5, the coefficients are integers averaging about 440 digits.




FIGURE 10.1. Roots in the closed upper half plane of a polynomial defining a degree 1200 Hurwitz number field

The polynomial is to some extent analyzable despite its large degree and large coefficients. From the factorization partitions (989, 208, 3) at 19 and (1181, 9, 6, 4) at 47, it has Galois group S_{1200} , in conformity with Principle B. From Newton polygons, it is wildly ramified at 2, 3, and 5, as predicted by Principle C. Figure 10.1 presents the roots of $f_{1200}(x)$ in the closed upper half plane, all of which lie in the drawn window $[-2.6, 2.6] \times [0, 2.4]$. There are 34 real roots and a general tendency of roots to cluster near the interval $[0, 1]$ connecting the special points 0 and 1.

Large degree Hurwitz number fields provide specific challenges to improve computational algorithms for general number fields. For example, from Newton polygons we have substantial information on how the field $K_{h,u}$ of this section factors over \mathbb{Q}_2 , \mathbb{Q}_3 , and \mathbb{Q}_5 . However we cannot go far enough to determine the exponents in its discriminant $-2^a 3^b 5^c$.

REFERENCES

1. José Bertin and Matthieu Romagny, *Champs de Hurwitz*, Mém. Soc. Math. Fr. (N.S.) (2011), no. 125-126, 219. MR 2920693
2. Manjul Bhargava, *The density of discriminants of quartic rings and fields*, Ann. of Math. (2) **162** (2005), no. 2, 1031–1063. MR 2183288
3. ———, *Mass formulae for extensions of local fields, and conjectures on the density of number field discriminants*, Int. Math. Res. Not. IMRN (2007), no. 17, Art. ID rnm052, 20. MR 2354798
4. ———, *The density of discriminants of quintic rings and fields.*, Ann. of Math. (2) **172** (2010), no. 3, 1559–1591.

5. Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265, Computational algebra and number theory (London, 1993). MR MR1484478
6. J. E. Cremona and M. P. Lingham, *Finding all elliptic curves with good reduction outside a given set of primes*, Experiment. Math. **16** (2007), no. 3, 303–312. MR 2367320
7. H. Davenport and H. Heilbronn, *On the density of discriminants of cubic fields. II*, Proc. Roy. Soc. London Ser. A **322** (1971), no. 1551, 405–420. MR 0491593
8. Claus Fieker and Jürgen Klüners, *Computation of Galois groups of rational polynomials*, LMS J. Comput. Math. **17** (2014), no. 1, 141–158. MR 3230862
9. John W. Jones and David P. Roberts, *A database of number fields*, LMS J. Comput. Math. **17** (2014), no. 1, 595–618, Database at <http://hobbes.la.asu.edu/NFDB/>. MR 3356048
10. Nicholas M. Katz, *Rigid Local Systems*, Annals of Mathematics Studies, vol. 139, Princeton University Press, Princeton, NJ, 1996. MR 1366651
11. Joachim Koenig, *Computation of Hurwitz spaces and new explicit polynomials for almost simple Galois groups*, Preprint, to appear in Math. Comp. 2015.
12. Gunter Malle, *Multi-parameter polynomials with given Galois group*, J. Symbolic Comput. **30** (2000), no. 6, 717–731. MR 1800034
13. Gunter Malle and B. Heinrich Matzat, *Inverse Galois theory*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1999. MR 1711577
14. Gunter Malle and David P. Roberts, *Number fields with discriminant $\pm 2^a 3^b$ and Galois group A_n or S_n* , LMS J. Comput. Math. **8** (2005), 80–101 (electronic). MR 2135031
15. David P. Roberts, *Hurwitz-Belyi maps*, Arxiv, August 30, 2016. Submitted.
16. ———, *Wild partitions and number theory*, J. Integer Seq. **10** (2007), no. 6, Article 07.6.6, 34. MR 2335791
17. ———, *Division polynomials with Galois group $SU_3(3).2 \cong G_2(2)$* , Advances in the theory of numbers, Fields Inst. Commun., vol. 77, Fields Inst. Res. Math. Sci., Toronto, ON, 2015, pp. 169–206. MR 3409329
18. ———, *Polynomials with prescribed bad primes*, Int. J. Number Theory **11** (2015), no. 4, 1115–1148. MR 3340686
19. David P. Roberts and Akshay Venkatesh, *Hurwitz monodromy and full number fields*, Algebra Number Theory **9** (2015), no. 3, 511–545. MR 3340543
20. Jean-Pierre Serre, *Topics in Galois theory*, second ed., Research Notes in Mathematics, vol. 1, A K Peters, Ltd., Wellesley, MA, 2008, With notes by Henri Darmon. MR 2363329
21. The PARI Group, Bordeaux, *PARI/GP version 2.7.5*, 2015, available from <http://pari.math.u-bordeaux.fr/>.
22. Helmut Völklein, *A transformation principle for covers of \mathbb{P}^1* , J. Reine Angew. Math. **534** (2001), 156–168. MR 1831635
23. Wolfram Research, Inc., *Mathematica 10.0.2.0*.

DIVISION OF SCIENCE AND MATHEMATICS, UNIVERSITY OF MINNESOTA MORRIS; MORRIS, MINNESOTA, 56267, USA

E-mail address: roberts@morris.umn.edu

URL: <http://cda.morris.umn.edu/~roberts>