

doi: 10.1515/umcsmath-2015-0007

---

ANNALES  
UNIVERSITATIS MARIAE CURIE-SKŁODOWSKA  
LUBLIN – POLONIA

VOL. LXVIII, NO. 2, 2014

SECTIO A

65–83

JAGANNATH PATEL and ASHOK KUMAR SAHOO

## On certain subclasses of analytic functions associated with the Carlson–Shaffer operator

ABSTRACT. The object of the present paper is to solve Fekete–Szegő problem and determine the sharp upper bound to the second Hankel determinant for a certain class  $\mathcal{P}^\lambda(a, c, A, B)$  of analytic functions in the unit disk. We also investigate several majorization properties for functions belonging to a subclass  $\tilde{\mathcal{P}}^\lambda(a, c, A, B)$  of  $\mathcal{P}^\lambda(a, c, A, B)$  and related function classes. Relevant connections of the main results obtained here with those given by earlier workers on the subject are pointed out.

**1. Introduction and preliminaries.** Let  $\mathcal{A}$  be the class of functions  $f$  of the form

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ . Also, let  $\mathcal{S}$  denote the subclass of  $\mathcal{A}$  consisting of functions of the form

$$(1.2) \quad g(z) = z - \sum_{n=2}^{\infty} b_n z^n \quad (b_n \geq 0).$$

A function  $f \in \mathcal{A}$  is said to be starlike function of order  $\alpha$  and convex function of order  $\alpha$ , respectively, if and only if  $\operatorname{Re}\{zf'(z)/f(z)\} > \alpha$  and  $\operatorname{Re}\{1 + (zf''(z)/f'(z))\} > \alpha$  for  $0 \leq \alpha < 1$  and for all  $z \in \mathbb{U}$ . By usual

2010 *Mathematics Subject Classification.* 30C45.

*Key words and phrases.* Analytic function, subordination, Hadamard product, majorization, Fekete–Szegő problem, Hankel determinant.

notations, we denote these classes of functions by  $\mathcal{S}^*(\alpha)$  and  $\mathcal{C}(\alpha)$  ( $0 \leq \alpha < 1$ ), respectively. We write  $\mathcal{S}^*(0) = \mathcal{S}^*$  and  $\mathcal{C}(0) = \mathcal{C}$ , the familiar subclasses of starlike functions and convex functions in  $\mathbb{U}$ .

Furthermore, a function  $f \in \mathcal{A}$  is said to be in the class  $\mathcal{R}(\alpha)$ , if it satisfies the inequality:

$$\operatorname{Re}\{f'(z)\} > \alpha \quad (0 \leq \alpha < 1; z \in \mathbb{U}).$$

Note that  $\mathcal{R}(\alpha)$  is a subclass of close-to-convex functions of order  $\alpha$  ( $0 \leq \alpha < 1$ ) in  $\mathbb{U}$ . We write  $\mathcal{R}(0) = \mathcal{R}$ , the familiar class functions whose derivatives have a positive real part in  $\mathbb{U}$ .

Let  $\mathcal{P}$  denote the class of analytic functions of the form

$$(1.3) \quad \phi(z) = 1 + p_1z + p_2z^2 + \cdots \quad (z \in \mathbb{U})$$

such that  $\operatorname{Re}\{\phi(z)\} > 0$  in  $\mathbb{U}$ .

For functions  $f$  and  $g$ , analytic in the unit disk  $\mathbb{U}$ , we say the  $f$  is said to be subordinate to  $g$ , written as  $f \prec g$  or  $f(z) \prec g(z)$  ( $z \in \mathbb{U}$ ), if there exists an analytic function  $\omega$  in  $\mathbb{U}$  with  $\omega(0) = 0$ ,  $|\omega(z)| \leq |z|$  ( $z \in \mathbb{U}$ ) and  $f(z) = g(\omega(z))$  for all  $z \in \mathbb{U}$ . In particular, if  $g$  is univalent in  $\mathbb{U}$ , then we have the following equivalence (see [20]):

$$f(z) \prec g(z) \quad (z \in \mathbb{U}) \iff f(0) = g(0) \text{ and } f(\mathbb{U}) \subset g(\mathbb{U}).$$

Following MacGregor [19], we say that  $f$  is majorized by  $g$  in  $\mathbb{U}$  and write

$$(1.4) \quad f(z) \ll g(z) \quad (z \in \mathbb{U}),$$

if there exists a function  $\psi$ , analytic in  $\mathbb{U}$  such that  $|\psi(z)| \leq 1$  and

$$(1.5) \quad f(z) = \psi(z)g(z) \quad (z \in \mathbb{U}).$$

For the functions  $f$  and  $g$  given by the power series

$$f(z) = \sum_{n=0}^{\infty} a_n z^n, \quad g(z) = \sum_{n=0}^{\infty} b_n z^n \quad (z \in \mathbb{U})$$

their Hadamard product (or convolution), denoted by  $f \star g$  is defined as

$$(f \star g)(z) = \sum_{n=0}^{\infty} a_n b_n z^n = (g \star f)(z) \quad (z \in \mathbb{U}).$$

We note that  $f \star g$  is analytic in  $\mathbb{U}$ .

For real or complex parameters  $a_1, a_2, \dots, a_p$  and  $b_1, b_2, \dots, b_q$  ( $b_j \notin \mathbb{Z}_0^- = \{\dots, -2, -1, 0\}$ ;  $j = 1, 2, \dots, q$ ), the generalized hypergeometric function  ${}_pF_q(a_1, a_2, \dots; b_1, b_2, \dots, b_q; z)$  is defined by the following infinite series (cf., e.g., [28]):

$$(1.6) \quad {}_pF_q(a_1, a_2, \dots; b_1, b_2, \dots, b_q; z) = \sum_{n=0}^{\infty} \frac{(a_1)_n (a_2)_n \cdots (a_p)_n}{(b_1)_n (b_2)_n \cdots (b_q)_n} \frac{z^n}{n!}$$

( $z \in \mathbb{U}$ ), where  $p \leq q + 1, p, q \in \mathbb{N}_0 = \mathbb{N} \cup \{0\} = \{0, 1, 2, 3, \dots\}$  and  $(x)_n$  is the Pochhammer symbol defined, in terms of the Gamma function  $\Gamma$ , by

$$(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)} = \begin{cases} 1, & (n = 0, x \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}), \\ x(x+1) \cdots (x+n-1), & (n \in \mathbb{N}, x \in \mathbb{C}). \end{cases}$$

It is easily seen that the radius of convergence  $\rho$  of the function  ${}_pF_q$  represented by the series (1.6) is

$$\rho = \begin{cases} \infty, & p < q + 1, \\ 1, & p = q + 1, \\ 0, & p > q + 1, \end{cases}$$

so that for  $p \leq q + 1$ , the function  ${}_pF_q$  is analytic in  $\mathbb{U}$ .

By making use of the Hadamard product, Carlson–Shaffer [3] defined the linear operator

$$\mathcal{L}(a, c) : \mathcal{A} \longrightarrow \mathcal{A}$$

in terms of the incomplete beta function  $\varphi$  by

$$(1.7) \quad \mathcal{L}(a, c)f(z) = \varphi(a, c; z) \star f(z) \quad (f \in \mathcal{A}; z \in \mathbb{U}),$$

where

$$\varphi(a, c; z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} z^{n+1} \quad (a \in \mathbb{C}, c \in \mathbb{C} \setminus \mathbb{Z}_0^-; z \in \mathbb{U}).$$

If  $f \in \mathcal{A}$  is given by (1.1), then it follows from (1.7) that

$$(1.8) \quad \mathcal{L}(a, c)f(z) = z + \sum_{n=1}^{\infty} \frac{(a)_n}{(c)_n} a_{n+1} z^{n+1} \quad (z \in \mathbb{U})$$

and

$$(1.9) \quad z(\mathcal{L}(a, c)f)'(z) = a\mathcal{L}(a+1, c)f(z) - (a-1)\mathcal{L}(a, c)f(z) \quad (z \in \mathbb{U}).$$

We note that for  $f \in \mathcal{A}$

- (i)  $\mathcal{L}(a, a)f(z) = f(z)$ ;
- (ii)  $\mathcal{L}(2, 1)f(z) = zf'(z)$ ;
- (iii)  $\mathcal{L}(3, 1)f(z) = zf'(z) + \frac{1}{2}z^2f''(z)$ ;
- (iv)  $\mathcal{L}(m+1, 1)f(z) = \mathcal{D}^m f(z) = \frac{z}{(1-z)^{m+1}} \star f(z)$  ( $m \in \mathbb{Z}; m > -1$ ), the Ruscheweyh derivative operator [26];

(v)  $\mathcal{L}(2, 2-\mu)f(z) = \Omega_z^\mu f(z)$  ( $0 \leq \mu < 1; z \in \mathbb{U}$ ), the well-known Owa–Srivastava fractional differential operator [25]. We also observe that  $\Omega_z^0 f(z) = f(z)$  and  $\Omega_z^1 f(z) = zf'(z)$ .

With the aid of the linear operator  $\mathcal{L}(a, c)$ , we introduce a subclass of  $\mathcal{A}$  as follows:

**Definition 1.1.** For the fixed parameters  $A, B$  ( $-1 \leq B < A \leq 1$ ),  $a > 0$  and  $c > 0$ , a function  $f \in \mathcal{A}$  is said to be in the class  $\mathcal{R}^\lambda(a, c, A, B)$ , if it satisfies the following subordination relation:

$$(1.10) \quad (1 - \lambda) \frac{\mathcal{L}(a, c)f(z)}{z} + \lambda \frac{\mathcal{L}(a + 1, c)f(z)}{z} \prec \frac{1 + Az}{1 + Bz} \quad (\lambda \geq 0; z \in \mathbb{U}).$$

Using the identity (1.9) in (1.10), it follows that

$$\left(1 - \frac{\lambda}{a}\right) \frac{\mathcal{L}(a, c)f(z)}{z} + \frac{\lambda}{a} (\mathcal{L}(a, c)f)'(z) \prec \frac{1 + Az}{1 + Bz} \quad (\lambda \geq 0; z \in \mathbb{U}).$$

By suitably specializing the parameters  $a, c, \lambda, A$  and  $B$ , we obtain the following subclasses of  $\mathcal{A}$ .

$$\begin{aligned} \text{(i)} \quad & \mathcal{R}^0(a, c, 1 - 2\alpha, -1) = \mathcal{R}_{a,c}(\alpha) \\ & = \left\{ f \in \mathcal{A} : \operatorname{Re} \left( \frac{\mathcal{L}(a, c)f(z)}{z} \right) > \alpha, 0 \leq \alpha < 1; z \in \mathbb{U} \right\}. \\ \text{(ii)} \quad & \mathcal{R}^2(2, 2 - \mu, \beta(1 - 2\alpha), -\beta) = \mathcal{R}(\mu, \alpha, \beta) \\ & = \left\{ f \in \mathcal{A} : \left| \frac{(\Omega_z^\mu f)'(z) - 1}{(\Omega_z^\mu f)'(z) + 1 - 2\alpha} \right| < \beta, 0 \leq \alpha < 1, 0 < \beta \leq 1, \right. \\ & \quad \left. 0 \leq \mu < 1; z \in \mathbb{U} \right\}. \end{aligned}$$

We note that  $\mathcal{R}(0, \alpha, \beta) = \mathcal{R}(\alpha, \beta)$  ( $0 \leq \alpha < 1, 0 < \beta \leq 1$ ), the class studied by Juneja and Mogra [10], which in turn give the class considered in [2] for  $\beta = 1$ .

$$\begin{aligned} \text{(iii)} \quad & \mathcal{R}^\lambda(m + 1, 1, 1 - 2\alpha, -1) = \mathcal{R}_m^\lambda(\alpha) \\ & = \left\{ f \in \mathcal{A} : \operatorname{Re} \left( (1 - \lambda) \frac{\mathcal{D}^m f(z)}{z} + \lambda \frac{\mathcal{D}^{m+1} f(z)}{z} \right) > \alpha, m \in \mathbb{N}_0, \right. \\ & \quad \left. 0 \leq \alpha < 1; z \in \mathbb{U} \right\}. \\ \text{(iv)} \quad & \mathcal{R}^\lambda(2, 1, 1 - 2\alpha, -1) = \mathcal{R}^\lambda(\alpha) \\ & = \left\{ f \in \mathcal{A} : \operatorname{Re} \left( f'(z) + \frac{\lambda}{2} z f''(z) \right) > \alpha, 0 \leq \lambda, 0 \leq \alpha < 1; z \in \mathbb{U} \right\}. \end{aligned}$$

Next, we define a subclass of  $\mathcal{F}$  as follows:

**Definition 1.2.** For the fixed parameters  $A, B$  ( $-1 \leq B < A \leq 1, 1 \leq B < 0$ ),  $a > 0$  and  $c > 0$ , a function  $f \in \mathcal{F}$  is said to be in the class  $\tilde{\mathcal{R}}^\lambda(a, c, A, B)$ , if it satisfies the following subordination relation:

$$(1.11) \quad (1 - \lambda) \frac{\mathcal{L}(a, c)f(z)}{z} + \lambda \frac{\mathcal{L}(a + 1, c)f(z)}{z} \prec \frac{1 + Az}{1 + Bz} \quad (\lambda \in \mathbb{C}, \operatorname{Re}(\lambda) \geq 0; z \in \mathbb{U}).$$

In view of (1.9), it is easily seen that the subordination relation (1.11) is equivalent to

$$\frac{\mathcal{L}(a, c)f(z)}{z} + \frac{\lambda}{a}z \left( \frac{\mathcal{L}(a, c)f(z)}{z} \right)' \prec \frac{1 + Az}{1 + Bz} \quad (z \in \mathbb{U}).$$

If we set  $h(z) = \mathcal{L}(a, c)f(z)/z$ , then the above expression further reduces to

$$h(z) + \frac{\lambda}{a}zh'(z) \prec \frac{1 + Az}{1 + Bz} \quad (\lambda \in \mathbb{C}, \operatorname{Re}(\lambda) \geq 0; z \in \mathbb{U}).$$

We write

$$\begin{aligned} \tilde{\mathcal{R}}^\lambda(1, 1, 1 - 2\alpha, -1) &= \tilde{\mathcal{R}}^\lambda(\alpha) \\ &= \left\{ f \in \mathcal{F} : \operatorname{Re} \left( \frac{f(z)}{z} + \lambda z \left( \frac{f(z)}{z} \right)' \right) > \alpha, z \in \mathbb{U} \right\} \end{aligned}$$

( $\lambda \in \mathbb{C}, \operatorname{Re}(\lambda) \geq 0, 0 \leq \alpha < 1$ ).

Noonan and Thomas [23] defined the  $q$ -th Hankel determinant of the function  $f \in \mathcal{A}$  given by (1.1) as

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \cdots & a_{n+q} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n+q-1} & a_{n+q} & \cdots & a_{n+2q-2} \end{vmatrix} \quad (a_1 = 1; n, q \in \mathbb{N}).$$

This determinant has been studied by several authors with the subject of inquiry ranging from the rate of growth of  $H_q(n)$  (as  $n \rightarrow \infty$ ) [24] to the determination of precise bounds with specific values of  $n$  and  $q$  for certain subclasses of analytic functions in the unit disc  $\mathbb{U}$ .

For  $n = 1, q = 2 = 1$  and  $n = q = 2$ , the Hankel determinant simplifies to

$$H_2(1) = |a_3 - a_2^2| \quad \text{and} \quad H_2(2) = |a_2a_4 - a_3^2|.$$

We refer to  $H_2(2)$  as the second Hankel determinant. It is known [4] that if  $f$  given by (1.1) is analytic and univalent in  $\mathbb{U}$ , then the sharp inequality  $H_2(1) = |a_3 - a_2^2| \leq 1$  holds. For a family  $\mathcal{F}$  of functions in  $\mathcal{A}$  of the form (1.1), the more general problem of finding the sharp upper bounds for the functionals  $|a_3 - \mu a_2^2|$  ( $\mu \in \mathbb{R}/\mathbb{C}$ ) is popularly known as Fekete–Szegő problem for the class  $\mathcal{F}$ . The Fekete–Szegő problem for the known classes of univalent functions, that is, starlike functions, convex functions and close-to-convex functions has been completely settled [5, 11, 12, 13]. Recently, Janteng et al. [8, 9] have obtained the sharp upper bounds to the second Hankel determinant  $H_2(2)$  for the family  $\mathcal{R}$ . For initial work on the class  $\mathcal{R}$ , one may refer to the paper by MacGregor [17].

A majorization properties for the class of starlike functions of complex order  $\gamma$  and the class of convex functions of complex order  $\gamma$  ( $\gamma \in \mathbb{C}^*$ ) has been investigated by Altıntaş et al. [1] and MacGregor [19] has also studied

the same problem for the classes  $\mathcal{S}^*$  and  $\mathcal{C}$ , respectively. Recently, Goyal and Goswami [6], and Goyal et al. [7] generalized these results for different function classes.

In the present article, by following the techniques devised by Libera and Złotkiewicz [14, 15], we solve the Fekete–Szegő problem and also determine the sharp upper bound to the second Hankel determinant for the class  $\mathcal{R}^\lambda(a, c, A, B)$ . We also investigate several majorization properties for certain subclasses of analytic functions in the unit disk  $\mathbb{U}$ . Relevant connections of the results presented here with those obtained in earlier works are also mentioned.

To establish our main results, we shall need the following lemmas.

**Lemma 1.1** ([4]). *Let the function  $\phi$ , given by (1.3) be a member of the class  $\mathcal{P}$ . Then*

$$|p_k| \leq 2 \quad (k \geq 1)$$

and the estimate is sharp for the function  $\phi(z) = (1+z)/(1-z)$ ,  $z \in \mathbb{U}$ .

**Lemma 1.2** ([16]). *If the function  $\phi$ , given by (1.3) belongs to the class  $\mathcal{P}$ , then for any  $\gamma \in \mathbb{C}$*

$$|p_2 - \gamma p_1^2| \leq 2 \max\{1, |2\gamma - 1|\}$$

and the result is sharp for the functions given by

$$\phi(z) = \frac{1+z^2}{1-z^2} \quad \text{and} \quad \phi(z) = \frac{1+z}{1-z} \quad (z \in \mathbb{U}).$$

**Lemma 1.3** ([15], see also [14]). *If the function  $\phi$ , given by (1.3) belongs to the class  $\mathcal{P}$ , then*

$$p_2 = \frac{1}{2} \{p_1^2 + (4 - p_1^2)x\}$$

and

$$p_3 = \frac{1}{4} \{p_1^3 + 2(4 - p_1^2)p_1x - (4 - p_1^2)p_1x^2 + 2(4 - p_1^2)(1 - |x|^2)z\}$$

for some complex numbers  $x, z$  satisfying  $|x| \leq 1$  and  $|z| \leq 1$ .

**2. Hankel determinant for the class  $\mathcal{R}^\lambda(a, c, A, B)$ .** Unless otherwise mentioned, we assume throughout the sequel that

$$a > 0, \quad c > 0, \quad \lambda \geq 0 \quad \text{and} \quad -1 \leq B < A \leq 1.$$

Now, we determine the sharp upper bound for the functional  $|a_3 - \gamma a_2^2|$  ( $\gamma \in \mathbb{R}$ ) for functions belonging to the class  $\mathcal{R}^\lambda(a, c, A, B)$ .

**Theorem 2.1.** *If  $\gamma \in \mathbb{R}$  and the function  $f$ , given by (1.1) belongs to the class  $\mathcal{R}^\lambda(a, c, A, B)$ , then*

$$(2.1) \quad |a_3 - \gamma a_2^2| \leq \begin{cases} -\frac{(c)_2(A-B)}{(a+1)(a+2\lambda)} \left\{ B + \frac{(a+1)(a+2\lambda)c(A-B)\gamma}{(a+\lambda)^2(c+1)} \right\}, & \gamma < \rho_1, \\ \frac{(c)_2(A-B)}{(a+1)(a+2\lambda)}, & \rho_1 \leq \gamma \leq \rho_2, \\ \frac{(c)_2(A-B)}{(a+1)(a+2\lambda)} \left\{ B + \frac{(a+1)(a+2\lambda)c(A-B)\gamma}{(a+\lambda)^2(c+1)} \right\}, & \gamma > \rho_2, \end{cases}$$

where

$$\rho_1 = -\frac{(1+B)(a+\lambda)^2(c+1)}{(a+1)(a+2\lambda)c(A-B)} \quad \text{and} \quad \rho_2 = \frac{(1-B)(a+\lambda)^2(c+1)}{(a+1)(a+2\lambda)c(A-B)}.$$

The estimate in (2.1) is sharp.

**Proof.** From (1.10), we have

$$(2.2) \quad (1-\lambda) \frac{\mathcal{L}(a, c)f(z)}{z} + \lambda \frac{\mathcal{L}(a+1, c)f(z)}{z} = \frac{1-A+(1+A)\phi(z)}{1-B+(1+B)\phi(z)}$$

( $z \in \mathbb{U}$ ), where the function  $\phi$ , given by (1.3) belongs to the class  $\mathcal{P}$ . Writing the series expansion of  $\mathcal{L}(a, c)f(z)/z$ ,  $\mathcal{L}(a+1, c)f(z)$ ,  $\phi(z)$  from (1.8) and (1.3) in (2.2), and comparing the like powers of  $z$  in the resulting equation, we deduce that

$$(2.3) \quad a_2 = \frac{c(A-B)}{2(a+\lambda)} p_1$$

$$(2.4) \quad a_3 = \frac{c(c+1)(A-B)}{2(a+1)(a+2\lambda)} \left\{ p_2 - \frac{1}{2}(1+B)p_1^2 \right\}$$

and

$$(2.5) \quad a_4 = \frac{c(c+1)(c+2)(A-B)}{2(a+1)(a+2)(a+3)} \left\{ p_3 - (1+B)p_1p_2 + \frac{1}{4}(1+B)^2p_1^3 \right\}.$$

Using (2.3) and (2.4), we obtain

$$|a_3 - \gamma a_2^2| = \frac{(c)_2(A-B)}{(a+1)(a+2\lambda)} \left| p_2 - \left\{ \frac{(a+1)(a+2\lambda)c(A-B)}{2(a+\lambda)^2(c+1)} \gamma + B + 1 \right\} p_1^2 \right|$$

and with the aid of Lemma 1.2, the above expression yields

$$(2.6) \quad \begin{aligned} & |a_3 - \gamma a_2^2| \\ & \leq \frac{2(c)_2(A-B)}{(a+1)(a+2\lambda)} \max \left\{ 1, \left| \frac{(a+1)(a+2\lambda)c(A-B)}{(a+\lambda)^2(c+1)} \gamma + B \right| \right\}. \end{aligned}$$

If  $\gamma < \rho_1$ , then

$$\frac{(a+1)(a+2\lambda)c(A-B)}{(a+\lambda)^2(c+1)} \gamma + B < -1,$$

which in view of (2.6) implies the first case of the estimate in (2.1). In the case  $\rho_1 \leq \gamma \leq \rho_2$ , we obtain

$$\left| \frac{(a+1)(a+2\lambda)c(A-B)}{(a+\lambda)^2(c+1)}\gamma + B \right| \leq 1.$$

Thus, from (2.6), we get the second case of the estimate in (2.1). Finally, for  $\gamma > \rho_2$ , we deduce that

$$\left| \frac{(a+1)(a+2\lambda)c(A-B)}{(a+\lambda)^2(c+1)}\gamma + B \right| > 1,$$

which again with the aid of (2.6) gives the third case of the estimate in (2.1).

It is easily seen that the estimate for the first and third cases in (2.1) are sharp for the function  $f$ , defined in  $\mathbb{U}$  by

$$f(z) = \begin{cases} \varphi(c, a; z) \star \frac{z(1+Az)}{1+Bz}, & \lambda = 0, \\ z {}_3F_2\left(1, \frac{a}{\lambda}, c; a, 1 + \frac{a}{\lambda}; z\right) \star \frac{z(1+Az)}{1+Bz}, & \lambda > 0. \end{cases}$$

The estimate for the second case in (2.1) is sharp for the function  $f$ , defined in  $\mathbb{U}$  by

$$(2.7) \quad f(z) = \begin{cases} \varphi(c, a; z) \star \frac{z(1+Az^2)}{1+Bz^2}, & \lambda = 0, \\ z {}_3F_2\left(1, \frac{a}{\lambda}, c; a, 1 + \frac{a}{\lambda}; z\right) \star \frac{z(1+Az^2)}{1+Bz^2}, & \lambda > 0, \end{cases}$$

where the function  ${}_3F_2$  is defined by (1.6). □

Setting  $a = 2$ ,  $c = 2 - \mu$  ( $0 \leq \mu < 1$ ),  $A = \beta(1 - 2\alpha)$ ,  $B = -\beta$  ( $0 \leq \alpha < 1$ ,  $0 < \beta \leq 1$ ) and  $\lambda = 2$  in Theorem 2.1, we obtain the following result.

**Corollary 2.1.** *If  $\gamma \in \mathbb{R}$  and the function  $f$ , given by (1.1) belongs to the class  $\mathcal{R}(\mu, \alpha, \beta)$ , then*

$$|a_3 - \gamma a_2^2| \leq \begin{cases} \frac{\beta(1-\alpha)(2-\mu)(3-\mu)}{9} \left\{ \beta - \frac{9\beta(1-\alpha)(2-\mu)\gamma}{4(3-\mu)} \right\}, & \gamma < \sigma_1, \\ \frac{\beta(1-\alpha)(2-\mu)(3-\mu)}{9}, & \sigma_1 \leq \gamma \leq \sigma_2, \\ \frac{\beta(1-\alpha)(2-\mu)(3-\mu)}{9} \left\{ \frac{9\beta(1-\alpha)(2-\mu)\gamma}{4(3-\mu)} - \beta \right\}, & \gamma > \sigma_2, \end{cases}$$

where

$$\sigma_1 = -\frac{4(1+\beta)(3-\mu)}{9\beta(1-\alpha)(2-\mu)} \quad \text{and} \quad \sigma_2 = \frac{4(1-\beta)(3-\mu)}{9\beta(1-\alpha)(2-\mu)}.$$

The estimate is sharp for the functions  $f$ , defined in  $\mathbb{U}$  by

$$f(z) = z {}_3F_2(1, 1, 2 - \mu; 2, 2; z) \star \frac{z\{1 + \beta(1 - 2\alpha)z\}}{1 - \beta z}$$

and

$$f(z) = z {}_3F_2(1, 1, 2 - \mu; 2, 2; z) \star \frac{z\{1 + \beta(1 - 2\alpha)z^2\}}{1 - \beta z^2}.$$



Letting  $a = 2, c = 1, A = 1 - 2\alpha$  ( $0 \leq \alpha < 1$ ) and  $B = -1$  in Theorem 1, we get

**Corollary 2.2.** *If  $\gamma \in \mathbb{R}$  and the function  $f$ , given by (1.1) belongs to the class  $\mathcal{R}^\lambda(\alpha)$ , then*

$$|a_3 - \gamma a_2^2| \leq \begin{cases} \frac{2(1-\alpha)}{3(1+\lambda)} \left\{ 1 - \frac{6(1+\lambda)(1-\alpha)\gamma}{(2+\lambda)^2} \right\}, & \gamma < 0, \\ \frac{2(1-\alpha)}{3(1+\lambda)}, & 0 \leq \gamma \leq \frac{(2+\lambda)^2}{3(1+\lambda)(1-\alpha)}, \\ \frac{2(1-\alpha)}{3(1+\lambda)} \left\{ \frac{6(1+\lambda)(1-\alpha)\gamma}{(2+\lambda)^2} - 1 \right\}, & \gamma > \frac{(2+\lambda)^2}{3(1+\lambda)(1-\alpha)}. \end{cases}$$

The estimate is sharp for the functions  $f$ , defined in  $\mathbb{U}$  by

$$f(z) = \begin{cases} \varphi(1, 2; z) \star \frac{z\{1+(1-2\alpha)z\}}{1-z}, & \lambda = 0, \\ z {}_3F_2(1, 1, 2 - \mu; 2, 2; z) \star \frac{z\{1+(1-2\alpha)z\}}{1-z}, & \lambda > 0 \end{cases}$$

and

$$f(z) = \begin{cases} \varphi(1, 2; z) \star \frac{z\{1+(1-2\alpha)z\}}{1-z}, & \lambda = 0, \\ z {}_3F_2(1, 1, 2 - \mu; 2, 2; z) \star \frac{z\{1+(1-2\alpha)z^2\}}{1-z^2}, & \lambda > 0. \end{cases}$$

In the following theorem, we find the sharp upper bound to the second Hankel determinant for the class  $\mathcal{R}^\lambda(a, c, A, B)$ .

**Theorem 2.2.** *Let  $a \geq c > 0, \lambda \geq 0, -1 \leq B < A \leq 1$  and*

$$(2.8) \quad 2(1+|B|)(a+2)(c+1)(a+\lambda)(a+3\lambda) \geq (1+2|B|)(a+1)(c+2)(a+2\lambda)^2.$$

*If the function  $f$ , given by (1.1) belongs to the class  $\mathcal{R}^\lambda(a, c, A, B)$ , then*

$$(2.9) \quad |a_2 a_4 - a_3^2| \leq \frac{(c)_2^2 (A - B)^2}{(a + 1)^2 (a + 2\lambda)^2}.$$

The estimate in (2.9) is sharp.

**Proof.** Assuming that  $f$ , given by (1.1) belongs to the class  $\mathcal{R}^\lambda(a, c, A, B)$  and using (2.3), (2.4) and (2.5), we deduce that

$$(2.10) \quad |a_2 a_4 - a_3^2| = \frac{c^2(c+1)(A-B)^2}{4(a+1)(a+\lambda)(a+3\lambda)} \times \left| K_1 p_1 p_3 - (1+B)(K_1 - K_2) p_1^2 p_2 + \frac{(1+B)^2(K_1 - K_2)}{4} p_1^4 \right|,$$

where

$$K_1 = \frac{c+2}{a+2} \quad \text{and} \quad K_2 = \frac{(c+1)(a+\lambda)(a+3\lambda)}{(a+1)(a+2\lambda)^2}.$$

Since the functions  $\phi(z)$  and  $\phi(e^{i\theta}z)$  ( $\theta \in \mathbb{R}$ ), defined by (1.3) are in the class  $\mathcal{P}$  simultaneously, we assume without loss of generality that  $p_1 > 0$ .

For convenience, we write  $p_1 = p$  ( $0 \leq p \leq 2$ ). Now, by using Lemma 1.3 in (2.10), we get

$$\begin{aligned}
 & |a_2a_4 - a_3^2| \\
 &= \frac{c^2(c+1)(A-B)^2}{4(a+1)(a+\lambda)(a+3\lambda)} \left| \left\{ \frac{K_1}{4}p^4 + \frac{K_1}{2}(4-p^2)p^2x \right. \right. \\
 &\quad \left. \left. - \frac{K_1}{4}(4-p^2)p^2x^2 + \frac{K_1}{2}(4-p^2)p(1-|x|^2)z \right\} \right. \\
 (2.11) \quad & \left. - \left\{ \frac{(1+B)(K_1-K_2)}{2}p^4 + \frac{(1+B)(K_1-K_2)}{2}(4-p^2)p^2 \right\} \right. \\
 & \left. - \left\{ \frac{K_2}{4}p^4 + \frac{K_2}{2}(4-p^2)p^2x + \frac{K_2}{4}(4-p^2)^2x^2 \right\} \right. \\
 & \left. + \frac{(1+B)^2(K_1-K_2)}{4}p^4 \right| \\
 &= \frac{c^2(c+1)(A-B)^2}{4(a+1)(a+\lambda)(a+3\lambda)} \left| \frac{B^2(K_1-K_2)}{4}p^4 - \frac{B(K_1-K_2)}{2}(4-p^2)p^2x \right. \\
 & \quad \left. - \frac{1}{4}(4-p^2) \{K_1p^2 + K_2(4-p^2)\}x^2 + \frac{K_1}{2}(4-p^2)p(1-|x|^2)z \right|
 \end{aligned}$$

for some complex numbers  $x$  ( $|x| \leq 1$ ) and  $z$  ( $|z| \leq 1$ ). Applying the triangle inequality in (2.11) and upon replacing  $|x|$  by  $y$  in the resulting expression, we get

$$\begin{aligned}
 (2.12) \quad |a_2a_4 - a_3^2| &\leq \frac{c^2(c+1)(A-B)^2}{4(a+1)(a+\lambda)(a+3\lambda)} \left\{ \frac{|B|^2(K_1-K_2)}{4}p^4 \right. \\
 &\quad \left. + \frac{|B|(K_1-K_2)}{2}(4-p^2)p^2y \right. \\
 &\quad \left. + \frac{1}{4}(4-p^2) \{K_1p^2 + K_2(4-p^2) - 2K_1p\}y^2 \right. \\
 &\quad \left. + \frac{K_1}{2}(4-p^2)p \right\} \\
 &=: \mathcal{G}(p, y)
 \end{aligned}$$

( $0 \leq p \leq 2$ ,  $0 \leq y \leq 1$ ). We next maximize the function  $\mathcal{G}(p, y)$  on the closed rectangle  $[0, 2] \times [0, 1]$ . Since

$$2K_2 > K_1 > K_2 > 0,$$

we have

$$\begin{aligned}
 \frac{\partial \mathcal{G}}{\partial y} &= \frac{c^2(c+1)(A-B)^2}{8(a+1)(a+\lambda)(a+3\lambda)}(4-p^2) \\
 &\quad \times [(K_1-K_2)|B|p^2 + (2-p)\{(2+p)K_2 - K_1p\}y] > 0
 \end{aligned}$$

for  $0 < p < 2$  and  $0 < y < 1$ . Thus,  $\mathcal{G}(p, y)$  cannot have a maximum in the interior on the closed rectangle  $[0, 2] \times [0, 1]$ . Therefore, for fixed  $p \in [0, 2]$

$$\max_{0 \leq y \leq 1} \mathcal{G}(p, y) = \mathcal{G}(p, 1) =: \mathcal{F}(p),$$

where

$$\begin{aligned} \mathcal{F}(p) = & \frac{c^2(c+1)(A-B)^2}{8(a+1)(a+\lambda)(a+3\lambda)} \left[ \frac{|B|^2(K_1 - K_2)}{4} p^4 \right. \\ & \left. + \frac{|B|(K_1 - K_2)}{2} (4 - p^2)p^2 + \frac{1}{4}(4 - p^2) \{K_1 p^2 + K_2(4 - p^2)\} \right]. \end{aligned}$$

A routine calculation yields

$$\mathcal{F}'(p) = p\{(|B|^2 - 2|B| - 1)(K_1 - K_2)p^2 + 2(1 + 2|B|)K_1 - 4(1 + |B|)K_2\}.$$

Thus,  $\mathcal{F}'(p) = 0$  implies that either  $p = 0$  or

$$p^2 = -\frac{2\{2(1 + |B|)K_2 - (1 + 2|B|)K_1\}}{\{2 - (1 - |B|^2)\}(K_1 - K_2)} < 0,$$

which is not true. We, further observe that

$$\mathcal{F}''(0) = 2\{(1 + 2|B|)K_1 - 2(1 + |B|)K_2\} < 0$$

by (2.8). Since  $\mathcal{F}(2) < \mathcal{F}(0)$ ,  $\max_{0 \leq p \leq 2} \mathcal{F}(p)$  occurs at  $p = 0$ . Thus, the upper bound of (2.12) corresponds to  $p = 0$  and  $y = 1$ , from which we get the estimate in (2.9).

It is easily seen that the estimate (2.9) is sharp for the function  $f$ , given by (2.7) and thus the proof of Theorem 2.2 is completed.  $\square$

Setting  $\lambda = 0$ ,  $A = 1 - 2\alpha$  ( $0 \leq \alpha < 1$ ) and  $B = -1$  in Theorem 2.2, we get the following result obtained by Mishra and Kund [21].

**Corollary 2.3.** *If  $a \geq c > 0$ ,  $ac - 2a + 5c + 2 \geq 0$  and the function  $f$ , given by (1.1) belongs to the class  $\mathcal{R}_{a,c}(\alpha)$ , then*

$$|a_2 a_4 - a_3^2| \leq \left\{ \frac{2(c)_2(1 - \alpha)}{(a)_2} \right\}^2.$$

The estimate is sharp for the function  $f$ , defined by

$$f(z) = \varphi(c, a; z) \star \frac{z\{1 + (1 - 2\alpha)z^2\}}{1 - z^2} \quad (0 \leq \alpha < 1; z \in \mathbb{U}).$$

Letting  $a = 2$ ,  $c = 2 - \mu$ ,  $A = \beta(1 - 2\alpha)$ ,  $B = -\beta$  and  $\lambda = 2$  in Theorem 2.2, we obtain

**Corollary 2.4.** *If the function  $f$ , given by (1.1) belongs to the class  $\mathcal{R}(\mu, \alpha, \beta)$ , then*

$$|a_2 a_4 - a_3^2| \leq \frac{\{\beta(1 - \alpha)(2 - \mu)(3 - \mu)\}^2}{9} \quad (0 \leq \mu < 1, 0 \leq \alpha < 1, 0 < \beta \leq 1)$$

and the estimate is sharp for the function  $f$ , defined by

$$f(z) = z {}_3F_2(1, 1, 2 - \mu; 2, 2; z) \star \frac{z\{1 + \beta(1 - 2\alpha)z^2\}}{1 - \beta z^2} \quad (z \in \mathbb{U}).$$

Taking  $a = 2$ ,  $c = 1$ ,  $A = 1 - 2\alpha$  and  $B = -1$  in Theorem 2.2, we get the following result, which in turn yields the corresponding work of Mishra and Kund [21] for  $\lambda = 0$ , and the work of Janteng et al. [8] for  $\lambda = \alpha = 0$ .

**Corollary 2.5.** *If the function  $f$ , given by (1.1) belongs to the class  $\mathcal{R}^\lambda(\alpha)$ , then*

$$|a_2 a_4 - a_3^2| \leq \frac{4(1 - \alpha)^2}{9(1 + \lambda)^2} \quad \left(0 \leq \alpha < 1, 0 \leq \lambda \leq \frac{5 + 2\sqrt{10}}{3}\right)$$

and the estimate is sharp for the function  $f$ , defined in  $\mathbb{U}$  by

$$f(z) = \begin{cases} \varphi(1, 2; z) \star \frac{z\{1 + (1 - 2\alpha)z^2\}}{1 - z^2}, & \lambda = 0 \\ z {}_3F_2\left(1, 1, \frac{2}{\lambda}; 2, 1 + \frac{2}{\lambda}; z\right) \star \frac{z\{1 + (1 - 2\alpha)z^2\}}{1 - z^2}, & \lambda > 0. \end{cases}$$

**3. Majorization properties.** We prove the following lemmas, which will be used in our investigation of majorization properties for the class  $\tilde{\mathcal{R}}^\lambda(a, c, A, B)$ .

**Lemma 3.1.** *If  $a \geq c > 0$  and the function  $g$ , given by (1.2) belongs to the class  $\tilde{\mathcal{R}}^\lambda(a, c, A, B)$ , then*

$$(3.1) \quad \sum_{n=1}^{\infty} b_{n+1} \leq \frac{c(A - B)}{\{a + \operatorname{Re}(\lambda)\}(1 - B)}.$$

**Proof.** It follows from (1.11) that

$$\operatorname{Re} \left\{ (1 - \lambda) \frac{\mathcal{L}(a, c)f(z)}{z} + \lambda \frac{\mathcal{L}(a + 1, c)f(z)}{z} \right\} > \frac{1 - A}{1 - B} \quad (z \in \mathbb{U})$$

which upon substituting the series expansion of  $\mathcal{L}(a, c)f(z)/z$  and  $\mathcal{L}(a + 1, c)f(z)/z$  gives

$$\operatorname{Re} \left\{ 1 - \sum_{n=1}^{\infty} \frac{(a)_n}{(c)_n} \left(1 + \frac{\lambda n}{a}\right) b_{n+1} z^n \right\} > \frac{1 - A}{1 - B} \quad (z \in \mathbb{U}).$$

Letting  $z \rightarrow 1^-$  through real values in the above expression, we find that

$$1 - \sum_{n=1}^{\infty} \frac{(a)_n}{(c)_n} \left\{1 + \frac{\operatorname{Re}(\lambda)n}{a}\right\} b_{n+1} > \frac{1 - A}{1 - B}.$$

Since  $a \geq c > 0$ ,  $b_{n+1} \geq 0$  and  $\operatorname{Re}(\lambda) \geq 0$ , the above inequality implies that

$$\frac{a}{c} \left\{1 + \frac{\operatorname{Re}(\lambda)}{a}\right\} \sum_{n=1}^{\infty} b_{n+1} \leq \sum_{n=1}^{\infty} \frac{(a)_n}{(c)_n} \left\{1 + \frac{\operatorname{Re}(\lambda)n}{a}\right\} b_{n+1} \leq \frac{A - B}{1 - B}.$$

This completes the proof of Lemma 3.1.  $\square$

**Lemma 3.2.** *Under the hypothesis of Lemma 3.1, we have for  $|z| = r < 1$*

$$(3.2) \quad 1 - \frac{c(A - B)r}{\{a + \operatorname{Re}(\lambda)\}(1 - B)} \leq \operatorname{Re} \left( \frac{g(z)}{z} \right) \\ \leq \left| \frac{g(z)}{z} \right| \leq 1 - \frac{c(A - B)r}{\{a + \operatorname{Re}(\lambda)\}(1 - B)}.$$

**Proof.** Since

$$\left| \frac{g(z)}{z} \right| \leq 1 + \left( \sum_{n=1}^{\infty} b_{n+1} \right) |z| \quad (z \in \mathbb{U}),$$

Lemma 3.1 leads to

$$\left| \frac{g(z)}{z} \right| \leq 1 + \frac{c(A - B)r}{\{a + \operatorname{Re}(\lambda)\}(1 - B)} \quad (|z| = r < 1).$$

Similarly, we have

$$\operatorname{Re} \left( \frac{g(z)}{z} \right) = 1 - \operatorname{Re} \left( \sum_{n=1}^{\infty} b_{n+1} z^n \right) \geq 1 - \left| \sum_{n=1}^{\infty} b_{n+1} z^n \right| \\ \geq 1 - \left( \sum_{n=1}^{\infty} b_{n+1} \right) |z| \geq 1 - \frac{c(A - B)r}{\{a + \operatorname{Re}(\lambda)\}(1 - B)} \quad (|z| = r < 1)$$

and the proof of Lemma 3.2 is completed.  $\square$

Now, we prove

**Theorem 3.1.** *Let the function  $g$  be in the class  $\mathcal{F}$ . If  $a \geq c > 0$ , the function  $h \in \mathcal{T}$  satisfies*

$$(3.3) \quad \frac{z^2 (\mathcal{L}(a, c)h)'(z)}{\mathcal{L}(a, c)h(z)} \in \tilde{\mathcal{K}}^\lambda(a, c, A, B)$$

and  $\mathcal{L}(a, c)g \ll \mathcal{L}(a, c)h$  in  $\mathbb{U}$ , then

$$(3.4) \quad |(\mathcal{L}(a, c)g)'(z)| \leq |(\mathcal{L}(a, c)h)'(z)| \quad (|z| \leq r(\lambda, a, c, A, B)),$$

where  $r(\lambda, a, c, A, B)$  is the root of the cubic equation

$$(3.5) \quad c(A - B)r^3 - \{a + \operatorname{Re}(\lambda)\}(1 - B)r^2 \\ - [2\{a + \operatorname{Re}(\lambda)\}(1 - B) + c(A - B)]r \\ + \{a + \operatorname{Re}(\lambda)\}(1 - B) = 0$$

in  $(0, 1)$ .

**Proof.** From (3.3), by using Lemma 3.2, we get for  $|z| = r < 1$

$$(3.6) \quad |\mathcal{L}(a, c)h(z)| \leq \frac{\{a + \operatorname{Re}(\lambda)\}(1 - B)r}{\{a + \operatorname{Re}(\lambda)\}(1 - B) - c(A - B)r} |(\mathcal{L}(a, c)h)'(z)|.$$

Since  $\mathcal{L}(a, c)g(z) \ll \mathcal{L}(a, c)h(z)$  in  $\mathbb{U}$ , we have by (1.5)

$$\mathcal{L}(a, c)g(z) = \psi(z)\mathcal{L}(a, c)h(z) \quad (z \in \mathbb{U}),$$

where the function  $\psi$  is analytic in  $\mathbb{U}$  and satisfies  $|\psi(z)| \leq 1$  in  $\mathbb{U}$ , so that

$$(3.7) \quad (\mathcal{L}(a, c)g)'(z) = \psi(z) (\mathcal{L}(a, c)h)'(z) + \psi'(z)\mathcal{L}(a, c)h(z) \quad (z \in \mathbb{U}).$$

Using the following estimate [22]

$$(3.8) \quad |\psi'(z)| \leq \frac{1 - |\psi(z)|^2}{1 - |z|^2} \quad (z \in \mathbb{U})$$

followed by (3.6) in (3.7), we obtain

$$\begin{aligned} & |(\mathcal{L}(a, c)g)'(z)| \\ & \leq \left[ |\psi(z)| + \frac{1 - |\psi(z)|^2}{1 - |z|^2} \frac{\{a + \operatorname{Re}(\lambda)\}(1 - B)r}{\{a + \operatorname{Re}(\lambda)\}(1 - B) - c(A - B)r} \right] |(\mathcal{L}(a, c)h)'(z)| \end{aligned}$$

which upon setting  $|\psi(z)| = x$  ( $0 \leq x \leq 1$ ) yields the inequality

$$(3.9) \quad |(\mathcal{L}(a, c)g)'(z)| \leq \left\{ \frac{\Psi(x)}{(1 - r^2) \{ (a + \operatorname{Re}(\lambda))(1 - B) - c(A - B)r \}} \right\} |(\mathcal{L}(a, c)h)'(z)|,$$

where

$$\begin{aligned} \Psi(x) &= \{a + \operatorname{Re}(\lambda)\}(1 - B)rx^2 \\ &+ (1 - r^2) [\{a + \operatorname{Re}(\lambda)\}(1 - B) - c(A - B)r]x \\ &+ \{a + \operatorname{Re}(\lambda)\}(1 - B)r. \end{aligned}$$

The function  $\Psi$  attains its maximum value at  $x = 1$  with  $r = r(\lambda, a, c, A, B)$ , the root of the equation (3.5) contained in  $(0, 1)$ . Furthermore, if  $0 \leq y \leq r(\lambda, a, c, A, B)$ , then the function

$$\begin{aligned} \Theta(x) &= -\{a + \operatorname{Re}(\lambda)\}(1 - B)yx^2 \\ &+ (1 - y^2) [\{a + \operatorname{Re}(\lambda)\}(1 - B) - c(A - B)y]x \\ &+ \{a + \operatorname{Re}(\lambda)\}(1 - B)y \end{aligned}$$

increases in the interval  $0 \leq x \leq 1$ , so that

$$\Theta(x) \leq \Theta(1) = (1 - y^2) [\{a + \operatorname{Re}(\lambda)\}(1 - B) - c(A - B)y].$$

Thus, by substituting  $x = 1$  in (3.9), we conclude that the inequality in (3.4) holds true for  $|z| \leq r(\lambda, a, c, A, B)$ , where  $r(\lambda, a, c, A, B)$  is given by (3.5). This completes the proof of Theorem 3.1.  $\square$

For  $a = c = 1$ ,  $A = 1 - 2\alpha$  ( $0 \leq \alpha < 1$ ) and  $B = -1$ , Theorem 3.1 gives the following result.

**Corollary 3.1.** *If the function  $g \in \mathcal{T}$  and  $g \ll h$  in  $\mathbb{U}$ , where  $z^2h'(z)/h(z) \in \tilde{\mathcal{R}}^\lambda(\alpha)$ , then*

$$|g'(z)| \leq |h'(z)| \quad (|z| \leq r(\lambda, \alpha)),$$

where  $r(\lambda, \alpha)$  is the root of the cubic equation

$$(1 - \alpha)r^3 - \{1 + \operatorname{Re}(\lambda)\}r^2 - \{3 - \alpha + 2\operatorname{Re}(\lambda)\}r + 1 + \operatorname{Re}(\lambda) = 0$$

in  $(0, 1)$ .

In the special case  $\lambda = 0$ , Corollary 3.1 simplifies to the following result.

**Corollary 3.2.** *Let the function  $g$  be in the class  $\mathcal{T}$ . If the function  $h \in \mathcal{S}^*(\alpha) \cap \mathcal{T}$  and  $g \ll h$  in  $\mathbb{U}$ , then*

$$|g'(z)| \leq |h'(z)| \quad (|z| \leq r(\alpha)),$$

where  $r(\alpha)$  is the root of the cubic equation  $(1 - \alpha)r^3 - r^2 - (3 - \alpha)r + 1 = 0$  in  $(0, 1)$ .

With the aid of the following inclusion relation [27, Theorem 7]

$$\mathcal{C}(\alpha) \cap \mathcal{T} \subset \mathcal{S}^*\left(\frac{2}{3 - \alpha}\right) \cap \mathcal{T} \quad (0 \leq \alpha < 1),$$

we get the following result from Corollary 3.2.

**Corollary 3.3.** *Let the function  $g$  be in the class  $\mathcal{T}$ . If the function  $h \in \mathcal{C}(\alpha) \cap \mathcal{T}$  and  $g \ll h$  in  $\mathbb{U}$ , then*

$$|g'(z)| \leq |h'(z)| \quad (|z| \leq \tilde{r}(\alpha)),$$

where  $\tilde{r}(\alpha)$  is the root of the cubic equation  $(1 - \alpha)r^3 - (3 - \alpha)r^2 - (7 - 3\alpha)r + (3 - \alpha) = 0$  in  $(0, 1)$ .

Finally, we prove

**Theorem 3.2.** *Let the function  $f$  be in the class  $\mathcal{A}$ . If the function  $g \in \mathcal{A}$  satisfies the subordination condition:*

$$(3.10) \quad \frac{z(\mathcal{L}(a, c)g)'(z)}{\mathcal{L}(a, c)g(z)} \prec \frac{1 + Az}{1 + Bz} \quad (-1 \leq B < A \leq 1; z \in \mathbb{U})$$

and  $\mathcal{L}(a, c)f \ll \mathcal{L}(a, c)g$  in  $\mathbb{U}$ , then

$$(3.11) \quad |\mathcal{L}(a + 1, c)f(z)| \leq |\mathcal{L}(a + 1, c)g(z)| \quad (|z| \leq r(a, A, B)),$$

where  $r(a, A, B)$  is the root of the cubic equation

$$(3.12) \quad |A + (a - 1)B|r^3 - (a + 2|B|)r^2 - (|A + (a - 1)B| + 2)r + a = 0$$

in  $(0, 1)$ .

**Proof.** From the definition of subordination, it follows from (3.10) that

$$(3.13) \quad \frac{z(\mathcal{L}(a, c)g)'(z)}{\mathcal{L}(a, c)g(z)} = \frac{1 + A\omega(z)}{1 + B\omega(z)} \quad (z \in \mathbb{U}),$$

where  $\omega$  is analytic in  $\mathbb{U}$  with  $\omega(0) = 0$  and  $|\omega(z)| < 1$  for all  $z \in \mathbb{U}$ . Now, making use of the identity (1.9) for the function  $g$  in (3.13), we deduce that

$$(3.14) \quad \begin{aligned} |\mathcal{L}(a, c)g(z)| &\leq \frac{a(1 + |B||\omega(z)|)}{a - |A + (a - 1)B||\omega(z)|} |\mathcal{L}(a + 1, c)g(z)| \\ &\leq \frac{a(1 + |B||z|)}{a - |A + (a - 1)B||z|} |\mathcal{L}(a + 1, c)g(z)| \quad (z \in \mathbb{U}). \end{aligned}$$

Since  $\mathcal{L}(a, c)f$  is majorized by  $\mathcal{L}(a, c)g$  in  $\mathbb{U}$ , we have

$$\mathcal{L}(a, c)f(z) = \psi(z)\mathcal{L}(a, c)g(z) \quad (z \in \mathbb{U}),$$

where  $\psi$  is analytic in  $\mathbb{U}$  and satisfies  $|\psi(z)| \leq 1$  in  $\mathbb{U}$ . Differentiating the above expression with respect to  $z$ , using the identity (1.9) for both the functions  $f$  and  $g$  in the resulting equation, we obtain

$$(3.15) \quad a \mathcal{L}(a + 1, c)f(z) = a \psi(z)\mathcal{L}(a + 1, c)g(z) + z\psi'(z)\mathcal{L}(a, c)g(z)$$

( $z \in \mathbb{U}$ ). Using the estimate (3.8) and (3.14) in (3.15), we get

$$\begin{aligned} &|\mathcal{L}(a + 1, c)f(z)| \\ &\leq \left\{ |\psi(z)| + \frac{1 - |\psi(z)|^2}{1 - |z|^2} \frac{(1 + |B||z|)|z|}{a - |A + (a - 1)B||z|} \right\} |\mathcal{L}(a + 1, c)g(z)| \end{aligned}$$

which upon setting  $|z| = r$  and  $|\psi(z)| = x$  ( $0 \leq x \leq 1$ ) yields the inequality

$$|\mathcal{L}(a + 1, c)f(z)| \leq \left[ \frac{\Psi(x)}{(1 - r^2)\{a - |A + (a - 1)B|r\}} \right] |\mathcal{L}(a + 1, c)g(z)|,$$

where

$$\Psi(x) = -r(1 + |B|r)x^2 + (1 - r^2)\{a - |A + (a - 1)B|r\}x + (1 + |B|r)r$$

which takes on its maximum value at  $x = 1$  with  $r = r(a, A, B)$ , where  $r(a, A, B)$  is the root of the equation (3.12) in  $(0, 1)$ .

The remaining part of the proof of Theorem 3.2 is much akin to that of Theorem 3.1, and so we omit the details.  $\square$

Letting  $a = c = 1$ ,  $A = \beta(1 - 2\alpha)$  and  $B = -\beta$  in Theorem 3.2, we get

**Corollary 3.4.** *Let the function  $f$  be in the class  $\mathcal{A}$ . If the function  $g \in \mathcal{A}$  satisfies*

$$\left| \frac{zg'(z)/g(z) - 1}{zg'(z)/g(z) + 1 - 2\alpha} \right| < \beta \quad (0 \leq \alpha < 1, 0 < \beta \leq 1; z \in \mathbb{U})$$

and  $f \ll g$  in  $\mathbb{U}$ , then

$$|f'(z)| \leq |g'(z)| \quad (|z| \leq r(\alpha, \beta)),$$

where

$$r(\alpha, \beta) = \begin{cases} \frac{3 + \beta|1 - 2\alpha| - \sqrt{\beta^2|1 - 2\alpha|^2 + 2\beta|1 - 2\alpha| + 9}}{2\beta|1 - 2\alpha|}, & \alpha \neq \frac{1}{2}, \\ \frac{1}{3}, & \alpha = \frac{1}{2}. \end{cases}$$



Using the following best possible inclusion relationship [18]

$$\mathcal{C}(\alpha) \subset \mathcal{S}^*(\varkappa(\alpha)),$$

where

$$(3.16) \quad \varkappa(\alpha) = \begin{cases} \frac{1-2\alpha}{2^{2(1-\alpha)}(1-2^{2\alpha-1})}, & \alpha \neq \frac{1}{2}, \\ \frac{1}{2 \ln 2}, & \alpha = \frac{1}{2}, \end{cases}$$

and taking  $\beta = 1$  in Corollary 3.4, we obtain

**Corollary 3.5.** *Let the function  $f$  be in the class  $\mathcal{A}$ . If the function  $g \in \mathcal{C}(\alpha)$  and  $f \ll g$  in  $\mathbb{U}$ , then*

$$|f'(z)| \leq |g'(z)| \quad (|z| \leq r(\varkappa(\alpha))),$$

where

$$r(\varkappa(\alpha)) = \begin{cases} \frac{3+|1-2\varkappa(\alpha)|-\sqrt{|1-2\varkappa(\alpha)|^2+2|1-2\varkappa(\alpha)|+9}}{2|1-2\varkappa(\alpha)|}, & \varkappa(\alpha) \neq \frac{1}{2}, \\ \frac{1}{3}, & \varkappa(\alpha) = \frac{1}{2} \end{cases}$$

and  $\varkappa(\alpha)$  is given by (3.16).

**Remark.** (i) For the choice  $\beta = 1$  in Corollary 3.4, we get the result due to Altıntaş et al. [1, Theorem 1], which in turn yields the corresponding work of MacGregor [19, Theorem 1B] for  $\alpha = 0$ .

(ii) Corollary 3.5 improves the corresponding result obtained by Altıntaş et al. [1, Theorem 2].

(iii) For the case  $\alpha = 0$ , Corollary 3.5 yields a result due to MacGregor [19, Theorem 1C].

**Acknowledgement.** The authors would like to thank the referee(s) for the suggestions/comments which improved the presentation of the paper.

#### REFERENCES

- [1] Altıntaş, O., Özkan, O., Srivastava, H. M., *Majorization by starlike functions of complex order*, Complex Var. **46** (2001), 207–218.
- [2] Caplinger, T. R., Causey, W. M., *A class of univalent functions*, Proc. Amer. Math. Soc. **39** (1973), 357–361.
- [3] Carlson, B. C., Shaffer, D. B., *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal. **15** (1984), 737–745.
- [4] Duren, P. L., *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, 259, Springer-Verlag, New York, USA, 1983.
- [5] Fekete, M., Szegő, G., *Eine Bemerkung über ungerede schlichte funktionen*, J. London Math. Soc. **8** (1933), 85–89.
- [6] Goyal, S. P., Goswami, P., *Majorization for certain subclass of analytic functions defined by linear operator using differential subordination*, Appl. Math. Letters **22** (2009), 1855–1858.
- [7] Goyal, S. P., Bansal, S. K., Goswami, P., *Majorization for certain classes of functions by fractional derivatives*, J. Appl. Math. Stat. Informatics **6**(2) (2010), 45–50.

- 
- [8] Janteng, A., Halim, S. A., Darus, M., *Coefficient inequality for a function whose derivative has a positive real part*, J. Inequal. Pure Appl. Math. **7** (2006), Art. 50 [<http://jipam.vu.edu.au/>].
- [9] Janteng, A., Halim, S. A., Darus, M., *Estimate on the second Hankel functional for functions whose derivative has a positive real part*, J. Quality Measurement and Analysis **4** (2008), 189–195.
- [10] Juneja, O. P., Mogra, M. L., *A class of univalent functions*, Bull. Sci. Math. (2) **103** (1979), 435–447.
- [11] Keogh, F. R., Merkes, E. P., *A coefficient inequality for certain classes of analytic functions*, Proc. Amer. Math. Soc. **20** (1969), 8–12.
- [12] Koepf, W., *On the Fekete-Szegő problem for close-to-convex functions. II*, Arch. Math. (Basel) **49** (1987), 420–433.
- [13] Koepf, W., *On the Fekete-Szegő problem for close-to-convex functions*, Proc. Amer. Math. Soc. **101** (1987), 89–95.
- [14] Libera, R. J., Zlotkiewicz, E. J., *Early coefficient of the inverse of a regular convex function*, Proc. Amer. Math. Soc. **85** (2) (1982), 225–230.
- [15] Libera, R. J., Zlotkiewicz, E. J., *Coefficient bounds for the inverse of a function with derivative in  $\mathcal{P}$* , Proc. Amer. Math. Soc. **87** (2) (1983), 251–257.
- [16] Ma, W. C., Minda, D., *A unified treatment of some special classes of univalent functions*, Proceedings of the Conference on Complex Analysis (Tianjin, 1992), Z. Li, F. Ren, L. Yang and S. Zhang (Eds.), Int. Press, Cambridge, MA, 1994, 157–169.
- [17] MacGregor, T. H., *Functions whose derivative have a positive real part*, Trans. Amer. Math. Soc. **104**(3) (1962), 532–537.
- [18] MacGregor, T. H., *The radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc. **14** (1963), 514–520.
- [19] MacGregor, T. H., *Majorization by univalent functions*, Duke Math. J. **34** (1967), 95–102.
- [20] Miller, S. S., Mocanu, P. T., *Differential Subordinations: Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics, Vol. 225, Marcel Dekker Inc., New York and Basel, 2000.
- [21] Mishra, A. K., Kund, S. N., *The second Hankel determinant for a class of analytic functions associated with the Carlson-Shaffer operator*, Tamkang J. Math. **44**(1) (2013), 73–82.
- [22] Nehari, Z., *Conformal Mapping*, McGraw-Hill Book Company, New York, Toronto and London, 1952.
- [23] Noonan, J. W., Thomas, D. K., *On the second Hankel determinant of areally mean  $p$ -valent functions*, Trans. Amer. Math. Soc. **223** (1976), 337–346.
- [24] Noor, K. I., *Hankel determinant problem for the class of functions with bounded boundary rotation*, Rev. Roum. Math. Pures Appl. **28** (1983), no. 8, 731–739.
- [25] Owa, S., Srivastava, H. M., *Univalent and starlike generalized hypergeometric functions*, Canad. J. Math. **39** (1987), 1057–1077.
- [26] Ruscheweyh, St., *New criteria for univalent functions*, Proc. Amer. Math. Soc. **49** (1975), 109–115.
- [27] Silverman, H., *Univalent functions with negative coefficients*, Proc. Amer. Math. Soc. **51** (1975), 109–116.
- [28] Srivastava, H. M., Karlson, P. W., Karlsson, Per W., *Multiple Gaussian Hypergeometric Series* (Mathematics and its Applications), A Halsted Press Book (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.

Jagannath Patel  
Department of Mathematics  
Utkal University  
Vani Vihar, Bhubaneswar-751004  
India  
e-mail: [jpatelmath@yahoo.co.in](mailto:jpatelmath@yahoo.co.in)

Ashok Kumar Sahoo  
Department of Mathematics  
Veer Surendra Sai University of Technology  
Sidhi Vihar, Burla-768018  
India  
e-mail: [ashokuamt@gmail.com](mailto:ashokuamt@gmail.com)

Received January 14, 2014