

MARCIN BORYC and ŁUKASZ KRUK

A multidimensional singular stochastic control problem on a finite time horizon

ABSTRACT. A singular stochastic control problem in n dimensions with time-dependent coefficients on a finite time horizon is considered. We show that the value function for this problem is a generalized solution of the corresponding HJB equation with locally bounded second derivatives with respect to the space variables and the first derivative with respect to time. Moreover, we prove that an optimal control exists and is unique.

Singular stochastic control is a class of problems in which one is allowed to change the drift of a Markov process (usually a diffusion) at a price proportional to the variation of the control used. Admissible controls do not have to be absolutely continuous with respect to the Lebesgue measure and they may have jumps. This setup is natural for many problems of practical interest, including portfolio selection in finance, control of queueing networks and spacecraft control, to mention just a few examples. The reader is referred to Chapter VIII of [5] for more information and basic references.

One-dimensional singular stochastic control problems are well understood by now, see, e.g., [2] and the references given there. In this case, if the running cost is convex, the optimal control makes the underlying process a reflected diffusion at the boundary of the so-called *nonaction region* \mathcal{C} . In the case of a diffusion with time-independent coefficients and discounted cost on the infinite time horizon, \mathcal{C} is just an interval and the value function enjoys

2010 *Mathematics Subject Classification.* Primary: 93E20; Secondary: 35Q93.

Key words and phrases. Singular stochastic control, generalized derivative, HJB equation, optimal control.

C^2 -regularity (*smooth fit*). Both C^2 -regularity of the value function and the characterization of the optimally controlled process have been extended to the case of singular control for the two-dimensional Brownian motion [14]. In $n \geq 3$ dimensions, except for “close to one-dimensional” cases of a single push direction [15, 16] and the radially symmetric running cost [9], only partial results are known. For example, for optimal control of the Brownian motion on the infinite time horizon, regularity of the boundary of \mathcal{C} away from some “corner points” was shown in [17] and a characterization of the optimal control as a solution of the corresponding modified Skorokhod problem was given in [8].

In this paper we consider a n -dimensional singular stochastic control problem on a finite time horizon in which state is governed by a linear stochastic differential equation with time-dependent coefficients, the running cost is convex and controls may act in any direction. We provide estimates for the corresponding value function. These estimates imply that the value function has locally bounded generalized derivatives of the second order with respect to the space variable and of the first order with respect to the time variable. These properties are needed to consider the value function as a solution of the corresponding parabolic Hamilton–Jacobi–Bellman (HJB) equation in some generalized sense and to show existence and uniqueness of an optimal control.

Similar results have been shown in Theorem 2.1 and Theorem 3.4 from [2] in the one-dimensional case with a single push direction. The corresponding results for a multidimensional singular stochastic control problem on the infinite time horizon with time-independent drift, covariance, cost (i.e., for the elliptic case) can be found in [11]. Our article contains a generalization (or adjustment) of the approach of [2, 11] to an n -dimensional parabolic problem. It turns out that while the main ideas from those papers may be applied in our case, a mathematically rigorous analysis of our problem is somewhat delicate and needs rather careful arguments.

Our motivation for pursuing this project is the hope that the results given here will allow for a characterization of the optimal policy in the parabolic case as a solution to the corresponding Skorokhod problem for a domain with time-dependent (moving) boundary, which would be an analog of the main theorem from [8]. Indeed, the analysis of [8] used the results from [11] as the starting point, so it is plausible that their analogs will be useful in proving the corresponding result on a finite time horizon. Such a characterization would address a long-standing open problem on the structure of the optimal control in the case under consideration. We hope to address this issue in a subsequent paper.

Existence results for multidimensional singular control problems closely related to our work may be found in [1, 3, 6]. Apparently, in spite of their considerable generality, none of them contains our existence result

as a special case. Indeed, in these papers optimal *weak solutions* to the corresponding SDEs are constructed, while we are concerned about finding an optimal *strong solution*, i.e., for the given (as opposed to some) filtration and underlying Brownian motion. Moreover, the problem considered in [1] is elliptic and the allowable control directions lie in a cone, the opening of which cannot be too wide. In [3, 6] the time horizon is finite, but the problem considered in [3] has the final cost instead of the running cost, while in [6] the drift of the controlled diffusion is bounded, which excludes its linear dependence on the state.

The structure of this paper is as follows. In Section 1 we pose the singular stochastic control problem, give definitions and prove lemmas needed in further considerations. In Section 2 we prove estimates for the value function. In Section 3 we consider the Bellman's dynamic programming principle (DPP) and the HJB equation related to this problem. Section 4 contains proofs of existence and uniqueness of an optimal control.

1. Notation, assumptions and lemmas. Let $\mathbb{M}^{n \times n}$ denote the set of matrices of dimension $n \times n$ with the operator norm, i.e. $\|A\| = \sup\{|Ax| : x \in \mathbb{R}^n, |x| = 1\}$. Let $T > 0$ be a fixed number representing our time horizon. For a function $u = u(x, t) : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}$ we denote the gradient and the Hessian of u with respect to the space variables (i.e., x_i) by Du and D^2u , respectively.

Let $(W_t, t \geq 0)$ be a standard n -dimensional Brownian motion defined on a complete probability space (Ω, \mathcal{F}, P) . Let $(\mathcal{F}_t, t \geq 0)$ be the augmentation of the filtration generated by W (see [7], p. 89). Denote by \mathcal{V} the set of controls v which are left-continuous, adapted to the filtration $(\mathcal{F}_t, t \geq 0)$ random processes acting from $[0, T]$ into \mathbb{R}^n , with P -a.s. bounded variation and s.t. $v(0) = 0$ P -a.s. We note that these processes are also progressively measurable (see [7], Th. 1.1.13). As it is customary in singular stochastic control theory (see, e.g., [8]), we write $v(t) = \int_0^t \gamma(s) d\xi(s)$, where $|\gamma(t)| = 1$ for every $t \in [0, T]$ and ξ is nondecreasing and left-continuous. In other words, $\xi(t)$ is the total variation of v on the time interval $[0, t]$ and $\gamma(t)$ is the Radon–Nikodym derivative of the vector-valued measure induced by v on $[0, T]$ with respect to its total variation ξ .

Consider the state process described by the stochastic integral equation

$$(1) \quad y_{xt}(s) = x + \int_t^s \left(a(r)y_{xt}(r) + b(r) \right) dr + \int_t^s \sigma(r) dW_{r-t} + v(s-t),$$

$s \in [t, T]$, where $t \in [0, T]$ is an initial time, $x \in \mathbb{R}^n$ is an initial position, $b : [0, T] \rightarrow \mathbb{R}^n$ and $a, \sigma : [0, T] \rightarrow \mathbb{M}^{n \times n}$ stand for the drift and the covariance terms. Note that $(y_{xt}(s))_{s \in [t, T]}$ is a random process adapted to $(\mathcal{F}_{s-t})_{s \in [t, T]}$.

To each control $v \in \mathcal{V}$, we associate a cost given by the payoff functional

$$(2) \quad J_{xt}(v) = \mathbb{E} \left\{ \int_t^T f(y_{xt}(s), s) e^{-\int_t^s \alpha(r) dr} ds + \int_t^T c(s) e^{-\int_t^s \alpha(r) dr} d\xi(s-t) \right\},$$

where f , α and c are respectively the running cost, the discount factor and the instantaneous cost per unit of “fuel”.

Our purpose is to characterize the optimal cost, the so-called value function

$$(3) \quad u(x, t) = \inf \{ J_{xt}(v) : v \in \mathcal{V} \}.$$

It is often convenient to consider the following penalized problem associated with (3):

$$(4) \quad u_\epsilon(x, t) = \inf \{ J_{xt}(v) : v \in \mathcal{V}_\epsilon \},$$

where $\epsilon > 0$ and \mathcal{V}_ϵ is the set of all controls $v \in \mathcal{V}$ which are Lipschitz continuous and $|\frac{dv}{dt}(t)| \leq \frac{1}{\epsilon}$ for almost every $t \in [0, T]$ almost surely.

Definition 1.1. We say that the finite time horizon stochastic control problem has the dynamic programming property in the weak sense if for every $x \in \mathbb{R}^n$, $t, t' \in [0, T]$ s.t. $t < t'$ and $y_{xt}^0(s)$ given by (1) with $v \equiv 0$ we have

$$(5) \quad u(x, t) \leq \mathbb{E} \left\{ \int_t^{t'} f(y_{xt}^0(s), s) e^{-\int_t^s \alpha(r) dr} ds + u(y_{xt}^0(t'), t') e^{-\int_t^{t'} \alpha(r) dr} \right\}.$$

Let us assume the following:

- α, c are Lipschitz continuous from $[0, T]$ into $[0, \infty)$ with constant $L > 0$,
- b is Lipschitz continuous from $[0, T]$ into \mathbb{R}^n with the same constant $L > 0$,
- a, σ are Lipschitz continuous from $[0, T]$ into $\mathbb{M}^{n \times n}$ with the same constant $L > 0$,
- there exists $c_0 > 0$ such that $c(t) \geq c_0$ for all $t \in [0, T]$,
- $f : \mathbb{R}^n \times [0, T] \rightarrow [0, \infty)$ and there exist constants $p > 1, C_0, \tilde{C}_0 > 0$ such that for all $t, t' \in [0, T]$, $x, x' \in \mathbb{R}^n$ and $\lambda \in (0, 1)$ we have

$$(6) \quad \tilde{C}_0 |x|^p - C_0 \leq f(x, t) \leq C_0(1 + |x|^p),$$

$$(7) \quad |f(x, t) - f(x + x', t)| \leq C_0(1 + f(x, t) + f(x + x', t))^{1-1/p} |x'|,$$

$$(8) \quad |f(x, t) - f(x, t')| \leq C_0(1 + |x|^p) |t - t'|,$$

$$(9) \quad \begin{aligned} 0 < f(x + \lambda x', t) - 2f(x, t) + f(x - \lambda x', t) \\ \leq C_0 \lambda^2 (1 + f(x, t))^q, \quad q = (1 - 2/p)^+. \end{aligned}$$

The last assumption implies strict convexity of the function f with respect to x .

Let us denote by c_{\max} and α_{\max} the maximum of the function c , α , respectively. Moreover, by a_{\max} , σ_{\max} , β_{\max} and b_{\max} we denote the maximum over $t \in [0, T]$ of the norms of the matrices $a(t)$, $\sigma(t)$, $\beta(t)$ and the vector $b(t)$ respectively, where $\beta(t) = \sigma(t)\sigma^T(t)$.

Now we give lemmas needed for the proofs of the Theorems 2.1 and 2.2. The first one is well known.

Lemma 1.2. *For all $x, y \geq 0$ we have*

$$\begin{aligned} x^p + y^p &\leq (x + y)^p \leq 2^{p-1}(x^p + y^p), \text{ if } p \geq 1, \\ 2^{p-1}(x^p + y^p) &\leq (x + y)^p \leq x^p + y^p, \text{ if } p \in (0, 1). \end{aligned}$$

Lemma 1.3 (See [10], Corollary 2.5.12). *Consider an n -dimensional process described by a stochastic integral equation*

$$x(t) = x_0 + \int_0^t g(x(s), s)ds + \int_0^t h(x(s), s)dW_s, \quad t \geq 0,$$

where $x_0 \in \mathbb{R}^n$, $g : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^n$ and $h : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{M}^{n \times n}$. We assume that there exists a constant C such that for all $x \in \mathbb{R}^n$ and $t \geq 0$

$$(10) \quad \|h(x, t)\| + |g(x, t)| \leq C(1 + |x|).$$

Then for every $q > 0$ there exists a constant $C_{11} > 0$ depending only on q, C such that for all $t \geq 0$

$$(11) \quad \mathbb{E} \sup_{0 \leq s \leq t} |x(s)|^q \leq C_{11} e^{C_{11}t} (1 + |x_0|)^q.$$

Remark 1.4. For the process y_{xt} defined by (1) with $v \equiv 0$ the assumption (10) holds. Indeed, σ is Lipschitz continuous, independent of x and defined on a finite time interval $[0, T]$, so it is bounded. We conclude the same about a, b , so $|g(x, t)| = |a(t) \cdot x + b(t)| \leq C(1 + |x|)$, where $C = \max\{\|a(t)\|, |b(t)| : t \in [0, T]\}$.

Lemma 1.5. *Let $x, x' \in \mathbb{R}^n$, $t \in [0, T]$ and $g(s) = y_{xt}(s) - y_{x't}(s)$ for $s \in [t, T]$. Then*

$$(12) \quad \frac{dg}{ds}(s) = a(s)g(s), \quad |g(s)| \leq C_{12}|x - x'|, \quad s \in [t, T],$$

where $C_{12} = (1 + a_{\max}Te^{a_{\max}T})$.

Proof. In view of (1) we have

$$g(s) = x - x' + \int_t^s a(r)(y_{xt}(r) - y_{x't}(r))dr = x - x' + \int_t^s a(r)g(r)dr.$$

Taking the derivative d/ds of both sides, we get the differential equation $\frac{dg}{ds}(s) = a(s)g(s)$ with initial data $g(t) = x - x'$. The solution of this problem satisfies

$$|g(s)| \leq |x - x'| + \int_t^s |a(r)g(r)|dr \leq |x - x'| + a_{\max} \int_t^s |g(r)|dr.$$

Using the Gronwall's inequality (see [4], p. 625), we get the second part of (12). \square

Lemma 1.6. *Suppose that for some $x \in \mathbb{R}^n$, $t \in [0, T]$, $v \in \mathcal{V}$ we have*

$$\mathbb{E} \int_t^T f(y_{xt}(s), s) e^{-\int_t^s \alpha(r) dr} ds \leq C(1 + |x|^p)$$

for a suitable constant $C > 0$ independent of x, t . Then

$$(13) \quad \mathbb{E} \int_t^T f(y_{xt}(s), s) ds \leq C_{13}(1 + |x|^p), \text{ where } C_{13} = C \cdot e^{\int_0^T \alpha(r) dr}.$$

Proof. Indeed, multiplying both sides of our assumption by $e^{\int_t^T \alpha(r) dr}$, we get

$$\mathbb{E} \int_t^T f(y_{xt}(s), s) e^{\int_s^T \alpha(r) dr} ds \leq C e^{\int_t^T \alpha(r) dr} (1 + |x|^p) \leq C_{13}(1 + |x|^p).$$

Of course, the left-hand side is not smaller than $\mathbb{E} \int_t^T f(y_{xt}(s), s) ds$. \square

Lemma 1.7 (Compare a statement in [17], p. 181). *The function $J_{xt}(v)$ is convex with respect to (x, v) , more precisely, for all $x_1, x_2 \in \mathbb{R}^n$, $t \in [0, T]$, $v_1, v_2 \in \mathcal{V}$ and $\theta \in [0, 1]$,*

$$J_{\theta x_1 + (1-\theta)x_2, t}(\theta v_1 + (1-\theta)v_2) \leq \theta J_{x_1, t}(v_1) + (1-\theta) J_{x_2, t}(v_2).$$

Proof. First, we note that the set \mathcal{V} is obviously convex. Let $y_{xt}^v(s)$ be the solution of (1) corresponding to a control v . Denote $v_0 = \theta v_1 + (1-\theta)v_2$ and $x_0 = \theta x_1 + (1-\theta)x_2$. In view of the definition of $J_{xt}(v)$, it suffices to prove two following inequalities

$$(14) \quad f(y_{x_0, t}^{v_0}(s), s) \leq \theta f(y_{x_1, t}^{v_1}(s), s) + (1-\theta) f(y_{x_2, t}^{v_2}(s), s), \quad s \in [t, T],$$

$$(15) \quad \int_t^T d\xi_0(s-t) \leq \theta \int_t^T d\xi_1(s-t) + (1-\theta) \int_t^T d\xi_2(s-t),$$

where ξ_0, ξ_1, ξ_2 are the total variations of v_0, v_1, v_2 respectively.

The latter inequality is a consequence of the fact that the variation of the sum of functions is not greater than the sum of their variations. So $\xi_0 \leq \theta \xi_1 + (1-\theta)\xi_2$. Because ξ_0, ξ_1, ξ_2 are nondecreasing and $\xi_0(0) = \xi_1(0) = \xi_2(0) = 0$ P -a.s., we conclude that (15) is true.

To prove (14) we show first that

$$(16) \quad y_{x_0, t}^{v_0}(s) = \theta y_{x_1, t}^{v_1}(s) + (1-\theta) y_{x_2, t}^{v_2}(s).$$

Indeed, using (1) we get

$$y_{x_i, t}^{v_i}(s) = x_i + \int_t^s (a(r) y_{x_i, t}^{v_i}(r) + b(r)) dr + \int_t^s \sigma(r) dW_{r-t} + v_i(s-t),$$

$i = 0, 1, 2$. Let $g(s) = y_{x_0,t}^{v_0}(s) - \theta y_{x_1,t}^{v_1}(s) - (1 - \theta)y_{x_2,t}^{v_2}(s)$. Then

$$g(s) = \int_t^s a(r) \left(y_{x_0,t}^{v_0}(r) - \theta y_{x_1,t}^{v_1}(r) - (1 - \theta)y_{x_2,t}^{v_2}(r) \right) dr = \int_t^s a(r)g(r)dr .$$

Taking the derivative d/ds of both sides, we get the differential equation $\frac{dg}{ds}(s) = a(s)g(s)$ with initial data $g(t) = x_0 - \theta x_1 - (1 - \theta)x_2 = 0$. The solution of this problem is $g(s) \equiv 0$, so (16) holds. Using (16) and convexity of f we have (14). \square

Lemma 1.8. *Suppose that for some $t' \in [0, T]$, $x \in \mathbb{R}^n$, $v \in \mathcal{V}$ we have*

$$\mathbb{E} \int_0^{T-t'} c(t' + s) e^{-\int_0^s \alpha(t'+r)dr} d\xi(s) \leq C(1 + |x|^p)$$

for a suitable constant $C > 0$ independent of x , t' . Then there exists a constant $C_{17} > 0$ independent of x , t' such that

$$(17) \quad \mathbb{E}\xi(T - t') \leq C_{17}(1 + |x|^p).$$

Proof. Indeed, multiplying both sides of our assumption by $e^{\int_0^{T-t'} \alpha(t'+r)dr}$ and using the lower bound of c , we get

$$\begin{aligned} c_0 \mathbb{E}\xi(T - t') &= c_0 \mathbb{E} \int_0^{T-t'} d\xi(s) \leq \mathbb{E} \int_0^{T-t'} c(t' + s) e^{\int_s^{T-t'} \alpha(t'+r)dr} d\xi(s) \\ &\leq C e^{\int_0^T \alpha(r)dr} (1 + |x|^p). \end{aligned} \quad \square$$

Lemma 1.9. *Suppose that for some $x \in \mathbb{R}^n$, $t \in [0, T]$, $v \in \mathcal{V}$ we have*

$$\mathbb{E} \int_0^{T-t} f(y_{xt}(t + s), t + s) e^{-\int_0^s \alpha(t+r)dr} ds \leq C(1 + |x|^p)$$

for a suitable constant $C > 0$ independent of x , t . Then there exists a constant $C_{18} > 0$ independent of x , t such that

$$(18) \quad \mathbb{E} \int_0^{T-t} (1 + |y_{xt}(t + s)|^p) ds \leq C_{18}(1 + |x|^p).$$

Proof. From Lemma 1.6 we know that

$$\mathbb{E} \int_0^{T-t} f(y_{xt}(t + s), t + s) ds \leq C_{13}(1 + |x|^p).$$

Using (6), we get

$$\mathbb{E} \int_0^{T-t} \left(\tilde{C}_0 |y_{xt}(t + s)|^p - C_0 \right) ds \leq C_{13}(1 + |x|^p).$$

Hence

$$\tilde{C}_0 \mathbb{E} \int_0^{T-t} |y_{xt}(t + s)|^p ds \leq (C_{13} + C_0 T)(1 + |x|^p)$$

and finally

$$\tilde{C}_0 \mathbb{E} \int_0^{T-t} (1 + |y_{xt}(t+s)|^p) ds \leq (C_{13} + C_0 T + \tilde{C}_0 T)(1 + |x|^p). \quad \square$$

Lemma 1.10. *Let $0 \leq t' \leq t \leq T$ and suppose that for some $x \in \mathbb{R}^n$, $v \in \mathcal{V}$ we have*

$$\mathbb{E} \int_0^{T-t'} f(y_{xt'}(t'+s), t'+s) e^{-\int_0^s \alpha(t'+r) dr} ds \leq C(1 + |x|^p)$$

for a suitable constant $C > 0$ independent of x , t , t' . Then there exists a constant $C_{19} > 0$ independent of x , t , t' such that

$$(19) \quad \mathbb{E} \int_0^{T-t} f(y_{xt'}(t'+s), t+s) ds \leq C_{19}(1 + |x|^p).$$

Proof. We observe that using (8) we have

$$\begin{aligned} f(y_{xt'}(t'+s), t+s) &\leq |f(y_{xt'}(t'+s), t+s) - f(y_{xt'}(t'+s), t'+s)| \\ &\quad + f(y_{xt'}(t'+s), t'+s) \\ &\leq C_0 |t - t'| (1 + |y_{xt'}(t'+s)|^p) + f(y_{xt'}(t'+s), t'+s). \end{aligned}$$

Hence, in view of Lemma 1.6 and Lemma 1.9, we get

$$\begin{aligned} \mathbb{E} \int_0^{T-t} f(y_{xt'}(t'+s), t+s) ds &\leq C_0 |t - t'| C_{18} (1 + |x|^p) + C_{13} (1 + |x|^p) \\ &\leq C_{19} (1 + |x|^p), \end{aligned}$$

where $C_{19} = C_0 T C_{18} + C_{13}$. \square

Lemma 1.11. *Let $0 \leq t' \leq t \leq T$, $x \in \mathbb{R}^n$, $v \in \mathcal{V}$. Assume that*

$$\mathbb{E} \int_0^{T-t'} f(y_{xt'}(t'+s), t'+s) e^{-\int_0^s \alpha(t'+r) dr} ds \leq C(1 + |x|^p)$$

for a suitable constant $C > 0$ independent of x , t' , t . Then there exists a constant $C_{20} > 0$ independent of x , t' , t such that for all $s \in [0, T - t]$ we have

$$(20) \quad \mathbb{E} |y_{xt'}(t'+s) - y_{xt}(t+s)|^p \leq C_{20} |t - t'|^p (1 + |x|^p).$$

Proof. For $s \in [0, T - t]$, we have

$$y_{xt}(t+s) = x + \int_0^s (a(t+r)y_{xt}(t+r) + b(t+r)) dr + \int_0^s \sigma(t+r) dW_r + v(s),$$

$$y_{xt'}(t'+s) = x + \int_0^s (a(t'+r)y_{xt'}(t'+r) + b(t'+r)) dr + \int_0^s \sigma(t'+r) dW_r + v(s),$$

so

$$(21) \quad y_{xt'}(t'+s) - y_{xt}(t+s) = A_s + B_s + M_s,$$

where

$$\begin{aligned} A_s &= \int_0^s (a(t' + r)y_{xt'}(t' + r) - a(t + r)y_{xt}(t + r))dr = A_s^1 + A_s^2, \\ A_s^1 &= \int_0^s a(t + r)(y_{xt'}(t' + r) - y_{xt}(t + r))dr, \\ A_s^2 &= \int_0^s (a(t' + r) - a(t + r))y_{xt'}(t' + r)dr, \\ B_s &= \int_0^s (b(t' + r) - b(t + r))dr, \\ M_s &= \int_0^s (\sigma(t' + r) - \sigma(t + r))dW_r. \end{aligned}$$

Recall that a, b, σ are Lipschitz continuous with the constant L . The process M_s is a martingale with quadratic variation

$$[M]_s = \int_0^s (\sigma(t' + r) - \sigma(t + r))^2 dr \leq L^2 |t - t'|^2 s.$$

This, together with the Burkholder–Davis–Gundy inequalities (see [7], Theorem 3.3.28), implies the existence of a constant C_p , depending only on p , such that

$$(22) \quad \mathbb{E} \sup_{0 \leq s \leq T-t} |M_s|^p \leq C_p L^p T^{\frac{p}{2}} |t - t'|^p.$$

Clearly,

$$(23) \quad \sup_{0 \leq s \leq T-t} |B_s| \leq LT |t - t'|.$$

By the Hölder's inequality, for $q = p/(p - 1)$ we have

$$(24) \quad |A_s^1|^p \leq (a_{\max}^q s)^{\frac{p}{q}} \int_0^s |y_{xt'}(t' + r) - y_{xt}(t + r)|^p dr,$$

$$(25) \quad |A_s^2|^p \leq ((L|t - t'|)^q s)^{\frac{p}{q}} \int_0^s |y_{xt'}(t' + r)|^p dr.$$

By Lemma 1.9, the inequality (18) holds for every $t \in [0, T]$. Lemma 1.2 and the relations (18), (21)–(25) imply that the random variable

$$\sup_{0 \leq s \leq T-t} |y_{xt'}(t' + s) - y_{xt}(t + s)|^p$$

is integrable and hence, by the Lebesgue dominated convergence theorem, the function $F(s) = \mathbb{E}|y_{xt'}(t' + s) - y_{xt}(t + s)|^p$ is continuous on $[0, T - t]$. From Lemma 1.2 and (18), (21)–(25) we also have, for each $s \in [0, T - t]$,

$$F(s) \leq c_1 |t - t'|^p (1 + |x|^p) + c_2 \int_0^s F(r) dr,$$

where $c_1 = 2^{2p-2}(L^p T^p + C_p L^p T^{\frac{p}{2}} + (LT)^{\frac{p}{q}} C_{18})$, $c_2 = 2^{2p-2} a_{\max}^p T^{\frac{p}{q}}$. This, together with the Gronwall's inequality (see, e.g., [7], Problem 5.2.7), implies that for all $s \in [0, T - t]$,

$$F(s) \leq c_1 |t - t'|^p (1 + |x|^p) \left(1 + c_2 \int_0^s e^{c_2(s-r)} dr \right).$$

We have obtained (20) with $C_{20} = c_1(1 + c_2 \int_0^T e^{c_2(T-r)} dr)$. \square

Lemma 1.12. *Suppose that for some $x \in \mathbb{R}^n$, $t \in [0, T]$, $v \in \mathcal{V}$ we have*

$$\mathbb{E} \int_t^T f(y_{xt}(s), s) e^{-\int_t^s \alpha(r) dr} ds \leq C(1 + |x|^p)$$

for a suitable constant $C > 0$ independent of x, t . Then there exists a constant $C_{26} > 0$ independent of x, x', t such that for every $x' \in \mathbb{R}^n$,

$$(26) \quad \mathbb{E} \int_t^T f(y_{x+x',t}(s), s) ds \leq C_{26}(1 + |x|^p + |x + x'|^p).$$

Proof. From (6) and Lemma 1.2 we have

$$\begin{aligned} \mathbb{E} \int_t^T f(y_{x+x',t}(s), s) ds &\leq \mathbb{E} \int_t^T C_0(1 + |y_{x+x',t}(s)|^p) ds \\ &\leq TC_0 + C_0 2^{p-1} \mathbb{E} \int_t^T |y_{x+x',t}(s) - y_{x,t}(s)|^p ds + C_0 2^{p-1} \mathbb{E} \int_t^T |y_{x,t}(s)|^p ds. \end{aligned}$$

Now using Lemma 1.5, Lemma 1.9 and Lemma 1.2 again, we get

$$\begin{aligned} \mathbb{E} \int_t^T f(y_{x+x',t}(s), s) ds &\leq TC_0 + C_0 2^{p-1} T \cdot C_{12}^p |x'|^p + C_0 2^{p-1} C_{18}(1 + |x|^p) \\ &\leq TC_0 + C_0 2^{2p-2} T \cdot C_{12}^p (|x' + x|^p + |x|^p) + C_0 2^{p-1} C_{18}(1 + |x|^p) \\ &\leq C_{26}(1 + |x|^p + |x + x'|^p), \end{aligned}$$

where $C_{26} = C_0(T + 2^{2p-2} T \cdot C_{12}^p + 2^{p-1} C_{18})$. \square

Lemma 1.13. *Suppose that for some $x \in \mathbb{R}^n$, $t' \in [0, T]$, $v \in \mathcal{V}$ we have*

$$\mathbb{E} \int_0^{T-t'} f(y_{xt'}(t' + s), t' + s) e^{-\int_0^s \alpha(t'+r) dr} ds \leq C(1 + |x|^p)$$

for a suitable constant $C > 0$ independent of x, t' . Then there exists a constant $C_{27} > 0$ independent of x, t', t such that for every $t \in [t', T]$,

$$(27) \quad \mathbb{E} \int_0^{T-t} f(y_{xt}(t + s), t + s) ds \leq C_{27}(1 + |x|^p).$$

Proof. Using (6) and Lemma 1.2, we have

$$\begin{aligned} \mathbb{E} \int_0^{T-t} f(y_{xt}(t+s), t+s) ds &\leq \mathbb{E} \int_0^{T-t} C_0(1 + |y_{xt}(t+s)|^p) ds \\ &\leq C_0 T + 2^{p-1} C_0 \mathbb{E} \int_0^{T-t} |y_{xt'}(t'+s)|^p ds \\ &\quad + 2^{p-1} C_0 \mathbb{E} \int_0^{T-t} |y_{xt}(t+s) - y_{xt'}(t'+s)|^p ds . \end{aligned}$$

In view of Lemma 1.9, the Fubini's theorem and Lemma 1.11, we get

$$\begin{aligned} \mathbb{E} \int_0^{T-t} f(y_{xt}(t+s), t+s) ds \\ \leq C_0 T + 2^{p-1} C_0 C_{18}(1 + |x|^p) + 2^{p-1} C_0 T C_{20} |t-t'|^p (1 + |x|^p) \\ \leq C_{27}(1 + |x|^p), \end{aligned}$$

where $C_{27} = C_0(T + 2^{p-1}C_{18} + 2^{p-1}T^{p+1}C_{20})$. \square

The next two definitions and lemma refer to mollification of a given function (see [4], p. 629–630).

Definition 1.14. Define $\eta \in C^\infty(\mathbb{R}^n)$ by

$$(28) \quad \eta(x) = \begin{cases} C_{28} \exp\left(\frac{1}{|x|^2-1}\right) & \text{if } |x| < 1 \\ 0 & \text{if } |x| \geq 1, \end{cases}$$

where the constant C_{28} is selected so that $\int_{\mathbb{R}^n} \eta(x) dx = 1$. For each $m \in \mathbb{N}$ set $\eta_m(x) = m^n \cdot \eta(mx)$. We call η the standard mollifier. The functions η_m belong to the class $C^\infty(\mathbb{R}^n)$ and satisfy $\int_{\mathbb{R}^n} \eta_m(x) dx = 1$.

Definition 1.15. Fix $t' \in [0, T]$. For each $m \in \mathbb{N}$ we define mollification of the function $u(\cdot, t')$ by

$$u_m(x) = \int_{B(0, \frac{1}{m})} \eta_m(y) u(x-y, t') dy, \quad x \in \mathbb{R}^n,$$

where $B(0, r) = \{x \in \mathbb{R}^n : |x| < r\}$.

Lemma 1.16. For each $m \in \mathbb{N}$ we have $u_m \in C^\infty(\mathbb{R}^n)$. Moreover, if $u(\cdot, t')$ is continuous, then $u_m(x) \rightarrow u(x, t')$ uniformly on compact subsets of \mathbb{R}^n as $m \rightarrow \infty$.

2. Estimates for the value function. Let the assumptions from Section 1 appearing immediately after Definition 1.1 hold.

Theorem 2.1. Let u be the value function defined by (3). Then for some positive constants C_{29}, C_{30}, C_{31} , the same $p > 1$ as in the assumptions (6)–(9) and every $t \in [0, T]$, $x, x' \in \mathbb{R}^n$ and $\lambda \in (0, 1)$, the following estimates

hold:

$$(29) \quad 0 \leq u(x, t) \leq C_{29}(1 + |x|^p),$$

$$(30) \quad |u(x, t) - u(x + x', t)| \leq C_{30}(1 + |x|^{p-1} + |x + x'|^{p-1})|x'|,$$

$$(31) \quad 0 \leq u(x + \lambda x', t) - 2u(x, t) + u(x - \lambda x', t) \leq C_{31}\lambda^2(1 + |x|)^{(p-2)^+}.$$

Proof: Proof of (29). Nonnegativity of u is the consequence of nonnegativity of f and c . Next, taking the control $v \equiv 0$ and using (6), the Fubini's theorem, Lemma 1.3 and Lemma 1.2, we get

$$\begin{aligned} u(x, t) &\leq J_{xt}(0) = \mathbb{E} \int_t^T f(y_{xt}^0(s), s) e^{-\int_t^s \alpha(r) dr} ds \\ &\leq \mathbb{E} \int_t^T C_0(1 + |y_{xt}^0(s)|^p) ds = C_0 \int_t^T \mathbb{E}(1 + |y_{xt}^0(s)|^p) ds \\ &\leq C_0 \int_t^T \mathbb{E} \left(1 + C_{11} e^{C_{11}(s-t)} (1 + |x|)^p \right) ds \\ &\leq C_0 \int_0^T (1 + C_{11} e^{C_{11}T} 2^{p-1}) (1 + |x|^p) ds \\ &= C_0 T (1 + C_{11} e^{C_{11}T} 2^{p-1}) (1 + |x|^p) \\ &= C_{29}(1 + |x|^p), \end{aligned}$$

where C_{29} depends only on C_0, T, C_{11}, p , so (29) is proved. \square

Proof of (30). Now we note that

$$\begin{aligned} u(x + x', t) - u(x, t) &= \inf_{v' \in \mathcal{V}} \sup_{v \in \mathcal{V}} \left(J_{x+x',t}(v') - J_{x,t}(v) \right) \\ &\leq \sup_{v \in \mathcal{V}} \left(J_{x+x',t}(v) - J_{x,t}(v) \right). \end{aligned}$$

Hence

$$\begin{aligned} u(x + x', t) - u(x, t) &\leq \sup_{v \in \mathcal{V}} |J_{xt}(v) - J_{x+x',t}(v)| \\ &\leq \sup_{v \in \mathcal{V}} \mathbb{E} \int_t^T |f(y_{xt}(s), s) - f(y_{x+x',t}(s), s)| e^{-\int_t^s \alpha(r) dr} ds. \end{aligned}$$

Applying (7), we can estimate the last expression from above by

$$\sup_{v \in \mathcal{V}} \mathbb{E} \int_t^T C_0(1 + f(y_{xt}(s), s) + f(y_{x+x',t}(s), s))^{1-1/p} \cdot |y_{xt}(s) - y_{x+x',t}(s)| ds.$$

Using Lemma 1.5, we have

$$\begin{aligned} u(x + x', t) - u(x, t) &\leq \sup_{v \in \mathcal{V}} C_0 C_{12} |x'| \cdot \mathbb{E} \int_t^T (1 + f(y_{xt}(s), s) + f(y_{x+x',t}(s), s))^{\frac{p-1}{p}} ds. \end{aligned}$$

We use the Hölder's inequality with exponent $\frac{p}{p-1}$ to estimate the last expression above by

$$(32) \quad \sup_{v \in \mathcal{V}} C_0 C_{12} |x'| \cdot \left(\mathbb{E} \int_t^T (1 + f(y_{xt}(s), s) + f(y_{x+x',t}(s), s)) ds \right)^{\frac{p-1}{p}} T^{\frac{1}{p}}.$$

By virtue of (29) we can consider only those controls v for which

$$\mathbb{E} \int_t^T f(y_{xt}(s), s) e^{-\int_t^s \alpha(r) dr} ds \leq (C_{29} + \epsilon)(1 + |x|^p)$$

for some arbitrary $\epsilon > 0$. From (32), Lemma 1.6 and Lemma 1.12 we see that

$$\begin{aligned} u(x + x', t) - u(x, t) &\leq C_0 C_{12} |x'| \left(T + C_{13}(1 + |x|^p) \right. \\ &\quad \left. + C_{26}(1 + |x|^p + |x + x'|^p) \right)^{\frac{p-1}{p}} T^{\frac{1}{p}} \\ &\leq C_{30} |x'| (1 + |x|^p + |x + x'|^p)^{\frac{p-1}{p}}, \end{aligned}$$

where $C_{30} = T^{1/p} \cdot C_0 C_{12} (T + C_{13} + C_{26})^{1-1/p}$. Finally using Lemma 1.2, we get

$$u(x + x', t) - u(x, t) \leq C_{30}(1 + |x|^{p-1} + |x + x'|^{p-1})|x'|.$$

In an analogous manner we get the same estimate for $u(x, t) - u(x + x', t)$. \square

Proof of (31). We observe that

$$\begin{aligned} &u(x + \lambda x', t) + u(x - \lambda x', t) - 2u(x, t) \\ &\leq \sup_{v \in \mathcal{V}} (J_{x+\lambda x',t}(v) + J_{x-\lambda x',t}(v) - 2J_{x,t}(v)) \\ &= \sup_{v \in \mathcal{V}} \mathbb{E} \int_t^T (f(y_{x+\lambda x',t}(s), s) + f(y_{x-\lambda x',t}(s), s) - 2f(y_{x,t}(s), s)) e^{-\int_t^s \alpha(r) dr} ds. \end{aligned}$$

In view of (12) we can apply (9) to get

$$\begin{aligned} &u(x + \lambda x', t) + u(x - \lambda x', t) - 2u(x, t) \\ &\leq \sup_{v \in \mathcal{V}} \mathbb{E} \int_t^T C_0 \lambda^2 (1 + f(y_{xt}(s), s))^{(1-2/p)^+} ds. \end{aligned}$$

If $p \leq 2$ we have $u(x + \lambda x', t) + u(x - \lambda x', t) - 2u(x, t) \leq C_0 T \lambda^2$. If $p > 2$ we use the Hölder inequality with exponent $\frac{p}{p-2}$ to get

$$\begin{aligned} &u(x + \lambda x', t) + u(x - \lambda x', t) - 2u(x, t) \\ &\leq \sup_{v \in \mathcal{V}} C_0 \lambda^2 \left(\mathbb{E} \int_t^T (1 + f(y_{xt}(s), s)) ds \right)^{1-2/p} T^{2/p}. \end{aligned}$$

By virtue of (29) we can consider only those controls v for which

$$\mathbb{E} \int_t^T f(y_{xt}(s), s) e^{-\int_t^s \alpha(r) dr} ds \leq (C_{29} + \epsilon)(1 + |x|^p)$$

for some arbitrary $\epsilon > 0$. From Lemma 1.6 and Lemma 1.2 we see that

$$\begin{aligned} u(x + \lambda x', t) + u(x - \lambda x', t) - 2u(x, t) \\ \leq C_0 \lambda^2 \left(T + C_{13}(1 + |x|^p) \right)^{1-2/p} T^{2/p} \\ \leq C_{31} \lambda^2 (1 + |x|^p)^{1-2/p} \\ \leq C_{31} \lambda^2 (1 + |x|)^{p-2}, \end{aligned}$$

where $C_{31} = T^{2/p} C_0 (T + C_{13})^{1-2/p}$. We have proved the upper bound of (31).

To prove the lower bound of (31), it clearly suffices to prove convexity of $u(x, t)$ with respect to the first variable. In view of the definition of u we know that for every $\epsilon > 0$, $x_1, x_2 \in \mathbb{R}^n$, $t \in [0, T]$, $\theta \in [0, 1]$ there exist $v_1, v_2 \in \mathcal{V}$ such that $J_{x_i, t}(v_i) \leq u(x_i, t) + \epsilon$, $i = 1, 2$.

Using Lemma 1.7, we get

$$\begin{aligned} u(\theta x_1 + (1 - \theta)x_2, t) &\leq J_{\theta x_1 + (1 - \theta)x_2, t}(\theta v_1 + (1 - \theta)v_2) \\ &\leq \theta J_{x_1, t}(v_1) + (1 - \theta) J_{x_2, t}(v_2) \\ &\leq \theta u(x_1, t) + (1 - \theta) u(x_2, t) + \epsilon. \end{aligned}$$

Because $\epsilon > 0$ is arbitrary, we get convexity of $u(x, t)$ with respect to the first variable. \square

Theorem 2.2. *Let the assumptions of Theorem 2.1 be satisfied. Assume that the dynamic programming property in the weak sense holds (Definition 1.1). Then for some constant $C_{33} > 0$ and every $t, t' \in [0, T]$, $x \in \mathbb{R}^n$, we have*

$$(33) \quad |u(x, t) - u(x, t')| \leq C_{33}(1 + |x|^p)|t - t'|.$$

Proof. We note that

$$\begin{aligned} u(x, t) - u(x, t') &= \inf_{v \in \mathcal{V}} \sup_{v' \in \mathcal{V}} \left(J_{xt}(v) - J_{xt'}(v') \right) \\ &\leq \sup_{v' \in \mathcal{V}} \left(J_{xt}(v') - J_{xt'}(v') \right). \end{aligned}$$

For $t' \leq t$ the difference $J_{xt}(v) - J_{xt'}(v)$ is equal to

$$\begin{aligned} \mathbb{E} \left\{ \int_0^{T-t} \left(f(y_{xt}(t+s), t+s) e^{-\int_0^s \alpha(t+r) dr} \right. \right. \\ \left. \left. - f(y_{xt'}(t'+s), t'+s) e^{-\int_0^s \alpha(t'+r) dr} \right) ds \right. \\ \left. + \int_0^{T-t} \left(c(t+s) e^{-\int_0^s \alpha(t+r) dr} - c(t'+s) e^{-\int_0^s \alpha(t'+r) dr} \right) d\xi(s) \right. \\ \left. - \int_{T-t}^{T-t'} f(y_{xt'}(t'+s), t'+s) e^{-\int_0^s \alpha(t'+r) dr} ds \right. \\ \left. - \int_{T-t}^{T-t'} c(t'+s) e^{-\int_0^s \alpha(t'+r) dr} d\xi(s) \right\}. \end{aligned}$$

Let us denote the expectations of the first two integrals in the last expression by A and B , respectively. Because the last two integrals are nonnegative we get

$$(34) \quad J_{xt}(v) - J_{xt'}(v) \leq A + B.$$

We can estimate B as follows:

$$B \leq \mathbb{E} \int_0^{T-t} \left| c(t+s) e^{-\int_0^s \alpha(t+r) dr} - c(t'+s) e^{-\int_0^s \alpha(t'+r) dr} \right| d\xi(s).$$

Adding and subtracting $c(t+s) e^{-\int_0^s \alpha(t'+r) dr}$ under the absolute value sign and using the triangle inequality and positivity of α , we get

$$B \leq \mathbb{E} \int_0^{T-t} \left(c_{\max} \left| e^{-\int_0^s \alpha(t+r) dr} - e^{-\int_0^s \alpha(t'+r) dr} \right| + |c(t+s) - c(t'+s)| \right) d\xi(s).$$

Because $|e^x - e^y| \leq |x - y|$ for $x, y \leq 0$ and c, α are Lipschitz continuous, we have

$$\begin{aligned} B &\leq \mathbb{E} \int_0^{T-t} \left(c_{\max} \int_0^s |\alpha(t+r) - \alpha(t'+r)| dr + |c(t+s) - c(t'+s)| \right) d\xi(s) \\ &\leq (c_{\max} T + 1)L|t - t'| \mathbb{E} \int_0^{T-t} d\xi(s) = (c_{\max} T + 1)L|t - t'| \mathbb{E} \xi(T - t). \end{aligned}$$

By virtue of (29) we can consider only those controls v for which

$$\mathbb{E} \int_0^{T-t'} c(t'+s) e^{-\int_0^s \alpha(t'+r) dr} d\xi(s) \leq (C_{29} + \epsilon)(1 + |x|^p)$$

for some arbitrary $\epsilon > 0$. Using Lemma 1.8, we get $\mathbb{E} \xi(T - t) \leq \mathbb{E} \xi(T - t') \leq C_{17}(1 + |x|^p)$ and

$$(35) \quad B \leq C_{35} |t - t'| (1 + |x|^p), \text{ where } C_{35} = (c_{\max} T + 1)LC_{17}.$$

Now we estimate A :

$$\begin{aligned}
A &\leq \mathbb{E} \int_0^{T-t} \left| f(y_{xt}(t+s), t+s) e^{-\int_0^s \alpha(t+r) dr} \right. \\
&\quad \left. - f(y_{xt}(t+s), t+s) e^{-\int_0^s \alpha(t'+r) dr} \right| ds \\
&\quad + \mathbb{E} \int_0^{T-t} \left| f(y_{xt}(t+s), t+s) e^{-\int_0^s \alpha(t'+r) dr} \right. \\
&\quad \left. - f(y_{xt'}(t'+s), t'+s) e^{-\int_0^s \alpha(t'+r) dr} \right| ds \\
&= A_1 + A_2 .
\end{aligned}$$

Using the inequality $|e^x - e^y| \leq |x - y|$ for $x, y \leq 0$ again, we get

$$\begin{aligned}
(36) \quad A_1 &\leq \mathbb{E} \int_0^{T-t} f(y_{xt}(t+s), t+s) \left(\int_0^s |\alpha(t+r) - \alpha(t'+r)| dr \right) ds \\
&\leq TL|t - t'| \mathbb{E} \int_0^{T-t} f(y_{xt}(t+s), t+s) ds .
\end{aligned}$$

By virtue of (29) we can consider only those controls v for which

$$(37) \quad \mathbb{E} \int_0^{T-t'} f(y_{xt'}(t'+s), t'+s) e^{-\int_0^s \alpha(t'+r) dr} ds \leq (C_{29} + \epsilon)(1 + |x|^p)$$

for some arbitrary $\epsilon > 0$. Using (36) and Lemma 1.13, we get

$$(38) \quad A_1 \leq C_{38} |t - t'| (1 + |x|^p) , \text{ where } C_{38} = TLC_{27} .$$

To estimate A_2 we use (7)–(8) and we have that A_2 is less than or equal to

$$\begin{aligned}
&\mathbb{E} \int_0^{T-t} \left| f(y_{xt}(t+s), t+s) - f(y_{xt'}(t'+s), t'+s) \right. \\
&\quad \left. + f(y_{xt'}(t'+s), t'+s) - f(y_{xt'}(t'+s), t'+s) \right| ds \\
&\leq \mathbb{E} \int_0^{T-t} C_0 \left(1 + f(y_{xt}(t+s), t+s) + f(y_{xt'}(t'+s), t'+s) \right)^{1-1/p} \\
&\quad \times |y_{xt'}(t'+s) - y_{xt}(t+s)| ds \\
&\quad + \mathbb{E} \int_0^{T-t} C_0 (1 + |y_{xt'}(t'+s)|^p) |t - t'| ds = A_3 + A_4 .
\end{aligned}$$

Using the Hölder's inequality and the Fubini's theorem, we get

$$\begin{aligned}
A_3 &\leq C_0 \left\{ \mathbb{E} \int_0^{T-t} \left(1 + f(y_{xt}(t+s), t+s) + f(y_{xt'}(t'+s), t'+s) \right) ds \right\}^{1-1/p} \\
&\quad \times \left\{ \int_0^{T-t} \mathbb{E} |y_{xt'}(t'+s) - y_{xt}(t+s)|^p ds \right\}^{1/p} .
\end{aligned}$$

From this together with (37), Lemma 1.13, Lemma 1.10 and Lemma 1.11 we have

$$A_3 \leq C_0 \{(T + C_{27} + C_{19})(1 + |x|^p)\}^{1-1/p} \cdot \left\{ TC_{20} |t - t'|^p (1 + |x|^p) \right\}^{1/p}.$$

Because $1 + |x|^p \leq (1 + |x|)^p$, we get

$$A_3 \leq C_0 (T + C_{27} + C_{19})^{1-1/p} (1 + |x|)^{p-1} \cdot (TC_{20})^{1/p} |t - t'| (1 + |x|).$$

Hence, from Lemma 1.2,

$$(39) \quad A_3 \leq C_{39} |t - t'| (1 + |x|^p),$$

where $C_{39} = C_0 (T + C_{27} + C_{19})^{1-1/p} (TC_{20})^{1/p} 2^{p-1}$. Furthermore, from Lemma 1.9 we get

$$(40) \quad A_4 \leq C_{40} |t - t'| (1 + |x|^p), \text{ where } C_{40} = C_0 C_{18}.$$

In view of (34)–(35) and (38)–(40) we get for $t' \leq t$,

$$(41) \quad u(x, t) - u(x, t') \leq C_{41} |t - t'| (1 + |x|^p),$$

where $C_{41} = C_{35} + C_{38} + C_{39} + C_{40}$.

To obtain a similar inequality for $t < t'$ we proceed as follows. Let $(y_{xt}^0(s))_{s \in [t, T]}$ be a solution of (1) with $v \equiv 0$. We can write the i -th coordinate of $y_{xt}^0(s)$ as follows

$$(42) \quad y_{xt}^0(s)_i = x_i + \int_t^s \left(\sum_{j=1}^n a_{ij}(r) y_{xt}^0(r)_j + b_i(r) \right) dr + \sum_{j=1}^n \int_t^s \sigma_{ij}(r) dW_{r-t}^j,$$

$i = 1, \dots, n$, where subscripts denote the corresponding coordinates. Let $\{u_m(\cdot)\}_{m \in \mathbb{N}}$ be a sequence of mollifications of the function $u(\cdot, t')$ (see Def. 1.15). Applying the Itô's formula ([7], Th. 3.3.6), we get

$$\begin{aligned} & \mathbb{E} u_m(y_{xt}^0(t')) \\ &= u_m(x) + \mathbb{E} \sum_{i=1}^n \int_t^{t'} \frac{\partial u_m(y_{xt}^0(s))}{\partial x_i} \left(\sum_{j=1}^n a_{ij}(s) y_{xt}^0(s)_j + b_i(s) \right) ds \\ (43) \quad &+ \mathbb{E} \sum_{i=1}^n \int_t^{t'} \frac{\partial u_m(y_{xt}^0(s))}{\partial x_i} \sum_{j=1}^n \sigma_{ij}(s) dW_{s-t}^j \\ &+ \frac{1}{2} \mathbb{E} \sum_{i,j=1}^n \int_t^{t'} \frac{\partial^2 u_m(y_{xt}^0(s))}{\partial x_i \partial x_j} d[y_{xt}^0(s)_i, y_{xt}^0(s)_j] \\ &= u_m(x) + \mathbb{A} + \mathbb{B} + \mathbb{C}. \end{aligned}$$

We need the following lemma.

Lemma 2.3. *We assume (29)–(31). Let $t' \in [0, T]$ be fixed. Then there exist constants $C_{45}, C_{46} > 0$ such that for all $x \in \mathbb{R}^n$, $m \in \mathbb{N}$ and $i, j \in \{1, \dots, n\}$,*

$$(44) \quad \lim_{m \rightarrow \infty} u_m(x) = u(x, t'),$$

$$(45) \quad \left| \frac{\partial u_m(x)}{\partial x_i} \right| \leq C_{45}(1 + |x|)^{p-1},$$

$$(46) \quad 0 \leq \frac{\partial^2 u_m(x)}{\partial x_i \partial x_j} \leq C_{46}(1 + |x|^p).$$

We estimate \mathbb{A} as follows

$$\begin{aligned} \mathbb{A} &\leq \mathbb{E} \sum_{i=1}^n \int_t^{t'} \left| \frac{\partial u_m(y_{xt}^0(s))}{\partial x_i} \right| \cdot \left| \sum_{j=1}^n a_{ij}(s) y_{xt}^0(s)_j + b_i(s) \right| ds \\ &\leq \mathbb{E} \sum_{i=1}^n \int_t^{t'} \left| \frac{\partial u_m(y_{xt}^0(s))}{\partial x_i} \right| \left(n \|a(s)\| \cdot |y_{xt}^0(s)| + |b(s)| \right) ds. \end{aligned}$$

Using Lemma 2.3 and Lemma 1.2, we see that \mathbb{A} is not greater than

$$(47) \quad \begin{aligned} &\sum_{i=1}^n \mathbb{E} \int_t^{t'} C_{45}(1 + |y_{xt}^0(s)|)^{p-1} n(a_{\max} + b_{\max})(1 + |y_{xt}^0(s)|) ds \\ &\leq C_{47} \mathbb{E} \int_t^{t'} (1 + |y_{xt}^0(s)|^p) ds, \end{aligned}$$

where $C_{47} = n^2 C_{45} (a_{\max} + b_{\max}) 2^{p-1}$.

Now we show that $\mathbb{B} = 0$. Indeed,

$$\mathbb{B} = \mathbb{E} \sum_{i,j=1}^n Z_{ij}(t'), \text{ where } Z_{ij}(s) = \int_t^s \frac{\partial u_m(y_{xt}^0(r))}{\partial x_i} \sigma_{ij}(r) dW_{r-t}^j \text{ for } s \in [t, t'].$$

From properties of the Itô's integrals (see [7], Section 3.2) the process $(Z_{ij}(s))_{s \in [t, t']}$ is a martingale provided that

$$\mathbb{E} \int_t^{t'} \left(\frac{\partial u_m(y_{xt}^0(s))}{\partial x_i} \sigma_{ij}(s) \right)^2 ds \leq \infty.$$

Using Lemma 2.3 and Lemma 1.2, we have

$$\begin{aligned} &\mathbb{E} \int_t^{t'} \left(\frac{\partial u_m(y_{xt}^0(s))}{\partial x_i} \sigma_{ij}(s) \right)^2 ds \leq \mathbb{E} \int_t^{t'} C_{45}^2 (1 + |y_{xt}^0(s)|)^{2p-2} \sigma_{\max}^2 ds \\ &\leq C_{45}^2 \sigma_{\max}^2 \mathbb{E} \int_t^{t'} (1 + |y_{xt}^0(s)|)^{2p} ds \\ &\leq C_{45}^2 \sigma_{\max}^2 2^{2p-1} \mathbb{E} \int_t^{t'} (1 + |y_{xt}^0(s)|^p) ds. \end{aligned}$$

Using the Fubini's theorem and Lemma 1.3, we get

$$\begin{aligned} & \mathbb{E} \int_t^{t'} \left(\frac{\partial u_m(y_{xt}^0(s))}{\partial x_i} \sigma_{ij}(s) \right)^2 ds \\ & \leq C_{45}^2 \sigma_{\max}^2 2^{2p-1} \int_t^{t'} \left(1 + C_{11} e^{C_{11}T} (1 + |x|)^{2p} \right) ds < \infty. \end{aligned}$$

Hence Z_{ij} is a martingale and $\mathbb{E}Z_{ij}(t') = \mathbb{E}Z_{ij}(t) = 0$. So

$$(48) \quad \mathbb{B} = \mathbb{E} \sum_{i,j=1}^n Z_{ij}(t') = 0.$$

Now, using the conventional “multiplication rules” (see [7], p. 154), we know that

$$ds ds = 0, \quad ds dW_s^i = 0, \quad dW_s^i dW_s^i = ds, \quad dW_s^i dW_s^j = 0 \quad \text{for } i \neq j.$$

So in view of (42) we can write

$$d[y_{xt}^0(s)_i, y_{xt}^0(s)_j] = \sum_{k=1}^n \sigma_{ik}(s) dW_{s-t}^k \cdot \sum_{l=1}^n \sigma_{jl}(s) dW_{s-t}^l = \sum_{k=1}^n \sigma_{ik}(s) \sigma_{jk}(s) ds.$$

From Lemma 2.3 we have

$$\begin{aligned} \mathbb{C} &= \frac{1}{2} \sum_{i,j=1}^n \mathbb{E} \int_t^{t'} \frac{\partial^2 u_m(y_{xt}^0(s))}{\partial x_i \partial x_j} \sum_{k=1}^n \sigma_{ik}(s) \sigma_{jk}(s) ds \\ (49) \quad &\leq \frac{1}{2} \sum_{i,j=1}^n \mathbb{E} \int_t^{t'} C_{46} (1 + |y_{xt}^0(s)|^p) n \sigma_{\max}^2 ds \\ &= C_{49} \mathbb{E} \int_t^{t'} (1 + |y_{xt}^0(s)|^p) ds, \quad \text{where } C_{49} = \frac{1}{2} C_{46} \sigma_{\max}^2 n^3. \end{aligned}$$

In summary, in view of (43) and (47)–(49)

$$\mathbb{E} u_m(y_{xt}^0(t')) \leq u_m(x) + (C_{47} + C_{49}) \mathbb{E} \int_t^{t'} (1 + |y_{xt}^0(s)|^p) ds.$$

Taking the limit as $n \rightarrow \infty$ and using the Fatou's lemma, we get

$$(50) \quad \mathbb{E} u(y_{xt}^0(t'), t') \leq u(x, t') + C_{50} \mathbb{E} \int_t^{t'} (1 + |y_{xt}^0(s)|^p) ds,$$

$C_{50} = C_{47} + C_{49}$. Furthermore, from Lemma 1.2 and Lemma 1.3 we have for each $s \in [t, T]$

$$(51) \quad \mathbb{E}(1 + |y_{xt}^0(s)|^p) \leq C_{51} (1 + |x|^p), \quad C_{51} = C_{11} e^{C_{11}T} 2^{p-1} + 1.$$

Next, from (5), (6), (50), (51) and the Fubini's theorem we conclude

$$\begin{aligned} u(x, t) &\leq \mathbb{E} \left\{ \int_t^{t'} f(y_{xt}^0(s), s) e^{-\int_t^s \alpha(r) dr} ds + u(y_{xt}^0(t'), t') e^{-\int_t^{t'} \alpha(r) dr} \right\} \\ &\leq C_0 \int_t^{t'} \mathbb{E}(1 + |y_{xt}^0(s)|^p) ds + \mathbb{E}u(y_{xt}^0(t'), t') \\ &\leq (C_0 C_{51} + C_{50} C_{51}) (|t - t'| (1 + |x|^p)) + u(x, t'). \end{aligned}$$

Hence, for $t < t'$

$$(52) \quad u(x, t) - u(x, t') \leq C_{52} |t - t'| (1 + |x|^p), \quad C_{52} = C_{51} (C_0 + C_{50}).$$

It is clear that (41) and (52) imply (33). \square

Now we give the proof of Lemma 2.3.

Proof: Proof of (44). The continuity of $u(\cdot, t')$ is a consequence of (30). So in view of Lemma 1.16 we conclude that $\lim_{m \rightarrow \infty} u_m(x) = u(x, t')$. \square

Proof of (45). Let $x \in \mathbb{R}^n$, $0 \leq |x'| < 1$. From Definitions 1.14, 1.15 and (30) we get

$$\begin{aligned} |u_m(x) - u_m(x + x')| &= \left| \int_{B(0, \frac{1}{m})} \eta_m(y) (u(x - y, t') - u(x + x' - y, t')) dy \right| \\ &\leq \int_{B(0, \frac{1}{m})} m^n \cdot \eta(my) |u(x - y, t') - u(x + x' - y, t')| dy \\ &\leq C_{28} C_{30} |x'| m^n \int_{B(0, \frac{1}{m})} (1 + |x - y|^{p-1} + |x + x' - y|^{p-1}) dy. \end{aligned}$$

Because $|x'| < 1$ and $|y| \leq \frac{1}{m} \leq 1$, we have

$$\begin{aligned} 1 + |x - y|^{p-1} + |x + x' - y|^{p-1} &\leq 1 + (1 + |x|)^{p-1} + (2 + |x|)^{p-1} \\ &\leq (2 + 2^{p-1})(1 + |x|)^{p-1}. \end{aligned}$$

Furthermore (see [4], p. 615),

$$\int_{B(0, \frac{1}{m})} dy = \frac{\Pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} \cdot \frac{1}{m^n}, \quad \text{where } \Gamma(t) = \int_0^\infty s^{t-1} e^{-s} ds, \quad \text{for } t > 0.$$

In summary,

$$\frac{|u_m(x) - u_m(x + x')|}{|x'|} \leq C_{28} C_{30} \frac{\Pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} (2 + 2^{p-1})(1 + |x|)^{p-1}.$$

Taking the limit as $|x'| \rightarrow 0$ on both sides, we conclude (45). \square

Proof of (46). Let $x' \in \mathbb{R}^n$ and $\lambda \in (0, 1)$. We have

$$\begin{aligned} & u_m(x + \lambda x') - 2u_m(x) + u_m(x - \lambda x') \\ &= \int_{B(0, \frac{1}{m})} \eta_m(y) (u(x + \lambda x', t') - 2u(x, t') + u(x - \lambda x', t')) dy. \end{aligned}$$

From (31) and nonnegativity of η_m we conclude that $\frac{\partial^2 u_m(x)}{\partial x_i \partial x_j} \geq 0$. On the other hand, using (31) and mimicking the proof of (45), we see that

$$\begin{aligned} & u_m(x + \lambda x') - 2u_m(x) + u_m(x - \lambda x') \\ & \leq \int_{B(0, \frac{1}{m})} m^n \cdot \eta(my) C_{31} \lambda^2 (1 + |x|)^{(p-2)^+} dy \\ & \leq \lambda^2 C_{28} C_{31} \frac{\Pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} (1 + |x|)^{(p-2)^+}. \end{aligned}$$

For $p \in (1, 2]$, $(1 + |x|)^{(p-2)^+} = 1 \leq (1 + |x|^p) \leq 2^{p-1}(1 + |x|^p)$. For $p > 2$, in view of Lemma 1.2, $(1 + |x|)^{(p-2)^+} = (1 + |x|)^{p-2} \leq (1 + |x|)^p \leq 2^{p-1}(1 + |x|^p)$. Thus, for all $p > 1$ we have

$$\frac{u_m(x + \lambda x') - 2u_m(x) + u_m(x - \lambda x')}{\lambda^2} \leq C_{28} C_{31} \frac{\Pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} 2^{p-1} (1 + |x|^p).$$

Taking the limit as $\lambda \rightarrow 0$, we can conclude (46). □

Remark 2.4. Theorems 2.1 and 2.2 are true for functions u_ϵ (see (4)) instead of u . Indeed, in view of the proofs we see that the constants $C_{29}, C_{30}, C_{31}, C_{33}$ do not depend on ϵ .

Remark 2.5. It follows from (44)–(46) that for every $t' \in [0, T]$ $Du_m(\cdot; t')$ converges to $Du(\cdot, t')$ the distributional gradient of u with respect to x almost uniformly as $m \rightarrow \infty$ (see the proof of Theorem 3.5, to follow, for a similar argument, with u_m replaced by u_{ϵ_m}). This implies differentiability of u with respect to x in the classical sense (see, e.g., Theorem 7.17 in [13]), so Du is the classical gradient of u with respect to x at any point $(x, t') \in \mathbb{R}^n \times [0, T]$. Moreover, by (46) Du_m are locally Lipschitz in x uniformly in m , so Du is also locally Lipschitz in x . Thus Theorems 2.1, 2.2 and their proofs imply that the value function $u(x, t)$ has generalized derivatives of the first order with respect to t and of the second order with respect to x . These generalized derivatives belongs to the space $L_{loc}^\infty(\mathbb{R}^n \times [0, T])$ of all functions essentially bounded on every open bounded subset of the domain.

Proposition 2.6. For all $x \in \mathbb{R}^n$ and $t \in [0, T]$ we have $u(x, t) \leq (c_{\max} + C_{29})(1 + |x|)$.

Proof. Let $x' \in \mathbb{R}^n$ be arbitrary. Consider controls for which $\lim_{s \rightarrow 0^+} v_s = x$. In view of (2) and (3) we have

$$\begin{aligned} u(x', t) &= \inf\{J_{x't}(v) : v \in \mathcal{V}\} \leq c(t)|x| + \inf\{J_{x+x',t}(v) : v \in \mathcal{V}\} \\ &= c(t)|x| + u(x + x', t). \end{aligned}$$

So $u(x', t) - u(x + x', t) \leq c(t)|x|$. Similarly $u(x + x', t) - u(x', t) \leq c(t)|x|$, so

$$(53) \quad |u(x + x', t) - u(x', t)| \leq c(t)|x|.$$

Taking $x' = 0$, we get $|u(x, t) - u(0, t)| \leq c(t)|x|$. From (29) we see that $u(0, t) \leq C_{29}$ so $u(x, t) \leq c(t)|x| + u(0, t) \leq c_{\max}|x| + C_{29} \leq (c_{\max} + C_{29})(1 + |x|)$. \square

Remark 2.7. The proof of Proposition 2.6 is not valid for u_ϵ instead of u , because if a control $v \in \mathcal{V}_\epsilon$, then it is continuous, so the condition $\lim_{s \rightarrow 0^+} v_s = x$ is invalid for $x \neq 0$.

Remark 2.8. The value function $u(x, t)$ satisfies $|Du(x, t)| \leq c(t)$ for all $(x, t) \in \mathbb{R}^n \times [0, T]$. Indeed, the gradient exists for all $(x, t) \in \mathbb{R}^n \times [0, T]$ in view of Remark 2.5. From (53) we see that the first derivative of $u(x, t)$ with respect to x in any direction is bounded by $c(t)$. Hence, the norm of the gradient $Du(x, t)$ is bounded by $c(t)$, too.

3. Dynamic Programming Principle and HJB equation. To consider the DPP and the HJB equation for our problem we will first prove the pointwise convergence of u_ϵ to u if $\epsilon \rightarrow 0^+$. For this purpose we need an integral form of the Gronwall's inequality with locally finite measures.

Lemma 3.1 (see [18]). *Let μ be a locally finite measure on the Borel σ -algebra of $[t, T]$, where $0 \leq t \leq T$. We consider a measurable function ϕ defined on $[t, T]$ such that $\int_t^T |\phi(r)|\mu(dr) < \infty$. We assume that there exists a Borel function $\psi \geq 0$ on $[t, T]$ such that for all $s \in [t, T]$,*

$$\phi(s) \leq \psi(s) + \int_{[t,s)} \phi(r)\mu(dr).$$

Then for all $s \in [t, T]$,

$$\phi(s) \leq \psi(s) + \int_{[t,s)} \psi(r)e^{\mu([r,s))}\mu(dr).$$

Theorem 3.2. *For all $(x, t) \in \mathbb{R}^n \times [0, T]$ we have $\lim_{\epsilon \rightarrow 0^+} u_\epsilon(x, t) = u(x, t)$.*

Proof. Fix $x \in \mathbb{R}^n$ and $t \in [0, T]$. Consider an arbitrary $v \in \mathcal{V}$ such that $J_{xt}(v) < \infty$.

Step 1. We show first that $v \in L^p(\Omega \times [0, T - t], P \otimes \mu_{Leb})$, where μ_{Leb} denotes the Lebesgue's measure. Since $J_{xt}(v) < \infty$, we have

$$\mathbb{E} \int_t^T f(y_{xt}(s), s) ds < \infty$$

and from (6) we get

$$(54) \quad \mathbb{E} \int_t^T |y_{xt}(s)|^p ds < \infty.$$

From (1) we can write for $s \in [t, T]$

$$(55) \quad v(s-t) = y_{xt}(s) - x - \int_t^s b(r)dr - \int_t^s \sigma(r)dW_{r-t} - \int_t^s a(r)y_{xt}(r)dr.$$

Using (54) and properties of the normal distribution, we know that each term from the line above, maybe except for the last one, belongs to the space $L^p(\Omega \times [0, T-t])$. But the last term belongs to this space, too. Indeed,

$$\mathbb{E} \int_t^T \left| \int_t^s a(r)y_{xt}(r)dr \right|^p ds \leq a_{\max}^p \mathbb{E} \int_t^T \left(\int_t^T |y_{xt}(r)|dr \right)^p ds.$$

Using the Hölder's inequality and (54), we can estimate the last expression above by

$$\begin{aligned} & a_{\max}^p \mathbb{E} \int_t^T \left(\int_t^T |y_{xt}(r)|^p dr \cdot |T-t|^{p/q} \right) ds \\ & \leq a_{\max}^p T^{1+p/q} \mathbb{E} \int_t^T |y_{xt}(r)|^p dr < \infty, \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$. Hence, from (55) we see that $v \in L^p(\Omega \times [0, T-t])$.

Step 2. Now we define a sequence of bounded controls $\{v_R, R > 0\}$ such that v_R is convergent to v in the space $L^p(\Omega \times [0, T-t])$ and the total variation of v_R is pointwise convergent to the total variation of v from below. Let

$$v_R(s) = \begin{cases} v(s), & |v(s)| \leq R \\ \frac{v(s)}{|v(s)|} \cdot R, & |v(s)| > R. \end{cases}$$

We see that for all $s \in [0, T-t]$ $\lim_{R \rightarrow \infty} v_R(s) = v(s)$ and $|v_R(s)| \leq |v(s)|$. Hence, from Lemma 1.2 and Step 1,

$$\mathbb{E} \int_0^{T-t} |v(s) - v_R(s)|^p ds \leq 2^p \mathbb{E} \int_0^{T-t} |v(s)|^p ds < \infty$$

and using the Lebesgue's dominated convergence theorem, we get

$$\lim_{R \rightarrow \infty} \mathbb{E} \int_0^{T-t} |v(s) - v_R(s)|^p ds = \mathbb{E} \int_0^{T-t} \lim_{R \rightarrow \infty} |v(s) - v_R(s)|^p ds = 0.$$

The convergence in L^p is proved. Moreover, if $\xi(s), \xi_R(s)$ denote the total variations on the interval $[0, s]$ of the functions v, v_R respectively, then for all $s \in [0, T-t]$,

$$(56) \quad \xi_R(s) \leq \xi(s) \text{ and } \lim_{R \rightarrow \infty} \xi_R(s) = \xi(s).$$

Step 3. Let y_{xt}^v, y_{xt}^{vR} denote the state processes (see (1)) corresponding to the controls v, v_R respectively. We want to show that $\{y_{xt}^{vR}\}$ is convergent to y_{xt}^v in the space $L^p(\Omega \times [t, T])$. First we observe that for $s \in [t, T]$,

$$y_{xt}^v(s) - y_{xt}^{vR}(s) = \int_t^s a(r)(y_{xt}^v(r) - y_{xt}^{vR}(r))dr + v(s-t) - v_R(s-t).$$

Denoting $z_R(s) = y_{xt}^v(s) - y_{xt}^{vR}(s)$ and $u_R(s) = v(s-t) - v_R(s-t)$, we can rewrite the last equality in the form $z_R(s) = \int_t^s a(r)z_R(r)dr + u_R(s)$. Hence $|z_R(s)| \leq \int_t^s |z_R(r)|a_{\max}dr + |u_R(s)|$. Using Lemma 3.1 with $\phi = |z_R|$, $\psi = |u_R|$ and $\mu = a_{\max} \cdot \mu_{Leb}$, we get

$$(57) \quad \begin{aligned} |z_R(s)| &\leq |u_R(s)| + \int_t^s |u_R(r)|e^{a_{\max}(s-r)}dr \\ &\leq |u_R(s)| + C_{57} \int_t^s |u_R(r)|dr, \end{aligned}$$

where $C_{57} = e^{a_{\max}T}$. So from Lemma 1.2 and the Hölder's inequality

$$(58) \quad \begin{aligned} |z_R(s)|^p &\leq 2^{p-1} \left\{ |u_R(s)|^p + C_{57}^p \left(\int_t^s |u_R(r)|dr \right)^p \right\} \\ &\leq 2^{p-1} \left\{ |u_R(s)|^p + C_{57}^p (s-t)^{p/q} \int_t^s |u_R(r)|^p dr \right\} \\ &\leq C_{58} \left\{ |u_R(s)|^p + \int_t^T |u_R(r)|^p dr \right\}, \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$ and $C_{58} = 2^{p-1}(1 + C_{57}^p T^{p/q})$. Finally, in view of Step 2 we have

$$\begin{aligned} \lim_{R \rightarrow \infty} \mathbb{E} \int_t^T |z_R(s)|^p ds &\leq \lim_{R \rightarrow \infty} C_{58} \mathbb{E} \int_t^T \left\{ |u_R(s)|^p + \int_t^T |u_R(r)|^p dr \right\} ds \\ &\leq \lim_{R \rightarrow \infty} \left\{ C_{58} \mathbb{E} \int_t^T |u_R(s)|^p ds + C_{58} T \mathbb{E} \int_t^T |u_R(r)|^p dr \right\} = 0. \end{aligned}$$

Step 4. The next step is to show that $J_{xt}(v_R) \rightarrow J_{xt}(v)$ if $R \rightarrow \infty$. Indeed,

$$\begin{aligned} |J_{xt}(v) - J_{xt}(v_R)| &\leq \left| \mathbb{E} \int_t^T \left(f(y_{xt}^v(s), s) - f(y_{xt}^{vR}(s), s) \right) e^{-\int_t^s \alpha(r)dr} ds \right| \\ &\quad + \left| \mathbb{E} \int_t^T c(s) e^{-\int_t^s \alpha(r)dr} d(\xi - \xi_R)(s-t) \right| = A_R + B_R. \end{aligned}$$

In view of (56),

$$B_R \leq c_{\max} \left| \mathbb{E} \int_t^T d(\xi - \xi_R)(s-t) \right| = c_{\max} \mathbb{E}(\xi(T-t) - \xi_R(T-t)).$$

Using (56) again and the assumption that $J_{xt}(v) < \infty$, we see that $\mathbb{E}(\xi(T-t) - \xi_R(T-t)) \leq \mathbb{E}\xi(T-t) < \infty$. Hence, from the Lebesgue's dominated convergence theorem we get

$$\begin{aligned} \lim_{R \rightarrow \infty} B_R &\leq \lim_{R \rightarrow \infty} c_{\max} \mathbb{E}(\xi(T-t) - \xi_R(T-t)) \\ &= c_{\max} \mathbb{E} \lim_{R \rightarrow \infty} (\xi(T-t) - \xi_R(T-t)) = 0. \end{aligned}$$

Using (7) and the Hölder's inequality, we have

$$\begin{aligned} A_R &\leq \mathbb{E} \int_t^T |f(y_{xt}^v(s), s) - f(y_{xt}^{vR}(s), s)| ds \\ &\leq \mathbb{E} \int_t^T (1 + f(y_{xt}^v(s), s) + f(y_{xt}^{vR}(s), s))^{1-1/p} |y_{xt}^v(s) - y_{xt}^{vR}(s)| ds \\ &\leq \left\{ \mathbb{E} \int_t^T (1 + f(y_{xt}^v(s), s) + f(y_{xt}^{vR}(s), s)) ds \right\}^{1-1/p} \left\{ \mathbb{E} \int_t^T |y_{xt}^v(s) - y_{xt}^{vR}(s)|^p ds \right\}^{1/p}. \end{aligned}$$

In view of Step 3, the second factor in the last expression goes to 0 if $R \rightarrow \infty$. We must show that the first factor is bounded. Indeed, from (6) and Lemma 1.2 we can write

$$\begin{aligned} &\mathbb{E} \int_t^T (1 + f(y_{xt}^v(s), s) + f(y_{xt}^{vR}(s), s)) ds \\ &\leq (1 + C_0) \mathbb{E} \int_t^T (2 + |y_{xt}^v(s)|^p + |y_{xt}^{vR}(s)|^p) ds \\ &\leq (1 + C_0) \mathbb{E} \int_t^T (2 + |y_{xt}^v(s)|^p + 2^{p-1} |y_{xt}^v(s)|^p + 2^{p-1} |y_{xt}^v(s) - y_{xt}^{vR}(s)|^p) ds. \end{aligned}$$

Using (54) and Step 3 again, we conclude that the last expression is bounded uniformly in R . Hence $\lim_{R \rightarrow \infty} A_R = 0$.

Summarizing Steps 1-4, we know that $J_{xt}(v_R)$ goes to $J_{xt}(v)$ if $R \rightarrow \infty$, so we can consider only bounded controls.

Step 5. Consider $v \in \mathcal{V}$ such that $\|v\|_\infty < R$ for some $R > 0$. We will construct a sequence of controls $\{v_n, n \in \mathbb{N}\}$ convergent to v in $L^p(\Omega \times [0, T-t])$ and such that $v_n \in V_{1/(2nR)}$ for all n . Besides we shall prove that the variation of v_n is pointwise convergent to the variation of v from below. Let $v_n(s) = n \int_{(s-1/n) \vee 0}^s v(r) dr$, $s \in [0, T-t]$. We observe that v_n is a progressively measurable continuous random process such that $\|v_n\|_\infty \leq R$, so $v_n \in L^p(\Omega \times [0, T-t])$. From left-continuity of v we know that

$$(59) \quad \forall \omega \in \Omega \quad \forall s \in [0, T-t] \quad \lim_{n \rightarrow \infty} v_n(s) = v(s).$$

Using the Lebesgue's dominated convergence theorem, we get

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^{T-t} |v(s) - v_n(s)|^p ds = \mathbb{E} \int_0^{T-t} \lim_{n \rightarrow \infty} |v(s) - v_n(s)|^p ds = 0,$$

so L^p -convergence is proved.

Now we want to check that $v_n \in V_{1/(2nR)}$. Indeed,

$$\left| \frac{d}{ds} v_n(s) \right| = \left| \frac{d}{ds} \left(n \int_{(s-1/n) \vee 0}^s v(r) dr \right) \right| = n |v(s) - v((s-1/n) \vee 0)| \leq 2nR.$$

Let $\xi_n(s), \xi(s)$ denote the variations on the interval $[0, s]$ of the functions v_n, v respectively. For convenience, we define $v(r) \equiv 0$ for $r < 0$. Then $v_n(s) = n \int_{s-1/n}^s v(r) dr$, $s \in [0, T-t]$. Fix $\omega \in \Omega$, $s \in (0, T-t]$. Let $\Pi = \{s_0, s_1, \dots, s_k\}$ be a partition of the interval $[0, s]$, where $0 = s_0 < s_1 < \dots < s_k = s$. Then

$$\begin{aligned} \sum_{i=1}^k |v_n(s_i) - v_n(s_{i-1})| &= n \sum_{i=1}^k \left| \int_{s_{i-1}/n}^{s_i} v(r) dr - \int_{s_{i-1}-1/n}^{s_{i-1}} v(r) dr \right| \\ &= n \sum_{i=1}^k \left| \int_0^{1/n} \left(v(s_i + r - 1/n) - v(s_{i-1} + r - 1/n) \right) dr \right| \\ &\leq n \int_0^{1/n} \sum_{i=1}^k |v(s_i + r - 1/n) - v(s_{i-1} + r - 1/n)| dr \\ &\leq n \int_0^{1/n} \xi(s) dr = \xi(s). \end{aligned}$$

Letting $|\Pi| \rightarrow 0$, we get

$$(60) \quad \xi_n(s) \leq \xi(s).$$

On the other hand, from (59) we see that

$$\begin{aligned} \sum_{i=1}^k |v(s_i) - v(s_{i-1})| &= \sum_{i=1}^k \left| \lim_{n \rightarrow \infty} v_n(s_i) - \lim_{n \rightarrow \infty} v_n(s_{i-1}) \right| \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^k |v_n(s_i) - v_n(s_{i-1})| \leq \liminf_{n \rightarrow \infty} \xi_n(s). \end{aligned}$$

Letting $|\Pi| \rightarrow 0$ and using (60), we have

$$\xi(s) \leq \liminf_{n \rightarrow \infty} \xi_n(s) \leq \limsup_{n \rightarrow \infty} \xi_n(s) \leq \xi(s) \quad \Leftrightarrow \quad \lim_{n \rightarrow \infty} \xi_n(s) = \xi(s).$$

Step 6. In view of Step 5 we can mimic Steps 3 and 4 to conclude that $J_{xt}(v_n) \rightarrow J_{xt}(v)$ if $n \rightarrow \infty$, where $\|v\|_\infty < R$ for some $R > 0$. From this

and Step 4, remembering that $v_n \in \mathcal{V}_{1/(2nR)}$, we can write

$$(61) \quad \inf_{v \in \mathcal{V}} J_{xt}(v) = \inf_{v \in \bigcup_{\epsilon > 0} \mathcal{V}_\epsilon} J_{xt}(v)$$

and $\lim_{\epsilon \rightarrow 0^+} u_\epsilon(x, t) = u(x, t)$. □

Theorem 3.3 (Bellman's dynamic programming principle). *Let $x \in \mathbb{R}^n$, $t \in [0, T]$ and let y_{xt}^v denote the state process corresponding to a control $v \in \mathcal{V}$. Let $\tau \in [0, T - t]$ be a Markov time with respect to $\{\mathcal{F}_t\}$. Then*

$$u(x, t) = \inf_{v \in \mathcal{V}} \mathbb{E} \left\{ \int_t^{t+\tau} f(y_{xt}^v(s), s) e^{-\int_t^s \alpha(r) dr} ds + \int_t^{t+\tau} c(s) e^{-\int_t^s \alpha(r) dr} d\xi(s-t) + u(y_{xt}^v(t+\tau), t+\tau) \right\}.$$

Proof. For convenience let us denote

$$J_{xt}(v, \tau) = \mathbb{E} \left\{ \int_t^{t+\tau} f(y_{xt}^v(s), s) e^{-\int_t^s \alpha(r) dr} ds + \int_t^{t+\tau} c(s) e^{-\int_t^s \alpha(r) dr} d\xi(s-t) \right\}.$$

It is known that DPP holds for regular stochastic control problems (see, e.g., [10], Th. 3.1.6). Hence we have for each $\epsilon > 0$,

$$(62) \quad u_\epsilon(x, t) = \inf_{v \in \mathcal{V}_\epsilon} \left\{ J_{xt}(v, \tau) + \mathbb{E} u_\epsilon(y_{xt}^v(t+\tau), t+\tau) \right\}.$$

Considering any $\tilde{v} \in \mathcal{V}_\epsilon$, we have

$$u_\epsilon(x, t) \leq J_{xt}(\tilde{v}, \tau) + \mathbb{E} u_\epsilon(y_{xt}^{\tilde{v}}(t+\tau), t+\tau).$$

If $\epsilon \rightarrow 0^+$, from Theorem 3.2 and the Lebesgue's dominated convergence theorem we get

$$u(x, t) \leq J_{xt}(\tilde{v}, \tau) + \mathbb{E} u(y_{xt}^{\tilde{v}}(t+\tau), t+\tau).$$

Because $\epsilon > 0$ and $\tilde{v} \in \mathcal{V}_\epsilon$ are arbitrary we can conclude that

$$(63) \quad u(x, t) \leq \inf_{v \in \bigcup_{\epsilon > 0} \mathcal{V}_\epsilon} \left\{ J_{xt}(v, \tau) + \mathbb{E} u(y_{xt}^v(t+\tau), t+\tau) \right\}.$$

On the other hand, from (62)

$$u_\epsilon(x, t) \geq \inf_{v \in \bigcup_{\epsilon > 0} \mathcal{V}_\epsilon} \left\{ J_{xt}(v, \tau) + \mathbb{E} u(y_{xt}^v(t+\tau), t+\tau) \right\}.$$

Letting $\epsilon \rightarrow 0^+$, we get

$$(64) \quad u(x, t) \geq \inf_{v \in \bigcup_{\epsilon > 0} \mathcal{V}_\epsilon} \left\{ J_{xt}(v, \tau) + \mathbb{E} u(y_{xt}^v(t+\tau), t+\tau) \right\}.$$

The inequalities (63), (64) and an argument similar to the proof of Theorem 3.2 (see (61)) imply that

$$u(x, t) = \inf_{v \in \mathcal{V}} \left\{ J_{xt}(v, \tau) + \mathbb{E} u(y_{xt}^v(t+\tau), t+\tau) \right\}. \quad \square$$

Corollary 3.4. *The dynamic programming property in the weak sense holds (see Definition 1.1) and hence the value function satisfies (33).*

Denote

$$Au(x, t) = \frac{-\partial u(x, t)}{\partial t} - \frac{1}{2}\beta(t) \circ D^2u(x, t) - \left(a(t)x + b(t)\right) \circ Du(x, t) + \alpha(t)u(x, t),$$

where \circ denotes the scalar product of vectors and matrices respectively.

Theorem 3.5 (The HJB equation). *The value function u satisfies almost everywhere (a.e.) the following second-order differential equation:*

$$(65) \quad \max \left\{ Au(x, t) - f(x, t), |Du(x, t)| - c(t) \right\} = 0.$$

Proof. An application of the DPP for regular stochastic control problems yields for $\epsilon > 0$ the following equation (see [5], Chapter IV.3):

$$(66) \quad Au_\epsilon(x, t) + \frac{1}{\epsilon} \left(|Du_\epsilon(x, t)| - c(t) \right)^+ = f(x, t) \quad a.e.$$

In view of Theorems 2.1, 2.2, Remark 2.4, Theorem 3.2, Corollary 3.4 and the Arzela–Ascoli’s theorem ([7], Th. 2.4.9) we see that $u_\epsilon \rightarrow u$ uniformly on every compact set if $\epsilon \rightarrow 0^+$.

Fix $t \in [0, T]$. From (31) and Remark 2.4 we see that $D^2u_\epsilon(\cdot, t)$ are locally uniformly bounded for all $\epsilon > 0$ in their domains, so using the Arzela–Ascoli’s theorem from every sequence $\{\epsilon_m\}_{m \in \mathbb{N}}$ convergent to 0, we can choose a subsequence $\{\tilde{\epsilon}_m\}_{m \in \mathbb{N}}$ such that

$$Du_{\tilde{\epsilon}_m}(\cdot, t) \rightarrow v = (v_1, \dots, v_n) \quad \text{almost uniformly if } m \rightarrow \infty.$$

But v must be equal to $Du(\cdot, t)$ in the distribution sense. Indeed, for any function $\phi \in C_c^\infty(\mathbb{R}^n)$ and any $k = 1, \dots, n$ we have

$$\int_{\mathbb{R}^n} \frac{\partial \phi(x)}{\partial x_k} u_{\tilde{\epsilon}_m}(x, t) dx = - \int_{\mathbb{R}^n} \phi(x) \frac{\partial u_{\tilde{\epsilon}_m}(x, t)}{\partial x_k} dx.$$

Letting $m \rightarrow \infty$, we get

$$\int_{\mathbb{R}^n} \frac{\partial \phi(x)}{\partial x_k} u(x, t) dx = - \int_{\mathbb{R}^n} \phi(x) v_k(x) dx,$$

so $v_k(\cdot) = \frac{\partial u(\cdot, t)}{\partial x_k}$ almost everywhere. Since $\frac{\partial u}{\partial x_k}$ and v_k are Lipschitz continuous, the equality holds for all $x \in \mathbb{R}^n$. Thus, v does not depend on the choice of the subsequence $\{\tilde{\epsilon}_m\}$, so

$$(67) \quad \forall t \in [0, T] \quad Du_\epsilon(\cdot, t) \rightarrow Du(\cdot, t) \quad \text{almost uniformly if } \epsilon \rightarrow 0^+.$$

Let $\psi = (1 + |x|)^{-2p-n-1}$. From (29)–(33) we conclude that $|Au_\epsilon(x, t)|$ is not greater than

$$(C_{33} + C_{29}\alpha_{\max})(1 + |x|^p) + \frac{1}{2}\beta_{\max}n^2C_{31}(1 + |x|)^{(p-2)^+} \\ + (a_{\max}|x| + b_{\max})nC_{30}(1 + 2|x|^{p-1})$$

for almost every $(x, t) \in \mathbb{R}^n \times [0, T]$. Using Lemma 1.2, we have the estimate

$$(68) \quad |Au_\epsilon(x, t)| \leq C_{68}(1 + |x|)^p \quad \text{a.e.}$$

for some constant $C_{68} > 0$ depending only on $C_{29}, C_{30}, C_{31}, C_{33}, n, p, a_{\max}, b_{\max}, \alpha_{\max}, \beta_{\max}$. Hence

$$|Au_\epsilon(x, t)|^2\psi(x) \leq \frac{C_{68}^2(1 + |x|)^{2p}}{(1 + |x|)^{2p+n+1}} = \frac{C_{68}^2}{(1 + |x|)^{n+1}} \quad \text{a.e.}$$

The same estimate holds for u instead u_ϵ . So $|Au_\epsilon|^2\psi, |Au|^2\psi \in L^1(\mathbb{R}^n \times [0, T])$. Moreover, Au_ϵ, Au are uniformly bounded in the space L_ψ^2 , where

$$L_\psi^2 = \left\{ v : v^2\psi \in L^1(\mathbb{R}^n \times [0, T]) \right\} = L_{\psi \cdot \mu_{Leb}}^2(\mathbb{R}^n \times [0, T]).$$

From the Banach–Alaoglu theorem ([12], Th. 3.15) we know that balls in the space L^2 are weakly compact. So for each sequence $\{\epsilon_m\}_{m \in \mathbb{N}}$ convergent to 0, there exists a subsequence $\{\tilde{\epsilon}_m\}_{m \in \mathbb{N}}$ such that $Au_{\tilde{\epsilon}_m} \rightharpoonup v$ in L_ψ^2 if $m \rightarrow \infty$. We will show that $v = Au$ in the distribution sense. Indeed, for any function ϕ belonging to the class $C_c^\infty(\mathbb{R}^n \times [0, T])$, we have

$$\int_0^T \int_{\mathbb{R}^n} (Au_{\tilde{\epsilon}_m})\phi dxdt = \int_0^T \int_{\mathbb{R}^n} \frac{\partial \phi}{\partial t} u_{\tilde{\epsilon}_m} dxdt - \frac{1}{2} \int_0^T \int_{\mathbb{R}^n} (\beta(t) \circ D^2\phi) u_{\tilde{\epsilon}_m} dxdt \\ + \int_0^T \int_{\mathbb{R}^n} \left((a(t)x + b(t)) \circ D\phi \right) u_{\tilde{\epsilon}_m} dxdt + \int_0^T \int_{\mathbb{R}^n} \text{tr}(a(t)) u_{\tilde{\epsilon}_m} \phi dxdt \\ + \int_0^T \int_{\mathbb{R}^n} \alpha(t)\phi u_{\tilde{\epsilon}_m} dxdt.$$

Letting $m \rightarrow \infty$, we get

$$\int_0^T \int_{\mathbb{R}^n} v\phi dxdt = \int_0^T \int_{\mathbb{R}^n} \frac{\partial \phi}{\partial t} u dxdt - \frac{1}{2} \int_0^T \int_{\mathbb{R}^n} (\beta(t) \circ D^2\phi) u dxdt \\ + \int_0^T \int_{\mathbb{R}^n} \left((a(t)x + b(t)) \circ D\phi \right) u dxdt + \int_0^T \int_{\mathbb{R}^n} \text{tr}(a(t)) u \phi dxdt \\ + \int_0^T \int_{\mathbb{R}^n} \alpha(t)\phi u dxdt = \int_0^T \int_{\mathbb{R}^n} (Au)\phi dxdt.$$

Hence $Au_{\tilde{\epsilon}_m} \rightharpoonup Au$ in L_ψ^2 if $m \rightarrow \infty$. From uniqueness of the limit we conclude

$$(69) \quad Au_\epsilon \rightharpoonup Au \quad \text{in } L_\psi^2 \text{ if } \epsilon \rightarrow 0^+.$$

In view of (66) we have $Au_\epsilon \leq f$ a.e. From this and (69) we see that $Au(x, t) \leq f(x, t)$ a.e. This, together with Remark 2.8 ensure us that

$$(70) \quad \max \left\{ Au(x, t) - f(x, t), |Du(x, t)| - c(t) \right\} \leq 0 \quad \text{a.e.}$$

Take a sequence $\epsilon_n \rightarrow 0$ and let \mathcal{D} be the set of $(x, t) \in \mathbb{R}^n \times [0, T]$ such that (66) holds at (x, t) for all ϵ_n . Then $\mu_{Leb}((\mathbb{R}^n \times [0, T]) \setminus \mathcal{D}) = 0$. Choose $t \in [0, T]$ such that for almost every $x \in \mathbb{R}^n$ we have $(x, t) \in \mathcal{D}$. Since $|Du_{\epsilon_n}(x, t)| \rightarrow |Du(x, t)|$ as $n \rightarrow \infty$ (see (67)), $\mathbb{I}_{\{|Du_{\epsilon_n}(x, t)| < c(t)\}} = \mathbb{I}_{\{|Du(x, t)| < c(t)\}}$ for n large enough (depending on (x, t)). We have from this and (66) that

$$(71) \quad \mathbb{I}_{\{|Du(x, t)| < c(t)\}} Au_{\epsilon_n}(x, t) \rightarrow \mathbb{I}_{\{|Du(x, t)| < c(t)\}} f(x, t) \quad \text{a.e.}$$

On the other hand, (69) yields

$$(72) \quad \mathbb{I}_{\{|Du(x, t)| < c(t)\}} Au_{\epsilon_n}(x, t) \rightharpoonup \mathbb{I}_{\{|Du(x, t)| < c(t)\}} Au(x, t) \quad \text{in } L^2_\psi,$$

so the sequence $\{\mathbb{I}_{\{|Du(x, t)| < c(t)\}} Au_{\epsilon_n}(x, t)\}$ is bounded in L^2_ψ and thus it is uniformly integrable in L^1_ψ . This, together with (71), implies that for every $\phi \in C_c^\infty(\mathbb{R}^n \times [0, T]) \subset L^2_\psi$,

$$(73) \quad \int_0^T \int_{\mathbb{R}^n} \mathbb{I}_{\{|Du(x, t)| < c(t)\}} (Au_{\epsilon_n} \phi \psi)(x, t) dx dt \\ \rightarrow \int_0^T \int_{\mathbb{R}^n} \mathbb{I}_{\{|Du(x, t)| < c(t)\}} (f \phi \psi)(x, t) dx dt,$$

which, together with (72) implies that

$$\mathbb{I}_{\{|Du(x, t)| < c(t)\}} Au(x, t) = \mathbb{I}_{\{|Du(x, t)| < c(t)\}} f(x, t) \quad \text{a.e.} \quad \square$$

4. Existence and uniqueness of the optimal control. The results of this section are analogous to Theorems 7 and 8 from [11].

Fix $(t, x) \in [0, T] \times \mathbb{R}^n$ (for $t = T$ the only admissible control is $v(0) = 0$ a.s.). Let m_t be the measure on $([t, T] \times \Omega, \mathcal{B}([t, T]) \otimes \mathcal{F})$ equal to the product of the Lebesgue's measure and P .

Remark 4.1. If a process X is a modification of a process Y and both processes have left-continuous sample paths a.s., then the processes X, Y are indistinguishable (compare Problem 1.1.5, [7]).

Theorem 4.2. *The optimal control $v^* \in \mathcal{V}$, if it exists, is unique up to the indistinguishability.*

Proof. Suppose there are $v_1, v_2 \in \mathcal{V}$ for which $u(x, t) = J_{xt}(v_1) = J_{xt}(v_2)$. Put $v_0 = (v_1 + v_2)/2$. Of course $v_0 \in \mathcal{V}$. From Lemma 1.7 we have

$$(74) \quad u(x, t) - J_{xt}(v_0) = \frac{1}{2} (J_{xt}(v_1) + J_{xt}(v_2)) - J_{xt}(v_0) \geq 0.$$

Let $y_{xt}^0, y_{xt}^1, y_{xt}^2$ be the solutions of (1) corresponding to $v = v_0, v_1, v_2$ respectively. In view of the proof of Lemma 1.7 and strict convexity of the running cost function f , we have

$$(75) \quad f(y_{xt}^0(s), s) < \frac{1}{2}f(y_{xt}^1(s), s) + \frac{1}{2}f(y_{xt}^2(s), s)$$

provided that $y_{xt}^1(s) \neq y_{xt}^2(s)$.

Assume that v_1, v_2 are not indistinguishable. Then there exists $s' \in (t, T]$ such that $P(A) > 0$, where $A = \{v_1(s') \neq v_2(s')\}$ (see Remark 4.1). Because v_1, v_2 have left-continuous sample paths a.s., there exists $s''(\omega) \in (t, s')$ such that $v_1(s) \neq v_2(s)$ for all $s \in [s''(\omega), s']$, $\omega \in A$. Thus, $y_{xt}^1(s) \neq y_{xt}^2(s)$ on some m_t -nonzero set. This fact together with (75) and the definition of J_{xt} imply that the inequality (74) is strict, so we get a contradiction. We conclude that v_1, v_2 must be indistinguishable. \square

Lemma 4.3. *Let $\{z_n\}_{n \in \mathbb{N}}$ be a sequence in $L^p(m_t)$. If $z_n \rightarrow 0$ in $L^p(m_t)$, then $\mathcal{T}z_n \rightarrow 0$ in $L^p(m_t)$, where*

$$\mathcal{T}z_n(s, \omega) = z_n(s, \omega) - \int_t^s a(r)z_n(r, \omega)dr.$$

Proof. By the Hölder's inequality, the function $g(s, \omega) = \int_t^s a(r)z_n(r, \omega)dr$ satisfies

$$\|g\|_{L^p}^p \leq a_{\max}^p \mathbb{E} \int_t^T \int_t^T |z_n(r, \omega)|^p \cdot T^{p/q} dr ds \leq a_{\max}^p \cdot T^{1+p/q} \cdot \|z_n\|_{L^p}^p,$$

so \mathcal{T} is a bounded operator from $L^p(m_t)$ into $L^p(m_t)$. \square

Theorem 4.4. *There exists an optimal control $v^* \in \mathcal{V}$.*

Proof. Let $\{v_k\}_{k \in \mathbb{N}}$ be a sequence of admissible controls such that $J_{xt}(v_k) \rightarrow u(x, t)$ as $k \rightarrow \infty$ and let y_{xt}^k be the solution of (1) corresponding to $v = v_k$. Then $J_{xt}(v_k)$ are uniformly bounded in k . By Lemma 1.9 the sequence $\{y_{xt}^k\}_{k \in \mathbb{N}}$ is bounded in $L^p(m_t)$ and hence, by the Banach–Alaoglu theorem ([12], Th. 3.15), there exists a subsequence (still denoted by $\{y_{xt}^k\}$) and a process y_{xt} such that $y_{xt}^k \rightharpoonup y_{xt}$ in $L^p(m_t)$.

Fix $k \in \mathbb{N}$. Since the sequence $\{y_{xt}^i\}_{i \geq k}$ is also convergent to y_{xt} , by the Mazur theorem there exists

$$z_{xt}^k = \sum_{i=k}^{n(k)} \alpha_{k,i} \cdot y_{xt}^i, \quad \alpha_{k,i} \geq 0, \quad \sum_{i=k}^{n(k)} \alpha_{k,i} = 1, \quad k \leq n(k) < \infty$$

such that $\|z_{xt}^k - y_{xt}\|_{L^p} \leq 1/k$. In particular $z_{xt}^k \rightarrow y_{xt}$ in $L^p(m_t)$. Let $\eta_k = \sum_{i=k}^{n(k)} \alpha_{k,i} \cdot v_i$ be the control corresponding to z_{xt}^k in (1). Then $\eta_k \in \mathcal{V}$. Moreover by Lemma 1.7,

$$u(x, t) \leq J_{xt}(\eta_k) \leq \sum_{i=k}^{n(k)} \alpha_{k,i} \cdot J_{xt}(v_i) \leq \max_{i=k, \dots, n(k)} J_{xt}(v_i) \xrightarrow{k \rightarrow \infty} u(x, t).$$

For $s \in [t, T]$ we have

$$z_{xt}^k(s) - z_{xt}^m(s) - \int_t^s a(r)(z_{xt}^k(r) - z_{xt}^m(r))dr = \eta_k(s-t) - \eta_m(s-t).$$

Because $\{z_{xt}^k\}$ is convergent in $L^p(m_t)$, $z_{xt}^k - z_{xt}^m$ goes to 0 in $L^p(m_t)$ as $k, m \rightarrow \infty$. Using Lemma 4.3 we conclude that $\{\eta_k(\cdot - t, \cdot)\}_{k \in \mathbb{N}}$ is a Cauchy sequence in $L^p(m_t)$ so it is convergent to a process $v \in L^p(m_t)$. Without loss of generality we may assume that $v(0) \equiv 0$.

Now we choose a subsequence (still denoted by k) such that $\eta_k(s, \omega) \rightarrow v(s, \omega)$ as $k \rightarrow \infty$ for $(s, \omega) \in \mathcal{A}$, where $(\mu_{Leb} \times P)(\mathcal{A}) = T - t$. For $\omega \in \Omega$ and $s \in [0, T - t]$, we define

$$\mathcal{A}_\omega = \{s \in [0, T - t] : (s, \omega) \in \mathcal{A}\}, \quad \mathcal{A}_s = \{\omega \in \Omega : (s, \omega) \in \mathcal{A}\}.$$

Note that $P(\mathcal{A}_0) = 1$ because $\eta_k(0) = v(0) = 0$ P -a.s. Furthermore, let

$$\tilde{\Omega} = \{\omega \in \Omega : \mu_{Leb}(\mathcal{A}_\omega) = T - t\}, \quad \mathcal{S} = \{s \in [0, T - t] : P(\mathcal{A}_s) = 1\}.$$

Then $P(\tilde{\Omega}) = 1$ and $\mu_{Leb}(\mathcal{S}) = T - t$. Let \mathcal{N} be a countable subset of \mathcal{S} , dense in $[0, T - t]$, including 0 and let $\mathcal{A}_{\mathcal{N}} = \bigcap_{s \in \mathcal{N}} \mathcal{A}_s$. We have $P(\mathcal{A}_{\mathcal{N}}) = 1$.

Let $\xi_k(s)$ denote the total variation of η_k on the interval $[0, s]$. Because $J_{xt}(\eta_k)$ are uniformly bounded in k , there exists a constant $C > 0$ such that $\mathbb{E}\xi_k(T - t) \leq C$ for all $k \in \mathbb{N}$. In view of the Fatou's lemma, $\mathbb{E} \liminf_{k \rightarrow \infty} \xi_k(T - t) \leq \liminf_{k \rightarrow \infty} \mathbb{E}\xi_k(T - t) \leq C$, so $\liminf_{k \rightarrow \infty} \xi_k(T - t)$ is finite a.s.

Fix $\omega \in \Omega$ and let $\Pi \subset \mathcal{A}_\omega$, $\Pi = \{t_0, t_1, \dots, t_m\}$, $0 = t_0 < t_1 < \dots < t_m \leq T - t$. Let $k_n = k_n(\omega) \rightarrow \infty$ be a sequence of natural numbers such that $\lim_{k_n \rightarrow \infty} \xi_{k_n}(T - t) = \liminf_{k \rightarrow \infty} \xi_k(T - t)$. Then

$$\begin{aligned} \sum_{i=0}^{m-1} |v(t_{i+1}) - v(t_i)| &= \lim_{k_n \rightarrow \infty} \sum_{i=0}^{m-1} |\eta_{k_n}(t_{i+1}) - \eta_{k_n}(t_i)| \\ &\leq \lim_{k_n \rightarrow \infty} \xi_{k_n}(T - t) = \liminf_{k \rightarrow \infty} \xi_k(T - t). \end{aligned}$$

Thus, $v|_{\mathcal{A}_\omega}$ has bounded variation and hence it has left-hand and right-hand limits at each point. Let $v^* = 0$ on the P -zero set $\Omega \setminus (\mathcal{A}_{\mathcal{N}} \cap \tilde{\Omega})$. On $\mathcal{A}_{\mathcal{N}} \cap \tilde{\Omega}$ let

$$v^*(s) = \begin{cases} 0 = v(0), & s = 0 \\ \lim_{\mathcal{A}_\omega \ni u \uparrow s} v(u) = \lim_{\mathcal{N} \ni u \uparrow s} v(u), & s \in (0, T - t]. \end{cases}$$

Then v^* is progressively measurable, left-continuous and $v^*(0) = 0$. Moreover, for $\omega \in \mathcal{A}_{\mathcal{N}} \cap \tilde{\Omega}$ and for each partition $\Pi = \{t_0, t_1, \dots, t_m\}$, $0 = t_0 < t_1 < \dots < t_m \leq T - t$, we can choose $\{t_i^k\}_{k \in \mathbb{N}} \subset \mathcal{A}_\omega$ such that $t_i^k \uparrow t_i$ as

$k \rightarrow \infty$, $i = 1, 2, \dots, m$. Therefore,

$$\begin{aligned} \sum_{i=0}^{m-1} |v^*(t_{i+1}) - v^*(t_i)| &= \lim_{k \rightarrow \infty} \sum_{i=0}^{m-1} |v(t_{i+1}^k) - v(t_i^k)| \\ &\leq \text{Var}(v, [0, T-t] \cap \mathcal{A}_\omega) \leq \liminf_{k \rightarrow \infty} \xi_k(T-t), \end{aligned}$$

so v^* has bounded variation. This ensures us that $v^* \in \mathcal{V}$.

For $\omega \in \mathcal{A}_\mathcal{N} \cap \tilde{\Omega}$, the set $A_\omega \cap \{s \in [0, T-t] : v(s, \omega) \neq v^*(s, \omega)\}$ is countable, so its Lebesgue's measure is equal to 0. Therefore $v = v^*$ m_t -a.e. In particular, $\eta_k \rightarrow v^*$ in $L^p(m_t)$. Proceeding as in Steps 3–4 in the proof of Theorem 3.2 we can show that $y_{xt}^{\eta_k} \rightarrow y_{xt}^{v^*}$ in $L^p(m_t)$ and hence

$$(76) \quad \lim_{k \rightarrow \infty} \mathbb{E} \int_t^T f(y_{xt}^{\eta_k}(s), s) e^{-\int_t^s \alpha(r) dr} ds = \mathbb{E} \int_t^T f(y_{xt}^{v^*}(s), s) e^{-\int_t^s \alpha(r) dr} ds.$$

To finish the proof we need to check that

$$(77) \quad \mathbb{E} \int_t^T c(s) e^{-\int_t^s \alpha(r) dr} d\xi^*(s-t) \leq \liminf_{k \rightarrow \infty} \mathbb{E} \int_t^T c(s) e^{-\int_t^s \alpha(r) dr} d\xi_k(s-t),$$

where ξ^* is the total variation of v^* . Fix $\omega \in \mathcal{A}_\mathcal{N} \cap \tilde{\Omega}$ and let $0 \leq s_1 \leq s_2 \leq T-t$. Let $\Pi = \{t_0, t_1, \dots, t_m\}$, $s_1 = t_0 < t_1 < \dots < t_m = s_2$ and let $\{t_i^k\}_{k \in \mathbb{N}} \subset \mathcal{A}_\omega$ be such that $t_i^k \uparrow t_i$ as $k \rightarrow \infty$, $i = 0, 1, \dots, m$. Then for every $k_0 \in \mathbb{N}$

$$\begin{aligned} \sum_{i=0}^{m-1} |v^*(t_{i+1}) - v^*(t_i)| &= \lim_{k \rightarrow \infty} \sum_{i=0}^{m-1} |v(t_{i+1}^k) - v(t_i^k)| \\ &\leq \text{Var}(v, [t_0^{k_0}, s_2] \cap \mathcal{A}_\omega). \end{aligned}$$

Letting $k_0 \rightarrow \infty$, we get $\sum_{i=0}^{m-1} |v^*(t_{i+1}) - v^*(t_i)| \leq \text{Var}(v, [s_1, s_2] \cap \mathcal{A}_\omega)$ and hence

$$(78) \quad \text{Var}(v^*, [s_1, s_2]) \leq \text{Var}(v, [s_1, s_2] \cap \mathcal{A}_\omega).$$

Let $\xi(s) = \text{Var}(v, [0, s] \cap \mathcal{A}_\omega)$, $s \in [0, T-t]$. Restricting $\Pi = \{t_0, t_1, \dots, t_m\}$, $s_1 = t_0 < t_1 < \dots < t_m = s_2$ so that $\Pi \subset \mathcal{A}_\omega$ (in particular assuming $s_1, s_2 \in \mathcal{A}_\omega$), we get

$$\begin{aligned} \sum_{i=0}^{m-1} |v(t_{i+1}) - v(t_i)| &= \lim_{k \rightarrow \infty} \sum_{i=0}^{m-1} |\eta_k(t_{i+1}) - \eta_k(t_i)| \\ &\leq \liminf_{k \rightarrow \infty} (\xi_k(s_2) - \xi_k(s_1)). \end{aligned}$$

As $\|\Pi\| \rightarrow 0$ we get

$$(79) \quad \xi(s_2) - \xi(s_1) \leq \liminf_{k \rightarrow \infty} (\xi_k(s_2) - \xi_k(s_1)).$$

Now take $\Pi = \{t_0, t_1, \dots, t_m\}$, $0 = t_0 < t_1 < \dots < t_m \leq T-t$, $\Pi \subset \mathcal{N}$. In particular, $\Pi \subset \mathcal{A}_\omega$ for all $\omega \in \mathcal{A}_\mathcal{N}$. For every interval $[t_i, t_{i+1}]$,

$i = 0, 1, \dots, m-1$, let $l_i = \min \left\{ c(s) e^{-\int_t^s \alpha(r) dr} : s \in [t_i + t, t_{i+1} + t] \right\}$. For $\omega \in \mathcal{A}_{\mathcal{N}} \cap \tilde{\Omega}$, by (78)–(79), we have

$$\begin{aligned} \sum_{i=0}^{m-1} l_i \cdot (\xi^*(t_{i+1}) - \xi^*(t_i)) &\leq \sum_{i=0}^{m-1} l_i \cdot (\xi(t_{i+1}) - \xi(t_i)) \\ &\leq \liminf_{k \rightarrow \infty} \sum_{i=0}^{m-1} l_i \cdot (\xi_k(t_{i+1}) - \xi_k(t_i)). \end{aligned}$$

This together with the Fatou's lemma and the fact that $P(\mathcal{A}_{\mathcal{N}} \cap \tilde{\Omega}) = 1$ yields

$$\begin{aligned} \mathbb{E} \sum_{i=0}^{m-1} l_i \cdot (\xi^*(t_{i+1}) - \xi^*(t_i)) &\leq \mathbb{E} \liminf_{k \rightarrow \infty} \sum_{i=0}^{m-1} l_i \cdot (\xi_k(t_{i+1}) - \xi_k(t_i)) \\ &\leq \liminf_{k \rightarrow \infty} \mathbb{E} \sum_{i=0}^{m-1} l_i \cdot (\xi_k(t_{i+1}) - \xi_k(t_i)) \\ &\leq \liminf_{k \rightarrow \infty} \mathbb{E} \int_t^T c(s) e^{-\int_t^s \alpha(r) dr} d\xi_k(s-t). \end{aligned}$$

Letting $\|\Pi\| \rightarrow 0$, $t_m \uparrow T - t$ so that each partition in the sequence is contained in the next one, by the monotone convergence theorem, we get (77).

From (76) and (77)

$$J_{xt}(v^*) \leq \liminf_{k \rightarrow \infty} J_{xt}(\eta_k) = u(x, t).$$

On the other hand, $J_{xt}(v^*) \geq u(x, t)$ because $v^* \in \mathcal{V}$ and hence $J_{xt}(v^*) = u(x, t)$ so v^* is an optimal control. \square

REFERENCES

- [1] Budhiraja, A., Ross, K., *Existence of optimal controls for singular control problems with state constraints*, Ann. Appl. Probab. **16**, No. 4 (2006), 2235–2255.
- [2] Chow, P. L., Menaldi, J. L., Robin, M., *Additive control of stochastic linear systems with finite horizon*, SIAM J. Control Optim. **23**, No.6 (1985), 858–899.
- [3] Dufour, F., Miller, B., *Singular stochastic control problems*, SIAM J. Control Optim. **43**, No. 2 (2004), 708–730.
- [4] Evans, L. C., *Partial Differential Equations*, American Mathematical Society, Providence, RI, 1998.
- [5] Fleming, W. H., Soner, H. M., *Controlled Markov Processes and Viscosity Solutions*, Springer, New York, 2006.
- [6] Haussman, U. G., Suo, W., *Singular optimal stochastic controls. I. Existence*, SIAM J. Control Optim. **33**, No. 3 (1995), 916–936.
- [7] Karatzas, I., Shreve, S. E., *Brownian Motion and Stochastic Calculus*, Springer-Verlag, New York, 1988.
- [8] Kruk, Ł., *Optimal policies for n-dimensional singular stochastic control problems, Part I: The Skorokhod problem*, SIAM J. Control Optim. **38**, No. 5 (2000), 1603–1622.

-
- [9] Kruk, Ł., *Optimal policies for n-dimensional singular stochastic control problems, Part II: The radially symmetric case*. *Ergodic control*, SIAM J. Control Optim. **39**, No. 2 (2000), 635–659.
 - [10] Krylov, N. V., *Controlled Diffusion Processes*, Springer-Verlag, New York, 1980.
 - [11] Menaldi, J. L., Taksar, M. I., *Optimal correction problem of a multidimensional stochastic system*, Automatica J. IFAC **25**, No. 2 (1989), 223–232.
 - [12] Rudin, W., *Functional Analysis*, McGraw-Hill Book Company, New York, 1991.
 - [13] Rudin, W., *Principles of Mathematical Analysis*, McGraw-Hill Book Company, New York, 1976.
 - [14] Soner, H. M., Shreve, S. E., *Regularity of the value function for a two-dimensional singular stochastic control problem*, SIAM J. Control Optim. **27** (1989), 876–907.
 - [15] Soner, H. M., Shreve, S. E., *A free boundary problem related to singular stochastic control*, *Applied stochastic analysis (London, 1989)*, Stochastics Monogr. 5, Gordon and Breach, New York, 1991, 265–301.
 - [16] Soner, H. M., Shreve, S. E., *A free boundary problem related to singular stochastic control: the parabolic case*, *Comm. Partial Differential Equations* **16** (1991), 373–424.
 - [17] S. A. Williams, P. L. Chow and J. L. Menaldi, *Regularity of the free boundary in singular stochastic control*, *J. Differential Equations* **111** (1994), 175–201.
 - [18] http://en.wikipedia.org/wiki/Gronwall's_inequality, 24.09.2013.

Marcin Boryc
Institute of Mathematics
Maria Curie-Skłodowska University
pl. M. Curie-Skłodowskiej 1
20-031 Lublin
Poland
e-mail: borycmarcin@gmail.com

Łukasz Kruk
Institute of Mathematics
Maria Curie-Skłodowska University
pl. M. Curie-Skłodowskiej 1
20-031 Lublin
Poland
e-mail: lkruk@hektor.umcs.lublin.pl

Received April 24, 2014