

doi: 10.17951/a.2016.70.2.9

ANNALES
UNIVERSITATIS MARIAE CURIE-SKŁODOWSKA
LUBLIN – POLONIA

VOL. LXX, NO. 2, 2016

SECTIO A

9–27

ALBO CARLOS CAVALHEIRO

Existence and uniqueness of solutions for a class of degenerate nonlinear elliptic equations

ABSTRACT. In this work we are interested in the existence and uniqueness of solutions for the Navier problem associated to the degenerate nonlinear elliptic equations

$$\begin{aligned} \Delta(v(x)|\Delta u|^{p-2}\Delta u) - \sum_{j=1}^n D_j[\omega_1(x)\mathcal{A}_j(x, u, \nabla u)] + b(x, u, \nabla u)\omega_2(x) \\ = f_0(x) - \sum_{j=1}^n D_j f_j(x), \quad \text{in } \Omega \end{aligned}$$

in the setting of the weighted Sobolev spaces.

1. Introduction. In this work we prove the existence and uniqueness of (weak) solutions in the weighted Sobolev space

$$X = W^{2,p}(\Omega, v) \cap W_0^{1,p}(\Omega, \omega_1, \omega_2)$$

(see Definition 2.4 and Definition 2.5) for the Navier problem

$$(P) \begin{cases} Lu(x) = f_0(x) - \sum_{j=1}^n D_j f_j(x), & \text{in } \Omega \\ u(x) = \Delta u(x) = 0, & \text{on } \partial\Omega \end{cases}$$

2010 *Mathematics Subject Classification.* 35J70, 35J60.

Key words and phrases. Degenerate nonlinear elliptic equations, weighted Sobolev spaces.

where L is the partial differential operator

$$Lu(x) = \Delta(v(x) |\Delta u|^{p-2} \Delta u) - \sum_{j=1}^n D_j [\omega_1(x) \mathcal{A}_j(x, u(x), \nabla u(x))] \\ + b(x, u, \nabla u) \omega_2(x)$$

where $D_j = \partial/\partial x_j$, Ω is a bounded open set in \mathbb{R}^n , ω_1 , ω_2 and v are three weight functions, Δ is the Laplacian operator, $1 < p < \infty$ and the functions $\mathcal{A}_j : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ ($j = 1, \dots, n$) and $b : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfy the following assumptions:

(H1) The function $x \mapsto \mathcal{A}_j(x, \eta, \xi)$ is measurable on Ω for all $(\eta, \xi) \in \mathbb{R} \times \mathbb{R}^n$. The function $(\eta, \xi) \mapsto \mathcal{A}_j(x, \eta, \xi)$ is continuous on $\mathbb{R} \times \mathbb{R}^n$ for almost all $x \in \Omega$.

(H2) There exists a constant $\theta_1 > 0$ such that

$$[\mathcal{A}(x, \eta, \xi) - \mathcal{A}(x, \tilde{\eta}, \tilde{\xi})] \cdot (\xi - \tilde{\xi}) \geq \theta_1 |\xi - \tilde{\xi}|^p,$$

whenever $\xi, \tilde{\xi} \in \mathbb{R}^n$, $\xi \neq \tilde{\xi}$, where $\mathcal{A}(x, \eta, \xi) = (\mathcal{A}_1(x, \eta, \xi), \dots, \mathcal{A}_n(x, \eta, \xi))$ (where a dot denotes here the Euclidean scalar product in \mathbb{R}^n).

(H3) $\mathcal{A}(x, \eta, \xi) \cdot \xi \geq \lambda_1 |\xi|^p + \Lambda_1 |\eta|^p$, where λ_1 and Λ_1 are nonnegative constants.

(H4) $|\mathcal{A}(x, \eta, \xi)| \leq K_1(x) + h_1(x) |\eta|^{p/p'} + h_2(x) |\xi|^{p/p'}$, where K_1, h_1 and h_2 are nonnegative functions, with h_1 and $h_2 \in L^\infty(\Omega)$, and $K_1 \in L^{p'}(\Omega, \omega_1)$ (with $1/p + 1/p' = 1$).

(H5) The function $x \mapsto b(x, \eta, \xi)$ is measurable on Ω for all $(\eta, \xi) \in \mathbb{R} \times \mathbb{R}^n$. The function $(\eta, \xi) \mapsto b(x, \eta, \xi)$ is continuous on $\mathbb{R} \times \mathbb{R}^n$ for almost all $x \in \Omega$.

(H6) There exists a constant $\theta_2 > 0$ such that

$$[b(x, \eta, \xi) - b(x, \tilde{\eta}, \tilde{\xi})](\eta - \tilde{\eta}) \geq \theta_2 |\eta - \tilde{\eta}|^p,$$

whenever $\eta, \tilde{\eta} \in \mathbb{R}$, $\eta \neq \tilde{\eta}$.

(H7) $b(x, \eta, \xi) \eta \geq \lambda_2 |\xi|^p + \Lambda_2 |\eta|^p$, where $\lambda_2 \geq 0$ and $\Lambda_2 > 0$ are constants.

(H8) $|b(x, \eta, \xi)| \leq K_2(x) + h_3(x) |\eta|^{p/p'} + h_4(x) |\xi|^a$, where K_2, h_3 and h_4 are nonnegative functions, with $K_2 \in L^{p'}(\Omega, \omega_2)$, h_3 and $h_4 \in L^\infty(\Omega)$, and $a = (p-1)/q'$, where $1 < q < \infty$ ($1/q + 1/q' = 1$).

(H9) $\lambda_1 + \lambda_2 > 0$.

By a *weight*, we shall mean a locally integrable function ω on \mathbb{R}^n such that $\omega(x) > 0$ for a.e. $x \in \mathbb{R}^n$. Every weight ω gives rise to a measure on the measurable subsets on \mathbb{R}^n through integration. This measure will be denoted by μ . Thus, $\mu(E) = \int_E \omega(x) dx$ for measurable sets $E \subset \mathbb{R}^n$.

In general, the Sobolev spaces $W^{k,p}(\Omega)$ without weights occur as spaces of solutions for elliptic and parabolic partial differential equations. For degenerate partial differential equations, i.e., equations with various types of singularities in the coefficients, it is natural to look for solutions in weighted Sobolev spaces (see [1], [2], [4], [8] and [13]).

A class of weights, which is particularly well understood, is the class of A_p -weights (or Muckenhoupt class) that was introduced by B. Muckenhoupt (see [10]). These classes have found many useful applications in harmonic analysis (see [12]). Another reason for studying A_p -weights is the fact that powers of the distance to submanifolds of \mathbb{R}^n often belong to A_p (see [9]). There are, in fact, many interesting examples of weights (see [8] for p -admissible weights).

In the non-degenerate case (i.e. with $\omega(x) \equiv 1$), for all $f \in L^p(\Omega)$ the Poisson equation associated with the Dirichlet problem

$$\begin{cases} -\Delta u = f(x), & \text{in } \Omega \\ u(x) = 0, & \text{on } \partial\Omega \end{cases}$$

is uniquely solvable in $W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)$ (see [7]), and the nonlinear Dirichlet problem

$$\begin{cases} -\Delta_p u = f(x), & \text{in } \Omega \\ u(x) = 0, & \text{on } \partial\Omega \end{cases}$$

is uniquely solvable in $W_0^{1,p}(\Omega)$ (see [3]), where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ is the p -Laplacian operator. In the degenerate case, the weighted p -Biharmonic operator has been studied by many authors (see [11] and the references therein), and the degenerated p -Laplacian has been studied in [4]. The problem with degenerated p -Laplacian and p -Biharmonic operators

$$\begin{cases} \Delta(\omega(x)|\Delta u|^{p-2} \Delta u) - \operatorname{div}[\omega(x)|\nabla u|^{p-2} \nabla u] = f(x) - \operatorname{div}(G(x)), & \text{in } \Omega \\ u(x) = \Delta u(x) = 0, & \text{in } \partial\Omega \end{cases}$$

has been studied by the author in [2].

The following theorem will be proved in Section 3.

Theorem 1.1. *Assume (H1)–(H8). If $\omega_1, \omega_2, v \in A_p$ (with $1 < p < \infty$), $\omega_1 \leq \omega_2$ a.e., $\omega_2/\omega_1 \in L^q(\Omega, \omega_1)$ ($1 < q < \infty$) $f_0/\omega_2 \in L^{p'}(\Omega, \omega_2)$ and $f_j/\omega_1 \in L^{p'}(\Omega, \omega_1)$ ($j = 1, \dots, n$), then the problem (P) has a unique solution $u \in X = W^{2,p}(\Omega, v) \cap W_0^{1,p}(\Omega, \omega_1, \omega_2)$. Moreover, we have*

$$\|u\|_X \leq \frac{1}{\gamma^{p'/p}} \left(\left\| \frac{f_0}{\omega_2} \right\|_{L^{p'}(\Omega, \omega_2)} + \sum_{j=1}^n \left\| \frac{f_j}{\omega_1} \right\|_{L^{p'}(\Omega, \omega_1)} \right)^{p'/p},$$

where $\gamma = \min\{\lambda_1 + \lambda_2, \Lambda_2, 1\}$.

Corollary 1.2. *Let the assumptions of Theorem 1.1 be fulfilled. If $u_1, u_2 \in X = W^{2,p}(\Omega, v) \cap W_0^{1,p}(\Omega, \omega_1, \omega_2)$ are solutions of*

$$(P_1) \begin{cases} Lu_1(x) = f_0(x) - \sum_{j=1}^n D_j f_j(x), & \text{in } \Omega \\ u_1(x) = \Delta u_1(x) = 0, & \text{on } \partial\Omega \end{cases}$$

$$(P_2) \begin{cases} Lu_2(x) = \tilde{f}_0(x) - \sum_{j=1}^n D_j \tilde{f}_j(x), & \text{in } \Omega \\ u_2(x) = \Delta u_2(x) = 0, & \text{on } \partial\Omega \end{cases}$$

then

$$\|u_1 - u_2\|_X \leq \frac{1}{\varepsilon^{p-1}} \left(\left\| \frac{f_0 - \tilde{f}_0}{\omega_2} \right\|_{L^{p'}(\Omega, \omega_2)} + \sum_{j=1}^n \left\| \frac{f_j - \tilde{f}_j}{\omega_1} \right\|_{L^{p'}(\Omega, \omega_1)} \right)^{p-1},$$

where $\varepsilon = \min\{\beta_p, \theta_1, \theta_2\}$ (with β_p as in Lemma 2.2 (b)).

2. Definitions and basic results. Let ω be a locally integrable nonnegative function in \mathbb{R}^n and assume that $0 < \omega(x) < \infty$ almost everywhere. We say that ω belongs to the Muckenhoupt class A_p , $1 < p < \infty$, or that ω is an A_p -weight, if there is a constant $C = C_{p,\omega}$ such that

$$\left(\frac{1}{|B|} \int_B \omega(x) dx \right) \left(\frac{1}{|B|} \int_B \omega^{1/(1-p)}(x) dx \right)^{p-1} \leq C$$

for all balls $B \subset \mathbb{R}^n$, where $|\cdot|$ denotes the n -dimensional Lebesgue measure in \mathbb{R}^n . If $1 < q \leq p$, then $A_q \subset A_p$ (see [6], [8] or [12] for more information about A_p -weights). The weight ω satisfies the doubling condition if there exists a positive constant C such that $\mu(B(x; 2r)) \leq C\mu(B(x; r))$ for every ball $B = B(x; r) \subset \mathbb{R}^n$, where $\mu(B) = \int_B \omega(x) dx$. If $\omega \in A_p$, then μ is doubling (see Corollary 15.7 in [8]).

As an example of A_p -weight, the function $\omega(x) = |x|^\alpha$, $x \in \mathbb{R}^n$, is in A_p if and only if $-n < \alpha < n(p-1)$ (see Corollary 4.4, Chapter IX in [12]).

If $\omega \in A_p$, then

$$\left(\frac{|E|}{|B|} \right)^p \leq C \frac{\mu(E)}{\mu(B)}$$

whenever B is a ball in \mathbb{R}^n and E is a measurable subset of B (see 15.5 *strong doubling property* in [8]). Therefore, if $\mu(E) = 0$, then $|E| = 0$.

Definition 2.1. Let ω be a weight, and let $\Omega \subset \mathbb{R}^n$ be open. For $0 < p < \infty$ we define $L^p(\Omega, \omega)$ as the set of measurable functions f on Ω such that

$$\|f\|_{L^p(\Omega, \omega)} = \left(\int_{\Omega} |f(x)|^p \omega(x) dx \right)^{1/p} < \infty.$$

If $\omega \in A_p$, $1 < p < \infty$, then $\omega^{-1/(p-1)}$ is locally integrable and we have $L^p(\Omega, \omega) \subset L^1_{\text{loc}}(\Omega)$ for every open set Ω (see Remark 1.2.4 in [13]). It thus makes sense to talk about weak derivatives of functions in $L^p(\Omega, \omega)$.

Definition 2.2. Let $\Omega \subset \mathbb{R}^n$ be open, k be a nonnegative integer and $\omega \in A_p$ ($1 < p < \infty$). We define the weighted Sobolev space $W^{k,p}(\Omega, \omega)$ as

the set of functions $u \in L^p(\Omega, \omega)$ with weak derivatives $D^\alpha u \in L^p(\Omega, \omega)$ for $1 \leq |\alpha| \leq k$. The norm of u in $W^{k,p}(\Omega, \omega)$ is defined by

$$(2.1) \quad \|u\|_{W^{k,p}(\Omega, \omega)} = \left(\int_{\Omega} |u(x)|^p \omega(x) dx + \sum_{1 \leq |\alpha| \leq k} \int_{\Omega} |D^\alpha u(x)|^p \omega(x) dx \right)^{1/p}.$$

We also define $W_0^{k,p}(\Omega, \omega)$ as the closure of $C_0^\infty(\Omega)$ with respect to the norm $\|\cdot\|_{W^{k,p}(\Omega, \omega)}$.

If $\omega \in A_p$, then $W^{k,p}(\Omega, \omega)$ is the closure of $C^\infty(\Omega)$ with respect to the norm (2.1) (see Theorem 2.1.4 in [13]). The spaces $W^{k,p}(\Omega, \omega)$ and $W_0^{k,p}(\Omega, \omega)$ are Banach spaces.

It is evident that the weight function ω which satisfies $0 < c_1 \leq \omega(x) \leq c_2$ for $x \in \Omega$ (c_1 and c_2 positive constants), gives nothing new (the space $W_0^{k,p}(\Omega, \omega)$ is then identical with the classical Sobolev space $W_0^{k,p}(\Omega)$). Consequently, we shall be interested above all in such weight functions ω which either vanish somewhere in $\Omega \cup \partial\Omega$ or increase to infinity (or both).

Definition 2.3. Let $\Omega \subset \mathbb{R}^n$ be open, $1 < p < \infty$, and let ω_1 and ω_2 be A_p -weights. We define the weighted Sobolev space $W^{1,p}(\Omega, \omega_1, \omega_2)$ as the set of functions $u \in L^p(\Omega, \omega_2)$ with weak derivatives $D_j u \in L^p(\Omega, \omega_1)$, for $j = 1, \dots, n$. The norm of u in $W^{1,p}(\Omega, \omega_1, \omega_2)$ is given by

$$(2.2) \quad \|u\|_{W^{1,p}(\Omega, \omega_1, \omega_2)} = \left(\int_{\Omega} |u(x)|^p \omega_2(x) dx + \sum_{j=1}^n \int_{\Omega} |D_j u(x)|^p \omega_1(x) dx \right)^{1/p}.$$

The space $W_0^{1,p}(\Omega, \omega_1, \omega_2)$ is the closure of $C_0^\infty(\Omega)$ with respect to the norm (2.2). The dual space of $W_0^{1,p}(\Omega, \omega_1, \omega_2)$ is the space

$$\begin{aligned} [W_0^{1,p}(\Omega, \omega_1, \omega_2)]^* &= W^{-1,p'}(\Omega, \omega_1, \omega_2) \\ &= \left\{ T = f_0 - \operatorname{div} F : F = (f_1, \dots, f_n), \frac{f_0}{\omega_2} \in L^{p'}(\Omega, \omega_2), \frac{f_j}{\omega_1} \in L^{p'}(\Omega, \omega_1) \right\}. \end{aligned}$$

In this article we use the following results.

Theorem 2.1. Let $\omega \in A_p$, $1 < p < \infty$, and let Ω be a bounded open set in \mathbb{R}^n . If $u_m \rightarrow u$ in $L^p(\Omega, \omega)$, then there exist a subsequence $\{u_{m_k}\}$ and a function $\Phi \in L^p(\Omega, \omega)$ such that

- (i) $u_{m_k}(x) \rightarrow u(x)$, $m_k \rightarrow \infty$, μ -a.e. on Ω ;
- (ii) $|u_{m_k}(x)| \leq \Phi(x)$, μ -a.e. on Ω ;
- (where $\mu(E) = \int_E \omega(x) dx$).

Proof. The proof of this theorem follows the lines of Theorem 2.8.1 in [5]. \square

Lemma 2.2. *Let $1 < p < \infty$.*

(a) *There exists a constant α_p such that*

$$\left| |x|^{p-2}x - |y|^{p-2}y \right| \leq \alpha_p |x - y| (|x| + |y|)^{p-2}, \forall x, y \in \mathbb{R}^n;$$

(b) *There exist two positive constants β_p, γ_p such that for every $x, y \in \mathbb{R}^n$*

$$\begin{aligned} \beta_p (|x| + |y|)^{p-2} |x - y|^2 &\leq (|x|^{p-2}x - |y|^{p-2}y)(x - y) \\ &\leq \gamma_p (|x| + |y|)^{p-2} |x - y|^2. \end{aligned}$$

Proof. See [3], Proposition 17.2 and Proposition 17.3. \square

Definition 2.4. We denote $X = W^{2,p}(\Omega, v) \cap W_0^{1,p}(\Omega, \omega_1, \omega_2)$ with the norm

$$\|u\|_X = \left(\int_{\Omega} |u|^p \omega_2 dx + \int_{\Omega} |\nabla u|^p \omega_1 dx + \int_{\Omega} |\Delta u|^p v dx \right)^{1/p}.$$

Definition 2.5. We say that an element

$$u \in X = W^{2,p}(\Omega, v) \cap W_0^{1,p}(\Omega, \omega_1, \omega_2)$$

is a (weak) solution of problem (P) if, for all $\varphi \in X$,

$$\begin{aligned} \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta \varphi v dx + \sum_{j=1}^n \int_{\Omega} \omega_1 \mathcal{A}_j(x, u(x), \nabla u(x)) D_j \varphi(x) dx \\ + \int_{\Omega} b(x, u, \nabla u) \varphi \omega_2 dx = \int_{\Omega} f_0(x) \varphi(x) dx + \sum_{j=1}^n \int_{\Omega} f_j(x) D_j \varphi(x) dx. \end{aligned}$$

3. Proof of Theorem 1.1. The basic idea is to reduce the problem (P) to an operator equation $Au = T$ and apply the theorem below.

Theorem 3.1. *Let $A : X \rightarrow X^*$ be a monotone, coercive and hemicontinuous operator on the real, separable, reflexive Banach space X . Then the following assertions hold:*

- (a) *For each $T \in X^*$ the equation $Au = T$ has a solution $u \in X$;*
- (b) *If the operator A is strictly monotone, then equation $Au = T$ is uniquely solvable in X .*

Proof. See Theorem 26.A in [15]. \square

To prove the existence of solutions, we define $B, B_1, B_2, B_3 : X \times X \rightarrow \mathbb{R}$ and $T : X \rightarrow \mathbb{R}$ by

$$\begin{aligned} B(u, \varphi) &= B_1(u, \varphi) + B_2(u, \varphi) + B_3(u, \varphi), \\ B_1(u, \varphi) &= \sum_{j=1}^n \int_{\Omega} \omega_1 \mathcal{A}_j(x, u, \nabla u) D_j \varphi dx = \int_{\Omega} \omega_1 \mathcal{A}(x, u, \nabla u) \cdot \nabla \varphi dx, \end{aligned}$$

$$\begin{aligned}
B_2(u, \varphi) &= \int_{\Omega} b(x, u, \nabla u) \varphi \omega_2 dx, \\
B_3(u, \varphi) &= \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta \varphi v dx, \\
T(\varphi) &= \int_{\Omega} f_0(x) \varphi(x) dx + \sum_{j=1}^n \int_{\Omega} f_j(x) D_j \varphi(x) dx.
\end{aligned}$$

Then $u \in X$ is a (weak) solution to problem (P) if for all $\varphi \in X$

$$B(u, \varphi) = B_1(u, \varphi) + B_2(u, \varphi) + B_3(u, \varphi) = T(\varphi).$$

Step 1. For $j = 1, \dots, n$ we define the operator $F_j : X \rightarrow L^{p'}(\Omega, \omega_1)$ by

$$(F_j u)(x) = \mathcal{A}_j(x, u(x), \nabla u(x)).$$

We now show that operator F_j is bounded and continuous.

(i) Using (H4) and $\omega_1 \leq \omega_2$, we obtain

$$\begin{aligned}
\|F_j u\|_{L^{p'}(\Omega, \omega_1)}^{p'} &= \int_{\Omega} |F_j u(x)|^{p'} \omega_1 dx \\
&= \int_{\Omega} |\mathcal{A}_j(x, u, \nabla u)|^{p'} \omega_1 dx \\
&\leq \int_{\Omega} \left(K_1 + h_1 |u|^{p/p'} + h_2 |\nabla u|^{p/p'} \right)^{p'} \omega_1 dx \\
&\leq C_p \int_{\Omega} \left[(K_1^{p'} + h_1^{p'} |u|^p + h_2^{p'} |\nabla u|^p) \omega_1 \right] dx \\
(3.1) \quad &\leq C_p \left[\int_{\Omega} K_1^{p'} \omega_1 dx + \int_{\Omega} h_1^{p'} |u|^p \omega_2 dx + \int_{\Omega} h_2^{p'} |\nabla u|^p \omega_1 dx \right],
\end{aligned}$$

where the constant C_p depends only on p . We have,

$$\begin{aligned}
\int_{\Omega} h_1^{p'} |u|^p \omega_2 dx &\leq \|h_1\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |u|^p \omega_2 dx \\
&\leq \|h_1\|_{L^\infty(\Omega)}^{p'} \|u\|_X^p,
\end{aligned}$$

and

$$\begin{aligned}
\int_{\Omega} h_2^{p'} |\nabla u|^p \omega_1 dx &\leq \|h_2\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |\nabla u|^p \omega_1 dx \\
&\leq \|h_2\|_{L^\infty(\Omega)}^{p'} \|u\|_X^p.
\end{aligned}$$

Therefore, by (3.1) we obtain

$$\|F_j u\|_{L^{p'}(\Omega, \omega_1)} \leq C_p \left(\|K_1\|_{L^{p'}(\Omega, \omega_1)} + (\|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Omega)}) \|u\|_X^{p/p'} \right),$$

and hence the boundedness.

(ii) Let $u_m \rightarrow u$ in X as $m \rightarrow \infty$. We need to show that $F_j u_m \rightarrow F_j u$ in $L^{p'}(\Omega, \omega_1)$. We will apply the Lebesgue Dominated Theorem. If $u_m \rightarrow u$ in X , then $u_m \rightarrow u$ in $L^p(\Omega, \omega_2)$ and $|\nabla u_m| \rightarrow |\nabla u|$ in $L^p(\Omega, \omega_1)$. Using Theorem 2.1, there exist a subsequence $\{u_{m_k}\}$ and two functions $\Phi_1 \in L^p(\Omega, \omega_1)$ and $\Phi_2 \in L^p(\Omega, \omega_2)$ such that

$$\begin{aligned} u_{m_k}(x) &\rightarrow u(x), \quad \mu_2\text{-a.e. in } \Omega, \\ |u_{m_k}(x)| &\leq \Phi_2(x), \quad \mu_2\text{-a.e. in } \Omega, \\ |\nabla u_{m_k}(x)| &\rightarrow |\nabla u(x)|, \quad \mu_1\text{-a.e. in } \Omega, \\ |\nabla u_{m_k}(x)| &\leq \Phi_1(x), \quad \mu_1\text{-a.e. in } \Omega. \end{aligned}$$

where $\mu_i = \int_E \omega_i(x) dx$ ($i = 1, 2$). Hence, using (H4) and $\omega_1 \leq \omega_2$, we obtain

$$\begin{aligned} \|F_j u_{m_k} - F_j u\|_{L^{p'}(\Omega, \omega_1)}^{p'} &= \int_{\Omega} |F_j u_{m_k}(x) - F_j u(x)|^{p'} \omega_1 dx \\ &= \int_{\Omega} |\mathcal{A}_j(x, u_{m_k}, \nabla u_{m_k}) - \mathcal{A}_j(x, u, \nabla u)|^{p'} \omega_1 dx \\ &\leq C_p \int_{\Omega} \left(|\mathcal{A}_j(x, u_{m_k}, \nabla u_{m_k})|^{p'} + |\mathcal{A}_j(x, u, \nabla u)|^{p'} \right) \omega_1 dx \\ &\leq C_p \left[\int_{\Omega} \left(K_1 + h_1 |u_{m_k}|^{p/p'} + h_2 |\nabla u_{m_k}|^{p/p'} \right)^{p'} \omega_1 dx \right. \\ &\quad \left. + \int_{\Omega} \left(K_1 + h_1 |u|^{p/p'} + h_2 |\nabla u|^{p/p'} \right)^{p'} \omega_1 dx \right] \\ &\leq 2 C_p \int_{\Omega} \left(K_1 + h_1 \Phi_2^{p/p'} + h_2 \Phi_1^{p/p'} \right)^{p'} \omega_1 dx \\ &\leq 2 C_p \left[\int_{\Omega} K_1^{p'} \omega_1 dx + \int_{\Omega} h_1^{p'} \Phi_2^p \omega_1 dx + \int_{\Omega} h_2^{p'} \Phi_1^p \omega_1 dx \right] \\ &\leq 2 C_p \left[\|K_1\|_{L^{p'}(\Omega, \omega_1)}^{p'} + \|h_1\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} \Phi_2^p \omega_2 dx \right. \\ &\quad \left. + \|h_2\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} \Phi_1^p \omega_1 dx \right] \\ &\leq 2 C_p \left[\|K_1\|_{L^{p'}(\Omega, \omega_1)}^{p'} + \|h_1\|_{L^\infty(\Omega)}^{p'} \|\Phi_2\|_{L^p(\Omega, \omega_2)}^p \right. \\ &\quad \left. + \|h_2\|_{L^\infty(\Omega)}^{p'} \|\Phi_1\|_{L^p(\Omega, \omega_1)}^p \right]. \end{aligned}$$

By condition (H1), we have

$$F_j u_{m_k}(x) = \mathcal{A}_j(x, u_{m_k}(x), \nabla u_{m_k}(x)) \rightarrow \mathcal{A}_j(x, u(x), \nabla u(x)) = F_j u(x),$$

as $m_k \rightarrow +\infty$. Therefore, by the Lebesgue Dominated Convergence Theorem, we obtain

$$\|F_j u_{m_k} - F_j u\|_{L^{p'}(\Omega, \omega_1)} \rightarrow 0,$$

that is,

$$F_j u_{m_k} \rightarrow F_j u \text{ in } L^{p'}(\Omega, \omega_1).$$

By the Convergence Principle in Banach spaces (see Proposition 10.13 in [14]), we have

$$(3.2) \quad F_j u_m \rightarrow F_j u \text{ in } L^{p'}(\Omega, \omega_1).$$

Step 2. We define the operator

$$\begin{aligned} G : X &\rightarrow L^{p'}(\Omega, v) \\ (Gu)(x) &= |\Delta u(x)|^{p-2} \Delta u(x). \end{aligned}$$

The operator G is continuous and bounded. In fact:

(i) We have

$$\begin{aligned} \|Gu\|_{L^{p'}(\Omega, v)}^{p'} &= \int_{\Omega} |\Delta u|^{p-2} \Delta u|^{p'} v \, dx \\ &= \int_{\Omega} |\Delta u|^{(p-2)p'} |\Delta u|^{p'} v \, dx \\ &= \int_{\Omega} |\Delta u|^p v \, dx \\ &\leq \|u\|_X^p. \end{aligned}$$

Hence, $\|Gu\|_{L^{p'}(\Omega, v)} \leq \|u\|_X^{p/p'}$.

(ii) If $u_m \rightarrow u$ in X , then $\Delta u_m \rightarrow \Delta u$ in $L^p(\Omega, v)$. By Theorem 2.1, there exist a subsequence $\{u_{m_k}\}$ and a function $\Phi_3 \in L^p(\Omega, v)$ such that

$$\begin{aligned} \Delta u_{m_k}(x) &\rightarrow \Delta u(x), \quad \mu_3\text{-a.e. in } \Omega \\ |\Delta u_{m_k}(x)| &\leq \Phi_3(x), \quad \mu_3\text{-a.e. in } \Omega, \end{aligned}$$

where $\mu_3(E) = \int_E v(x) \, dx$. Hence, using Lemma 2.2 (a), we obtain, if $p \neq 2$

$$\begin{aligned} \|Gu_{m_k} - Gu\|_{L^{p'}(\Omega, v)}^{p'} &= \int_{\Omega} |Gu_{m_k} - Gu|^{p'} v \, dx \\ &= \int_{\Omega} \left| |\Delta u_{m_k}|^{p-2} \Delta u_{m_k} - |\Delta u|^{p-2} \Delta u \right|^{p'} v \, dx \\ &\leq \int_{\Omega} \left[\alpha_p |\Delta u_{m_k} - \Delta u| (|\Delta u_{m_k}| + |\Delta u|)^{(p-2)} \right]^{p'} v \, dx \\ &\leq \alpha_p^{p'} \int_{\Omega} |\Delta u_{m_k} - \Delta u|^{p'} (2\Phi_3)^{(p-2)p'} v \, dx \end{aligned}$$

$$\begin{aligned} &\leq \alpha_p^{p'} 2^{(p-2)p'} \left(\int_{\Omega} |\Delta u_{m_k} - \Delta u|^p v \, dx \right)^{\frac{p'}{p}} \left(\int_{\Omega} \Phi_3^{\frac{(p-2)p p'}{p-p'}} v \, dx \right)^{\frac{p-p'}{p}} \\ &\leq \alpha_p^{p'} 2^{(p-2)p'} \|u_{m_k} - u\|_X^{p'} \|\Phi\|_{L^p(\Omega, v)}^{p-p'}, \end{aligned}$$

since $(p-2)pp'/(p-p') = p$ if $p \neq 2$. If $p = 2$, we have

$$\begin{aligned} \|Gu_{m_k} - Gu\|_{L^2(\Omega, v)}^2 &= \int_{\Omega} |\Delta u_{m_k} - \Delta u|^2 v \, dx \\ &\leq \|u_{m_k} - u\|_X^2. \end{aligned}$$

Therefore (for $1 < p < \infty$), by the Lebesgue Dominated Convergence Theorem, we obtain

$$\|Gu_{m_k} - Gu\|_X \rightarrow 0,$$

that is, $Gu_{m_k} \rightarrow Gu$ in $L^{p'}(\Omega, v)$. By the Convergence Principle in Banach spaces (see Proposition 10.13 in [14]), we have

$$(3.3) \quad Gu_m \rightarrow Gu \text{ in } L^{p'}(\Omega, v).$$

Step 3. We define the operator $H : X \rightarrow L^{p'}(\Omega, \omega_2)$ by

$$(Hu)(x) = b(x, u(x), \nabla u(x)).$$

The operator H is continuous and bounded. In fact:

(i) Using (H8) and $a = (p-1)/q'$, we obtain

$$\begin{aligned} \|Hu\|_{L^{p'}(\Omega, \omega_2)}^{p'} &= \int_{\Omega} |Hu|^{p'} \omega_2 \, dx \\ &= \int_{\Omega} |b(x, u, \nabla u)|^{p'} \omega_2 \, dx \\ &\leq \int_{\Omega} \left(K_2 + h_3 |u|^{p/p'} + h_4 |\nabla u|^a \right)^{p'} \omega_2 \, dx \\ &\leq C_p \int_{\Omega} \left[(K_2^{p'} + h_3^{p'} |u|^p + h_4^{p'} |\nabla u|^{ap'}) \omega_2 \right] dx \\ &= C_p \left[\int_{\Omega} K_2^{p'} \omega_2 \, dx + \int_{\Omega} h_3^{p'} |u|^p \omega_2 \, dx + \int_{\Omega} h_4^{p'} |\nabla u|^{ap'} \omega_2 \, dx \right]. \end{aligned}$$

We have

$$\begin{aligned} \int_{\Omega} h_3^{p'} |u|^p \omega_2 &\leq \|h_3\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |u|^p \omega_2 \, dx \\ &\leq \|h_3\|_{L^\infty(\Omega)}^{p'} \|u\|_X^p, \end{aligned}$$

and

$$\int_{\Omega} h_4^{p'} |\nabla u|^{ap'} \omega_2 \, dx \leq \|h_4\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |\nabla u|^{p/q'} \omega_2 \, dx$$

$$\begin{aligned}
&= \|h_4\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |\nabla u|^{p/q'} \frac{\omega_2}{\omega_1} \omega_1 dx \\
&\leq \|h_4\|_{L^\infty(\Omega)}^{p'} \left(\int_{\Omega} |\nabla u|^p \omega_1 dx \right)^{1/q'} \left(\int_{\Omega} \left(\frac{\omega_1}{\omega_2} \right)^q \omega_1 dx \right)^{1/q} \\
&\leq \|h_4\|_{L^\infty(\Omega)}^{p'} \|u\|_X^{p/q'} \|\omega_2/\omega_1\|_{L^q(\Omega, \omega_1)}.
\end{aligned}$$

Hence,

$$\begin{aligned}
\|Hu\|_{L^{p'}(\Omega, \omega_2)} &\leq C_p \left[\|K_2\|_{L^{p'}(\Omega, \omega_2)} + \|h_3\|_{L^\infty(\Omega)} \|u\|_X^{p/p'} \right. \\
&\quad \left. + \|h_4\|_{L^\infty(\Omega)} \|\omega_2/\omega_1\|_{L^q(\Omega, \omega_1)}^{1/p'} \|u\|_X^{(p-1)/q'} \right].
\end{aligned}$$

(ii) By the same argument used in Step 1(ii) (and condition (H5)), we obtain analogously, if $u_m \rightarrow u$ in X , then

$$(3.4) \quad Hu_m \rightarrow Hu, \quad \text{in } L^{p'}(\Omega, \omega_2).$$

Step 4. We also have

$$\begin{aligned}
|T(\varphi)| &\leq \int_{\Omega} |f_0| |\varphi| dx + \sum_{j=1}^n \int_{\Omega} |f_j| |D_j \varphi| dx \\
&= \int_{\Omega} \frac{|f_0|}{\omega_2} |\varphi| \omega_2 dx + \sum_{j=1}^n \int_{\Omega} \frac{|f_j|}{\omega_1} |D_j \varphi| \omega_1 dx \\
&\leq \|f_0/\omega_2\|_{L^{p'}(\Omega, \omega_2)} \|\varphi\|_{L^p(\Omega, \omega_2)} \\
&\quad + \sum_{j=1}^n \|f_j/\omega_1\|_{L^{p'}(\Omega, \omega_1)} \|D_j \varphi\|_{L^p(\Omega, \omega_1)} \\
&\leq \left(\|f_0/\omega_2\|_{L^{p'}(\Omega, \omega_2)} + \sum_{j=1}^n \|f_j/\omega_1\|_{L^{p'}(\Omega, \omega_1)} \right) \|\varphi\|_X.
\end{aligned}$$

Moreover, using (H4), (H8) and the Hölder inequality, we also have

$$\begin{aligned}
|B(u, \varphi)| &\leq |B_1(u, \varphi)| + |B_2(u, \varphi)| + |B_3(u, \varphi)| \\
&\leq \sum_{j=1}^n \int_{\Omega} |\mathcal{A}_j(x, u, \nabla u)| |D_j \varphi| \omega_1 dx \\
(3.5) \quad &\quad + \int_{\Omega} |\Delta u|^{p-2} |\Delta u| |\Delta \varphi| v dx \\
&\quad + \int_{\Omega} |b(x, u, \nabla u)| |\varphi| \omega_2 dx.
\end{aligned}$$

In (3.5) we have

$$\begin{aligned}
& \int_{\Omega} |\mathcal{A}(x, u, \nabla u)| |\nabla \varphi| \omega_1 dx \\
& \leq \int_{\Omega} \left(K_1 + h_1 |u|^{p/p'} + h_2 |\nabla u|^{p/p'} \right) |\nabla \varphi| \omega_1 dx \\
& \leq \|K_1\|_{L^{p'}(\Omega, \omega_1)} \|\nabla \varphi\|_{L^p(\Omega, \omega_1)} + \|h_1\|_{L^\infty(\Omega)} \|u\|_{L^p(\Omega, \omega_2)}^{p/p'} \|\nabla \varphi\|_{L^p(\Omega, \omega_1)} \\
& \quad + \|h_2\|_{L^\infty(\Omega)} \|\nabla u\|_{L^p(\Omega, \omega_1)}^{p/p'} \|\nabla \varphi\|_{L^p(\Omega, \omega_1)} \\
& \leq \left(\|K_1\|_{L^{p'}(\Omega, \omega_1)} + (\|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Omega)}) \|u\|_X^{p/p'} \right) \|\varphi\|_X,
\end{aligned}$$

and

$$\begin{aligned}
\int_{\Omega} |\Delta u|^{p-2} |\Delta u| |\Delta \varphi| v dx &= \int_{\Omega} |\Delta u|^{p-1} |\Delta \varphi| v dx \\
&\leq \left(\int_{\Omega} |\Delta u|^p v dx \right)^{1/p'} \left(\int_{\Omega} |\Delta \varphi|^p v dx \right)^{1/p} \\
&\leq \|u\|_X^{p/p'} \|\varphi\|_X,
\end{aligned}$$

and

$$\begin{aligned}
\int_{\Omega} |b(x, u, \nabla u)| |\varphi| \omega_2 dx &\leq \int_{\Omega} \left(K_2 + h_3 |u|^{p/p'} + h_4 |\nabla u|^a \right) |\varphi| \omega_2 dx \\
&\leq \int_{\Omega} K_2 |\varphi| \omega_2 dx + \|h_3\|_{L^\infty(\Omega)} \int_{\Omega} |u|^{p/p'} |\varphi| \omega_2 dx \\
&\quad + \|h_4\|_{L^\infty(\Omega)} \int_{\Omega} |\nabla u|^a |\varphi| \omega_2 dx \\
&\leq \left(\|K_2\|_{L^{p'}(\Omega, \omega_2)} + \|h_3\|_{L^\infty(\Omega)} \|u\|_X^{p/p'} \right) \|\varphi\|_X \\
&\quad + \|h_4\|_{L^\infty(\Omega)} \left(\int_{\Omega} |\nabla u|^{ap'} \omega_2 dx \right)^{1/p'} \left(\int_{\Omega} |\varphi|^p \omega_2 dx \right)^{1/p} \\
&\leq \left(\|K_2\|_{L^{p'}(\Omega, \omega_2)} + \|h_3\|_{L^\infty(\Omega)} \|u\|_X^{p/p'} \right) \|\varphi\|_X \\
&\quad + \|h_4\|_{L^\infty(\Omega)} \left(\int_{\Omega} |\nabla u|^{p/q'} \frac{\omega_2}{\omega_1} dx \right)^{1/p'} \|\varphi\|_X \\
&\leq \left(\|K_2\|_{L^{p'}(\Omega, \omega_2)} + \|h_3\|_{L^\infty(\Omega)} \|u\|_X^{p/p'} \right) \|\varphi\|_X \\
&\quad + \|h_4\|_{L^\infty(\Omega)} \left(\int_{\Omega} |\nabla u|^p \omega_1 dx \right)^{1/(p'q')} \|\omega_2/\omega_1\|_{L^q(\Omega, \omega_1)}^{1/p'} \|\varphi\|_X
\end{aligned}$$

$$\begin{aligned} &\leq \left(\|K_2\|_{L^{p'}(\Omega, \omega_2)} + \|h_3\|_{L^\infty(\Omega)} \|u\|_X^{p/p'} \right. \\ &\quad \left. + \|h_4\|_{L^\infty(\Omega)} \|\omega_2/\omega_1\|_{L^q(\Omega, \omega_1)}^{1/p'} \|u\|_X^{p/(q'p')} \right) \|\varphi\|_X. \end{aligned}$$

Therefore, in (3.5) we obtain, for all $u, \varphi \in X$,

$$\begin{aligned} |B(u, \varphi)| &\leq \left[\|K_1\|_{L^{p'}(\Omega, \omega_1)} + \|K_2\|_{L^{p'}(\Omega, \omega_2)} \right. \\ &\quad + (\|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Omega)} + \|h_3\|_{L^\infty(\Omega)} + 1) \|u\|_X^{p/p'} \\ &\quad \left. + \|h_4\|_{L^\infty(\Omega)} \|\omega_2/\omega_1\|_{L^q(\Omega, \omega_1)}^{1/p'} \|u\|_X^{p/(p'q')} \right] \|\varphi\|_X. \end{aligned}$$

Since $B(u, \cdot)$ is linear, for each $u \in X$, there exists a linear and continuous operator $A : X \rightarrow X^*$ such that $\langle Au, \varphi \rangle = B(u, \varphi)$, for all $u, \varphi \in X$ (where $\langle f, x \rangle$ denotes the value of the linear functional f at the point x) and

$$\begin{aligned} \|Au\|_* &\leq \|K_1\|_{L^{p'}(\Omega, \omega_1)} + \|K_2\|_{L^{p'}(\Omega, \omega_2)} \\ &\quad + (\|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Omega)} + \|h_3\|_{L^\infty(\Omega)} + 1) \|u\|_X^{p/p'} \\ &\quad + \|h_4\|_{L^\infty(\Omega)} \|\omega_2/\omega_1\|_{L^q(\Omega, \omega_1)}^{1/p'} \|u\|_X^{p/(p'q')}. \end{aligned}$$

Consequently, problem (P) is equivalent to the operator equation

$$Au = T, \quad u \in X.$$

Step 5. Using condition (H2), (H6) and Lemma 2.2 (b), we have

$$\begin{aligned} \langle Au_1 - Au_2, u_1 - u_2 \rangle &= B(u_1, u_1 - u_2) - B(u_2, u_1 - u_2) \\ &= \int_{\Omega} \omega_1 \mathcal{A}(x, u_1, \nabla u_1) \cdot \nabla(u_1 - u_2) \, dx + \int_{\Omega} |\Delta u_1|^{p-2} \Delta u_1 \Delta(u_1 - u_2) v \, dx \\ &\quad + \int_{\Omega} b(x, u_1, \nabla u_1)(u_1 - u_2) \omega_2 \, dx - \int_{\Omega} \omega_1 \mathcal{A}(x, u_2, \nabla u_2) \cdot \nabla(u_1 - u_2) \, dx \\ &\quad - \int_{\Omega} |\Delta u_2|^{p-2} \Delta u_2 \Delta(u_1 - u_2) v \, dx - \int_{\Omega} b(x, u_2, \nabla u_2)(u_1 - u_2) \omega_2 \, dx \\ &= \int_{\Omega} \omega_1 \left(\mathcal{A}(x, u_1, \nabla u_1) - \mathcal{A}(x, u_2, \nabla u_2) \right) \cdot \nabla(u_1 - u_2) \, dx \\ &\quad + \int_{\Omega} (|\Delta u_1|^{p-2} \Delta u_1 - |\Delta u_2|^{p-2} \Delta u_2) \Delta(u_1 - u_2) v \, dx \\ &\quad + \int_{\Omega} (b(x, u_1, \nabla u_1) - b(x, u_2, \nabla u_2))(u_1 - u_2) \omega_2 \, dx \\ &\geq \theta_1 \int_{\Omega} \omega_1 |\nabla(u_1 - u_2)|^p \, dx + \beta_p \int_{\Omega} (|\Delta u_1| + |\Delta u_2|)^{p-2} |\Delta u_1 - \Delta u_2|^2 v \, dx \end{aligned}$$

$$\begin{aligned}
& + \theta_2 \int_{\Omega} |u_1 - u_2|^p \omega_2 dx \\
& \geq \theta_1 \int_{\Omega} \omega_1 |\nabla(u_1 - u_2)|^p dx + \beta_p \int_{\Omega} (|\Delta u_1 - \Delta u_2|)^{p-2} |\Delta u_1 - \Delta u_2|^2 v dx \\
& \quad + \theta_2 \int_{\Omega} |u_1 - u_2|^p \omega_2 dx \\
& = \theta_1 \int_{\Omega} \omega_1 |\nabla(u_1 - u_2)|^p dx + \beta_p \int_{\Omega} |\Delta u_1 - \Delta u_2|^p v dx + \theta_2 \int_{\Omega} |u_1 - u_2|^p \omega_2 dx \\
& \geq \theta \|u_1 - u_2\|_X^p
\end{aligned}$$

where $\theta = \min \{\theta_1, \theta_2, \beta_p\}$.

Therefore, the operator A is strongly monotone, and this implies that the operator A is strictly monotone. Moreover, using (H3), (H7), (H9) and $\omega_1 \leq \omega_2$, we obtain

$$\begin{aligned}
\langle Au, u \rangle & = B(u, u) = B_1(u, u) + B_2(u, u) + B_3(u, u) \\
& = \int_{\Omega} \omega_1 \mathcal{A}(x, u, \nabla u) \cdot \nabla u dx + \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta u v dx \\
& \quad + \int_{\Omega} b(x, u, \nabla u) u \omega_2 dx \\
& \geq \int_{\Omega} (\lambda_1 |\nabla u|^p + \Lambda_1 |u|^p) \omega_1 dx + \int_{\Omega} |\Delta u|^p v dx \\
& \quad + \int_{\Omega} (\lambda_2 |\nabla u|^p + \Lambda_2 |u|^p) \omega_2 dx \\
& \geq \int_{\Omega} (\lambda_1 |\nabla u|^p + \Lambda_1 |u|^p) \omega_1 dx + \int_{\Omega} |\Delta u|^p v dx \\
& \quad + \lambda_2 \int_{\Omega} |\nabla u|^p \omega_1 dx + \Lambda_2 \int_{\Omega} |u|^p \omega_2 dx \\
& = (\lambda_1 + \lambda_2) \int_{\Omega} |\nabla u|^p \omega_1 dx + \Lambda_2 \int_{\Omega} |u|^p \omega_2 dx + \int_{\Omega} |\Delta u|^p v dx \\
& \geq \gamma \|u\|_X^p
\end{aligned}$$

where $\gamma = \min \{\lambda_1 + \lambda_2, \Lambda_2, 1\}$. Hence, since $p > 1$, we have

$$\frac{\langle Au, u \rangle}{\|u\|_X} \rightarrow +\infty, \text{ as } \|u\|_X \rightarrow +\infty,$$

that is, A is coercive.

Step 6. We need to show that the operator A is continuous. Let $u_m \rightarrow u$ in X as $m \rightarrow \infty$. We have,

$$|B_1(u_m, \varphi) - B_1(u, \varphi)| \leq \sum_{j=1}^n \int_{\Omega} |\mathcal{A}_j(x, u_m, \nabla u_m) - \mathcal{A}_j(x, u, \nabla u)| |D_j \varphi| \omega_1 dx$$

$$\begin{aligned}
&= \sum_{j=1}^n \int_{\Omega} |F_j u_m - F_j u| |D_j \varphi| \omega_1 dx \\
&\leq \sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega_1)} \|D_j \varphi\|_{L^p(\Omega, \omega_1)} \\
&\leq \sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega_1)} \|\varphi\|_X,
\end{aligned}$$

and

$$\begin{aligned}
&|B_3(u_m, \varphi) - B_3(u, \varphi)| \\
&= \left| \int_{\Omega} |\Delta u_m|^{p-2} \Delta u_m \Delta \varphi v dx - \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta \varphi v dx \right| \\
&\leq \int_{\Omega} \left| |\Delta u_m|^{p-2} \Delta u_m - |\Delta u|^{p-2} \Delta u \right| |\Delta \varphi| v dx \\
&= \int_{\Omega} |G u_m - G u| |\Delta \varphi| v dx \\
&\leq \|G u_m - G u\|_{L^{p'}(\Omega, v)} \|\varphi\|_X,
\end{aligned}$$

and

$$\begin{aligned}
|B_2(u_m, \varphi) - B_2(u, \varphi)| &\leq \int_{\Omega} |b(x, u_m, \nabla u_m) - b(x, u, \nabla u)| |\varphi| \omega_2 dx \\
&= \int_{\Omega} |H u_m - H u| |\varphi| \omega_2 dx \\
&\leq \|H u_m - H u\|_{L^{p'}(\Omega, \omega_2)} \|\varphi\|_X,
\end{aligned}$$

for all $\varphi \in X$. Hence,

$$\begin{aligned}
|B(u_m, \varphi) - B(u, \varphi)| &\leq |B_1(u_m, \varphi) - B_1(u, \varphi)| + |B_2(u_m, \varphi) - B_2(u, \varphi)| \\
&\quad + |B_3(u_m, \varphi) - B_3(u, \varphi)| \\
&\leq \left[\sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega_1)} + \|G u_m - G u\|_{L^{p'}(\Omega, v)} \right. \\
&\quad \left. + \|H u_m - H u\|_{L^{p'}(\Omega, \omega_2)} \right] \|\varphi\|_X.
\end{aligned}$$

Then we obtain

$$\begin{aligned}
\|A u_m - A u\|_* &\leq \sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega_1)} + \|G u_m - G u\|_{L^{p'}(\Omega, v)} \\
&\quad + \|H u_m - H u\|_{L^{p'}(\Omega, \omega_2)}.
\end{aligned}$$

Therefore, using (3.2), (3.3) and (3.4) we have $\|Au_m - Au\|_* \rightarrow 0$ as $m \rightarrow +\infty$, that is, A is continuous (and this implies that A is hemicontinuous).

Therefore, by Theorem 3.1, the operator equation $Au = T$ has a unique solution $u \in X$ and it is the unique solution for problem (P).

Step 7. In particular, by setting $\varphi = u$ in Definition 2.5, we have

$$(3.6) \quad B(u, u) = B_1(u, u) + B_2(u, u) + B_3(u, u) = T(u).$$

Hence, using (H3), (H7), (H9), $\omega_1 \leq \omega_2$ and $\gamma = \min\{\lambda_1 + \lambda_2, \Lambda_2, 1\}$, we obtain

$$\begin{aligned} B_1(u, u) + B_2(u, u) + B_3(u, u) &= \int_{\Omega} \omega_1 \mathcal{A}(x, u, \nabla u) \nabla u \, dx \\ &\quad + \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta u \, dx + \int_{\Omega} b(x, u, \nabla u) u \omega_2 \, dx \\ &\geq \int_{\Omega} (\lambda_1 |\nabla u|^p + \Lambda_1 |u|^p) \omega_1 \, dx + \int_{\Omega} |\Delta u|^p \, dx \\ &\quad + \int_{\Omega} (\Lambda_2 |u|^p + \Lambda_1 |\nabla u|^p) \omega_2 \, dx \\ &\geq \int_{\Omega} (\lambda_1 |\nabla u|^p + \Lambda_1 |u|^p) \omega_1 \, dx + \int_{\Omega} |\Delta u|^p \, dx \\ &\quad + \Lambda_2 \int_{\Omega} |u|^p \omega_2 \, dx + \lambda_1 \int_{\Omega} |\nabla u|^p \omega_1 \, dx \\ &\geq \gamma \|u\|_X^p \end{aligned}$$

and

$$\begin{aligned} T(u) &= \int_{\Omega} f_0 u \, dx + \sum_{j=1}^n \int_{\Omega} f_j D_j u \, dx \\ &\leq \|f_0/\omega_2\|_{L^{p'}(\Omega, \omega_2)} \|u\|_{L^p(\Omega, \omega_2)} + \sum_{j=1}^n \|f_j/\omega_1\|_{L^{p'}(\Omega)} \|D_j u\|_{L^p(\Omega, \omega_1)} \\ &\leq \left(\|f_0/\omega_2\|_{L^{p'}(\Omega, \omega_2)} + \sum_{j=1}^n \|f_j/\omega_1\|_{L^{p'}(\Omega, \omega_1)} \right) \|u\|_X. \end{aligned}$$

Therefore, in (3.6), we have

$$\gamma \|u\|_X^p \leq \left(\|f_0/\omega_2\|_{L^{p'}(\Omega, \omega_2)} + \sum_{j=1}^n \|f_j/\omega_1\|_{L^{p'}(\Omega, \omega_1)} \right) \|u\|_X,$$

and we obtain

$$\|u\|_X \leq \frac{1}{\gamma^{p'/p}} \left(\|f_0/\omega_2\|_{L^{p'}(\Omega, \omega_2)} + \sum_{j=1}^n \|f_j/\omega_1\|_{L^{p'}(\Omega, \omega_1)} \right)^{p'/p}.$$

Proof of Corollary 1.2. If $u_1, u_2 \in X$ are solutions of (P_1) and (P_2) respectively, then for all $\varphi \in X$ we have

$$\begin{aligned} & \int_{\Omega} |\Delta u_1|^{p-2} \Delta u_1 \Delta \varphi v \, dx + \sum_{j=1}^n \int_{\Omega} \omega_1 \mathcal{A}_j(x, u_1(x), \nabla u_1(x)) D_j \varphi(x) \, dx \\ & + \int_{\Omega} b(x, u_1, \nabla u_1) \varphi \omega_2 \, dx = \int_{\Omega} f_0(x) \varphi(x) \, dx + \sum_{j=1}^n \int_{\Omega} f_j(x) D_j \varphi(x) \, dx. \end{aligned}$$

and

$$\begin{aligned} & \int_{\Omega} |\Delta u_2|^{p-2} \Delta u_2 \Delta \varphi v \, dx + \sum_{j=1}^n \int_{\Omega} \omega_1 \mathcal{A}_j(x, u_2(x), \nabla u_2(x)) D_j \varphi(x) \, dx \\ & + \int_{\Omega} b(x, u_2, \nabla u_2) \varphi \omega_2 \, dx = \int_{\Omega} \tilde{f}_0(x) \varphi(x) \, dx + \sum_{j=1}^n \int_{\Omega} \tilde{f}_j(x) D_j \varphi(x) \, dx. \end{aligned}$$

In particular, for $\varphi = u_1 - u_2$ we obtain

$$\begin{aligned} & \int_{\Omega} \left(|\Delta u_1|^{p-1} \Delta u_1 - |\Delta u_2|^{p-1} \Delta u_2 \right) (\Delta u_1 - \Delta u_2) v \, dx \\ & + \int_{\Omega} \left(\mathcal{A}(x, u_1, \nabla u_1) - \mathcal{A}(x, u_2, \nabla u_2) \right) \cdot (\nabla u_1 - \nabla u_2) \omega_1 \, dx \\ (3.7) \quad & + \int_{\Omega} \left(b(x, u_1, \nabla u_1) - b(x, u_2, \nabla u_2) \right) (u_1 - u_2) \omega_2 \, dx \\ & = \int_{\Omega} (f_0 - \tilde{f}_0)(u_1 - u_2) \, dx + \sum_{j=1}^n \int_{\Omega} (f_j - \tilde{f}_j) D_j (u_1 - u_2) \, dx. \end{aligned}$$

(i) Using Lemma 2.2 (b), we obtain

$$\begin{aligned} & \int_{\Omega} \left(|\Delta u_1|^{p-1} \Delta u_1 - |\Delta u_2|^{p-1} \Delta u_2 \right) (\Delta u_1 - \Delta u_2) v \, dx \\ & \geq \beta_p \int_{\Omega} \left(|\Delta u_1| + |\Delta u_2| \right)^{p-2} |\Delta u_1 - \Delta u_2|^2 v \, dx \\ & \geq \beta_p \int_{\Omega} |\Delta u_1 - \Delta u_2|^{p-2} |\Delta u_1 - \Delta u_2|^2 v \, dx \\ & = \beta_p \int_{\Omega} |\Delta(u_1 - u_2)|^p v \, dx. \end{aligned}$$

(ii) By (H2), we have

$$\begin{aligned} & \int_{\Omega} \left(\mathcal{A}(x, u_1, \nabla u_1) - \mathcal{A}(x, u_2, \nabla u_2) \right) \cdot (\nabla u_1 - \nabla u_2) \omega_1 \, dx \\ & \geq \theta_1 \int_{\Omega} |\nabla u_1 - \nabla u_2|^p \omega_1 \, dx. \end{aligned}$$

(iii) By (H6) we have

$$\int_{\Omega} \left(b(x, u_1, \nabla u_1) - b(x, u_2, \nabla u_2) \right) (u_1 - u_2) \omega_2 dx \geq \theta_2 \int_{\Omega} |u_1 - u_2|^p \omega_2 dx.$$

Hence, if $\varepsilon = \min\{\beta_p, \theta_1, \theta_2\}$, in (3.7) we obtain

$$\begin{aligned} \varepsilon \|u_1 - u_2\|_X^p &\leq \beta_p \int_{\Omega} |\Delta(u_1 - u_2)|^p v dx + \theta_1 \int_{\Omega} |\nabla(u_1 - u_2)|^p \omega_1 dx \\ &\quad + \theta_2 \int_{\Omega} |u_1 - u_2|^p \omega_2 dx \\ &\leq \left(\|(f_0 - \tilde{f}_0)/\omega_2\|_{L^{p'}(\Omega, \omega_1)} + \sum_{j=1}^n \|(f_j - \tilde{f}_j)/\omega_1\|_{L^{p'}(\Omega, \omega_1)} \right) \|u_1 - u_2\|_X. \end{aligned}$$

Therefore,

$$\begin{aligned} &\|u_1 - u_2\|_X \\ &\leq \frac{1}{\varepsilon^{p-1}} \left(\|(f_0 - \tilde{f}_0)/\omega_2\|_{L^{p'}(\Omega, \omega_1)} + \sum_{j=1}^n \|(f_j - \tilde{f}_j)/\omega_1\|_{L^{p'}(\Omega, \omega_1)} \right)^{p-1}. \quad \square \end{aligned}$$

Example 1. Let $\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 1\}$. Consider the weight functions $\omega_1(x, y) = (x^2 + y^2)^{-1/4}$, $\omega_2(x, y) = (x^2 + y^2)^{-1/2}$ and $v(x, y) = (x^2 + y^2)^{-1/3}$ (we have $\omega_1, \omega_2, v \in A_2$, $p = 2$ and $q = 2$), and the functions $\mathcal{A} : \Omega \times \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $b : \Omega \times \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\begin{aligned} \mathcal{A}((x, y), \eta, \xi) &= h_2(x, y) \xi, \\ b((x, y), \eta, \xi) &= \eta (\cos^2(xy) + 1), \end{aligned}$$

where $h(x, y) = 2e^{(x^2+y^2)}$. Let us consider the partial differential operator

$$\begin{aligned} Lu(x, y) &= \Delta((x^2 + y^2)^{-1/3} |\Delta u| \Delta u) - \operatorname{div}((x^2 + y^2)^{-1/4} \mathcal{A}((x, y), u, \nabla u)) \\ &\quad + (x^2 + y^2)^{-1/2} b(x, u, \nabla u). \end{aligned}$$

Therefore, by Theorem 1.1, the problem

$$(P) \begin{cases} Lu(x) = \frac{\cos(xy)}{\sqrt{x^2 + y^2}} - \frac{\partial}{\partial x} \left(\frac{\sin(xy)}{\sqrt{x^2 + y^2}} \right) - \frac{\partial}{\partial y} \left(\frac{\sin(xy)}{\sqrt{x^2 + y^2}} \right), & \text{in } \Omega \\ u(x) = \Delta u(x) = 0, & \text{on } \partial\Omega \end{cases}$$

has a unique solution $u \in X = W^{2,2}(\Omega, v) \cap W_0^{1,2}(\Omega, \omega_1, \omega_2)$.

REFERENCES

- [1] Cavalheiro, A. C., *Existence and uniqueness of solutions for some degenerate nonlinear Dirichlet problems*, J. Appl. Anal. **19** (2013), 41–54.
- [2] Cavalheiro, A. C., *Existence results for Dirichlet problems with degenerated p -Laplacian and p -Biharmonic operators*, Appl. Math. E-Notes **13** (2013), 234–242.
- [3] Chipot, M., *Elliptic Equations: An Introductory Course*, Birkhäuser, Berlin, 2009.

-
- [4] Drábek, P., Kufner, A., Nicolosi, F., *Quasilinear Elliptic Equations with Degenerations and Singularities*, Walter de Gruyter, Berlin, 1997.
 - [5] Fučík, S., John, O., Kufner, A., *Function Spaces*, Noordhoff International Publ., Leyden, 1977.
 - [6] Garcia-Cuerva, J., Rubio de Francia, J. L., *Weighted Norm Inequalities and Related Topics*, North-Holland, Amsterdam, 1985.
 - [7] Gilbarg, D., Trudinger, N. S., *Elliptic Partial Equations of Second Order*, 2nd Ed., Springer, New York, 1983.
 - [8] Heinonen, J., Kilpeläinen, T., Martio, O., *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford University Press, Inc., New York, 1993.
 - [9] Kufner, A., *Weighted Sobolev Spaces*, John Wiley & Sons, 1985.
 - [10] Muckenhoupt, B., *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. **165** (1972), 207–226.
 - [11] Talbi, M., Tsouli, N., *On the spectrum of the weighted p -Biharmonic operator with weight*, Mediterr. J. Math. **4** (2007), 73–86.
 - [12] Torchinsky, A., *Real-Variable Methods in Harmonic Analysis*, Academic Press, San Diego, 1986.
 - [13] Turesson, B. O., *Nonlinear Potential Theory and Weighted Sobolev Spaces*, Springer-Verlag, Berlin–Heidelberg–New York, 2000.
 - [14] Zeidler, E., *Nonlinear Functional Analysis and Its Applications. Vol. I*, Springer-Verlag, New York, 1990.
 - [15] Zeidler, E., *Nonlinear Functional Analysis and Its Applications. Vol. II/B*, Springer-Verlag, New York, 1990.

Albo Carlos Cavalheiro
Department of Mathematics
State University of Londrina
Londrina - PR - Brazil, 86057-970
e-mail: accava@gmail.com

Received July 5, 2016