

Presentations

11-3-2013

Skill assessment of multiple models in the Chesapeake Bay

I. D. Irby

Virginia Institute of Marine Science

M. Friedrichs

Virginia Institute of Marine Science

C. Feng

R. Hood

Follow this and additional works at: <https://scholarworks.wm.edu/presentations>




Part of the [Environmental Sciences Commons](#)

Recommended Citation

Irby, I. D.; Friedrichs, M.; Feng, C.; and Hood, R.. "Skill assessment of multiple models in the Chesapeake Bay". 11-3-2013. 22nd Biennial Conference of the Coastal & Estuarine Research Federation, San Diego, CA.

This Presentation is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Presentations by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



Skill Assessment of Multiple Models in the Chesapeake Bay

Isaac (Ike) Irby¹, Marjorie Friedrichs¹, Yang Feng¹, Raleigh Hood², Jeremy Testa², Carl Friedrichs¹

¹Virginia Institute of Marine Science, The College of William and Mary, Gloucester Point VA, USA

²Center of Environmental Science, University of Maryland, Cambridge MD, USA



ABSTRACT

The Environmental Protection Agency (EPA) has developed a set of Total Maximum Daily Load (TMDL) allotments of nutrients and sediments for the six states that make up the Chesapeake Bay Watershed in order to elevate the health of the Bay, primarily in regards to dissolved oxygen concentrations. In developing the TMDLs, the EPA employed a coupled watershed-estuarine numerical modeling system together with an extensive set of monitoring data. Utilization of a multiple model approach when evaluating the status and recovery of the Bay system could enhance the overall confidence in model projections and better define model uncertainty. Open-source modeling systems such as the Regional Ocean Modeling System (ROMS) offer a cost effective way of utilizing the knowledge base of a large group of people from multiple institutions to address management issues within a single system. This study compares the relative skill of a set of ROMS-based models to the EPA regulatory model in terms of the seasonal variability of the Chesapeake Bay. Throughout the main stem of the Bay both model types achieve a similar model skill score in regards to dissolved oxygen (DO), the primary indicator of Bay health by the EPA, but vary significantly in terms of their ability to reproduce chlorophyll and nitrate.



INTRODUCTION

Chesapeake Bay and its surrounding watershed play host to an extensive suite of commercial and recreational fishing, agriculture, shipping, and tourism industries that have an estimated value upwards of one trillion dollars and is home to more than 16 million people. Ensuring the health of the Bay has become a major priority for the six states that make up the watershed and together they have committed to the implementation of a set of Total Maximum Daily Loads (TMDLs). The goal of the TMDLs is to improve water quality throughout the Bay by decreasing the levels of nutrients and sediment derived from the watershed. The EPA measures Bay water quality improvement primarily via increased dissolved oxygen (DO) concentrations. TMDL levels were determined using the EPA's coupled watershed-estuarine model, CH3D-ICM. A multiple model approach can be used to gauge uncertainty and elevate confidence in the regulatory model projections.

OBJECTIVE

Statistically compare output from a set of three open source estuarine models of varying biological complexity to the biologically sophisticated EPA regulatory model in terms of reproducing the mean and seasonal variability of temperature, salinity, stratification, dissolved oxygen, chlorophyll-a, and nitrate.



METHODS

- Simulations from the EPA regulatory model and three ROMS-based models were analyzed (Table 1):
 - CH3D – ICM: EPA
 - ROMS – RCA: UMCES
 - ChesNENA: VIMS
 - ChesROMS – BGC: UMCES
- Model output was compared to Chesapeake Bay Program monitoring data using a best time match system for roughly 17 cruises at 10 main stem station in 2004 (Fig. 1).
- Analyzed variables included surface and bottom temperature, salinity, dissolved oxygen, chlorophyll-a, and nitrate, as well as stratification.
- Model ability to reproduce the mean and seasonal variability of each variable was evaluated via Target Diagrams (Fig. 2).

	CH3D - ICM	ChesNENA	ChesROMS - BGC	ROMS - RCA
Nutrients Simulated	N, P, Si	C, N	N	N, P, Si
Biogeochemical Sediment Component	Yes	No	No	Yes
Number of Algal Groups	3	1	1	2
Horizontal Resolution	0.25 - 1km ²	~1km ²	~1km ²	~1km ²
Vertical Grid	z: ~5ft	σ: 20 layers	σ: 20 layers	σ: 20 layers

Table 1. Characteristics of the individual models.

ANALYSIS



	CH3D - ICM	ChesNENA	ChesROMS - BGC	ROMS - RCA
Temperature Surface	0.14	0.09	0.18	0.09
Temperature Bottom	0.21	0.33	0.21	0.19
Salinity Surface	0.23	0.31	0.43	0.36
Salinity Bottom	0.33	0.51	0.53	0.82
Stratification	1.11	1.36	1.34	1.33
DO Surface	0.71	0.51	0.74	0.66
DO Bottom	0.46	0.52	0.77	0.48
Chlorophyll Surface	1.17	1.25	1.70	2.08
Chlorophyll Bottom	0.88	1.05	1.28	1.54
Nitrate Surface	0.76	0.61	1.49	0.43
Nitrate Bottom	0.72	0.51	1.98	0.54

Table 2. Total normalized RMSD computed for each model for multiple variables using observations from cruises in 2004 at the 10 main stem stations shown in Figure 1. Shading indicates model results that perform better than the mean of the observations.

RESULTS

- The skill of all four models are similar to each other in terms of temperature, salinity, stratification, and DO, but the models vary significantly in terms of their chlorophyll and nitrate (Fig. 3, Table 2).
- All models consistently underestimate both the mean and standard deviation of maximum stratification, particularly at the northern stations (Fig. 4A).
- Despite the models' inability to resolve stratification, the models reproduce the mean and variability of DO quite well. All models perform better at the southern stations than the northernmost stations (Fig. 4B).
- Model skill for surface chlorophyll varies significantly between models, with the regulatory CH3D – ICM model performing best (Fig. 4C). ROMS – RCA is particularly challenged at the southern stations.
- Model skill for surface nitrate varies significantly between models, with two of the ROMS-based models performing as well or better than CH3D – ICM (Fig. 4D). ChesROMS is particularly challenged at the southern stations.

CONCLUSIONS

- Overall, models with lower biological complexity and lower resolution achieve similar skills scores as the EPA regulatory model in terms of seasonal variability along the main stem of the Chesapeake Bay.
- Multiple variables exhibit latitudinal dependence of model skill that is consistent throughout all four models, e.g. mean stratification is underestimated most and model skill for DO is lowest in the north.
- All four models do substantially better at resolving bottom DO than they do at resolving three variables that are primary influences on DO: stratification, chlorophyll, and nitrate. This follows because DO's variability is sensitive to temperature as a result of the solubility effect, and the models reproduce temperature very well.
- In terms of TMDL development, these findings offer a greater confidence in CH3D – ICM predictions of seasonal variability since a model does not necessarily need to perform well in terms of stratification, chlorophyll, or nitrate in order to resolve the mean and seasonal variation of DO.**

FUTURE WORK

Examine the skill of these multiple models in terms of interannual variability, with the goal of formulating a ROMS-based model that performs as well as the EPA regulatory model for both seasonal and interannual variability of DO and hypoxic volume.

ACKNOWLEDGEMENTS

This work was funded by the NOAA NOS IOOS as part of the Coastal Modeling Testbed (NA13NOS0120139) and the NASA Interdisciplinary Science Program as part of the USECoS project (NNX11AD47G). Special thanks to Aaron Bever and Ping Wang.

