

Reports

7-26-2013

Final Report An Assessment of Sea Scallop Abundance and Distribution in a Selected Closed Area: Nantucket Lightship Closed Area

David Rudders
Virginia Institute of Marine Science

William D. DuPaul
Virginia Institute of Marine Science

Jessica Bergeron
Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Rudders, D., DuPaul, W. D., & Bergeron, J. (2013) Final Report An Assessment of Sea Scallop Abundance and Distribution in a Selected Closed Area: Nantucket Lightship Closed Area. Marine Resource Report No. 2013-5. Virginia Institute of Marine Science, William & Mary. <http://dx.doi.org/doi:10.21220/m2-kqgf-ag37>

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Final Report

An Assessment of Sea Scallop Abundance and Distribution in a Selected Closed Area: Nantucket Lightship Closed Area

Award Number: 11-SCA-12

Submitted to:

National Marine Fisheries Service
Northeast Fisheries Science Center
Cooperative Research Program
166 Water Street
Woods Hole, Massachusetts 02543-1026

Submitted by:

David B. Rudders
William D. DuPaul
Jessica Bergeron

Virginia Institute of Marine Science
College of William and Mary
Gloucester Point, Virginia 23062

VIMS Marine Resource Report No. 2013-5
July 26, 2013

Project Summary

As the spatial and temporal dynamics of marine ecosystems have recently become better understood, the concept of entirely closing or limiting activities in certain areas has gained support as a method to conserve and enhance marine resources. In the last decade, the sea scallop resource has benefited from measures that have closed specific areas to fishing effort. As a result of closures on both Georges Bank and in the mid-Atlantic region, biomass of scallops in those areas has expanded. As the time approaches for the fishery to harvest scallops from the closed areas, quality, timely and detailed stock assessment information is required for managers to make informed decisions about the re-opening.

During May 2012, a survey was conducted in the access area of the Nantucket Lightship Closed Area (NLCA) aboard a commercial sea scallop vessel. At pre-determined sampling stations within the NLCA, both a NMFS survey dredge and a Coonamessett Farm Turtle Deflector Dredge (CFTDD) were simultaneously towed. From this trip, fine scale survey data was used to assess scallop abundance and distribution in the closed area. This data will also provide a comparison of the utility of using two different gears as survey tools in the context of industry based surveys.

Results indicate that the resource in NLCA is relatively abundant with sufficient exploitable biomass to support a commercial opening in 2013 and potentially in 2014. Of concern was the lack of observed recruitment that has the potential to impact the abundance of the resource in that area during subsequent years. Gear performance of the CFTDD was observed to differ with prior results with respect to the size of animals captured, although the relative efficiency of the CFTDD was consistent with prior surveys.

Project Background

The sea scallop, *Placopecten magellanicus*, supports a fishery that in the 2011 fishing year landed 58.1 million pounds of meats with an ex-vessel value of over US \$580 (Lowther, 2012). These landings resulted in the sea scallop fishery being the most valuable single species fishery along the East Coast of the United States. While historically subject to extreme cycles of productivity, the fishery has benefited from recent management measures intended to bring stability and sustainability. These measures include: limiting the number of participants, total effort (days-at-sea), gear and crew restrictions and most recently, a strategy to improve yield by protecting scallops through rotational area closures.

Amendment #10 to the Sea Scallop Fishery Management Plan officially introduced the concept of area rotation to the fishery. This strategy seeks to increase the yield and reproductive potential of the sea scallop resource by identifying and protecting discrete areas of high densities of juvenile scallops from fishing mortality. By delaying capture, the rapid growth rate of scallops is exploited to realize substantial gains in yield over short time periods. In addition to the formal attempts found in Amendment #10 to manage discrete areas of scallops for improved yield, specific areas on Georges Bank are also subject to area closures. In 1994, 17,000 km² of bottom were closed to any fishing gears capable of capturing groundfish. This closure was an attempt to aid in the rebuilding of severely depleted species in the groundfish complex. Since scallop dredges are capable of capturing groundfish, scallopers were also excluded from these areas. Since 1999, however, limited access to the three closed areas on Georges Bank has been allowed to harvest the dense beds of scallops that have accumulated in the absence of fishing pressure.

In order to effectively regulate the fishery and carry out a robust rotational area management strategy, current and detailed information regarding the abundance and distribution of sea scallops is essential. Currently, abundance and distribution information gathered by surveys comes from a variety of sources. The annual NMFS sea scallop survey provides a comprehensive and synoptic view of the resource from Georges Bank to Virginia. In contrast to the NMFS survey that utilizes a dredge as the sampling gear, the resource is also surveyed optically. Researchers from the School for Marine Science and Technology (SMAST) and the Woods Hole Oceanographic Institute (WHOI) are able to enumerate sea scallop abundance and distribution from images taken by both a still camera and a towed camera system (Stokesbury, *et. al.*, 2004; Stokesbury, 2002). Prior to the utilization of the optical surveys and in addition to the annual information supplied by the NMFS annual survey, commercial vessels were

contracted to perform surveys. Dredge surveys of the scallop access areas have been successfully completed by the cooperative involvement of industry, academic and governmental partners. The additional information provided by these surveys was vital in the determination of appropriate Total Allowable Catches (TAC) in the subsequent re-openings of the closed areas. This type of survey, using commercial fishing vessels, provides an excellent opportunity to gather required information and also involve stakeholders in the management of the resource.

The passing of Amendment #10 has set into motion changes to the sea scallop fishery that are designed to ultimately improve yield and create stability. This stability is an expected result of a spatially explicit rotational area management strategy where areas of juvenile scallops are identified and protected from harvest until they reach an optimum size. Implicit to the institution of the new strategy, is the highlighted need for further information to both assess the efficacy of an area management strategy and provide that management program with current and comprehensive information. In addition to rotational management areas, access to the scallop biomass encompassed by the Nantucket Lightship Closed Area is vital to the continued prosperity of the fishery.

In addition to collecting data to assess the abundance and distribution of sea scallops in the NLCA, the operational characteristics of commercial scallop vessels allow for the simultaneous towing of two dredges. As in past surveys, we towed two dredges at each station. One dredge was a NMFS sea scallop survey dredge and the other was a Coonamessett Farm Turtle Deflector Dredge (CFTDD). This paired design allowed for the estimation of the size selective characteristics of CFTDD equipped with turtle excluder chains. Gear performance (i.e. size selectivity and relative efficiency) information is limited for this dredge design and understanding how this dredge impacts the scallop resource will be beneficial for two reasons. First, it will be an important consideration for the stock assessment for scallops in that it provides the size selectivity characteristics of the most recent gear configuration and second, this information will support the use of this gear configuration to sample closed areas prior to re-openings. In addition, selectivity analyses using the SELECT method provide insight to the relative efficiency of the two gears used in the study (Millar, 1992). The relative efficiency measure from this experiment can be used to refine existing absolute efficiency estimates for the New Bedford style scallop dredge.

One of the stated advantages of a dredge sea scallop survey is that one can access and sample the target species. One parameter routinely measured is the shell height:meat weight relationship. While this relationship is used to determine swept area biomass for the area surveyed at that time, it can also be used as an indicator of seasonal shifts in biomass due to

the influence of spawning. For this reason, data on the shell height:meat weight relationship is routinely gathered by both the NMFS and VIMS scallop surveys. While this relationship may not be a direct indicator of animal health in and of itself, long term data sets may be useful in evaluating changing environmental conditions, food availability and density dependent interactions.

For this study, we pursued multiple objectives. The primary objective was to collect information to characterize the abundance and distribution of sea scallops within the access area of NLCA. Utilizing the same catch data with a different analytical approach, we estimated the size selectivity characteristics of the commercial sea scallop dredge. An additional component of the selectivity analysis allows for supplementary information regarding the efficiency of the commercial dredge relative to the NMFS survey dredge. As a third objective of this study, we collected biological samples to estimate a time and area specific shell height:meat weight relationship as well as additional samples were to assess product quality of the adult resource in the NLCA.

Methods

Survey Area and Sampling Design

The access area NLCA was surveyed during the course of this project. The boundary coordinates of the surveyed areas can be found in Table 1. Sampling stations for this study were selected within the context of a systematic random grid. With the patchy distribution of sea scallops determined by some unknown combination of environmental gradients (i.e. latitude, depth, hydrographic features, etc.), a systematic selection of survey stations results in an even dispersion of samples across the entire sampling domain. This sampling design has been successfully implemented during industry-based surveys since 1998.

The methodology to generate the systematic random grid entailed the decomposition of the domain (in this case a closed area) into smaller sampling cells. The dimensions of the sampling cells were primarily determined by a sample size analysis conducted using the catch data from survey trips conducted in the same areas during prior years. Since closed areas are of different dimensions and the total number of stations sampled per survey remains fairly constant, the distance between the stations varies. Generally, the distance between stations is roughly 3-4 nautical miles. Once the cell dimensions were set, a point within the most northwestern cell was randomly selected. This point served as the starting point and all of the other stations in the grid were based on its coordinates. The station locations for the 2012 NLCA survey are shown in Figure 1.

Sampling Protocols

While at sea, the vessel simultaneously towed two dredges. A NMFS survey dredge, 8 feet in width equipped with 2-inch rings, 4-inch diamond twine top and a 1.5-inch diamond mesh liner was towed on one side of the vessel. On the other side of the vessel, a 15 foot Coonamessett Farm Turtle Deflector Dredge (CFTDD) equipped with 4-inch rings, a 10-inch diamond mesh twine top and no liner was utilized. Turtle chains were used in configurations as dictated by the area surveyed and current regulations. In this paired design, it is assumed that the dredges cover a similar area of substrate and sample from the same population of scallops. The dredges were switched to opposite sides of the vessel mid-way throughout the trip to help minimize any bias.

For each survey tow, the dredges were fished for 15 minutes with a towing speed of approximately 3.8-4.0 kts. High-resolution navigational logging equipment was used to accurately determine and record vessel position. A Star-Oddi™ DST sensor was used on the dredge to measure and record dredge tilt angle as well as depth and depth (Figure 2). With these measurements, the start and end of each tow was estimated. Synchronous time stamps on both the navigational log and DST sensor were used to estimate the linear distance for each tow. A histogram depicting the estimated linear distances covered per tow over the entire survey is shown in Figure 3.

Sampling of the catch was performed using the protocols established by DuPaul and Kirkley, 1995 and DuPaul *et. al.* 1989. For each survey tow, the entire scallop catch was placed in baskets. Depending on the total volume of the catch, a fraction of these baskets were measured for sea scallop length frequency. The shell height of each scallop in the sampled fraction was measured on NMFS sea scallop measuring boards in 5 mm intervals. This protocol allows for the estimation of the size frequency for the entire catch by expanding the catch at each shell height by the fraction of total number of baskets sampled. Finfish and invertebrate bycatch were quantified, with finfish being sorted by species and measured to the nearest 1 cm.

Samples were taken to determine area specific shell height:meat weight relationships. At roughly 25 randomly selected stations the shell height of 10 randomly selected scallops were measured to the nearest 0.1 mm. These scallops were then carefully shucked and the adductor muscle individually packaged and frozen at sea. Upon return, the adductor muscle was weighed to the nearest 0.1 gram. The relationship between shell height and meat weight was estimated using a generalized linear mixed model (gamma distribution, log link) incorporating

depth as an explanatory variable using PROC GLIMMIX in SAS v. 9.2. The relationship was estimated with the following models:

$$MW = \exp(\alpha + \beta \cdot \ln SH + \gamma \cdot \ln \text{Depth})$$

$$MW = \exp(\alpha + \beta \cdot \ln SH)$$

where MW=meat weight (grams), SH=shell height (millimeters), Depth=depth (meters). α , β and γ are parameters to be estimated.

The standard data sheets in service since the 1998 Georges Bank survey were used. Data recorded on the bridge log included GPS location, tow-time (break-set/haul-back), tow speed, water depth, catch, bearing, weather and comments relative to the quality of the tow. The deck log maintained by the scientific personnel recorded detailed catch information on scallops, finfish, invertebrates and trash.

Data Analysis

The catch and navigation data were used to estimate swept area biomass within the area surveyed. The methodology to estimate biomass is similar to that used in previous survey work by VIMS. In essence, we estimate a mean abundance from the point estimates and scale that value up to the entire area of the domain sampled. This calculation is given:

$$TotalBiomass = \sum_j \left(\frac{\left(\frac{CatchWtperTowinSubarea_j}{AreaSweptperTow} \right)}{Efficiency} \right) SubArea_j$$

Catch weight per tow of exploitable scallops was calculated from the raw catch data as an expanded size frequency distribution with an area and depth appropriate shell height:meat weight relationship applied (length-weight relationships were obtained from SARC 50 document as well as the actual relationship taken during the cruise) (NEFSC, 2010). Exploitable biomass, defined as that fraction of the population vulnerable to capture by the currently regulated commercial gear, was calculated using two approaches. The observed catch at length data from the NMFS survey dredge (assumed to be non-size selective) was adjusted based upon the size selectivity characteristics of the commercial gear (Yochum and DuPaul, 2008). The

observed catch-at-length data from the commercial dredge was not adjusted due to the fact that these data already represent that fraction of the population that is subject to exploitation by the currently regulated commercial gear.

Utilizing the information obtained from the high resolution GPS, an estimate of area swept per tow was calculated. Throughout the cruise, the location of the ship was logged every three seconds. By determining the start and end of each tow based on the recorded times as delineated by the tilt sensor data, a survey tow can be represented by a series of consecutive coordinates (latitude, longitude). The linear distance of the tow is calculated by:

$$TowDist = \sum_{i=1}^n \sqrt{(long_2 - long_1)^2 + (lat_2 - lat_1)^2}$$

The linear distance of the tow is multiplied by the width of the gear (either 15 or 8 ft.) to result in an estimate of the area swept during a given survey tow.

The final two components of the estimation of biomass are constants and not determined from experimental data obtained on these cruises. Estimates of survey dredge gear efficiency have been calculated from a prior experiment using a comparison of optical and dredge catches (NEFSC, 2010). Based on this experiment, an efficiency value for the NMFS survey dredge of 38% was estimated for the rocky substrate areas on Georges Bank and a value of 44% was estimated for the smoother (sand, silt) substrates of some portions of Georges Bank and the entire mid-Atlantic. Estimates of commercial sea scallop dredge gear efficiency have been calculated from prior experiments using a variety of approaches (Gedamke *et. al.*, 2005, Gedamke *et. al.*, 2004, D. Hart, pers. comm.). The efficiency of the commercial dredge is generally considered to be higher and based on the prior work as well as the relative efficiency from the data generated from this study; an efficiency value of 60% was used for the NLCA. To scale the estimated mean scallop catch to the full domain, the total area of the NLCA was calculated in ArcGIS v. 10.0.

Size Selectivity

The estimation of size selectivity of the CFTDD equipped with 4" rings, a 10" twine top and turtle chains was based on a comparative analysis of the catches from the two dredges used in the survey. For this analysis, the NMFS survey dredge is assumed to be non-selective (i.e. a scallop that enters the dredge is retained by the dredge). Catch at length from the selective gear (commercial dredge) were compared to the non-selective gear via the SELECT

method (Millar, 1992). With this analytical approach, the selective properties (i.e. the length based probability of retention) of the commercial dredge were estimated. In addition to estimates of the length based probabilities of capture by the commercial dredge, the SELECT method characterizes a measure of relative fishing intensity. Assuming a known quantity of efficiency for one of the two gears (in this case the survey dredge at 38%), insight into the efficiency of the other gear (commercial dredge) can be attained.

Prior to analysis, all comparative tows were evaluated. Any tows that were deemed to have had problems during deployment or at any point during the tow (flipped, hangs, crossed towing wires, etc.) were removed from the analysis. In addition, tows where zero scallops were captured by both dredges were also removed from the analysis. The remaining tow pairs were then used to analyze the size selective properties of the commercial dredge with the SELECT method.

The SELECT method has become the preferred method to analyze size-selectivity studies encompassing a wide array of fishing gears and experimental designs (Millar and Fryer, 1999). This analytical approach conditions the catch of the selective gear at length l to the total catch (from both the selective gear variant and small mesh control).

$$\Phi_c(l) = \frac{p_c r_c(l)}{p_c r_c(l) + (1 - p_c)}$$

Where $r(l)$ is the probability of a fish at length l being retained by the gear given contact and p is the split parameter (measure of relative efficiency). Traditionally, selectivity curves have been described by the logistic function. This functional form has symmetric tails. In certain cases, other functional forms have been utilized to describe size selectivity of fishing gears. Examples of different functional forms include Richards, log-log and complimentary log-log. Model selection is determined by an examination of model deviance (the likelihood ratio statistic for model goodness of fit) as well as Akaike Information Criterion (AIC) (Xu and Millar, 1993, Sala, *et. al.*, 2008). For towed gears, however, the logistic function is the most common functional form observed in towed fishing gears. Given the logistic function:

$$r(l) = \left(\frac{\exp(a + bl)}{1 + \exp(a + bl)} \right)$$

by substitution:

$$\Phi(L) = \frac{pr(L)}{(1-p) + pr(L)} = \frac{p \frac{e^{a+bL}}{1+e^{a+bL}}}{(1-p) + p \frac{e^{a+bL}}{1+e^{a+bL}}} = \frac{pe^{a+bL}}{(1-p) + e^{a+bL}}$$

Where a , b , and p are parameters estimated via maximum likelihood. Based on the parameter estimates, L_{50} and the selection range (SR) are calculated.

$$L_{50} = \frac{-a}{b} \qquad SR = \frac{2 * \ln(3)}{b}$$

Where L_{50} defines the length at which an animal has a 50% probability of being retained, given contact with the gear and SR represents the difference between L_{75} and L_{25} which is a measure of the slope of the ascending portion of the logistic curve.

In situations where catch at length data from multiple comparative tows is pooled to estimate an average selectivity curve for the experiment, tow by tow variation is often ignored. Millar *et al.* (2004) developed an analytical technique to address this between-haul variation and incorporate that error into the standard error of the parameter estimates. Due to the inherently variable environment that characterizes the operation of fishing gears, replicate tows typically show high levels of between-haul variation. This variation manifests itself with respect to estimated selectivity curves for a given gear configuration (Fryer 1991, Millar *et al.*, 2004). If not accounted for, this between-haul variation may result in an underestimate of the uncertainty surrounding estimated parameters increasing the probability of spurious statistical significance (Millar *et al.*, 2004).

Approaches developed by Fryer (1991) and Millar *et al.*, (2004) address the issue of between-haul variability. One approach formally models the between-haul variability using a hierarchical mixed effects model (Fryer 1991). This approach quantifies the variability in the selectivity parameters for each haul estimated individually and may be more appropriate for complex experimental designs or experiments involving more than one gear. For more straightforward experimental designs, or studies that involve a single gear, a more intuitive combined-haul approach may be more appropriate.

This combined-hauls approach characterizes and then calculates an overdispersion correction for the selectivity curve estimated from the catch data summed over all tows, which is

identical to a curve calculated simultaneously to all individual tows. Given this identity, a replication estimate of between-haul variation (REP) can be calculated and used to evaluate how well the expected catch using the selectivity curve calculated from the combined hauls fits the observed catches for each individual haul (Millar *et. al.* 2004).

REP is calculated as the Pearson chi-square statistic for model goodness of fit divided by the degrees of freedom.

$$REP = \frac{Q}{d}$$

Where Q is equal to the Pearson chi-square statistic for model goodness of fit and d is equal to the degrees of freedom. The degrees of freedom are calculated as the number of terms in the summation, minus the number of estimated parameters. The calculated replicate estimate of between-haul variation was used to calculate observed levels of extra Poisson variation by multiplying the estimated standard errors by \sqrt{REP} . This correction is only performed when the data is not overdispersed (Millar, 1993).

A significant contribution of the SELECT model is the estimation of the split parameter which estimates the probability of an animal “choosing” one gear over another (Holst and Revill, 2009). This measure of relative efficiency, while not directly describing the size selectivity properties of the gear, is insightful relative to both the experimental design of the study as well as the characteristics of the gears used. A measure of relative efficiency (on the observational scale) can be calculated in instances where the sampling intensity is unequal. In this case, the sampling intensity is unequal due to differences in dredge width. Relative efficiency can be computed for each individual trip by the following formula:

$$RE = \frac{p/(1-p)}{p_0/(1-p_0)}$$

Where p is equal to the observed (estimated p value) and p_0 represents the expected value of the split parameter based upon the dredge widths in the study (Park *et. al.*, 2007). For this study, a 15 ft. commercial dredge was used with expected split parameter of 0.6521. The computed relative efficiency values were then used to scale the estimate of the NMFS survey dredge efficiency obtained from the optical comparisons (38%). Computing efficiency for the estimated p value from Yochum and DuPaul (2008) yields a commercial dredge efficiency of

64%. Preliminary observations suggest a slightly higher efficiency of the CFTDD relative to the standard New Bedford style scallop dredge that was used in Yochum and DuPaul (2008). This selectivity analysis will provide an additional piece of evidence related to the efficiency of the CFTDD.

Results

Abundance and distribution

The survey cruise to the access area of the NLCA was completed in May 2012. Summary statistics for the cruise is shown in Table 2. Length frequency distributions for the scallops captured during the NLCA survey is shown in Figure 4. Maps depicting the spatial distribution of the catches of pre-recruit (<70 mm shell height), and fully recruited (≥ 70 mm shell height) scallops from both the commercial and survey dredges are shown in Figures 5-8. Mean total and mean exploitable scallop densities for both the survey and commercial dredge are shown in Table 3. This information expanded to the area of the entire NLCA and representing an estimate of the total number of animals in the area is shown in Table 4. The mean estimated scallop meat weight for both the commercial and survey dredges for all of the shell height:meat weight relationships used is shown in Table 5. Mean catch (in grams of scallop meat) for the two dredge configurations as well as the four shell height: meat weight relationships are shown in Table 6. Total and exploitable biomass for both shell height:meat weight relationships and levels of assumed gear efficiency are shown in Tables 7-8 (total biomass is not estimated due to the selective properties of the commercial gear). Shell height:meat weight relationships were generated for the area. The resulting parameters as well as the parameters from SARC 50 (both a NLCA specific as well as a general Georges Bank relationship) are shown in Table 9. A comparative plot of the four curves is shown in Figure 9. Catch per unit of effort for finfish and invertebrate bycatch is shown in Table 10.

Size selectivity

The catch data was evaluated by the SELECT method with a variety of functional forms (logistic, Richards, log-log) in an attempt to characterize the most appropriate model. Examination of residual patterns model deviance and AIC values indicated that the logistic curve provided the best fit to the data. An additional model run was conducted to determine whether the hypotheses of equal fishing intensity (i.e. the two gears fished equally) were supported. Output for model runs for the logistic function with the split parameter (p) both held fixed at the expected value based on gear width and with p being estimated is shown in Table

11. Visual examination of residuals and values of model deviance and AIC indicated that the model with an estimated split parameter provided the best fit to the data. A fitted curve and deviance residuals for the NLCA cruise is shown in Figure 10. Estimated parameters for the final model run excluding tows with less than 50 total scallops caught is shown in Table 12. The estimated L_{50} value was 87.7 mm and the selection range was 34.26 mm. A final selectivity curve for this data set is shown in Figure 11.

The analysis that estimated the relative efficiency of the two gears based upon the expected and observed split parameter values resulted in an estimate relative efficiency value of 1.4202. Assuming the survey dredge operates with a 38% efficiency, the expected value for the efficiency of the commercial dredge was 54.0%. These results are similar to Yochum and DuPaul (2008) and are consistent with the 60% efficiency value in the previously calculated estimates of total and exploitable biomass.

In response to concerns from industry related to the product quality of some of the older animals in the NLCA, we qualitatively assessed scallop meats based on color and texture criteria. The phenomenon of “grey meats” is well established as well as stringy meats that tear easily. Based on our observations, the quality of the scallop meats in the NLCA during May was excellent and did not appear to have any detrimental characteristics associated with color or texture issues. We suspect that these issues can be ephemeral and are the result of factors that vary in time and space. This topic merits additional research to not only document its spatial extent and intensity, but to understand the underlying process.

As part of the outreach component of this project, a special data summary detailing the spatial distribution of scallops and bycatch species in NLCA was compiled. These data were shared with SMAST and were used to provide a starting point for their Bycatch Avoidance Program for 2013. The intent of sharing this data was to inform the sea scallop industry about the abundance and distribution of scallops in the area as well as potential areas of high finfish bycatch concentrations in an effort to direct effort away from these areas. It was hoped that by distributing this information, effort could be focused on areas that contained high densities of scallops while minimizing finfish bycatch. This is one potential strategy to reduce the rate of finfish bycatch.

Discussion

Fine scale surveys of closed areas are an important endeavor. These surveys provide information about subsets of the resource that may not have been subject to intensive sampling by other efforts. Additionally, the timing of industry-based surveys can be tailored to give

managers current information to guide important management decisions. This information can help time access to closed areas and help set Total Allowable Catches (TAC) for the re-opening. Finally, this type of survey is important in that it involves the stakeholders of the fishery in the management of the resource.

Our results suggest that for the NLCA sufficient biomass exists to support an opening in 2013 and perhaps 2014. For an area that had been dominated by a large size class, there appears to have been some limited recruitment in the area and that the age distribution of the resource is broader relative to prior years. These pre-recruits represent an important cohort and have the ability to realize year over year increases in growth as well as the potential to sustain openings in subsequent years. These animals, however, were spatially limited and their overall extent was not remarkable.

The use of commercial scallop vessels in a project of this magnitude presents some interesting challenges. One such challenge is the use of the commercial gear. This gear is not designed to be a survey gear; it is designed to be efficient in a commercial setting. The design of this current experiment however provides insight into the utility of using a commercial gear as a survey tool. One advantage of the use of this gear is that the catch from this dredge represent exploitable biomass and no further correction is needed. A disadvantage lies in the fact that there is very little ability of this gear to detect recruitment events. However, since this survey is designed to estimate exploitable biomass, this is not a critical issue.

The concurrent use of two different dredge configurations provides a means to not only test for agreement of results between the two gears, but also simultaneously conduct size selectivity experiments. In this instance, our experiment provided information regarding a recently mandated change to the commercial gear (CFTDD). While the expectation was that these changes should not affect the size selectivity characteristics of the gear (i.e. L_{50} and SR), as these characteristics are primarily determined by ring and mesh sizes, the possibility exists that the overall efficiency will be altered by different dredge frame design. Our results differed from Yochum and DuPaul (2008) with respect to L_{50} and SR. The estimate of L_{50} was lower by roughly 15 mm. This suggests that a number of tows in which high volumes of trash or scallops may have interfered with the passage of small scallops through the rings. This phenomenon is also reflected in a broad SR. Our estimated p value was very close to what was reported in Yochum and DuPaul (2008). This suggests a similar relative efficiency between the two dredge frames (Yochum and DuPaul (2008) used a New Bedford style dredge frame). These results, while different from other data sets, need to be taken in a broader context that includes different vessels, seasons and geographic regions. Given the major role that dredge efficiency plays in

the estimates of biomass from dredge surveys, it is clear that this topic is of critical importance and its refinement should be a high priority.

Biomass estimates are sensitive to other assumptions made about the biological characteristics of the resource; specifically, the use of appropriate shell height:meat weight parameters. Parameters generated from data collected during the course of the study were appropriate for the area and time sampled. There is however, a large variation in this relationship as a result of many factors. Seasonal and inter-annual variation can result in some of the largest differences in shell height:meat weight values. Traditionally, when the sea scallop undergoes its annual spawning cycle, metabolic energy is directed toward the production of gametes and the somatic tissue of the scallop is still recovering and is at some of their lowest levels relative to shell size (Serchuk and Smolowitz, 1989). While accurately representative for the month of the survey, biomass has the potential to be different relative to other times of the year. For comparative purposes, our results were also shown using the parameters from SARC 50 (NEFSC, 2010). These parameters reflect larger geographic regions (Georges Bank as well as NLCA) and are collected during the summer months. This allowed a comparison of results that may be reflective of some of the variations in biomass due to the fluctuations in the relationship between shell height and adductor muscle weight. Area and time specific shell height:meat weight parameters are another topic that merits consideration.

The survey of the NLCA during the late spring of 2012 provided a high-resolution view of the resource in this area. The NLCA is unique in that it will play a critical role in the spatial management strategy of the sea scallop resource over the next few years. With the other rotational areas of the mid-Atlantic (Hudson Canyon ,Elephant Trunk and DelMarVa) being either closed or nearing the end of their rotational cycles, the NLCA may have to carry some additional fishing pressure. While these data and subsequent analyses provide an additional source of information on which to base management decisions, it also highlights the need for further refinement of some of the components of industry based surveys. The use of industry based cooperative surveys provides an excellent mechanism to obtain the vital information to effectively regulate the sea scallop fishery in the context of an area management strategy.

Literature Cited

- DuPaul, W.D., E.J. Heist, and J.E. Kirkley, 1989. Comparative analysis of sea scallop escapement/retention and resulting economic impacts. College of William & Mary, Virginia Institute of Marine Science, Gloucester Point, VA. VIMS Marine Resource Report 88-10. 70 pp.
- DuPaul, W.D. and J.E. Kirkley, 1995. Evaluation of sea scallop dredge ring size. Contract report submitted to NOAA, National Marine Fisheries Service. Grant # NA36FD0131.
- Fryer, R.J. 1991. A model of between haul variation in selectivity. ICES J. of Mar. Sci. **48**: 281-290.
- Gedamke, T., W.D. DuPaul, and J.M. Hoenig. 2004. A spatially explicit open-ocean DeLury analysis to estimate gear efficiency in the dredge fishery for sea scallop *Placopecten magellanicus*. North American Journal of Fisheries Management 24:335-351.
- Gedamke, T., W.D. DuPaul, and J.M. Hoenig. 2005. Index-removal estimates of dredge Efficiency for sea scallops on Georges Bank. North American Journal of Fisheries Management 25:1122-1129.
- Holst, R. and A. Revill. 2009. A simple statistical method for catch comparison studies. Fisheries Research. **95**: 254-259.
- Lowther, A.. (Editor). 2012. Fisheries of the United States, 2011 NMFS Office of Science and Technology, Fisheries Statistics Division, Silver Spring, Maryland.
- Millar, R. B. 1992. Estimating the size-selectivity of fishing gear by conditioning on the total catch. Journal of the American Statistical Association. **87**: 962-968.
- Millar, R. B. 1993. Analysis of trawl selectivity studies (addendum): implementation in SAS. Fisheries Research. **17**: 373-377.
- Millar, R.B., M.K. Broadhurst, W.G. Macbeth. 2004. Modeling between-haul variability in the size selectivity of trawls. Fisheries Research. **67**:171-181.
- Millar, R.B. and R.J. Fryer. 1999. Estimating the size-selection curves of towed gears, traps, nets and hooks. Reviews Fish. Bio. Fish. **9**:89-116.
- Northeast Fisheries Science Center. 2010. 50th Northeast Regional Stock Assessment Workshop (45th SAW). 2007. 45th SAW assessment summary report. US Dep Commer, Northeast Fish Sci Cent Ref Doc. 10-09; 57 p.
- Park, H.H., R.B. Millar, H.C. An and H.Y. Kim. 2007. Size selectivity of drum-net traps for whelk (*Buccinum opisoplectum dall*) in the Korean coastal waters of the East Sea. Fish. Res. **86**: 113-119.
- Lowther, A.. (Editor). 2011. Fisheries of the United States, 2010. NMFS Office of Science and Technology, Fisheries Statistics Division, Silver Spring, Maryland.

- Sala, A., A. Lucchetti, C. Piccinetti and M. Ferretti. 2008. Size selection by diamond- and square-mesh codends in multi-species Mediterranean demersal trawl fisheries. *Fish. Res.* **93**:8-21.
- Serchuk, F.M. and Smolowitz, R.J. 1989. Seasonality in sea scallop somatic growth and reproductive cycles. *J. Shellfish Res.* 8:435.
- Stokesbury, K.D.E., B.P. Harris, M.C. Marino II, and J.I. Nogueira. 2004. Estimation of sea scallop abundance using a video survey in off-shore US waters. *Journal of Shellfish Research.* **23**:33-44.
- Stokesbury, K.D. 2002. Estimation of sea scallop abundance in closed areas of Georges Bank, USA. *Trans. of the Amer. Fish. Soc.* 131:1081-1092.
- Yochum, N. and DuPaul, W.D. 2008. Size-selectivity of the northwest Atlantic sea scallop (*Placopecten magellanicus*) dredge. *Journal of Shellfish Research* 27(2): 265-271.
- Xu, X and R.B. Millar. 1993. Estimation of trap selectivity for male snow crab (*Chionoecetes opilio*) using the SELECT modeling approach with unequal sampling effort. *Can J. Fish Aquat. Sci.* **50**: 2485-2490.

Table 1 Boundary coordinates of the access Area of Nantucket Lightship Closed Area.

NLCA	Latitude	Longitude
NLCA -1	40° 50' N	69° 30' W
NLCA -2	40° 30' N	69° 00' W
NLCA -3	40° 30' N	69° 14.5' W
NLCA -4	40° 50' N	69° 29.5' W
NLCA -5	40° 50' N	69° 30' W

Table 2 Summary statistics for the survey cruise.

Area	Cruise dates	Number of stations included in biomass estimate (survey dredge)	Number of stations included in biomass estimate (comm. dredge)
Nantucket Lightship Closed Area	May 24-28, 2012	108	108

Table 3 Mean total and mean exploitable scallop densities observed during the 2012 cooperative sea scallop survey of NLCA.

	Efficiency	Average Total Density (scallops/m²)	SE	Average Density of Exploitable Scallops (scallops/m²)	SE
Commercial	60%			0.167	0.021
Survey	38%	0.243	0.030	0.166	0.021

Table 4 Estimated number of scallops in the area surveyed. The estimate is based upon the estimated density of scallops at commercial dredge efficiency of 60% and survey dredge efficiency of 38%. The total area surveyed was estimated at 1,153.2 km².

	Efficiency	Estimated Total	Estimated Total Exploitable
NLCA			
Commercial	60%		192,316,171
Survey	38%	279,905,507	191,400,092

Table 5 Estimated average scallop meat weights for the area surveyed. Estimated weights are for the total size distribution of animals as represented by the catch from the NMFS survey dredge as well as the mean weight of exploitable scallops in the area as represented by the catches from both the survey and commercial dredge. Length:weight relationships from both SARC 50 as well as that observed from the cruise are shown.

	SH:MW	Mean Meat Weight (g) Total scallops	Mean Meat Weight (g) Exploitable scallops
Commercial	SARC 50 NCLA		38.99
Survey	SARC 50 NCLA	34.41	42.74
Commercial	SARC 50 W/ LAT		37.81
Survey	SARC 50 W/ LAT	33.26	41.63
Commercial	VIMS DEPTH WEIGHTED		45.43
Survey	VIMS DEPTH WEIGHTED	40.37	49.39
Commercial	VIMS		45.36
Survey	VIMS	40.29	49.34

Table 6 Mean catch of sea scallops observed during the 2012 VIMS-Industry cooperative surveys. Mean catch is depicted as a function of various shell height:meat weight relationships, either an area specific relationships derived from samples taken during the survey, or relationships from SARC 50. The top table depicts mean grams per tow of all scallops caught by the survey dredge. The bottom table depicts mean grams per tow for exploitable scallops caught by each gear.

	Samples	SH:MW	Mean Total (grams/tow)	Standard Error
Survey	108	SARC 50 NLCA	14,487.57	1,762.31
Survey	108	SARC 50 W/ LAT	14,004.61	1,689.57
Survey	108	VIMS DEPTH WEIGHTED	16,994.23	2,074.27
Survey	108	VIMS	16,961.48	2,073.57

	Samples	SH:MW	Mean Exploitable (grams/tow)	Standard Error
Commercial	108	SARC 50 NLCA	33,389.74	3,991.91
Survey	108	SARC 50 NLCA	12,283.64	1,503.43
Commercial	108	SARC 50 W/ LAT	32,380.87	3,844.35
Survey	108	SARC 50 W/ LAT	11,966.22	1,450.18
Commercial	108	VIMS DEPTH WEIGHTED	38,905.40	4,672.20
Survey	108	VIMS DEPTH WEIGHTED	14,195.42	1,743.44
Commercial	108	VIMS	38,844.16	4,667.78
Survey	108	VIMS	14,182.99	1,746.79

Table 7 Estimated total biomass of sea scallops observed during the 2012 VIMS-Industry cooperative survey. Biomass is presented as a function of different shell height:meat weight relationships, either an area specific relationship derived from samples taken during the actual survey or relationships from SARC 50.

	SH:MW	Efficiency	Total Biomass (mt)	95% CI	Lower Bound 95% CI	Upper Bound 95%CI
Survey	SARC 50 NCLA	38%	9,722.29	1,428.91	8,293.38	11,151.19
Survey	SARC 50 W/ LAT	38%	9,398.18	1,369.92	8,028.26	10,768.11
Survey	VIMS DEPTH WEIGHTED	38%	11,404.45	1,681.85	9,722.60	13,086.30
Survey	VIMS	38%	11,382.47	1,681.28	9,701.20	13,063.75

Table 8 Estimated exploitable biomass of sea scallops observed during the 2012 VIMS-Industry cooperative survey. Biomass is presented as a function of different shell height:meat weight relationships, either an area specific relationship derived from samples taken during the actual survey or relationships from SARC 50.

	SH:MW	Efficiency	Exploitable Biomass (mt)	95% CI	Lower Bound 95% CI	Upper Bound 95%CI
Commercial	SARC 50 NLCA	60%	7,568.63	1,373.78	6,194.85	8,942.40
Survey	SARC 50 NLCA	38%	8,243.28	1,219.00	7,024.28	9,462.28
Commercial	SARC 50 W/ LAT	60%	7,339.94	1,323.00	6,016.94	8,662.93
Survey	SARC 50 W/ LAT	38%	8,030.27	1,175.82	6,854.44	9,206.09
Commercial	VIMS DEPTH WEIGHTED	60%	8,818.89	1,607.89	7,211.00	10,426.78
Survey	VIMS DEPTH WEIGHTED	38%	9,526.23	1,413.61	8,112.62	10,939.84
Commercial	VIMS	60%	8,805.01	1,606.37	7,198.63	10,411.38
Survey	VIMS	38%	9,517.89	1,416.32	8,101.57	10,934.21

Table 9 Summary of area specific shell height:meat weight parameters used in the analyses. Parameters were obtained from two sources: (1) samples collected during the course of the surveys, and (2) SARC 50 (NEFSC, 2010).

	Date	α	β	γ	δ
Survey Data					
VIMS Depth Weighted	May, 2012	-6.8474	2.3440	-0.1590	
VIMS 2 parameter	May, 2012	-7.5356	2.3486		
SARC 50					
Georges Bank W/ Depth and Lat	-	9.6771	2.8387	-0.5084	-4.7629
NLCA specific	-	-8.1709	2.6454	-0.2298	

*The length weight relationship for sea scallops from data collected on the cruise is modeled as:

$$W = \exp(\alpha + \beta \ln(L))$$

For the relationship from SARC50 and the cruise that incorporates depth as a covariate

$$W = \exp(\alpha + \beta \ln(L) + \gamma \ln(D))$$

For SARC 50 (Georges Bank) depth and latitude term are included in the model as follows:

$$W = \exp(\alpha + \beta \ln(SH) + \gamma \ln(D) + \delta \ln(L))$$

Where W is meat weight in grams, SH is scallop shell height in millimeters (measured from the umbo to the ventral margin), D is depth in meters and L is latitude in decimal degrees.

Table 10 Catch per unit effort (a unit of effort is represented by one standard survey tow of 15 minute duration at 3.8 kts.) of finfish bycatch encountered during the survey of the Nantucket Lightship Closed Area during May 2012. In total, finfish bycatch was measured and recorded for 108 survey tows.

Common Name	Scientific Name	Commercial Dredge	Survey Dredge
Unclassified Skates	Raja spp.	76.99	27.98
Barndoor Skate	Raja laevis	1.91	0.58
Atlantic Cod	Gadus morhua	0.01	0.02
Haddock	Melanogrammus aeglefinus	0	0.01
Summer Flounder	Paralichthys dentatus	1.30	0.56
Fourspot Flounder	Paralichthys oblongotus	1.04	3.94
Yellowtail Flounder	Limanda ferruginea	0.59	2.13
Blackback Flounder	Psuedopleuronectes americana	0.85	0.72
Witch Flounder	Glyptocephalus cynoglossus	0.25	0.34
Windowpane Flounder	Scophthalmus aquasus	4.23	7.41
Monkfish	Lophius americanus	4.17	2.25

Table 11 Selection curve parameter estimates and hypotheses test. Selectivity data for each cruise was evaluated by a logistic curve with and without the split parameter (p) estimated. Improvements with respect to model fit were assessed by an examination of model deviance and AIC values.

	NLCA	
	Fixed p	Estimated p
a	-7.824	-5.625
b	0.133	0.064
p	0.500	0.728
L₂₅	42.190	53.500
L₅₀	58.670	87.750
L₇₅	75.140	122.030
Selection Range (SR)	16.470	34.280
Model Deviance	224.150	14.900
Degrees of Freedom	31	32
AIC	352.780	143.540

Table 12 Estimated logistic SELECT model fit for tows with total catch of greater than 50 scallops . Estimated parameters a , b and p as well as the length at 50% retention (L_{50}) and Selection Range (SR) are shown. The number of valid tows, as well as the replication estimate of between-haul variation (REP) is shown. This data set was determined to not be overdispersed and did not require an adjustment to the standard errors.

	NLCA	
Length Classes	27.5-182.5	
a	-5.616	1.067
b	1.064	0.014
p	0.727	0.020
L₅₀	87.66	26.00
Selection Range	34.26	7.81
REP	N/A	
# of tows in analysis	92	

Figure 1 Locations of sampling stations in the access area of Nantucket Lightship Closed Area survey by the F/V *Celtic* during the cruise conducted in May, 2012.




Figure 2 An example of the output Star-Oddi™ DST sensor. Arrows indicate the interpretation of the start and end of the dredge tow




Figure 3 Histogram of calculated tow lengths from the 2012 survey of the NLCA. Mean tow length was 1851.8 m with a standard deviation of 68.6 m.




Figure 4 Shell height frequencies for the two dredge configurations used to survey the access area of Nantucket Lightship Closed Area during May 2012. The frequencies represent the expanded but unadjusted catches of the two gears for all sampled tows.




Figure 5 Spatial distribution of sea scallop catches on the survey cruise of the access area of the Nantucket Lightship Closed Area during May 2012 by the NMFS survey dredge. This figure represents the catch of pre-recruit sea scallops (<70mm).




Figure 6 Spatial distribution of sea scallop catches on the survey cruise of the access area of the Nantucket Lightship Closed Area during May 2012 by the NMFS survey dredge. This figure represents the catch of recruit sea scallops (>70 mm).




Figure 7 Spatial distribution of sea scallop catches on the survey cruise of the access area of the Nantucket Lightship Closed Area during May 2012 by the CFTDD. This figure represents the catch of pre-recruit sea scallops (<70mm).




Figure 8 Spatial distribution of sea scallop catches on the survey cruise the access area of the Nantucket Lightship Closed Area during May 2012 by the CFTDD. This figure represents the catch of recruit sea scallops (>70 mm).




Figure 9 Shell height:meat weight relationships used in the study. The SARC-50 curve is an area specific curve for NLCA or a general relationship for Georges Bank. The VIMS 2012 curve is based on samples taken during the survey and is specific for the NLCA during May 2012 assuming an average depth for the area.




Figure 10 Top Panel: Logistic SELECT curve fit to the proportion of the total catch in the commercial dredge relative to the total catch (survey and commercial) for 2012 cruise to the NLCA. Bottom Panel: Deviance residuals for the model fit.





Figure 11 Estimated selectivity curve for the CFTDD based on data from the 2012 survey of the NLCA. The dashed line represents the length at 50% retention probability.

