



W&M ScholarWorks

Reports

11-1-1994

Button Bush Cephalanthus occidentalis L.

Gene Silberhorn Virginia Institute of Marine Science

Follow this and additional works at: https://scholarworks.wm.edu/reports



Part of the Plant Sciences Commons

Recommended Citation

Silberhorn, G. (1994) Button Bush Cephalanthus occidentalis L.. Wetland Flora Technical Reports, Wetlands Program, Virginia Institute of Marine Science. Virginia Institute of Marine Science, College of William and Mary. http://dx.doi.org/doi:10.21220/m2-9xjm-rh51

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Technical Report — Quetlands Program Wetland Flora



No. 94-10 / November 1994

Gene Silberhorn

Button Bush-

Cephalanthus occidentalis L.

Growth Habit and Diagnostic Characteristics ————

Button bush is a broad-leaved, deciduous shrub that grows up to 2 meters tall with an open spreading canopy. The simple, smooth marginate leaves are usually oppositely arranged throughout the lower branches, and are typically whorled (3 or 4 leaves at a node) just below the terminal borne fruit or flowering heads. Leaf petioles are often red during the peak flowering period when the white globose heads develop in July and August. In late August to September, the flowering heads mature to become compact, greenish, ball-like fruiting heads. By late autumn to early winter, the fruiting heads dry, turn brown, and eventually disintegrate into minute dry fruits called achenes. Button bush should not be confused with other wetland shrubs because of the combined diagnostic characteristics of whorled leaves, globose flowering and fruiting heads. Silky dogwood (Cornus amomum) and swamp loosestrife (Decodon verticillatus) have opposite leaves, but do not have ball-like flowers or fruits. Also, the latter is only semiwoody and does not branch.

Distribution -

Cephalanthus occidentalis is distributed throughout most of the eastern half of the United States from Florida to the Maritime Provinces of Canada, west to the Great Lake States and south to Mexico.

Button bush may occupy several different types of wetland habitats, including tidal and nontidal marshes, scrub/shrub and forested wetlands, and the margins of lakes, ponds, ditches and streams. In bottomland, hardwood forests dominated by tupelo (Nyssa aquatica) and bald cypress, Cephalanthus occidentalis is often associated with other hydrophytic shrubs such as swamp rose (*Rosa palustris*) and alder (Alnus serrulata). Cephalanthus frequently predominates in scrub/shrub wetlands where it may be associated with other shrubs such as swamp loosestrife, silky dogwood, swamp rose, alder and Virginia willow (Itea virginica).

Ecological Value / Benefits ————

Button bush frequently occupies the shrub strata of several different wetland types, therefore the broad range of ecological values attributed to wetlands in general may be accrued to this species.

Specifically, button bush flowers have high value for nectar gathering insects, the seeds are eaten by mallards and deer browse the foliage.

Hydrophytic Factor / Wetland Indicator Status —

According to the National List of Plant Species that Occur in Wetlands: Virginia (1988), Cephalanthus occidentalis is classified as an obligate wetland plant (OBL). OBLs are plants that almost always occur in wetlands (<99% probability).

Cephalanthus occidentalis L.



Wetlands Program School of Marine Science Virginia Institute of Marine Science College of William and Mary Gloucester Point, Virginia 23062 Dr. Carl Hershner, Program Director This report was funded, in part, by the Department of Environmental Quality's Coastal Resources Management Program through Grant No. NA470Z0287-01 of the National Oceanic and Atmospheric Administration, Office of Ocean and Coastal Resource Management, under the Zone Management Act of 1972, as amended.

Illustration by Kent Forrest

Printed on recycled paper.