



2014

Aberrant *myst4/brpf1* Signaling Misdirects Regional Neurogenesis Programs, Sustaining Expression of Self-Renewal Genes in Pediatric Brain Cancers

Timothy E. Van Meter

Asadullah Khan

Kevin Fitzmartin

Timothy E. Van Meter

College of William and Mary

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Van Meter, T. E., Khan, A., Fitzmartin, K., Taylor, M. D., Mack, S., & Cote, J. (2014). Eg-16aberrant *Myst4/brpf1* Signaling Misdirects Regional Neurogenesis Programs, Sustaining Expression Of Self-renewal Genes In Pediatric Brain Cancers. *Neuro-oncology*, 16(Suppl 5), v78.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Abstracts

EG-16. ABERRANT *Myst4*/Brpf1 SIGNALING MISDIRECTS REGIONAL NEUROGENESIS PROGRAMS, SUSTAINING EXPRESSION OF SELF-RENEWAL GENES IN PEDIATRIC BRAIN CANCERS

Timothy E. Van Meter^{1,2}, Asadullah Khan¹, Kevin Fitzmartin¹, Michael D. Taylor³, Stephen Mack³, and Jacques Cote⁴; ¹VCU School of Medicine, Richmond, VA, USA; ²College of William and Mary, Williamsburg, VA, USA; ³Hospital for Sick Children, Toronto, ON, Canada; ⁴University of Laval, Quebec, Canada

The *Myst* family of acetyltransferase proteins has been shown to play important role in development and self-renewal, primarily through lysine acetylation on Histones H3 and H4. In neural development, *Myst4* (*Morf/KAT6B*) has been found to be critical for self-renewal and for neuron generation in the

developing nervous system and during adult neurogenesis. This chromatin modifier exists in a complex with *Myst3* (*Moz/KAT6A*), the bromodomain factor *Brpf1*, which acts as a protein scaffold, targeting histone acetyltransferases to chromatin, as well as Inhibitor of Growth 5 (*ING5*) and *Esa1*-associated factor 6 (*EAF6*). Our laboratories have found alteration of these elements in pediatric brain cancers, suggesting a pathological role in abnormal neural progenitor growth. Exon-specific microarrays, DNA methylation studies and functional perturbation were performed to study the impact on tumor behavior. RNA interference in tumors and neural progenitors led to loss of H3K4acetylation in target genes and altered expression, including *Spondin-1*. Conversely, exogenous over-expression of targeting factors, such as *Brpf1*, enhanced expression of target genes, assessed by quantitative PCR analysis. Furthermore, expression of *Myst3/4* and *Brpf1* positively correlated with tumor malignancy markers in large patient cohorts, including *Ki67*, *PCNA*, and *MELK*, and with decreased overall survival. Assessment of direct functional relationships with tumor markers using ChIP-Seq approaches is currently being pursued to investigate direct *Myst3/4*/*Brpf1*-mediated promoter activity at known and novel target genes. These studies aim to elucidate the role of an important epigenetic mechanism in neurogenesis, the alteration of which may underlie global chromatin changes that contribute to tumor growth or initiation. Sustained progenitor growth may be suppressed by targeted therapies that disrupt these factors.