



W&M ScholarWorks

Reports

1-9-1981

A Definition of a Natural Oyster Rock

Dexter S. Haven Virginia Institute of Marine Science

Herbert M. Austin Virginia Institute of Marine Science

Follow this and additional works at: https://scholarworks.wm.edu/reports



Part of the Aquaculture and Fisheries Commons

Recommended Citation

Haven, D. S., & Austin, H. M. (1981) A Definition of a Natural Oyster Rock. Marine Resource Report No. 81-2. Virginia Institute of Marine Science, College of William and Mary. https://dx.doi.org/doi:10.25773/ v5-xrfz-mk53

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

A DEFINITION OF A NATURAL OYSTER ROCK

Prepared for
The Offices of Legislative Services
Virginia General Assembly

Ву

D. S. Haven, H. M. Austin, and

W. J. Hargis, Jr.

Virginia Institute of Marine Science and

School of Marine Science,
The College of William and Mary
Gloucester Point, Virginia 23062

January 9, 1981

VIRGINIA INSTITUTE OF MARINE SCIENCE SCHOOL OF MARINE SCIENCE

Gloucester Point, Virginia 23062

January 9, 1981

Phone (804) 642-21.11

12

MAL

Mr. Bernie Caton Legislative Services 2nd Floor P. O. Box 3-AG 910 Capitol Street Richmond, Virginia 23208

Dear Bernie:

Just prior to the Christmas holiday you asked Dexter to prepare a definition of a naturally productive oyster bottom. After considerable thought, we arrived at the definition which is enclosed with this letter. As you will see, this definition utilizes concepts derived from our four-year study of Baylor bottoms. This study is now in its final stages. When it is completed copies of the charts showing bottom types, as well as a text discussing other aspects will be forwarded to various State agencies. If desired, copies can be prepared for Legislative purposes.

While we believe that the definition submitted to you is accurate, we must point out that we have previously submitted to your office a study entitled, "A Classification Baylon Bottoms in Virginia." This is dated October 9 3 To divides many of the State's Baylor bottoms inco while closes with Class I being the most useful and ${f V}$ being Alexa of value. The option given in this report is the stative branch may decide on areas which might as saleted as a redefinition of Baylor bottoms.

Perhaps at some future data our definition which obtained is enclosed has a helpful in a further intergradation of these two studie

If we can be of further help, please let Dexter or me know.

Sincerely,

Herbert M. Austin

Assistant Director

HMA/gbr

Orglosure

ែក ខែក្រឡុងថ

A Definition of a Natural Oyster Rock

by

D. S. Haven, H. M. Austin, and W. J. Hargis, Jr.

Virginia Institute of Marine Science and School of Marine Science, The College of William and Mary Gloucester Point, Virginia 23062

January 9, 1981

A natural oyster rock is defined as a productive area of bottom which has a functional set which will support a reasonable level of commercial harvest for market or seed oysters. A set may typically occur each year, but the annual interval between sets may be as long as 15 years.

Such bottoms would necessarily have substrate classed as rock, mud-shell, sand-shell or buried shell as defined by the Baylor Ground study to be completed by the Virginia Institute of Marine Science in April 1981.

Moreover, the total volumes of shell, oysters and cinder in the substrate in any designated areas should equal or exceed on the average 10% by volume of the material in the upper five inches of the bottom.