



## W&M ScholarWorks

---

Arts & Sciences Articles

Arts and Sciences

---

2014

### First result from Q(weak)

David S. Armstrong  
*College of William and Mary*

Qweak Collaboration  
*College of William and Mary*

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

---

#### Recommended Citation

Armstrong, David S. and Qweak Collaboration, First result from Q(weak) (2014).  
10.1051/epjconf/20147307008

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).

## First result from $Q_{\text{weak}}$

David S. Armstrong<sup>a</sup> (for the  $Q_{\text{weak}}$  Collaboration)

Physics Dept., College of William & Mary, Williamsburg, VA 23187, USA

**Abstract.** Initial results are presented from the recently-completed  $Q_{\text{weak}}$  experiment at Jefferson Lab. The goal is a precise measurement of the proton's weak charge  $Q_w^p$ , to yield a test of the standard model and to search for evidence of new physics. The weak charge is extracted from the parity-violating asymmetry in elastic  $\vec{e}p$  scattering at low momentum transfer,  $Q^2 = 0.025 \text{ GeV}^2$ . A  $180 \mu\text{A}$  longitudinally-polarized 1.16 GeV electron beam was scattered from a 35 cm long liquid hydrogen at small angles,  $6^\circ < \theta < 12^\circ$ . Scattered electrons were analyzed in a toroidal magnetic field and detected using an array of eight Cerenkov detectors arranged symmetrically about the beam axis. The initial result, from 4% of the complete data set, is  $Q_w^p = 0.064 \pm 0.012$ , in excellent agreement with the standard model expectation. Full analysis of the data is expected to yield a value for the weak charge to about 5% precision.

### 1. Introduction

It is conventional wisdom that the enormously successful  $SU(3)_C \times SU(2)_L \times U(1)_Y$  standard model is actually a low-energy effective theory of some more complete higher-mass scale physics. Attempts to probe such possible new physics include direct searches at colliders (the energy frontier), and indirect searches (the precision or intensity frontier). The hallmark of the latter method is the use of observables which are well-predicted within the standard model, but for which new physics could alter those predictions, eg. via loop corrections or the exchange of new particles. One such example is the weak charge of the proton, which can be accessed in neutral-current processes such as parity-violating electron scattering (PVES) [1, 2].

The tree-level standard model expression for the proton's weak charge is

$$Q_w^p = 1 - 4 \sin^2(\theta_W) \quad (1)$$

where  $\theta_W$  is the electroweak mixing angle. It can also be expressed in terms of the weak vector couplings of the light quarks,  $Q_w^p = -2(2C_{1u} + C_{1d})$  which are again functions of the mixing angle. These expressions are modified beyond tree level by loop corrections [3], however after recent theoretical work on the troublesome  $\gamma Z$  box term [4–8], these are now sufficiently under control, and the standard model predictions for  $Q_w^p$  and for  $C_{1u}$  and  $C_{1d}$  are robust. The fortuitous modest suppression of the

---

<sup>a</sup>e-mail: [armd@physics.wm.edu](mailto:armd@physics.wm.edu)

numerical value of  $Q_w^p$ , since  $\sin^2(\theta_W)$  is near 0.23, increases the sensitivity of the measurement to potential new physics effects.

The experimental observable is the parity-violating asymmetry, the difference over the sum of the elastic scattering cross section for electrons with positive and negative helicity,

$$A_{ep} = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-}. \quad (2)$$

At tree level, in terms of electromagnetic, neutral-weak, and axial form factors, this is

$$A_{ep} = A_0 \left[ \frac{\varepsilon G_E^\gamma G_E^Z + \tau G_M^\gamma G_M^Z - (1 - 4 \sin^2 \theta_W) \varepsilon' G_M^\gamma G_A^Z}{\varepsilon (G_E^\gamma)^2 + \tau (G_M^\gamma)^2} \right] \quad (3)$$

where  $A_0 = \frac{-G_F Q^2}{4\pi\alpha\sqrt{2}}$ ,  $\varepsilon = [1 + 2(1 + \tau) \tan^2 \frac{\theta}{2}]^{-1}$ ,  $\varepsilon' = \sqrt{\tau(1 + \tau)(1 - \varepsilon^2)}$ ,  $G_F$  is the Fermi constant,  $-Q^2$  the four-momentum transfer squared,  $\alpha$  the fine structure constant,  $\tau = Q^2/4M^2$ ,  $M$  the proton mass,  $\theta$  the laboratory electron scattering angle and the weak charge appears as the  $Q^2 \rightarrow 0$  limit of  $G_E^Z$ .

Following [9], we can recast Eq. 3 for  $\theta \rightarrow 0$  as

$$A_{ep}/A_0 = Q_W^p + Q^2 B(Q^2, \theta), \quad (4)$$


wherein all the hadron structure dependence is subsumed in the  $B(Q^2, \theta)$  factor. Thus, a measurement of  $A_{ep}$  at forward angles and low  $Q^2$  gives access to the weak charge.

## 2. Experiment


The experiment built on techniques developed at Jefferson Lab over the last two decades for conducting precision parity-violating electron scattering measurements [10]. Of particular importance is the superbly small level of helicity-correlated fluctuations in beam properties available (through close cooperation with the electron-gun group and the accelerator operations group). Innovations for this experiment included the use of a very rapid reversal of the beam helicity (960 Hz) and the highest intensity beam yet used at JLab (180  $\mu$ A). The measurement took place during a two year period, and was conducted in Hall C, with a dedicated apparatus [11]. A toroidal-field magnetic spectrometer with an octagonal array of radiation-hard quartz Cherenkov detectors isolated electrons elastically scattered from a high-power (3 kW) thick liquid hydrogen target [12].

The kinematics (1.16 GeV incident energy, scattering angle  $\sim 8^\circ$ ,  $Q^2 = 0.025 \text{ GeV}^2$ ) were chosen to suppress the hadron-structure contribution to the asymmetry (especially the axial form-factor) while maintaining an acceptable figure of merit for the expected asymmetry ( $\sim 200$  ppb). The anode current signals from two PMTs mounted on each Cherenkov detector were integrated over each  $\sim 1$  ms helicity state and digitized using 18 bit ADCs sampling at 500 kHz. The kinematics were verified in special low beam-current calibration measurements obtained using a particle tracking system based on drift chambers located before and after the spectrometer magnet.

During the experiment the beam polarization ( $\sim 89\%$ ) was periodically measured using the Hall C Møller polarimeter [13] and, for the latter part of the experiment, continuously measured with a newly-developed Compton polarimeter based on a circularly polarized green laser in a low-gain cavity. The agreement between the results from the two polarimeters is excellent. For the initial results presented here, only Møller polarimeter data were available.



**Figure 1.** World data for reduced asymmetries from forward-angle PVES on the proton, including the present result, for measurements up to  $Q^2 = 0.63 (GeV)^2$ , presented in the forward angle limit (see text). The solid line is the global fit based on these data as well as on forward-angle  $^4\text{He}$  and deuterium data. The additional uncertainty arising from the rotation is indicated by outer error bars. The shaded region indicates the uncertainty in the fit.  $Q_W^p(\text{expt})$  is the intercept of the fit. The standard model (SM) prediction is also shown (arrow).



**Figure 2.** The running of the weak mixing angle with energy scale  $Q$ . The curve is the standard model predicted running in the MS scheme [2]. Data include collider results (LEP [17], SLD [18], Tevatron [19, 20]) at or near the Z-pole, the SLAC E158 electron-electron parity violation experiment [21], the Colorado atomic parity violation (APV) experiment on  $^{133}\text{Cs}$  [22, 23], NuTeV [24] (this latter is controversial, we show the value from [2]), and the present result; the inset shows the anticipated final precision of our experiment.

### 3. Results

We report here results from initial “commissioning” data, which represent about 4% of our full data set; these results appeared in the literature the same week as this conference [14]. After all corrections, the measured asymmetry was  $A_{ep} = -279 \pm 35$  (statistics)  $\pm 31$  (systematics) ppb.

In order to extract  $Q_W^p$  from the measured asymmetry  $A_{ep}$ , one needs to subtract the hadronic form factor piece in Eq. (4), *i.e.* the  $Q^2 B(Q^2, \theta)$  term, which vanishes at sufficiently low  $Q^2$ . This was done by exploiting previously-measured PVES asymmetries from the SAMPLE, HAPPEX, G0, and PVA4 experiments (see Ref. [10] and references therein), all of which were at significantly higher  $Q^2$  than the present result. These included data for hydrogen, deuterium and  $^4\text{He}$ , which were combined with our result in a global fit following the method outlined in [9]. All PVES data up to  $0.63 \text{ GeV}^2$  were used. The fit had five free parameters: the light quark weak charges  $C_{1u}$  and  $C_{1d}$ , the strange charge radius  $\rho_s$  and magnetic moment  $\mu_s$ , and the isovector axial form factor  $G_A^Z(T=1)$ . The isoscalar  $G_A^Z(T=0)$  was constrained by theory [15]. For the electromagnetic form factors  $G_E^\gamma$  and  $G_M^\gamma$  the parameterization of Kelley [16] was adopted, and the data were corrected for the  $\gamma Z$  box diagram contributions.

To display the quality of the fit, the  $\theta$  dependence of the proton-target data was removed using Eq. (3), and the asymmetries were divided by  $A_0$ , and plotted vs.  $Q^2$ ; see Fig. 1. The fitted intercept is  $Q_W^p(\text{expt}) = 0.064 \pm 0.012$ , in excellent agreement with the standard model (SM) expectation  $Q_W^p(\text{SM}) = 0.0710 \pm 0.0007$ , and represents the first direct extraction of the proton’s weak charge.

The “running” value of weak mixing angle at our kinematics can be extracted from  $Q_W^p$  [25], and is shown in Fig. 2, along with other precision determinations; again, our present result conforms well with standard model expectations.

The full analysis of the data is well underway. We anticipate a final precision on the weak charge of about 5%, which will probe certain classes of new physics at the multi-TeV scale.

I thank all the members of the  $Q_{\text{weak}}$  collaboration for their efforts and camaraderie. This work was supported by DOE contract No. DE-AC05-06OR23177, under which Jefferson Science Associates, LLC operates Thomas Jefferson National Accelerator Facility, as well as by NSERC (Canada) and the NSF (USA). The W&M group is grateful for support from the NSF, including grants PHY-1068667, PHY-0758068 and PHY-0320942.

## References

- [1] M.J. Musolf, T.W. Donnelly, J. Dubach, S.J. Pollock, S. Kowalski, and E.J. Beise, *Phys. Rep.* **239**, 1 (1994)
- [2] J. Erler and S. Su, *Prog. Part. Nucl. Phys.* **71**, 119 (2013)
- [3] K.S. Kumar, Sonny Mantry, W.J. Marciano, and P.A. Souder, *Annu. Rev. Nucl. Part. Sci.* **63**, 237 (2013)
- [4] M. Gorchtein and C.J. Horowitz, *Phys. Rev. Lett.* **102**, 091806 (2009)
- [5] A. Sibirtsev, P.G. Blunden, W. Melnitchouk, and A.W. Thomas, *Phys. Rev. D* **82**, 013011 (2010)
- [6] B.C. Rislow and C.E. Carlson, *Phys. Rev. D* **83**, 113007 (2011)
- [7] M. Gorchtein, C.J. Horowitz, and M.J. Ramsey-Musolf, *Phys. Rev. C* **84**, 015502 (2011)
- [8] N.L. Hall, P.G. Blunden, W. Melnitchouk, A.W. Thomas and R.D. Young, *Phys. Rev. D* **88**, 013011 (2014)
- [9] R.D. Young, R.D. Carlini, A.W. Thomas, and J. Roche, *Phys. Rev. Lett.* **99**, 122003 (2007)
- [10] D.S. Armstrong and R.D. McKeown *Annu. Rev. Nucl. Part. Sci.* **62**, 337 (2012)
- [11] R. D. Carlini et al., JLAB-PHY-12-1478, [arXiv:1202.1255](https://arxiv.org/abs/1202.1255) (2007)
- [12] G.R. Smith, *Nuovo Cimento* **35C**, No. 4, 159 (2012)
- [13] M. Hauer et al., *Nucl. Instrum. Meth. A* **462**, 382 (2001)
- [14] D. Androic et al. (Qweak Collaboration), *Phys. Rev. Lett.* **111**, 141803 (2013)
- [15] S.L. Zhu, S.J. Puglia, B.R. Holstein and M.J. Ramsey-Musolf, *Phys. Rev. D* **62**, 033008 (2000)
- [16] J.J. Kelly, *Phys. Rev. C* **70**, 068202 (2004)
- [17] ALEPH Collaboration et al., [arXiv:1012.2367](https://arxiv.org/abs/1012.2367) (2010)
- [18] K. Abe et al. (SLD Collaboration), *Phys. Rev. Lett.* **84**, 141803 (2013)
- [19] V.M. Abazov et al. (D0 Collaboration), *Phys. Rev. D* **84**, 012007 (2011)
- [20] J. Han et al., (CDF Collaboration), [arXiv:1110.0153](https://arxiv.org/abs/1110.0153) (2011)
- [21] P.L. Anthony et al. (SLAC E158 Collaboration), *Phys. Rev. Lett.* **95**, 081601 (2005)
- [22] C.S. Wood, S.C. Bennett, D. Cho, B.P. Masterson, J.L. Roberts, C.E. Tanner, and C.E. Wieman, *Science* **275**, 1759 (1997)
- [23] V.A. Dzuba, J.C. Berengut, V.V. Flambaum and B. Roberts, *Phys. Rev. Lett.* **109**, 203003 (2012)
- [24] G. Zeller et al. (NuTeV Collaboration), *Phys. Rev. Lett.* **88**, 091802 (2002)
- [25] J. Erler, A. Kurylov, and M.J. Ramsey-Musolf, *Phys. Rev. D* **68**, 016006 (2003)