

W&M ScholarWorks

Arts & Sciences Articles

Arts and Sciences

2014

# Existence of positive solutions to Kirchhoff type problems with zero mass

Yuhua Li

Fuyi Li

Junping Shi

Junping Shi College of William and Mary

Follow this and additional works at: https://scholarworks.wm.edu/aspubs

### **Recommended Citation**

Li, Y., Li, F., & Shi, J. (2014). Existence of positive solutions to Kirchhoff type problems with zero mass. Journal of Mathematical Analysis and Applications, 410(1), 361-374.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa



## Existence of positive solutions to Kirchhoff type problems with zero mass $^{\updownarrow}$



Yuhua Li<sup>a</sup>, Fuyi Li<sup>a,\*</sup>, Junping Shi<sup>a,b</sup>

<sup>a</sup> School of Mathematical Sciences, Shanxi University, Taiyuan 030006, Shanxi, PR China

<sup>b</sup> Department of Mathematics, College of William and Mary, Williamsburg, VA 23187-8795, USA

#### ARTICLE INFO

Article history: Received 16 May 2013 Available online 26 August 2013 Submitted by V. Radulescu

Keywords: Kirchhoff type problem Zero mass Variational method Cut-off functional

#### ABSTRACT

The existence of positive solutions depending on a nonnegative parameter  $\lambda$  to Kirchhoff type problems with zero mass is proved by using variational method, and the new result does not require usual compactness conditions. A priori estimate and a Pohozaev type identity are used to obtain the bounded Palais–Smale sequences for constant coefficient nonlinearity, while a cut-off functional and Pohozaev type identity are utilized to obtain the bounded Palais–Smale sequences for the variable-coefficient case.

© 2013 Elsevier Inc. All rights reserved.

#### 1. Introduction

In this paper, we consider positive solutions to the following nonlinear Kirchhoff type problem

$$\begin{cases} -\left(a+\lambda\int\limits_{\mathbb{R}^{N}}|\nabla u|^{2}\right)\Delta u=K(x)f(u), & \text{in } \mathbb{R}^{N},\\ u\in\mathcal{D}^{1,2}(\mathbb{R}^{N}), \end{cases}$$
(1.1)

where  $N \ge 3$ , *a* is a positive constant,  $\lambda \ge 0$  is a parameter, and *K* is a potential function. Kirchhoff type problem on a bounded domain  $\Omega \subset \mathbb{R}^N$ 

$$\begin{cases} -\left(a+b\int_{\Omega}|\nabla u|^{2}\right)\Delta u = f(u), & \text{in }\Omega, \\ u = 0, & \text{on }\partial\Omega \end{cases}$$
(1.2)

has been studied by many authors, for example [5-7,10,18-20,24,27,28]. Many solvability conditions on the nonlinearity f near zero and infinity for the problem (1.2) have been considered, such as the superlinear case [19]; and asymptotical linear case [24]. In addition, the following growth condition on f is often assumed:

(f)  $f(t)t \ge 4F(t)$  for |t| large, where  $F(t) = \int_0^t f(s) ds$ ,

\* Corresponding author. Fax: +86 0351 7011733.
E meil address: filesum odu ep (E Li)

<sup>\*</sup> Partially supported by National Natural Science Foundation of China (Grant No. 11071149, 11301313), and Science Council of Shanxi Province (2010011001-1, 2012011004-2, 2013021001-4) and Shanxi 100 Talent program.

E-mail address: fyli@sxu.edu.cn (F. Li).

<sup>0022-247</sup>X/\$ – see front matter @ 2013 Elsevier Inc. All rights reserved. http://dx.doi.org/10.1016/j.jmaa.2013.08.030

which assures the boundedness of any Palais-Smale (PS) or Cerami sequence. Indeed the condition (f) may appear in different forms as follows:

- (f<sub>0</sub>) there exists  $\theta \ge 1$  such that  $\theta G(t) \ge G(st)$  for all  $t \in \mathbb{R}$  and  $s \in [0, 1]$ , where G(t) = tf(t) 4F(t) (see [24]);
- (f<sub>1</sub>)  $\lim_{|t|\to\infty} G(t) = \infty$  (see [27]); or
- (f<sub>2</sub>)  $\lim_{|t|\to\infty} G(t) = \infty$  and there exists  $\sigma > \max\{1, N/2\}$  such that  $|f(t)|^{\sigma} \leq CG(t)|t|^{\sigma}$  for |t| large (see [19]).

In the papers above, each of the conditions  $(f_0)-(f_2)$  implies that the condition (f) holds. On the other hand, the condition (f) is sufficient to show the boundedness of any (PS) or Cerami sequence, which has been proved in [26].

There are few papers considering Kirchhoff type problems on  $\mathbb{R}^N$  except [8.9.12,17,25,26]. In [26], the author studied the problem

$$-\left(a+b\int_{\mathbb{R}^N}|\nabla u|^2\right)\Delta u+V(x)u=f(u),\quad\text{in }\mathbb{R}^N.$$
(1.3)

The existence of nontrivial solutions was proved in [26] under the condition (f) and

- (V)  $V \in C(\mathbb{R}^N, \mathbb{R})$ ,  $\inf_{x \in \mathbb{R}^N} V(x) > 0$  and for each M > 0, meas  $\{x \in \mathbb{R}^N : V(x) \leq M\} < \infty$ ; (f<sub>3</sub>)  $f \in C(\mathbb{R}_+, \mathbb{R}_+)$  and  $|f(t)| \leq C(|t| + |t|^{p-1})$  for all  $t \in \mathbb{R}_+ = [0, \infty)$  and some  $p \in (1, 2^*)$ , where  $2^* = 2N/(N-2)$  for (1)  $N \ge 3;$ (f<sub>4</sub>)  $\lim_{t\to 0} \frac{f(t)}{t} = 0;$ (f<sub>5</sub>)  $\lim_{t\to\infty} \frac{F(t)}{t^4} = \infty.$

In [26], the space  $\{u \in H^1(\mathbb{R}^N): \int_{\mathbb{R}^N} [|\nabla u|^2 + V(x)u^2] < \infty\}$  is compactly embedded into  $L^p(\mathbb{R}^N)$ , which makes the existence problem easier. In [8,9,25], the existence, multiplicity and concentration behavior of positive solutions of (1.3) were considered by relating the number of solutions with the topology of the set where V attains its minimum.

In [17], we proved the existence of a positive solution to the problem

$$\left(a + \lambda \int_{\mathbb{R}^N} |\nabla u|^2 + \lambda b \int_{\mathbb{R}^N} u^2\right) [-\Delta u + bu] = f(u), \quad \text{in } \mathbb{R}^N.$$
(1.4)

We assumed that a > 0, b > 0, and f satisfies  $(f_3)$ ,  $(f_4)$  and the following condition

(f<sub>6</sub>) 
$$\lim_{t\to\infty}\frac{f(t)}{t} = \infty$$
.

The result in [17] does not assume the condition (f) (or any of  $(f_0)$ - $(f_2)$ ), as the space  $H_r^1(\mathbb{R}^N)$  of radial functions can be compactly imbedded into  $L^p(\mathbb{R}^N)$ , which provides some compactness in the problem for the convergence. However, for the case of b = 0 in (1.4) (that is (1.1) with  $K \equiv 1$ ), one has to search for a positive solution in the space  $\mathcal{D}_{r}^{1,2}(\mathbb{R}^{N})$ , which does not possess compactness as  $H_r^1(\mathbb{R}^N)$ . Because of this difficulty, there are very few works up to now studying Kirchhoff problems with zero mass, i.e. the problem (1.1).

In this paper we consider the existence of positive solutions to (1.1), and we assume the following conditions which are considerably weaker than the ones in the previous works:

- (K<sub>0</sub>)  $K(x) \equiv 1$  for  $x \in \mathbb{R}^N$ ;
- (K<sub>1</sub>) let  $K : \mathbb{R}^N \to \mathbb{R}$  be a nonnegative continuous function and  $K \in [L^s(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)] \setminus \{0\}$  for some  $s \ge 2N/(N+2)$ ;
- (K<sub>2</sub>)  $|x \cdot \nabla K(x)| \leq \alpha K(x)$  for a.e.  $x \in \mathbb{R}^N$  and some  $\alpha \in (0, 2)$ ;
- (H<sub>1</sub>)  $f \in C(\mathbb{R}_+, \mathbb{R}_+)$  and  $\lim_{t\to 0^+} \frac{f(t)}{t^{2^*-1}} = 0;$
- (H<sub>2</sub>)  $\lim_{t\to\infty} \frac{f(t)}{t^{2^*-1}} = 0;$ (H<sub>3</sub>)  $\lim_{t\to\infty} \frac{f(t)}{t} = \infty.$

Our first result is for (1.1) with a constant potential function K(x), or equivalently for Eq. (1.4) with b = 0.

**Theorem 1.1.** Assume that  $N \ge 3$ , a is a positive constant, and  $\lambda \ge 0$  is a parameter. If the conditions  $(K_0)$ ,  $(H_1)$ ,  $(H_2)$  and  $(H_3)$  hold, then there exists  $\lambda_0 > 0$  such that for any  $\lambda \in [0, \lambda_0)$ , (1.1) has at least one positive solution.

Theorem 1.1 appears to be the first existence result for the problem (1.1). We remark also that the condition  $(H_3)$  is weaker than the ones in the papers mentioned above, in which  $\lim_{t\to\infty} f(t)/t^3 = \infty$  or a positive constant (which implies (H<sub>3</sub>)) was assumed. From Theorem 1.1, one can also have the following classical result (see [4]) if we let  $\lambda = 0$ .

**Corollary 1.2.** Assume that  $N \ge 3$ . If the conditions (H<sub>1</sub>), (H<sub>2</sub>) and (H<sub>3</sub>) hold, then the equation  $-\Delta u = f(u)$  in  $\mathbb{R}^N$  has at least one positive solution.

In our second result, we consider the case of non-constant potential function K(x), and we obtain the following result:

**Theorem 1.3.** Assume that  $N \ge 3$ , a is a positive constant, and  $\lambda \ge 0$  is a parameter. If the conditions  $(K_1)$ ,  $(K_2)$ ,  $(H_1)$ ,  $(H_2)$  and  $(H_3)$  hold, then there exists  $\lambda_0 > 0$  such that for any  $\lambda \in [0, \lambda_0)$ , (1.1) has at least one positive solution.

An example of K satisfying  $(K_1)$  and  $(K_2)$  is

$$K(x) = \frac{1}{1+|x|^{\alpha}},$$

where  $\alpha \in (1, 2)$ . One can easily verify that  $|x \cdot \nabla K(x)| = \frac{\alpha |x|^{\alpha}}{(1+|x|^{\alpha})^2}$ . Therefore, *K* satisfies the conditions (K<sub>1</sub>) for  $s \ge N$  and (K<sub>2</sub>).

The case of  $\lambda = 0$  in (1.1) corresponds to the following well-known nonlinear Poisson equation:

$$-\Delta u = K(x) f(u), \quad x \in \mathbb{R}^N.$$
(1.5)

Positive solutions of (1.5) with radially symmetric potential function have been investigated in, for example, [1,3,2,14–16]. For example, in [15], the case of  $K(x) = 1/(1 + |x|^2)$  and  $f(u) = u^p$  was studied and asymptotic expansions of solutions were obtained. In [14], the author assumed that  $f(u) = u^p$  and K satisfies some conditions similar to (K<sub>1</sub>) and (K<sub>2</sub>). The zero mass case for (1.5) was considered in [3,2]. Our result here implies the following corollary for (1.5):

**Corollary 1.4.** Assume that  $N \ge 3$ . If the conditions  $(K_1)$ ,  $(K_2)$ ,  $(H_1)$ ,  $(H_2)$  and  $(H_3)$  hold, then the equation  $-\Delta u = K(x) f(u)$  in  $\mathbb{R}^N$  has at least one positive solution.

In [1], the existence of a positive solution was proved for  $K \in L^{s}(\mathbb{R}^{N})$  with an additional condition

(H<sub>4</sub>) H(t) = tf(t) - 2F(t) is increasing in t and H(0) = 0.

In Corollary 1.4, we do not assume this monotonicity condition.

As in [17], we prove in this paper the existence of positive solutions to (1.1) without the condition (f) (or any of (f<sub>0</sub>)–(f<sub>2</sub>)). We use *a priori* estimate, a cut-off functional and a variable-coefficient Pohozaev type identity to obtain bounded (PS) sequences, and then we apply some known variational techniques to prove the existence of a positive solution. The Pohozaev identity with variable-coefficient proved in Lemma 2.2 seems to be the first of this kind for (1.1), which is of independent interest. Similar Pohozaev identities with variable-coefficient have also been obtained in [21,22] for *p*-Laplace equations with singular weight. Another difficulty in (1.1) is caused by the nonlocal term  $\int_{\mathbb{R}^N} |\nabla u|^2$ , which leads to some convergence difficulties, that is, if  $u_n$  converges weakly to u, then one cannot conclude that u is a weak solution of (1.1). So we must obtain strong convergence of the (PS) sequence, and this makes (1.1) more difficult to deal with than other similar elliptic equations.

In this paper the problem (1.1) is considered in the Sobolev space  $\mathcal{D}^{1,2}(\mathbb{R}^N) = \{u \in L^{2^*}(\mathbb{R}^N): |\nabla u| \in L^2(\mathbb{R}^N)\}$ . The space  $\mathcal{D}^{1,2}(\mathbb{R}^N)$  is equipped with the standard inner product and norm

$$(u, v) = \int_{\mathbb{R}^N} \nabla u \cdot \nabla v, \qquad \|u\| = (u, u)^{1/2}$$

When *K* is a constant, for obtaining the convergence, we consider the problem (1.1) in the subspace  $\mathcal{D}_r^{1,2}(\mathbb{R}^N)$  of  $\mathcal{D}^{1,2}(\mathbb{R}^N)$  consisting of radial functions. Then we have that  $\mathcal{D}_r^{1,2}(\mathbb{R}^N) \hookrightarrow L^{2^*}(\mathbb{R}^N)$  continuously. We denote by  $|\cdot|_q$  the usual  $L^q(\mathbb{R}^N)$  norm. In this paper, we consider only positive solutions to (1.1), so we assume that f(t) = 0 for t < 0. We recall some preliminaries and prove some lemmas in Section 2, and we give proofs of Theorems 1.1 and 1.3 in Sections 3 and 4, respectively.

#### 2. Preliminaries

Define a functional  $J_{\lambda}$  on the space  $\mathcal{D}^{1,2}(\mathbb{R}^N)$  by

$$J_{\lambda}(u) = \frac{1}{2}a\|u\|^2 + \frac{1}{4}\lambda\|u\|^4 - \int_{\mathbb{R}^N} KF(u), \quad u \in \mathcal{D}^{1,2}(\mathbb{R}^N).$$

It follows from  $(K_0)$  or  $(K_1)$ ,  $(H_1)$  and  $(H_2)$  that there exists C > 0 such that

(2.1)

$$|K(x)f(t)t| \leq C|t|^{2^*}, \quad |K(x)F(t)| \leq C|t|^{2^*}, \quad (x,t) \in \mathbb{R}^N \times \mathbb{R}.$$

Then we have that  $J_{\lambda}$  is well defined on  $\mathcal{D}^{1,2}(\mathbb{R}^N)$ , it is of  $C^1$  class for all  $\lambda \ge 0$ , and

$$\left\langle J_{\lambda}'(u), v \right\rangle = a(u, v) + \lambda \|u\|^{2}(u, v) - \int_{\mathbb{R}^{N}} Kf(u)v, \quad u, v \in \mathcal{D}^{1,2}(\mathbb{R}^{N}).$$

It is standard to verify that weak solutions of (1.1) correspond to critical points of the functional  $J_{\lambda}$ .

Next we recall a monotonicity method due to Struwe [23] and Jeanjean [10], which will be used in our proof. The version here is from [10].

**Theorem 2.1.** Let  $(X, \|\cdot\|)$  be a Banach space and  $I \subset \mathbb{R}_+$  an interval. Consider the family of  $C^1$  functionals on X

$$J_{\mu}(u) = A(u) - \mu B(u), \quad \mu \in I,$$

with B nonnegative and either  $A(u) \to \infty$  or  $B(u) \to \infty$  as  $||u|| \to \infty$  and such that  $J_{\mu}(0) = 0$ . For any  $\mu \in I$  we set

$$\Gamma_{\mu} = \{ \gamma \in C([0,1], X) \colon \gamma(0) = 0, J_{\mu}(\gamma(1)) < 0 \}.$$
(2.2)

If for every  $\mu \in I$  the set  $\Gamma_{\mu}$  is nonempty and

$$c_{\mu} = \inf_{\gamma \in \Gamma_{\mu}} \max_{t \in [0,1]} J_{\mu}(\gamma(t)) > 0, \tag{2.3}$$

then for almost every  $\mu \in I$  there is a sequence  $\{u_n\} \subset X$  such that

(i)  $\{u_n\}$  is bounded;

(ii)  $J_{\mu}(u_n) \to c_{\mu} \text{ as } n \to \infty$ ; (iii)  $J'_{\mu}(u_n) \to 0 \text{ as } n \to \infty$ , in the dual space  $X^{-1}$  of X.

To prove the boundedness of sequence of critical points in the proof later, we next introduce a Pohozaev type identity with variable-coefficient as follows.

**Lemma 2.2.** Assume that K satisfies  $(K_0)$  or  $(K_1)$ , and f satisfies  $(H_1)$  and  $(H_2)$ . If  $u \in \mathcal{D}^{1,2}(\mathbb{R}^N)$  is a weak solution of

$$-(a+\lambda ||u||^2)\Delta u = \mu K(x)f(u), \quad x \in \mathbb{R}^N,$$
(2.4)

then the following Pohozaev type identity holds

$$\frac{N-2}{2}\left(a+\lambda\|u\|^2\right)\int_{\mathbb{R}^N}|\nabla u|^2 = \mu N \int_{\mathbb{R}^N} KF(u) + \mu \int_{\mathbb{R}^N} F(u)(x \cdot \nabla K).$$
(2.5)

**Proof.** Since  $u \in \mathcal{D}^{1,2}(\mathbb{R}^N)$  is a weak solution of (2.4), by (K<sub>0</sub>) or (K<sub>1</sub>), (H<sub>1</sub>), (H<sub>2</sub>), and the standard regularity results, then  $u \in L^p_{loc}(\mathbb{R}^N)$  for all  $p \in [1, \infty)$ . Hence by the  $L^p$  estimate of elliptic equations, we know that  $u \in W^{2,p}_{loc}(\mathbb{R}^N)$  for all  $p \in [1, \infty)$ . Thus  $u \in C^{1,\beta}_{loc}(\mathbb{R}^N)$  for some  $\beta \in (0, 1)$ . It follows from (2.4) that

 $-(a+\lambda ||u||^2)\Delta u(x\cdot \nabla u) = \mu K f(u)(x\cdot \nabla u).$ 

We can calculate that

$$\begin{split} Kf(u)(x \cdot \nabla u) &= \operatorname{div} \big( xKF(u) \big) - NKF(u) - F(u)(x \cdot \nabla K), \\ \Delta u(x \cdot \nabla u) &= \operatorname{div} \big( \nabla u(x \cdot \nabla u) \big) - |\nabla u|^2 - x \cdot \nabla \left( \frac{|\nabla u|^2}{2} \right) \\ &= \operatorname{div} \left( \nabla u(x \cdot \nabla u) - x \frac{|\nabla u|^2}{2} \right) + \frac{N-2}{2} |\nabla u|^2. \end{split}$$

Therefore, for any R > 0,

364

Y. Li et al. / J. Math. Anal. Appl. 410 (2014) 361-374

$$\int_{B(0,R)} \left[ \Delta u(x \cdot \nabla u) - \frac{N-2}{2} |\nabla u|^2 \right] = \int_{\partial B(0,R)} \left[ \nabla u(x \cdot \nabla u) - x \frac{|\nabla u|^2}{2} \right] \cdot \frac{x}{R}$$
$$= \int_{\partial B(0,R)} \left[ \frac{1}{R} |x \cdot \nabla u|^2 - \frac{R}{2} |\nabla u|^2 \right],$$
$$\int_{B(0,R)} Kf(u)(x \cdot \nabla u) = \int_{\partial B(0,R)} xKF(u) \cdot \frac{x}{R} - \int_{B(0,R)} \left[ NKF(u) + F(u)x \cdot \nabla K \right]$$
$$= R \int_{\partial B(0,R)} KF(u) - \int_{B(0,R)} \left[ NKF(u) + F(u)x \cdot \nabla K \right].$$

Following [4], we show that the boundary terms above converge to 0 for at least a suitably chosen sequence  $R_n \rightarrow \infty$ . Note that

$$\int_{\mathbb{R}^{N}} \left[ KF(u) + |\nabla u|^{2} \right] \mathrm{d}x = \int_{0}^{\infty} \left\{ \int_{\partial B(0,R)} \left[ KF(u) + |\nabla u|^{2} \right] \mathrm{d}S \right\} \mathrm{d}R < \infty,$$

hence there exists a sequence  $\{R_n\}$  with  $R_n \to \infty$  such that

$$R_n \int_{\partial B(0,R_n)} \left[ KF(u) + |\nabla u|^2 \right] \mathrm{d}S \to 0.$$

Therefore, we can get the conclusion with the choice  $R = R_n$  and  $n \to \infty$ .  $\Box$ 

We conclude this section by proving a nonexistence result for larger  $\lambda$  for either  $N \ge 5$  or  $N \ge 6$ . This shows that the existence results in Theorems 1.1 and 1.3 for  $\lambda \in [0, \lambda_0)$  are somehow optimal. For the nonexistence result, we first observe the following lower bound for a positive solution:

**Lemma 2.3.** If *u* is a nontrivial weak solution of (1.1), then  $||u|| \ge r$  for some r > 0.

**Proof.** Since u is a weak solution of (1.1), then from (2.1),

$$a||u||^{2} \leq (a + \lambda ||u||^{2})||u||^{2} = \int_{\mathbb{R}^{N}} Kf(u)u \leq C ||u||^{2^{*}}$$

The proof is completed.  $\Box$ 

The nonexistence result is as follows:

#### Theorem 2.4.

- 1. If K satisfies (K<sub>0</sub>) and  $N \ge 5$ , then there exists a  $\lambda_1 > 0$  such that (1.1) has no positive solutions with nonnegative energy for  $\lambda \in (\lambda_1, \infty)$ .
- 2. If K satisfies  $(K_1)$  and  $(K_2)$  and  $N \ge 6$ , then there exists a  $\lambda_2 > 0$  such that (1.1) has no positive solutions with nonnegative energy for  $\lambda \in (\lambda_2, \infty)$ .

**Proof.** We may assume that *u* is a positive solution of (1.1) and  $J_{\lambda}(u) = c \ge 0$ . Then

$$\frac{1}{2}a\|u\|^2 + \frac{1}{4}\lambda\|u\|^4 - \int\limits_{\mathbb{R}^N} KF(u) = c.$$

If  $K \equiv 1$  and  $N \ge 5$ , then according to Lemma 2.2,

$$\frac{N-2}{2}(a+\lambda\|u\|^2)\int\limits_{\mathbb{R}^N}|\nabla u|^2=N\int\limits_{\mathbb{R}^N}F(u).$$

Hence

$$a||u||^{2} = \frac{1}{4}\lambda(N-4)||u||^{4} + cN \ge \frac{1}{4}\lambda||u||^{4}.$$

Therefore, we have  $\lambda r^2 \leq 4a$ , where *r* is defined in Lemma 2.3. If  $N \geq 6$  and *K* satisfies (K<sub>1</sub>) and (K<sub>2</sub>), then according to Lemma 2.2,

$$\frac{N-2}{2}(a+\lambda\|u\|^2)\int_{\mathbb{R}^N}|\nabla u|^2=N\int_{\mathbb{R}^N}KF(u)+\int_{\mathbb{R}^N}F(u)x\cdot\nabla K\leqslant (N+\alpha)\int_{\mathbb{R}^N}KF(u).$$

So

$$\frac{2+\alpha}{2}a\|u\|^{2} \ge \frac{1}{4}\lambda(N-4-\alpha)\|u\|^{4} + c(N+\alpha) \ge \frac{1}{4}\lambda(2-\alpha)\|u\|^{4},$$

thus we obtain  $\lambda r^2 \leq 2(2+\alpha)/(2-\alpha)$ , where *r* is defined in Lemma 2.3.

#### 3. The case that K is a constant

In this section, we consider the case that  $K(x) \equiv 1$ , and assume that conditions (H<sub>1</sub>)-(H<sub>3</sub>) are satisfied. First we recall the following estimate of the decay rate of radial functions in  $\mathcal{D}^{1,2}(\mathbb{R}^N)$  (see [4]).

**Lemma 3.1.** Suppose that  $N \ge 3$ . Then every radial function u in  $\mathcal{D}^{1,2}(\mathbb{R}^N)$  is almost everywhere equal to an even function  $U: \mathbb{R}^N \to \mathbb{R}$ , continuous for  $x \neq 0$ , such that

$$\left| U(x) \right| \leqslant C_N |x|^{(2-N)/2} \|u\|_{\mathcal{D}^{1,2}(\mathbb{R}^N)}, \quad |x| \ge 1,$$

where  $C_N$  only depends on N.

In this section, for the notation in Theorem 2.1, the space  $X = \mathcal{D}_r^{1,2}(\mathbb{R}^N)$ , and related functionals on  $\mathcal{D}_r^{1,2}(\mathbb{R}^N)$  are

$$A(u) = \frac{1}{2}a||u||^2 + \frac{1}{4}\lambda||u||^4, \qquad B(u) = \int_{\mathbb{R}^N} F(u)$$

So the perturbed functional which we will study is

$$J_{\lambda,\mu}(u) = \frac{1}{2}a\|u\|^2 + \frac{1}{4}\lambda\|u\|^4 - \mu \int_{\mathbb{R}^N} F(u).$$

and

$$\langle (J_{\lambda,\mu})'(u), v \rangle = a(u, v) + \lambda ||u||^2 (u, v) - \mu \int_{\mathbb{R}^N} f(u)v.$$
 (3.1)

To overcome the problem of lacking compactness, we need to consider the functional  $J_{\lambda,\mu}$  in the radial function space  $\mathcal{D}_r^{1,2}(\mathbb{R}^N)$ . We shall prove that  $J_{\lambda,\mu}$  satisfies the conditions of Theorem 2.1 in the next several lemmas. In the following two lemmas, we assume that  $(H_1)-(H_3)$  are satisfied. We choose a radial function  $\phi \in C_0^{\infty}(\mathbb{R}^N, \mathbb{R}_+)$  with  $\|\phi\| = 1$  and  $\operatorname{supp}(\phi) \subset B(0, R)$  for some R > 0. By  $(H_3)$ , we have that

for any  $C_1 > 0$  with  $C_1 \int_{B(0,R)} \phi^2 > a$ , there exists  $C_2 > 0$  such that

$$F(t) \ge C_1 |t|^2 - C_2, \quad t \in \mathbb{R}_+.$$
 (3.2)

Let

$$\lambda_0 = \frac{(C_1 \int_{B(0,R)} \phi^2 - a)^2}{4C_3},\tag{3.3}$$

where  $C_3 = C_2|B(0, R)|$ . Then we have the following lemma.

**Lemma 3.2.** Let  $\Gamma_{\mu}$  be the set of paths defined in (2.2). Then for  $\lambda \in [0, \lambda_0)$ ,  $\Gamma_{\mu} \neq \emptyset$  for  $\mu \in I = [1/2, 1]$ .

366

**Proof.** According to (3.2) and the definition of  $\lambda_0$ ,

$$J_{\lambda,\mu}(t\phi) = \frac{1}{2}at^2 + \frac{1}{4}\lambda t^4 - \mu \int_{\mathbb{R}^N} F(t\phi)$$
  
$$\leqslant \frac{1}{2}at^2 - \frac{1}{2}C_1t^2 \int_{B(0,R)} \phi^2 + C_3 + \frac{1}{4}\lambda t^4$$

If  $\lambda = 0$ , we can choose  $t_0 > 0$  large such that  $J_{\lambda,\mu}(t_0\phi) < 0$ . If  $\lambda \in (0, \lambda_0)$ , then by using

$$\left(C_1\int\limits_{B(0,R)}\phi^2-a\right)^2-4C_3\lambda>0,$$

we can choose  $t_0 > 0$  properly so that  $J_{\lambda,\mu}(t_0\phi) < 0$ . The proof is completed.  $\Box$ 

**Lemma 3.3.** Let  $c_{\mu}$  be defined as in (2.3). Then there exists a constant c > 0 such that  $c_{\mu} \ge c$  for all  $\mu \in I = [1/2, 1]$ .

**Proof.** For any  $\mu \in I$  and  $u \in \mathcal{D}_r^{1,2}(\mathbb{R}^N)$ , by using (H<sub>1</sub>) and (H<sub>2</sub>), we have

$$J_{\lambda,\mu}(u) \ge \frac{1}{2} a ||u||^2 + \frac{1}{4} \lambda ||u||^4 - C \int_{\mathbb{R}^N} |u|^{2^*}$$
$$\ge \frac{1}{2} a ||u||^2 - C \int_{\mathbb{R}^N} |u|^{2^*}.$$

From Sobolev's embedding theorem, we conclude that there exists  $\rho > 0$  such that  $J_{\lambda,\mu}(u) > 0$  for any  $\mu \in I$  and  $u \in \mathcal{D}_r^{1,2}(\mathbb{R}^N)$  with  $||u|| \in (0, \rho]$ . In particular, for  $||u|| = \rho$ , we have  $J_{\lambda,\mu}(u) \ge c > 0$ .

Fix  $\mu \in I$  and for any  $\gamma \in \Gamma_{\mu}$ , by the definition of  $\Gamma_{\mu}$ , we have  $\|\gamma(1)\| > \rho$ . Since  $\gamma(0) = 0$ , then from intermediate value theorem we deduce that there exists  $t_{\gamma} \in (0, 1)$  such that  $\|\gamma(t_{\gamma})\| = \rho$ . Therefore, for any  $\mu \in I$ ,

$$c_{\mu} \ge \inf_{\gamma \in \Gamma_{\mu}} J_{\lambda,\mu}(\gamma(t_{\gamma})) \ge c.$$
  $\Box$ 

Next we prove that the functional  $J_{\lambda,\mu}$  can achieve the critical value at  $c_{\mu}$  for any  $\mu \in I$ .

**Lemma 3.4.** For any  $\mu \in I$ , each bounded (PS) sequence of the functional  $J_{\lambda,\mu}$  in  $\mathcal{D}_r^{1,2}(\mathbb{R}^N)$  admits a convergent subsequence.

**Proof.** For any given  $\mu \in I$ , let  $\{u_n\}$  be a bounded (PS) sequence of  $J_{\lambda,\mu}$ , that is,  $\{u_n\}$  and  $\{J_{\lambda,\mu}(u_n)\}$  are bounded,  $(J_{\lambda,\mu})'(u_n) \to 0$  in  $\mathcal{D}'$ , where  $\mathcal{D}'$  is the dual space of  $\mathcal{D}_r^{1,2}(\mathbb{R}^N)$ . Since  $\{u_n\}$  is bounded, there exists a subsequence of  $\{u_n\}$  (still denoted by  $\{u_n\}$ ),  $u \in \mathcal{D}_r^{1,2}(\mathbb{R}^N)$  such that as  $n \to \infty$ ,

$$u_n \rightarrow u, \quad \text{in } \mathcal{D}_r^{1,2}(\mathbb{R}^N),$$
  

$$u_n \rightarrow u, \quad \text{in } L^p_{\text{loc}}(\mathbb{R}^N), \ p \in (1, 2^*),$$
  

$$u_n(x) \rightarrow u(x), \quad \text{a.e. } x \in \mathbb{R}^N.$$

According to Lemma 3.1, we may assume that

**\*** \*

$$|u_n(x)| \leqslant C|x|^{(2-N)/2}, \quad |x| \ge 1, \ n \ge 1.$$

$$(3.4)$$

From the conditions (H<sub>1</sub>) and (H<sub>2</sub>), for any  $\varepsilon > 0$ , there exist  $\delta > 0$  and  $C_{\varepsilon} > 0$  such that

$$\left|f(t)\right| \leqslant \varepsilon |t|^{2^{*}-1}, \quad |t| \leqslant \delta, \tag{3.5}$$

and

$$|f(t)| \le \varepsilon |t|^{2^*-1} + C_{\varepsilon} |t|^{2^*/2}, \quad t \in \mathbb{R}.$$
(3.6)

By (3.4), there exists an R > 0 such that  $|u_n(x)| \leq \delta$  for all  $|x| \geq R$  and all *n*. Therefore, from (3.5),

$$\left|f\left(u_{n}(x)\right)\right| \leqslant \varepsilon \left|u_{n}(x)\right|^{2^{*}-1}, \quad |x| \geqslant R, \ n \ge 1.$$

$$(3.7)$$

So it follows from (3.6) and (3.7) that

$$\left| \int_{\mathbb{R}^{N}} f(u_{n})(u_{n}-u) \right| \leq \varepsilon \int_{|x|$$

Hence we obtain that

$$\int_{\mathbb{R}^N} f(u_n)(u_n-u) \to 0, \quad n \to \infty.$$

Thus,

$$\langle (J_{\lambda,\mu})'(u_n), u_n - u \rangle = a(u_n, u_n - u) + \lambda ||u_n||^2 (u_n, u_n - u) - \mu \int_{\mathbb{R}^N} f(u_n)(u_n - u)$$
  
=  $[a + \lambda ||u_n||^2](u_n, u_n - u) + o(1),$ 

and then

 $\left[a+\lambda\|u_n\|^2\right](u_n,u_n-u)\to 0.$ 

It follows that  $||u_n|| \to ||u||$ . This together with  $u_n \rightharpoonup u$  shows that  $u_n \to u$  in  $\mathcal{D}_r^{1,2}(\mathbb{R}^N)$ . The proof is completed.  $\Box$ 

Now we are in the position to show that the modified functional  $J_{\lambda,\mu}$  has a nontrivial critical point.

**Lemma 3.5.** Let  $\lambda \in [0, \lambda_0)$ . For almost every  $\mu \in I$ , there exists  $u^{\mu} \in \mathcal{D}_r^{1,2}(\mathbb{R}^N) \setminus \{0\}$  such that  $(J_{\lambda,\mu})'(u^{\mu}) = 0$  and  $J_{\lambda,\mu}(u^{\mu}) = c_{\mu}$ .

**Proof.** From Theorem 2.1, for almost every  $\mu \in I$ , there exists a bounded sequence  $\{u_n^{\mu}\} \subset \mathcal{D}_r^{1,2}(\mathbb{R}^N)$  such that  $J_{\lambda,\mu}(u_n^{\mu}) \to c_{\mu}$  and  $(J_{\lambda,\mu})'(u_n^{\mu}) \to 0$  as  $n \to \infty$ . According to Lemma 3.4, we may assume that there exists  $u^{\mu} \in \mathcal{D}_r^{1,2}(\mathbb{R}^N)$  such that  $u_n^{\mu} \to u^{\mu}$  in  $\mathcal{D}_r^{1,2}(\mathbb{R}^N)$ . Then it follows that  $(J_{\lambda,\mu})'(u^{\mu}) = 0$ ,  $J_{\lambda,\mu}(u^{\mu}) = c_{\mu}$  and  $u^{\mu} \neq 0$  from Lemma 3.3.  $\Box$ 

According to Lemma 3.5, there exist a sequence  $\{\mu_n\} \subset I$  with  $\mu_n \to 1^-$  and an associated sequence  $\{u_n\} \subset D_r^{1,2}(\mathbb{R}^N)$  such that

$$J_{\lambda,\mu_n}(u_n) = c_{\mu_n}, \qquad (J_{\lambda,\mu_n})'(u_n) = 0.$$
(3.8)

The following lemma shows that  $\{u_n\}$  is bounded, which is a key for this paper.

**Lemma 3.6.** Let  $u_n$  be a critical point of  $J_{\lambda,\mu_n}$  at the level  $c_{\mu_n}$  as defined in (3.8).

- 1. If  $\lambda = 0$ , then there exists a constant C > 0 such that  $||u_n|| \leq C$  for all n.
- 2. There exists C > 0 such that for every  $\lambda \in (0, \lambda_0)$ , we have  $||u_n|| \leq C/\sqrt{\lambda}$  for all n.

**Proof.** Firstly, since  $(J_{\lambda,\mu_n})'(u_n) = 0$ , from Lemma 2.2,  $u_n$  satisfies the following Pohozaev type identity

$$\left(a+\lambda\|u_n\|^2\right)\frac{N-2}{2}\int\limits_{\mathbb{R}^N}|\nabla u_n|^2 = \mu_n N \int\limits_{\mathbb{R}^N} F(u_n).$$
(3.9)

Since also  $J_{\lambda,\mu_n}(u_n) = c_{\mu_n}$ , we have that

$$\frac{1}{2}aN\|u_n\|^2 + \frac{1}{4}\lambda N\|u_n\|^4 - \mu_n N \int_{\mathbb{R}^N} F(u_n) = c_{\mu_n} N.$$
(3.10)

Therefore, by (3.9) and (3.10), we obtain that

$$a \int_{\mathbb{R}^N} |\nabla u_n|^2 = c_{\mu_n} N + \frac{1}{4} \lambda (N-4) \|u_n\|^4.$$
(3.11)

We now estimate the right hand side of (3.11). By the min-max definition of the mountain pass level  $c_{\mu_n}$ , Lemma 3.2 and (3.2), we have that

$$c_{\mu_{n}} \leq \max_{t \in [0,t_{0}]} J_{\lambda,\mu_{n}}(t\phi)$$

$$\leq \max_{t \in [0,t_{0}]} \left\{ \frac{1}{2}at^{2} - \mu_{n} \int_{\mathbb{R}^{N}} F(t\phi) \right\} + \max_{t \in [0,t_{0}]} \frac{1}{4}\lambda t^{4}$$

$$\leq \max_{t \in [0,t_{0}]} \left\{ \frac{1}{2}at^{2} - \frac{1}{2}C_{1}t^{2} \int_{B(0,R)} \phi^{2} + \frac{1}{2}C_{3} \right\} + \max_{t \in [0,t_{0}]} \frac{1}{4}\lambda t^{4}$$

$$= \frac{1}{2}C_{3} + \frac{1}{4}\lambda t_{0}^{4}.$$

If  $\lambda = 0$  then it follows from (3.11) that  $2a||u_n||^2 \leq C_3 N$ . If  $\lambda \in (0, \lambda_0)$  and  $N \leq 4$ , then  $4a||u_n||^2 \leq 2C_3 N + \lambda_0 t_0^4$ . If  $N \geq 5$ , then by (3.11) we have

$$\|u_n\|^2 \leq 4a/\lambda.$$

Then the conclusion holds.  $\Box$ 

Now we are ready to prove Theorem 1.1.

**Proof of Theorem 1.1.** Let  $\lambda_0$  be defined as in (3.3), and let  $u_n$  be a critical point for  $J_{\lambda,\mu_n}$  at the level  $c_{\mu_n}$ . Then we may assume from Lemma 3.6 that for all n,

$$\|u_n\| \leqslant C_{\lambda},$$

where  $C_{\lambda}$  is defined by

$$C_{\lambda} = \begin{cases} C, & \lambda = 0, \\ C/\sqrt{\lambda}, & \lambda \in (0, \lambda_0). \end{cases}$$

Since  $\mu_n \to 1$ , we can show that  $\{u_n\}$  is a (PS) sequence of  $J_{\lambda}$ . Indeed, the boundedness of  $\{u_n\}$  implies that  $\{J_{\lambda}(u_n)\}$  is bounded. Also

$$\langle J'_{\lambda}(u_n), \mathbf{v} \rangle = \langle (J_{\lambda,\mu_n})'(u_n), \mathbf{v} \rangle + (\mu_n - 1) \int_{\mathbb{R}^N} f(u_n) \mathbf{v}, \quad \mathbf{v} \in \mathcal{D}^{1,2}_r(\mathbb{R}^N).$$

Thus  $J'_{\lambda}(u_n) \to 0$ , and consequently  $\{u_n\}$  is a bounded (PS) sequence of  $J_{\lambda}$ . By Lemma 3.4,  $\{u_n\}$  has a convergent subsequence, hence without loss of generality we may assume that  $u_n \to u$ . Consequently  $J'_{\lambda}(u) = 0$ . According to Lemma 3.3, we have that  $J_{\lambda}(u) = \lim_{n\to\infty} J_{\lambda}(u_n) = \lim_{n\to\infty} J_{\lambda,\mu_n}(u_n) \ge c > 0$  and u is a positive solution by the condition (H<sub>1</sub>). The proof is completed.  $\Box$ 

#### 4. The case $K \in L^{s}(\mathbb{R}^{N})$

In this section, we assume that *K* satisfies conditions (K<sub>1</sub>) and (K<sub>2</sub>), and *f* satisfies (H<sub>1</sub>)–(H<sub>3</sub>). For the non-constant *K* case, we need to use a cut-off functional to obtain the boundedness of { $u_n$ }. So following [11,13], we choose a cut-off function  $\psi \in C^{\infty}(\mathbb{R}_+, [0, 1])$  satisfying

$$\begin{cases} \psi(t) = 1, & t \in [0, 1], \\ \psi(t) = 0, & t \in [2, \infty), \\ |\psi'|_{\infty} \leq 2, & t \in [0, \infty), \end{cases}$$

and study the following modified functional  $J_{\lambda}^{T}: \mathcal{D}^{1,2}(\mathbb{R}^{N}) \to \mathbb{R}$  defined by

$$J_{\lambda}^{T}(u) = \frac{1}{2}a||u||^{2} + \frac{1}{4}\lambda h_{T}(u)||u||^{4} - \int_{\mathbb{R}^{N}} KF(u), \quad u \in \mathcal{D}^{1,2}(\mathbb{R}^{N}),$$

where for every T > 0,

$$h_T(u) = \psi\left(\frac{\|u\|^2}{T^2}\right).$$

With this penalization, for *T* sufficiently large and  $\lambda$  sufficiently small, we are able to find a critical point *u* of  $J_{\lambda}^{T}$  such that  $||u|| \leq T$  and so *u* is also a critical point of  $J_{\lambda}$ .

In this section, we consider the problem (1.1) in the space  $\mathcal{D}^{1,2}(\mathbb{R}^N)$  because that the function *K* is not assumed to be radial. For the setting of Theorem 2.1, the space  $X = \mathcal{D}^{1,2}(\mathbb{R}^N)$ ,

$$A(u) = \frac{1}{2}a||u||^2 + \frac{1}{4}\lambda h_T(u)||u||^4, \qquad B(u) = \int_{\mathbb{R}^N} KF(u)$$

So the perturbed functional which we will study is

$$J_{\lambda,\mu}^{T}(u) = \frac{1}{2}a\|u\|^{2} + \frac{1}{4}\lambda h_{T}(u)\|u\|^{4} - \mu \int_{\mathbb{R}^{N}} KF(u),$$

and

$$\left\langle \left(J_{\lambda,\mu}^{T}\right)'(u),\nu\right\rangle = a(u,\nu) + \lambda h_{T}(u) \|u\|^{2}(u,\nu) + \frac{\lambda}{2T^{2}}\psi'\left(\frac{\|u\|^{2}}{T^{2}}\right) \|u\|^{4}(u,\nu) - \mu \int_{\mathbb{R}^{N}} Kf(u)\nu.$$
(4.1)

Again we establish parallel steps as Lemmas 3.2 and 3.3 as follows.

**Lemma 4.1.** Let  $\Gamma_{\lambda,\mu}^{T}$  be defined by (2.2) for the functional  $J_{\lambda,\mu}^{T}$ . Then  $\Gamma_{\lambda,\mu}^{T} \neq \emptyset$  for all  $\mu \in I = [1/2, 1]$  and  $\lambda, T > 0$ .

**Proof.** By the condition (K<sub>1</sub>), we may assume that  $\int_{B(0,R)} K > 0$  for some R > 0. We now choose a function  $\phi \in C_0^{\infty}(\mathbb{R}^N, \mathbb{R}_+)$  with  $\|\phi\| = 1$ ,  $\operatorname{supp}(\phi) \subset B(0, R)$  and  $\int_{B(0,R)} K\phi^2 > 0$ . By the condition (H<sub>3</sub>), for  $C_1 = 2a(\int_{B(0,R)} K\phi^2)^{-1} > 0$ , there exists  $C_2 > 0$  such that

$$F(t) \ge C_1 |t|^2 - C_2, \quad t \in \mathbb{R}_+.$$
 (4.2)

For  $t^2 \ge 2T^2$ , we have from (4.2) that

$$J_{\lambda,\mu}^{T}(t\phi) = \frac{1}{2}at^{2} + \frac{1}{4}\lambda\psi\left(\frac{t^{2}}{T^{2}}\right)t^{4} - \mu\int_{\mathbb{R}^{N}}KF(t\phi)$$
$$\leq \frac{1}{2}at^{2} - \frac{1}{2}C_{1}t^{2}\int_{B(0,R)}K\phi^{2} + C_{3},$$

where  $C_3 = C_2 \int_{B(0,R)} K > 0$ . Thus we can choose t > 0 large such that  $J_{\lambda,\mu}^T(t\phi) < 0$ . The proof is completed.  $\Box$ 

**Lemma 4.2.** Let  $c_{\lambda,\mu}^T$  be defined by (2.3) for the functional  $J_{\lambda,\mu}^T$ . Then there exists a constant c > 0 such that  $c_{\lambda,\mu}^T \ge c$  for all  $\mu \in I$  and  $\lambda, T > 0$ .

**Proof.** Let  $\mu \in I$  and  $\lambda, T > 0$ . For any  $u \in \mathcal{D}^{1,2}(\mathbb{R}^N)$ , by (2.1), we have that

$$J_{\lambda,\mu}^{T}(u) \ge \frac{1}{2} a ||u||^{2} + \frac{1}{4} \lambda h_{T}(u) ||u||^{4} - C \int_{\mathbb{R}^{N}} |u|^{2}$$
$$\ge \frac{1}{2} a ||u||^{2} - C \int_{\mathbb{R}^{N}} |u|^{2^{*}}.$$

By Sobolev's embedding theorem, we conclude that there exists  $\rho > 0$  such that  $J_{\lambda,\mu}^{T}(u) > 0$  for  $u \in \mathcal{D}^{1,2}(\mathbb{R}^{N})$  with  $0 < ||u|| \leq \rho$ . In particular, for  $||u|| = \rho$ , it follows  $J_{\lambda,\mu}^{T}(u) \geq c > 0$ . For every  $\gamma \in \Gamma_{\lambda,\mu}^{T}$ , by the definition of  $\Gamma_{\lambda,\mu}^{T}$ , we have  $||\gamma(1)|| > \rho$ . Then from the intermediate value theorem, there exists  $t_{\gamma} \in (0, 1)$  such that  $||\gamma(t_{\gamma})|| = \rho$ . Therefore,

$$c_{\lambda,\mu}^T \ge \inf_{\gamma \in \Gamma_{\lambda,\mu}^T} J_{\lambda,\mu}^T (\gamma(t_{\gamma})) \ge c > 0.$$

The proof is completed.  $\Box$ 

The next lemma shows that the critical level defined in Lemma 4.2 can be achieved if  $\lambda$  is small.

**Lemma 4.3.** Let  $4\lambda T^2 < a$  and  $\mu \in I = [1/2, 1]$ . Then each bounded (PS) sequence of the functional  $J_{\lambda,\mu}^T$  admits a convergent subsequence.

**Proof.** Let  $4\lambda T^2 < a$  and  $\mu \in I$ . Let  $\{u_n\}$  be a bounded (PS) sequence of  $J_{\lambda,\mu}^T$ , that is,  $\{u_n\}$  and  $\{J_{\lambda,\mu}^T(u_n)\}$  are bounded, and  $(J_{\lambda,\mu}^T)'(u_n) \to 0$  in  $\mathcal{D}'$ , where  $\mathcal{D}'$  is the dual space of  $\mathcal{D}^{1,2}(\mathbb{R}^N)$ . Since  $\{u_n\}$  is bounded, we may assume that there exists  $u \in \mathcal{D}^{1,2}(\mathbb{R}^N)$  such that

$$u_n \to u, \quad \text{in } \mathcal{D}^{1,2}(\mathbb{R}^N),$$
  

$$u_n \to u, \quad \text{in } L^p_{\text{loc}}(\mathbb{R}^N), \ p \in (1, 2^*),$$
  

$$u_n(x) \to u(x), \quad \text{a.e. } x \in \mathbb{R}^N.$$

By the conditions (H<sub>1</sub>) and (H<sub>2</sub>), for any  $\varepsilon > 0$ , there exists  $C_{\varepsilon} > 0$  such that

$$\left|f(u)\right| \leq \varepsilon |u|^{2^{*}-1} + C_{\varepsilon} \chi_{(\tau_{1} \leq |u| \leq \tau_{2})}, \quad u \in D^{1,2}(\mathbb{R}^{N}),$$

$$(4.3)$$

where  $\tau_1$  and  $\tau_2$  are two positive constants. Let  $E_n = \{x \in \mathbb{R}^N : \tau_1 \leq |u_n(x)| \leq \tau_2\}$ . Then

$$au_1^{2^*}|E_n| \leq \int_{E_n} |u_n|^{2^*} \leq \int_{\mathbb{R}^N} |u_n|^{2^*} \leq C,$$

where  $|E_n|$  is Lebesgue's measure of  $E_n$ . This implies that  $|E_n| \leq C\tau_1^{-2^*}$ . So it follows from (4.3) and (K<sub>1</sub>) that for r > 0,

$$\begin{split} \left| \int_{|x| \ge r} Kf(u_n)(u_n - u) \right| &\leq \varepsilon |K|_{\infty} \int_{|x| \ge r} |u_n|^{2^* - 1} |u_n - u| + C_{\varepsilon} \int_{E_n \setminus B(0, r)} K|u_n - u| \\ &\leq \varepsilon |K|_{\infty} |u_n|^{2^* - 1} |u_n - u|_{2^*} + C_{\varepsilon} |E_n|^{1/s'} \left( \int_{E_n \setminus B(0, r)} K^s \right)^{1/s} |u_n - u|_{2^*} \end{split}$$

where  $s' \in (0, \infty]$  with  $1/s' + 1/s + 1/2^* = 1$ . Hence,

$$\limsup_{r\to\infty}\left|\int_{|x|\geqslant r}Kf(u_n)(u_n-u)\right|\leqslant\varepsilon C|K|_{\infty}.$$

On the other hand, we have that

$$\int_{|x| < r} Kf(u_n)(u_n - u) \to 0, \quad \text{as } n \to +\infty$$

for every r > 0. This implies that

$$\int_{\mathbb{R}^N} Kf(u_n)(u_n - u) \to 0, \quad \text{as } n \to +\infty.$$
(4.4)

Thus, by (4.4), we get that

$$\begin{split} \left\langle \left(J_{\lambda,\mu}^{T}\right)'(u_{n}), u_{n} - u \right\rangle &= a(u_{n}, u_{n} - u) + \lambda h_{T}(u_{n}) \|u_{n}\|^{2}(u_{n}, u_{n} - u) \\ &+ \frac{\lambda}{2T^{2}} \psi' \left(\frac{\|u_{n}\|^{2}}{T^{2}}\right) \|u_{n}\|^{4}(u_{n}, u_{n} - u) - \mu \int_{\mathbb{R}^{N}} Kf(u_{n})(u_{n} - u) \\ &= \left(a + \lambda h_{T}(u_{n}) \|u_{n}\|^{2} + \frac{\lambda}{2T^{2}} \psi' \left(\frac{\|u_{n}\|^{2}}{T^{2}}\right) \|u_{n}\|^{4}\right) (u_{n}, u_{n} - u) + o(1), \end{split}$$

and then as  $n \to \infty$ ,

$$\left(a + \lambda h_T(u_n) \|u_n\|^2 + \frac{\lambda}{2T^2} \psi'\left(\frac{\|u_n\|^2}{T^2}\right) \|u_n\|^4\right) (u_n, u_n - u) \to 0.$$

If  $||u_n||^2 > 2T^2$ , then  $|\psi'(\frac{||u_n||^2}{T^2})||u_n||^4| = 0$ . If  $||u_n||^2 \leq 2T^2$ , then  $|\psi'(\frac{||u_n||^2}{T^2})| \leq 2$  and  $|\psi'(\frac{||u_n||^2}{T^2})||u_n||^4| \leq 8T^4$ . According to  $4\lambda T^2 < a$ , then it follows that  $||u_n|| \to ||u||$ . This together with  $u_n \rightharpoonup u$  shows that  $u_n \to u$  in  $\mathcal{D}^{1,2}(\mathbb{R}^N)$ . The proof is completed.  $\Box$ 

Now we are in the position to show that the modified functional  $J_{\lambda,\mu}^T$  has a nontrivial critical point.

**Lemma 4.4.** Let  $4\lambda T^2 < a$ . Then for almost every  $\mu \in I$ , there exists  $u^{\mu} \in \mathcal{D}^{1,2}(\mathbb{R}^N)$  such that  $(J^T_{\lambda,\mu})'(u^{\mu}) = 0$  and  $J^T_{\lambda,\mu}(u^{\mu}) = c^T_{\lambda,\mu}$ .

**Proof.** By Theorem 2.1, for almost every  $\mu \in I$ , there exists a bounded sequence  $\{u_n^{\mu}\} \subset \mathcal{D}^{1,2}(\mathbb{R}^N)$  such that  $J_{\lambda,\mu}^T(u_n^{\mu}) \to c_{\lambda,\mu}^T$ and  $(J_{\lambda,\mu}^T)'(u_n^{\mu}) \to 0$  as  $n \to \infty$ . According to Lemma 4.3, we can suppose that there exists  $u^{\mu} \in \mathcal{D}^{1,2}(\mathbb{R}^N)$  such that  $u_n^{\mu} \to u^{\mu}$  in  $\mathcal{D}^{1,2}(\mathbb{R}^N)$ , then the assertion follows from Lemma 4.2.  $\Box$ 

According to Lemma 4.4, there exist a sequence  $\{\mu_n\} \subset I$  with  $\mu_n \to 1^-$  and a sequence  $\{u_n\} \subset D^{1,2}(\mathbb{R}^N)$  such that

$$J_{\lambda,\mu_n}^T(u_n) = c_{\mu_n}, \qquad (J_{\lambda,\mu_n}^T)'(u_n) = 0.$$
(4.5)

The following lemma shows that  $||u_n|| \leq T$  for all *n*, which is a key step for the proof.

**Lemma 4.5.** Let  $u_n$  be a critical point of  $J_{\lambda,\mu_n}^T$  at the level  $c_{\mu_n}$  as defined in (4.5). Then there exist positive constants

$$\lambda_0 = \frac{(2-\alpha)^2 a^2}{16(N-\alpha)(6N-10)C_3},\tag{4.6}$$

and

$$T_0^2 = \frac{4(N-\alpha)C_3}{(2-\alpha)a},$$
(4.7)

which satisfy

$$4(6N-10)(2-\alpha)^{-1}\lambda_0 T_0^2 = a,$$
(4.8)

such that for any  $\lambda \in [0, \lambda_0]$  and any  $T \ge T_0$ ,  $||u_n|| \le T$ .

**Proof.** Since  $(J_{\lambda,u_n}^T)'(u_n) = 0$ , then from Lemma 2.2,  $u_n$  satisfies the following Pohozaev type identity

$$\left(a + \lambda h_T(u_n) \|u_n\|^2 + \frac{\lambda}{2T^2} \psi'\left(\frac{\|u_n\|^2}{T^2}\right) \|u_n\|^4 \right) \frac{N-2}{2} \int_{\mathbb{R}^N} |\nabla u_n|^2 = \mu_n N \int_{\mathbb{R}^N} KF(u_n) + \mu_n \int_{\mathbb{R}^N} F(u_n)(x \cdot \nabla K)$$

$$\ge \mu_n (N-\alpha) \int_{\mathbb{R}^N} KF(u_n).$$

On the other hand, by using  $J_{\lambda,\mu_n}^T(u_n) = c_{\mu_n}$ , we have that

$$\frac{1}{2}a(N-\alpha)\|u_n\|^2 + \frac{1}{4}\lambda(N-\alpha)h_T(u_n)\|u_n\|^4 - \mu_n(N-\alpha)\int_{\mathbb{R}^N} KF(u_n) = c_{\mu_n}(N-\alpha).$$
(4.9)

Thus, we can obtain that

$$\frac{2-\alpha}{2}a\int_{\mathbb{R}^{N}}|\nabla u_{n}|^{2} \leq c_{\mu_{n}}(N-\alpha) + \frac{1}{4}\lambda(N-4+\alpha)h_{T}(u_{n})\|u_{n}\|^{4} + \frac{\lambda(N-2)}{4T^{2}}\psi'\left(\frac{\|u_{n}\|^{2}}{T^{2}}\right)\|u_{n}\|^{6}.$$
(4.10)

We will estimate the right hand side of (4.10). By the min-max definition of the mountain pass level  $c_{\mu_n}$ , Lemma 4.1 and (4.2), we have that

$$\begin{aligned} c_{\mu_n} &\leq \max_{t \in [0,\infty)} J^{I}_{\lambda,\mu_n}(t\phi) \\ &\leq \max_{t \in [0,\infty)} \left\{ \frac{1}{2}at^2 - \mu_n \int_{\mathbb{R}^N} KF(t\phi) \right\} + \max_{t \in [0,\infty)} \frac{1}{4}\lambda \psi\left(\frac{t^2}{T^2}\right) t^4 \\ &\leq \max_{t \in [0,\infty)} \left\{ \frac{1}{2}at^2 - \frac{1}{2}C_1t^2 \int_{B(0,R)} K\phi^2 + C_3 \right\} + \max_{t \in [0,\infty)} \frac{1}{4}\lambda \psi\left(\frac{t^2}{T^2}\right) t^4 \\ &\leq C_3 + \lambda T^4. \end{aligned}$$

We have also that

$$\frac{|N-4+\alpha|}{4}h_T(u_n)||u_n||^4 \leq (N-2+\alpha)T^4,$$
$$\frac{N-2}{4T^2} \left|\psi'\left(\frac{||u_n||^2}{T^2}\right)\right|||u_n||^6 \leq 4(N-2)T^4.$$

Then it follows that

$$\frac{2-\alpha}{2}a\int\limits_{\mathbb{R}^N}|\nabla u_n|^2 \leq (N-\alpha)C_3 + \lambda(6N-10)T^4$$

or equivalently

$$\int_{\mathbb{R}^N} |\nabla u_n|^2 \leqslant \frac{2(N-\alpha)}{(2-\alpha)a} C_3 + \lambda \frac{2(6N-10)}{(2-\alpha)a} T^4.$$
(4.11)

Now for  $\lambda_0$  and  $T_0$  defined in (4.6) and (4.7), since the equality (4.8) holds, we obtain that

$$2(N-\alpha)(2-\alpha)^{-1}a^{-1}C_3 + 2\lambda(2-\alpha)^{-1}(6N-10)a^{-1}T_0^4 \leq 2(N-\alpha)(2-\alpha)^{-1}a^{-1}C_3 + T_0^2/2 = T_0^2 \leq T^2.$$

Thus it follows from (4.11) that  $||u_n|| \leq T$  for all *n*, and the stated conclusion holds.  $\Box$ 

Now we can complete the proof of Theorem 1.3.

**Proof of Theorem 1.3.** Let  $\lambda_0$  and  $T_0$  be defined as in Lemma 4.5, and let  $u_n$  be a critical point for  $J_{\lambda,\mu_n}^T$  at the level  $c_{\mu_n}$ . Then we may assume that for all n,  $||u_n|| \leq T$  holds. Hence

$$J_{\lambda,\mu_n}^T(u_n) = \frac{1}{2}a\|u_n\|^2 + \frac{1}{4}\lambda\|u_n\|^4 - \mu_n \int_{\mathbb{R}^N} KF(u_n).$$

Since  $\mu_n \to 1$ , we will show that  $\{u_n\}$  is a (PS) sequence of  $J_{\lambda}$ . Indeed, the boundedness of  $\{u_n\}$  implies that  $\{J_{\lambda}(u_n)\}$  is bounded. Also

$$\left\langle J_{\lambda}'(u_n), \nu \right\rangle = \left\langle \left( J_{\lambda,\mu_n}^T \right)'(u_n), \nu \right\rangle + (\mu_n - 1) \int_{\mathbb{R}^N} Kf(u_n)\nu, \quad \nu \in \mathcal{D}^{1,2}(\mathbb{R}^N)$$

Thus  $J'_{\lambda}(u_n) \to 0$ , and  $\{u_n\}$  is a bounded (PS) sequence of  $J_{\lambda}$ . By Lemma 4.3,  $\{u_n\}$  has a convergent subsequence, and without loss of generality, we may assume that  $u_n \to u$ . Consequently  $J'_{\lambda}(u) = 0$ . According to Lemma 4.2, we have that  $J_{\lambda}(u) = \lim_{n\to\infty} J_{\lambda}(u_n) = \lim_{n\to\infty} J^T_{\lambda,\mu_n}(u_n) \ge c > 0$  and u is a positive solution by the condition (H<sub>1</sub>). The proof is completed.  $\Box$ 

#### Acknowledgment

The authors thank anonymous referees for careful readings and some helpful comments, which greatly improve the manuscript.

#### References

- Claudianor O. Alves, Marco A.S. Souto, Marcelo Montenegro, Existence of solution for two classes of elliptic problems in R<sup>N</sup> with zero mass, J. Differential Equations 252 (10) (2012) 5735–5750.
- [2] Antonio Azzollini, Alessio Pomponio, On a "zero mass" nonlinear Schrödinger equation, Adv. Nonlinear Stud. 7 (4) (2007) 599-627.
- [3] Antonio Azzollini, Alessio Pomponio, Compactness results and applications to some "zero mass" elliptic problems, Nonlinear Anal. 69 (10) (2008) 3559–3576.
- [4] Henry Berestycki, P.-L. Lions, Nonlinear scalar field equations. I. Existence of a ground state, Arch. Ration. Mech. Anal. 82 (4) (1983) 313–345.
- [5] Ching-Yu Chen, Yueh-Cheng Kuo, Tsung-Fang Wu, The Nehari manifold for a Kirchhoff type problem involving sign-changing weight functions, J. Differential Equations 250 (4) (2011) 1876–1908.
- [6] Xiaoming He, Wenming Zou, Infinitely many positive solutions for Kirchhoff-type problems, Nonlinear Anal. 70 (3) (2009) 1407–1414.
- [7] Xiaoming He, Wenming Zou, Multiplicity of solutions for a class of Kirchhoff type problems, Acta Math. Appl. Sin. Engl. Ser. 26 (3) (2010) 387–394.
- [8] Xiaoming He, Wenming Zou, Existence and concentration behavior of positive solutions for a Kirchhoff equation in R<sup>3</sup>, J. Differential Equations 252 (2) (2012) 1813–1834.
- [9] Xiaoming He, Wenming Zou, Ground states for nonlinear Kirchhoff equations with critical growth, Ann. Mat. (2012) 1–28.
- [10] Louis Jeanjean, On the existence of bounded Palais-Smale sequences and application to a Landesman-Lazer-type problem set on R<sup>N</sup>, Proc. Roy. Soc. Edinburgh Sect. A 129 (4) (1999) 787–809.
- [11] Louis Jeanjean, Stefan Le Coz, An existence and stability result for standing waves of nonlinear Schrödinger equations, Adv. Differential Equations 11 (7) (2006) 813–840.

- [12] Jiahua Jin, Xian Wu, Infinitely many radial solutions for Kirchhoff-type problems in  $\mathbb{R}^N$ , J. Math. Anal. Appl. 369 (2) (2010) 564–574.
- [13] Hiroaki Kikuchi, Existence and stability of standing waves for Schrödinger-Poisson-Slater equation, Adv. Nonlinear Stud. 7 (3) (2007) 403-437.
- [14] Yi Li, Asymptotic behavior of positive solutions of equation  $\Delta u + K(x)u^p = 0$  in  $\mathbb{R}^n$ , J. Differential Equations 95 (2) (1992) 304–330.
- [15] Yi Li, On the positive solutions of the Matukuma equation, Duke Math. J. 70 (3) (1993) 575-589.
- [16] Yi Li, Wei-Ming Ni, On conformal scalar curvature equations in  $\mathbb{R}^n$ , Duke Math. J. 57 (3) (1988) 895–924.
- [17] Yuhua Li, Fuyi Li, Junping Shi, Existence of a positive solution to Kirchhoff type problems without compactness conditions, J. Differential Equations 253 (7) (2012) 2285–2294.
- [18] Zhanping Liang, Fuyi Li, Junping Shi, Positive solutions to Kirchhoff type equations with nonlinearity having prescribed asymptotic behavior, Ann. Inst. H. Poincaré Anal. Non Linéaire (2013), http://dx.doi.org/10.1016/j.anihpc.2013.01.006.
- [19] Anmin Mao, Zhitao Zhang, Sign-changing and multiple solutions of Kirchhoff type problems without the P.S. condition, Nonlinear Anal. 70 (3) (2009) 1275–1287.
- [20] Kanishka Perera, Zhitao Zhang, Nontrivial solutions of Kirchhoff-type problems via the Yang index, J. Differential Equations 221 (1) (2006) 246–255.
- [21] Patrizia Pucci, Raffaella Servadei, Existence, non-existence and regularity of radial ground states for *p*-Laplacian equations with singular weights, Ann. Inst. H. Poincaré Anal. Non Linéaire 25 (3) (2008) 505–537.
- [22] Patrizia Pucci, Raffaella Servadei, Nonexistence for p-Laplace equations with singular weights, Commun. Pure Appl. Anal. 9 (5) (2010) 1421–1438.
- [23] Michael Struwe, Variational Methods: Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems, fourth edition, Ergeb. Math. Grenzgeb. (3), vol. 34, Springer-Verlag, Berlin, 2008.
- [24] Ji-Jiang Sun, Chun-Lei Tang, Existence and multiplicity of solutions for Kirchhoff type equations, Nonlinear Anal. 74 (4) (2011) 1212–1222.
- [25] Jun Wang, Lixin Tian, Junxiang Xu, Fubao Zhang, Multiplicity and concentration of positive solutions for a Kirchhoff type problem with critical growth, J. Differential Equations 253 (7) (2012) 2314–2351.
- [26] Xian Wu, Existence of nontrivial solutions and high energy solutions for Schrödinger-Kirchhoff-type equations in R<sup>N</sup>, Nonlinear Anal. Real World Appl. 12 (2) (2011) 1278-1287.
- [27] Yang Yang, Jihui Zhang, Nontrivial solutions of a class of nonlocal problems via local linking theory, Appl. Math. Lett. 23 (4) (2010) 377-380.
- [28] Zhitao Zhang, Kanishka Perera, Sign changing solutions of Kirchhoff type problems via invariant sets of descent flow, J. Math. Anal. Appl. 317 (2) (2006) 456–463.