



W&M ScholarWorks

Arts & Sciences Articles

Arts and Sciences

12-12-2016

How gauge covariance of the fermion and boson propagators in QED constrain the effective fermion-boson vertex

Shaoyang Jia

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Jia, Shaoyang, How gauge covariance of the fermion and boson propagators in QED constrain the effective fermion-boson vertex (2016).

10.1103/PhysRevD.94.116004

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

How gauge covariance of the fermion and boson propagators in QED constrain the effective fermion-boson vertex

Shaoyang Jia and M. R. Pennington

Phys. Rev. D **94**, 116004 — Published 12 December 2016

DOI: [10.1103/PhysRevD.94.116004](https://doi.org/10.1103/PhysRevD.94.116004)

How gauge covariance of the fermion and boson propagators in QED constrain the effective fermion-boson vertex

Shaoyang Jia*

Physics Department, College of William & Mary, Williamsburg, VA 23187, USA

M.R. Pennington†

*Physics Department, College of William & Mary, Williamsburg, VA 23187, USA and
Theory Center, Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA*

We derive the gauge covariance requirement imposed on the QED fermion-photon three-point function within the framework of a spectral representation for fermion propagators. When satisfied, such requirement ensures solutions to the fermion propagator Schwinger-Dyson equation (SDE) in any covariant gauge with arbitrary numbers of spacetime dimensions to be consistent with the Landau-Khalatnikov-Fradkin transformation (LKFT). The general result has been verified by the special cases of three and four dimensions. Additionally, we present the condition that ensures the vacuum polarization is independent of the gauge parameter. As an illustration, we show how the Gauge Technique dimensionally regularized in 4D does not satisfy the covariance requirement.

I. INTRODUCTION

The infinite set of Schwinger-Dyson equations constitutes the field equations of any theory. They relate Green's functions to each other. In QED and QCD, they relate, for instance, the fermion propagator to the gauge boson propagator and the fermion-boson interaction. The expansion of the Schwinger-Dyson equation for each Green's function in powers of the coupling yields the well-known perturbative series. However, most of the phenomena in hadron and nuclear physics are controlled by QCD in the regime of strong coupling [1, 2]. Then alternative truncations of the Schwinger-Dyson equations are required to capture the essence of the physics. A simple truncation, much used in QCD for the fermion propagator equation, combines the fermion-boson vertex and the gluon propagator function in a construction proportional to γ^μ . This ansatz from Maris and Tandy [3] is especially successful phenomenologically, and presumed to be appropriate in the Landau gauge. Then the fermion mass function in the strong coupling regime can be shown to have a characteristic momentum dependence illustrated in Fig. 2 of Ref. [4] (also see Fig. 1 of Ref. [5]). One realizes that whether the current quark mass (defined at some appropriate large momentum) is 5 or 100 MeV, the mass at low momenta is ~ 350 MeV heavier: an infrared behavior that matches constituent quark masses. It has been argued that such behavior of this gauge-dependent quantity is directly correlated with the momentum dependence of physical observables such as the electromagnetic formfactors of the pion and the proton [6–9]. It would then seem natural to check how the mass functions shown in Ref. [4] change with gauge.

As a simple illustration let us consider QED, in which a purely bare fermion-boson vertex is assumed. This pro-




FIG. 1. The dependence of the fermion propagator mass function $M(p^2)$ on ξ . The black solid line is the parametric form of $M(p^2)$ given by Eq. (2.1) of Ref. [10] with $M_0 = 3$ MeV, $c = 1.239$ and $\Lambda_{\text{QCD}} = 401$ MeV. The red dashed line and the blue dash-dot line correspond to what the mass function should be when $\alpha\xi = 3$ and $\alpha\xi = 6$ respectively. The red dashed line and the blue dash-dot line are obtained by the LKFT for the fermion propagator in 4D within the $\overline{\text{MS}}$ renormalization scheme at the scale $\mu = \Lambda$, with $F(p^2) = 1$ and $M(p^2)$ given by the black line as the initial conditions in the Landau gauge. Notice that gauge covariance produces a node-like feature, in this case at $p^2 \simeq -0.3$ GeV². Though this example is motivated by QCD in the choice of parameters, the calculations are from QED.

duces a mass function in the Landau gauge shown as the solid line in Fig. 1. Though this is QED, the qualitative behavior of the mass function is very like that for a light quark in QCD in the Maris-Tandy model. Solving the QED Schwinger-Dyson equation for the fermion in 4D with the same bare γ^μ vertex in the Feynman gauge (for instance) changes the mass function as in Fig. 11 of Ref. [11]. The corresponding 3D results are illustrated

* sjia@email.wm.edu

† michaelp@jlab.org

by Fig. 3.2 of Ref. [12].

However, the gauge covariance of the fermion propagator is exactly specified by the Landau-Khalatnikov-Fradkin transformation (LKFT) [13]. As we will remind the reader this relates the propagator functions in one gauge to those in another. If one applies this to the 4D fermion mass functions shown in Fig. 2 of Ref. [4] for the Landau gauge, one obtains the behavior in two other covariant gauges plotted here in Fig. 1. One sees that the mass function moves between gauges to produce what appears to be a *node*. This in fact ensures that any fermion condensate is gauge independent. The corresponding 3D results are given by Fig. 3.4 of Ref. [12]. However, solving the fermion Schwinger-Dyson equation in different gauges as shown in Fig. 11 of Ref. [11] gives mass functions that simply move up or down as the gauge changes. There is no hint of the nodal behavior required by the LKFT and seen in Fig. 1. This not surprisingly indicates that a vertex as simple as γ^μ cannot be appropriate in both the Landau and Feynman gauges in QED. Indeed, it may not hold in any covariant gauge.

The purpose of this paper is to present the conditions that ensure the solutions of the Schwinger-Dyson equation (SDE) for the fermion propagator are gauge covariant [14]. We study this in QED, where particles having a physical mass-shell means it is natural that the fermion propagator satisfies a spectral representation. This allows the SDE to be investigated at all momenta, and we are not restricted to spacelike momenta (or nearby time-like momenta) as in QCD studies.

This article is organized as the following. In Section II, the spectral representation is introduced for the fermion propagator. Then an abstract version of the SDE for the fermion propagator spectral functions is obtained in terms of the distribution Ω . Section III briefly reviews results of the LKFT for the fermion propagator. In Section IV, the gauge covariance requirements for the fermion propagator and the photon propagator SDEs are derived. In Section V, with known contributions to Ω calculated, the consistency requirement for the fermion equation is written for the unknown terms of Ω . As an example, the Gauge Technique ansatz of Salam, Delbourgo and Strathdee [15–18] translates into an Ω that is shown explicitly not to satisfy the consistency requirement in the quenched calculation in 4D. Section VI is the summary.

II. SPECTRAL REPRESENTATION OF FERMION PROPAGATOR AND ITS SDE

A. Spectral representation of fermion propagator as a bijective mapping

The fermion propagator $S_F(p)$ can be decomposed into Dirac vector and Dirac scalar components defined by

$$S_F(p) = S_1(p^2)\not{p} + S_2(p^2)\mathbb{1}. \quad (1)$$




FIG. 2. The illustration of analytic functions with branch cuts along the positive real axis. The contour can be used to prove Eq. (2) using Cauchy’s integral formula. When used to prove Eq. (3), z stands for the complex p^2 .

Each component function is similar to a scalar propagator function. Therefore, the spectral representation of fermion propagator requires two scalar spectral functions;

$$S_j(p^2; \xi) = \int_{m^2}^{+\infty} ds \frac{\rho_j(s; \xi)}{p^2 - s + i\epsilon}, \quad (2)$$

where $j = 1, 2$ and the dependence on the covariant gauge parameter ξ has been made explicit. The Feynman prescription of a momentum space propagator is denoted by the $i\epsilon$ term in the denominator, while the normal writing of ϵ is reserved for how far away the number of space-time dimensions is from 4 by $d = 4 - 2\epsilon$. The spectral integral given in Eq. (2) is convergent if $S_F(p)$ vanishes when $p^2 \rightarrow \infty$. That the integrals converge without the need for subtractions is assured by the renormalizability of QED in $d < 4$ dimensions.

Apparently when the fermion propagator takes its free-particle form, the spectral functions are given by $\rho_1(s) = \delta(s - m^2)$ and $\rho_2(s) = m\delta(s - m^2)$. When interactions are present, the fermion propagator is modified by quantum loop corrections and therefore develops branch cuts starting at the particle production thresholds. Such corrections add θ -functions to the spectral functions.

The existence of a spectral representation, Eq. (2), is determined by the analytic structure of the propagator functions $S_j(p^2)$ in the complex momentum plane. We expect that apart from free-particle poles and branch cuts along the positive real axis, propagator functions are holomorphic everywhere else. In this scenario, the spectral functions uniquely determine the propagator functions in the complex momentum plane.

Meanwhile, when the analytic structure of the fermion propagator meets such requirements, the inverse of Eq. (2) is given by

$$\rho_j(s; \xi) = -\frac{1}{\pi} \text{Im}\{S_j(s + i\epsilon; \xi)\}. \quad (3)$$




FIG. 3. The diagrammatic representation of the SDE for the fermion propagator in momentum space. The fermion-photon vertex is unknown and an ansatz is required to solve this equation.

The Feynman prescription combined with the limiting form of the δ -function,

$$\lim_{\varepsilon \rightarrow 0} \varepsilon / (x^2 + \varepsilon^2) = \pi \delta(x),$$

specifies that any simple pole structure of the propagator function corresponds to a δ -function term in its spectral function. In addition since $S_j(p^2 + i\varepsilon) = S_j^*(p^2 - i\varepsilon)$, functions calculated by Eq. (3) are indeed the spectral functions occurring in Eq. (2), which can be verified using Cauchy's integral formula with the contour in Fig. 2. Therefore we have shown that the spectral representation given by Eq. (2) and its inverse Eq. (3) specify a bijective mapping between the propagators as functions in the complex momentum plane and their spectral functions.

B. SDE for fermion propagator spectral functions

The SDE for the fermion propagator in momentum space is represented by Fig. 3. It has been solved extensively using specific ansätze for the fermion photon vertex [19, 20] (also see Ref. [11, 21–31]). Solutions in the Minkowski space have been obtained by [32] (also see Ref. [33–37]). Alternatively, these equations can be solved using complex conjugate poles to represent the propagator functions [5, 38–40].

Each diagram in Fig. 3 is not obviously linear in the spectral functions $\rho_j(s; \xi)$. However, an easier way to solve for the spectral functions $\rho_j(s; \xi)$ directly from the propagator SDE is by multiplying each term of the equation depicted in Fig. 3 by $S_F(p; \xi)$. After this multiplication, we obtain Fig. 4, where the first term on the right-hand side is clearly linear in $\rho_j(s; \xi)$. After decomposing this equation into its two Dirac components, the identity in Fig. 4 becomes

$$p^2 S_1(p^2) - m S_2(p^2) + \sigma_1(p^2) = 1 \quad (4a)$$

$$S_2(p^2) - m S_1 + \sigma_2(p^2) = 0, \quad (4b)$$

where $\sigma_j(p^2)$ are the Dirac scalar and vector components of the loop integral. For the second term on the right-hand side of Fig. 4, recall the Ward identity states that for QED, $Z_1 = Z_2$ [41]. Therefore the fermion propagator $S_F(p; \xi)$ shares the same renormalization constant with the fermion-photon vertex structure defined as $S_F(k)\Gamma^\mu(k, p)S_F(p)$, which indicates that the latter is also linear in $\rho_j(s; \xi)$.




FIG. 4. The diagrammatic representation of the SDE for the fermion propagator spectral functions.




FIG. 5. The diagrammatic representation of the SDE for the photon propagator.

The gauge covariance of the solutions to the fermion and boson propagator Schwinger-Dyson equations will constrain the allowed forms of the fermion-boson vertex $\Gamma^\mu(k, p)$. However the vertex in its full complexity with its 11 non-zero components is not required. Only the projections implied by the Schwinger-Dyson equation of Figs. 3, 4, and the corresponding equation for the inverse photon propagator (*i.e.* for the vacuum polarization) Fig. 5 are constrained. Thus it is this *effective* vertex that is restricted.

One specific spectral construction of the vertex structure linear in $\rho_j(s; \xi)$ and satisfying the longitudinal Ward-Green-Takahashi identity is the Gauge Technique [18], which makes the ansatz

$$S_F(p)\Gamma^\mu(k, p)S_F(p) = \int dW \frac{1}{\not{k} - W} \gamma^\mu \frac{1}{\not{p} - W} \rho(W), \quad (5)$$

where $\rho(W) = \text{sign}(W)[\rho_1(W^2) + W\rho_1(W^2)]$. In this particular case

$$\begin{aligned} \sigma_1(p^2) + \not{p}\sigma_2(p^2) &= ie^2 \int d\underline{k} \int dW \rho(W) \\ &\times \gamma^\nu \frac{1}{\not{k} - W} \gamma^\mu \frac{1}{\not{p} - W} D_{\mu\nu}(q). \end{aligned} \quad (6)$$

Transverse supplements to the Gauge Technique are required to meet various principles of QED, including renormalizability [42, 43], gauge covariance [44] and transverse Ward-Green-Takahashi identities [45–48]. However, from the equality $Z_1 = Z_2$ [41] we can further assume that such modifications are also linear in $\rho_j(s; \xi)$, and once known, allow us to calculate the loop integral in Fig. 4, resulting in a function of p as a linear functional of $\rho_j(s; \xi)$. Since this one-loop integral reduces to corrections to the fermion propagator in perturbative calculations, such p^2 dependences must be linearly generated from the free-particle propagator. Therefore after

taking the imaginary part of Fig. 4, or equivalently that of Eq. (4), we obtain

$$s\rho_1(s; \xi) - m_B\rho_2(s; \xi) - \frac{1}{\pi}\text{Im}\{\sigma_1(s + i\varepsilon; \xi)\} = 0, \quad (7a)$$

$$\rho_2(s; \xi) - m_B\rho_1(s; \xi) - \frac{1}{\pi}\text{Im}\{\sigma_2(s + i\varepsilon; \xi)\} = 0. \quad (7b)$$

The real constant term on the left-hand side disappears. After dividing Eq. (7a) by s , Eq. (7) can be rewritten as

$$\int ds' \begin{pmatrix} \Omega_{11}(s, s'; \xi) & \Omega_{12}(s, s'; \xi) \\ \Omega_{21}(s, s'; \xi) & \Omega_{22}(s, s'; \xi) \end{pmatrix} \begin{pmatrix} \rho_1(s'; \xi) \\ \rho_2(s'; \xi) \end{pmatrix} + \begin{pmatrix} \rho_1(s; \xi) \\ \rho_2(s; \xi) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (8)$$

where the $\Omega_{ij}(s, s'; \xi)$ encode all required linear operations on the spectral functions $\rho_j(s; \xi)$, which are obtained by functional derivatives similar to

$$\Omega(s, s') = -\frac{\delta}{\delta\rho(s')} \frac{1}{\pi}\text{Im}\{\sigma(s + i\varepsilon)\}. \quad (9)$$

The bare mass coupling in Eq. (7) is explicitly included in the off-diagonal terms of $\Omega_{ij}(s, s'; \xi)$. When the fermion-photon vertex is given by the Gauge Technique the resulting σ_j is given by Eq. (6). Then m_B is the only coupling between equations for ρ_1 and ρ_2 . However, when dimension-odd operators are allowed to enter the expression for $S_F(k)\Gamma^\mu(k, p)S_F(p)$, they will contribute additionally to off-diagonal elements of Ω_{ij} .

For a given ansatz for the fermion-photon vertex that ensures $S_F(k)\Gamma^\mu(k, p)S_F(p)$ being linear in $\rho_j(s; \xi)$, there is a corresponding Ω . It is the matrix Ω that is constrained by gauge covariance. Regardless of the photon being quenched or not, the SDE for fermion propagator spectral functions takes the form of Eq. (8). Solutions to Eq. (8) found in different covariant gauges are, of course, different because the fermion propagator is not a physical observable. However any ansatz for the fermion-photon vertex that respects Eq. (8) is expected to be gauge covariant. Satisfying the Ward-Green-Takahashi identity, a consequence of gauge invariance, however, is not sufficient to ensure the gauge covariance of solutions to Eq. (8), as we will see explicitly later on. In order to explore the conditions on the $\Omega_{ij}(s, s'; \xi)$ that ensure gauge covariance of solutions to Eq. (8), the LKFT for the fermion propagator needs to be solved first.

III. LKFT FOR FERMION PROPAGATOR

Detailed discussion of the LKFT for the fermion propagator has been made elsewhere [49]. Consequently in the present article, only crucial intermediate steps are included. Because of the existence of bijective relations among the fermion propagator in coordinate space, in momentum space and in spectral representation, LKFT manifests itself as isomorphic representations for the

fermion propagator in these spaces. Being linear in the coordinate representation suggests that LKFT in the spectral representation should also be a linear transform. Therefore without loss of generality,

$$\rho_j(s; \xi) = \int ds' \mathcal{K}_j(s, s'; \xi) \rho_j(s'; 0), \quad (10)$$

where distributions $\mathcal{K}_j(s, s'; \xi)$ represent linear operations that encode ξ dependences of $\rho_j(s; \xi)$ to be determined by LKFT. These operations observe closure, associativity, the existence of the identity element and the inverse elements.

Because the LKFT is independent of the initial conditions for $\rho_j(s; \xi)$, one can obtain the following differential equations for $\mathcal{K}_j(s, s'; \xi)$ by taking the ξ derivative of the coordinate space LKFT and subsequently taking the Fourier transform,

$$\frac{\partial}{\partial\xi} \int ds \frac{\mathcal{K}_j(s, s'; \xi)}{p^2 - s + i\varepsilon} = -\frac{\alpha}{4\pi} \int ds \frac{\Xi_j(p^2, s)}{p^2 - s + i\varepsilon} \mathcal{K}_j(s, s'; \xi), \quad (11)$$

where $z = p^2/s$. Explicit calculations reveal that

$$\frac{\Xi_1}{p^2 - s} = \frac{\Gamma(\epsilon)}{s} \left(\frac{4\pi\mu^2}{s} \right)^\epsilon \frac{(-2) {}_2F_1(\epsilon + 1, 3; 3 - \epsilon; z)}{(1 - \epsilon)(2 - \epsilon)} \quad (12a)$$

$$\frac{\Xi_2}{p^2 - s} = \frac{\Gamma(\epsilon)}{s} \left(\frac{4\pi\mu^2}{s} \right)^\epsilon \frac{-1}{1 - \epsilon} {}_2F_1(\epsilon + 1, 2; 2 - \epsilon; z). \quad (12b)$$

where recall ϵ is defined by $d = 4 - 2\epsilon$. Eq. (11) is solved by

$$\mathcal{K}_j = \exp\left(-\frac{\alpha\xi}{4\pi}\Phi_j\right), \quad (13)$$

where Φ_j are distributions independent of ξ and can be solved from

$$\int ds \frac{\Phi_j(s, s')}{p^2 - s + i\varepsilon} = \frac{\Xi_j(p^2, s')}{p^2 - s' + i\varepsilon}. \quad (14)$$

Eq. (14) is solved once we have established how to generate the z dependences in Eq. (12) from linear operations on the free-particle propagator with respect to the variable s alone. To do so requires the Riemann-Liouville definition of fractional calculus [50];

$$I^\alpha f(z) = \frac{1}{\Gamma(\alpha)} \int_0^z dz' (z - z')^{\alpha-1} f(z'). \quad (15)$$

For $\alpha > 0$, the Riemann-Liouville fractional derivative is defined as

$$D^\alpha f(z) = \left(\frac{d}{dz} \right)^{[\alpha]} I^{[\alpha] - \alpha} f(z), \quad (16)$$

where $[\alpha]$ is the ceiling function. Specifically for $\alpha \in (0, 1)$, $[\alpha] = 1$ and

$$D^\alpha f(z) = \frac{1}{\Gamma(1 - \alpha)} \frac{d}{dz} \int_0^z dz' (z - z')^{-\alpha} f(z'). \quad (17)$$

Furthermore, we define the dimensionless operator ϕ such that at the operator level $\int ds' \Phi = \phi$, then

$$\phi_n = \Gamma(\epsilon) \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} z^{2\epsilon+2-n} D^\epsilon z^{n-1} D^\epsilon z^{\epsilon-1}. \quad (18)$$

Distributions ϕ_n in Eq. (18) correspond to Φ_j with $n = 3, 2$ for $j = 1, 2$; see Eqs. (12a, 12b) respectively.

The exponential form of \mathcal{K}_j given by Eq. (13) remains illusive even with ϕ_n explicitly written as Eq. (18). To see how \mathcal{K}_j works explicitly, consider any function of the spectral variable that can be written as a linear combination of z^β , we can show that

$$\begin{aligned} \mathcal{K}_j z^\beta &= \exp(-\bar{\alpha}\bar{\phi}_n) z^\beta = \sum_{m=0}^{+\infty} \frac{(-\bar{\alpha})^m}{m!} \bar{\phi}_n^m z^\beta \\ &= \sum_{m=0}^{+\infty} \frac{(-\bar{\alpha})^m}{m!} \frac{\Gamma(n+\beta+(m-1)\epsilon-1)\Gamma(\beta+m\epsilon)}{\Gamma(n+\beta-\epsilon-1)\Gamma(\beta)} z^{\beta+m\epsilon}, \end{aligned} \quad (19)$$

where

$$\bar{\alpha} \equiv \frac{\alpha\xi}{4\pi} \frac{\Gamma(\epsilon)\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon. \quad (20)$$

Specifically for small ϵ , the operations given by Eq. (19) reduce to

$$\begin{aligned} \mathcal{K}_j &= \left(\frac{\mu^2 z}{p^2} \right)^{-\nu} \exp \left\{ -\nu \left[\frac{1}{\epsilon} + \gamma_E + \ln 4\pi + \mathcal{O}(\epsilon^1) \right] \right\} \\ &\times z^{2-n} I^\nu z^{n-1-\nu} I^\nu z^{-\nu-1}, \end{aligned} \quad (21)$$

where $\nu = \alpha\xi/(4\pi)$.

IV. GAUGE COVARIANCE REQUIREMENTS FOR THE PROPAGATOR SDES

A. Gauge covariance requirement on the fermion propagator SDE

For notational convenience, when two distributions are multiplied together, the integration over the spectral variable is implied. After adopting this notation, only dependences on ξ are required to be written explicitly. Therefore Eq. (8) becomes

$$\begin{pmatrix} \rho_1(\xi) \\ \rho_2(\xi) \end{pmatrix} + \begin{pmatrix} \Omega_{11}(\xi) & \Omega_{12}(\xi) \\ \Omega_{21}(\xi) & \Omega_{22}(\xi) \end{pmatrix} \begin{pmatrix} \rho_1(\xi) \\ \rho_2(\xi) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (22)$$

Since LKFT does not couple ρ_1 with ρ_2 , we have the following abbreviated versions of Eq. (10),

$$\rho_j(\xi) = \mathcal{K}_j(\xi)\rho_j(0). \quad (23)$$

Substituting Eq. (23) into Eq. (22) gives

$$\begin{pmatrix} \mathcal{K}_1(\xi) & \\ & \mathcal{K}_2(\xi) \end{pmatrix} \begin{pmatrix} \rho_1(0) \\ \rho_2(0) \end{pmatrix} + \begin{pmatrix} \Omega_{11}(\xi) & \Omega_{12}(\xi) \\ \Omega_{21}(\xi) & \Omega_{22}(\xi) \end{pmatrix} \begin{pmatrix} \mathcal{K}_1(\xi) & \\ & \mathcal{K}_2(\xi) \end{pmatrix} \begin{pmatrix} \rho_1(0) \\ \rho_2(0) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (24)$$

Since obviously

$$(\text{diag}\{\mathcal{K}_1(\xi), \mathcal{K}_2(\xi)\})^{-1} = \text{diag}\{\mathcal{K}_1(-\xi), \mathcal{K}_2(-\xi)\}$$

with matrix inversion defined by regular matrix multiplication and distribution inversion defined by distribution multiplication that gives a δ -function. Combining this result with Eq. (22) in the Landau gauge,

$$\begin{pmatrix} \rho_1(0) \\ \rho_2(0) \end{pmatrix} + \begin{pmatrix} \Omega_{11}(0) & \Omega_{12}(0) \\ \Omega_{21}(0) & \Omega_{22}(0) \end{pmatrix} \begin{pmatrix} \rho_1(0) \\ \rho_2(0) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (25)$$

yields

$$\begin{pmatrix} \Omega_{11}(0) & \Omega_{12}(0) \\ \Omega_{21}(0) & \Omega_{22}(0) \end{pmatrix} = \begin{pmatrix} \mathcal{K}_1(-\xi) & \\ & \mathcal{K}_2(-\xi) \end{pmatrix} \begin{pmatrix} \Omega_{11}(\xi) & \Omega_{12}(\xi) \\ \Omega_{21}(\xi) & \Omega_{22}(\xi) \end{pmatrix} \begin{pmatrix} \mathcal{K}_1(\xi) & \\ & \mathcal{K}_2(\xi) \end{pmatrix}. \quad (26)$$

Since for different ansatz the Landau gauge solutions $\rho(s; 0)$ are allowed to be different, Eq. (26) is the necessary condition for solutions to the SDE for the fermion propagator to be consistent with its LKFT.

Meanwhile, when $\Omega(0)$ is given by Eq. (26), Eq. (25) becomes Eq. (24), which, when viewed as equations for $\mathcal{K}_1(\xi)\rho_1(0)$ and $\mathcal{K}_2(\xi)\rho_2(0)$, is identical to Eq. (22). Therefore Eq. (26) is also the sufficient condition for solutions to the fermion propagator SDE to be consistent with LKFT. Therefore solutions of the SDE for fermion propagator are consistent with LKFT if and only if Eq. (26) is satisfied.

B. Gauge covariance requirement on the photon propagator SDE

After gauge fixing, the photon propagator becomes

$$D^{\mu\nu}(q) = \Delta^{\mu\nu}(q) + \xi \frac{q^\mu q^\nu}{q^4 + i\varepsilon}, \quad (27)$$

where

$$\Delta^{\mu\nu}(q) = \frac{G(q^2)}{q^2 + i\varepsilon} \left(g^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right) \quad (28)$$

is the Landau gauge photon propagator. The dressing function $G(q^2)$ is determined by the SDE for the photon propagator.

As illustrated in Fig. 5, the same vertex structure $S_F(k)\Gamma^\mu(k, p)S_F(p)$ appears in the SDE for photon propagator. This allows us to derive the gauge covariance

requirement on the photon propagator SDE. Meanwhile, the spectral representation ensures the transversality of the vacuum polarization tensor through the translational invariance of the loop momentum. To start with, the dependence of the photon propagator $D^{\mu\nu}(q)$ on the covariant gauge parameter ξ is completely specified by the $\xi q^\mu q^\nu / q^4$ term, as a direct consequence of which, $G(q^2)$ of Eq. (28) and the transverse vacuum polarization tensor $\Pi^{\mu\nu}(q^2) = (g^{\mu\nu} q^2 - q^\mu q^\nu) \Pi(q^2)$ are required to be independent of ξ .

Next, one should expect the analytic structure of the photon propagator to differ from that for the fermion propagator, with singularities in distinct (but related) positions. Nevertheless, we can still proceed by keeping the external momentum dependence explicit without introducing a spectral function for the photons. Therefore the consistency requirement for the photon propagator SDE is simply given by $\partial_\xi \Pi(q^2) = 0$. Since the vacuum polarization function $\Pi(q^2)$ is linear in the fermion propagator spectral functions. One can write

$$\begin{aligned} \Pi(q^2) &= \int dW \Omega^\gamma(q^2, W; \xi) \rho(W; \xi) \\ &= \int ds (\Omega_1^\gamma(q^2, s; \xi), \Omega_2^\gamma(q^2, s; \xi)) \begin{pmatrix} \rho_1(s; \xi) \\ \rho_2(s; \xi) \end{pmatrix}. \end{aligned} \quad (29)$$

With the ξ dependence of $\rho_j(s; \xi)$ given by Eq. (13), the ξ independence of $\Pi(q^2)$ specifies

$$\Omega_j^\gamma(q^2, s; \xi) = \int ds' \Omega_j^\gamma(q^2, s'; 0) \exp \left[\frac{\alpha\xi}{4\pi} \Phi_j(s', s) \right], \quad (30)$$

or at the operator level $\Omega_\xi^\gamma = \Omega_0^\gamma e^{\nu\Phi}$. This is the consistency requirement between the photon SDE and the LKFT.

V. THE DECOMPOSITION OF Ω

The operator Ω can be decomposed into components from the fermion mass, and the longitudinal and transverse parts of the photon propagator. Some of these can be calculated exactly. In the quenched approximation,

$G(q^2) = 1$ and the photon propagator is known exactly. When photons are unquenched, the vacuum polarization produces a nontrivial $G(q^2)$ in Eq. (28). In this case the introduction of a spectral representation for the photon propagator is required. Meanwhile, since the longitudinal part of the fermion-photon vertex is fixed by the Ward-Green-Takahashi identity, contributions from the $\xi q^\mu q^\nu / q^4$ term to Ω are known exactly regardless of either the dressing of the photon propagator or the transverse part of the fermion-photon vertex.

While the bare mass m_B contributes to off-diagonal terms of Ω_{ij} containing terms at most linear in m_B , allowing the following decomposition of Ω ,

$$\Omega = \Omega^m + \Omega^\xi + \Omega^\Delta, \quad (31)$$

where

$$\Omega^m(s, s') = \begin{pmatrix} & -\frac{m_B}{s} \delta(s - s') \\ -m_B \delta(s - s') & \end{pmatrix} \quad (32)$$

stands for the operation linear in m_B that is also independent of ξ . Furthermore, denoting by Ω^ξ the contribution from the longitudinal component of the photon propagator $\xi q^\mu q^\nu / q^4$, this can be readily computed exactly. While Ω^Δ is calculated with the $\Delta^{\mu\nu}(q)$ term of the photon propagator in Eq. (27), which remains unknown without either the photon dressing function or the transverse part of the fermion-photon vertex.

Ω^ξ , being linear in ξ , vanishes in the Landau gauge. While Ω^Δ depends on the gauge because of the transverse aspects of $S_F(k) \Gamma^\mu(k, p) S_F(p)$. These need not be zero in the Landau gauge, despite this being commonly assumed.

A. Exact expressions for Ω^ξ

In order to calculate Ω^ξ in any dimensions, based on Eq. (9) we need to calculate the contribution to $\sigma_j(p^2; \xi)$ as functionals of $\rho_j(s; \xi)$ with explicit dependence on the number of spacetime dimensions $d = 4 - 2\epsilon$. We denote by σ_j^ξ the contribution to σ_j from the longitudinal component of the photon propagator. After replacing $D_{\mu\nu}(q)$ by $\xi q_\mu q_\nu / q^4$, we have

$$\begin{aligned} \sigma_1^\xi(p^2) + \not{p} \sigma_2^\xi(p^2) &= ie^2 \xi \int dW \int d\underline{k} \not{k} \frac{1}{\not{k} - W} \not{k} \frac{1}{q^4} \frac{\rho(W)}{\not{p} - W} \\ &= \frac{-\alpha\xi}{4\pi} \int dW \int_0^1 dy \Gamma(\epsilon) \left(\frac{4\pi\mu^2}{s} \right)^\epsilon \left\{ \frac{-\epsilon(1-y)y}{(1-yz)\mathcal{D}^\epsilon} \not{p} + \frac{y[3\epsilon - 4 + (3 - 2\epsilon)y]}{\mathcal{D}^\epsilon} \not{p} + \frac{W}{\mathcal{D}^\epsilon} \right\} \frac{\rho(W)}{\not{p} - W}, \end{aligned} \quad (33)$$

with y being the Feynman parameter, $z = p^2/s$ and the combined denominator given by $\mathcal{D} = (1-y)(1-yz)$. After applying the integral definition of hypergeometric functions [51], we have

$$\int_0^1 dy \frac{y^p (1-y)^q}{(1-yz)^a} = \frac{\Gamma(p+1)\Gamma(q+1)}{\Gamma(p+q+2)} {}_2F_1(a, p+1; p+q+2; z). \quad (34)$$

Then the loop-integral factor of Eq. (33) becomes

$$ie^2 \int d\underline{k} \not{k} \frac{1}{\not{k} - W} \not{q} \frac{1}{q^4} = -\frac{\alpha}{4\pi} \Gamma(\epsilon) \left(\frac{4\pi\mu^2}{s} \right)^\epsilon \left\{ \frac{-\epsilon \not{p}}{(3-\epsilon)(2-\epsilon)} {}_2F_1(1+\epsilon, 2; 4-\epsilon; z) + \frac{(3\epsilon-4)\not{p}}{(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon, 2; 3-\epsilon; z) + \frac{2(3-2\epsilon)\not{p}}{(3-\epsilon)(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon, 3; 4-\epsilon; z) + \frac{W}{1-\epsilon} {}_2F_1(\epsilon, 1; 2; z) \right\}. \quad (35)$$

Since σ_j^ξ are properly formulated Feynman diagrams corresponding to loop-corrections to the fermion propagator where the $\rho_j(s; \xi)$ are given by δ -functions, one expects that linear combinations of hypergeometric functions in Eq. (35) are finite (at least in 4D) when $z \rightarrow 1$ such that there are contributions to fermion propagator functions no more singular than those of a free particle.

After numerous applications of contiguous relations for hypergeometric functions ${}_2F_1(a, b; c; z)$, Eq. (33) becomes

$$\sigma_1^\xi(p^2) = \frac{\alpha\xi}{4\pi} \int ds \left(\frac{4\pi\mu^2}{s} \right)^\epsilon \frac{\Gamma(\epsilon)}{1-\epsilon} \times {}_2F_1(\epsilon, 2; 2-\epsilon; z) \rho_1(s), \quad (36a)$$

$$\sigma_2^\xi(p^2) = \frac{\alpha\xi}{4\pi} \int ds \left(\frac{4\pi\mu^2}{s} \right)^\epsilon \frac{\epsilon\Gamma(\epsilon)}{(2-\epsilon)(1-\epsilon)} \times {}_2F_1(\epsilon+1, 2; 3-\epsilon; z) \frac{1}{s} \rho_2(s). \quad (36b)$$

Details of the intermediate steps can be found in Appendix A.

Next, since Ω^ξ is only linear in ξ , we define Θ as

$$\Theta = -\nu^{-1} \Omega^\xi, \quad (37)$$

(recalling $\nu \equiv \alpha\xi/4\pi$) such that the distribution Θ is independent of ξ . Apparently only diagonal elements of Θ_{ij} survive, therefore

$$-\nu\Theta_{11}(s, s'; \xi) = -\frac{1}{s} \frac{\delta}{\delta\rho_1(s'; \xi)} \frac{1}{\pi} \text{Im}\{\sigma_1^\xi(s+i\varepsilon; \xi)\}. \quad (38a)$$

$$-\nu\Theta_{22}(s, s'; \xi) = -\frac{\delta}{\delta\rho_2(s'; \xi)} \frac{1}{\pi} \text{Im}\{\sigma_2^\xi(s+i\varepsilon; \xi)\}. \quad (38b)$$

Let us define at the operator level $\int ds\Theta = \text{diag}\{\theta_1, \theta_2\}$. Since

$$\frac{1}{p^2 - s + i\varepsilon} = -\frac{1}{p^2} \sum_{\beta=1}^{+\infty} z^\beta,$$

Eqs. (36, 38) imply

$$-\nu\theta_1 \frac{1}{p^2 - s + i\varepsilon} = \nu\Gamma(\epsilon) \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{1}{p^2} \frac{1}{1-\epsilon} \times \sum_{\beta=1}^{+\infty} \frac{(\epsilon)_{\beta-1} (2)_{\beta-1}}{(2-\epsilon)_{\beta-1} (\beta-1)!} z^{\beta+\epsilon} \quad (39a)$$

$$-\nu\theta_2 \frac{1}{p^2 - s + i\varepsilon} = \nu\Gamma(\epsilon) \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{1}{p^2} \frac{\epsilon}{(2-\epsilon)(1-\epsilon)} \times \sum_{\beta=1}^{+\infty} \frac{(\epsilon+1)_{\beta-1} (2)_{\beta-1}}{(3-\epsilon)_{\beta-1} (\beta-1)!} z^{\beta+\epsilon}. \quad (39b)$$

Therefore we have the following identities for θ_j ,

$$\theta_1 z^\beta = \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{\Gamma(1-\epsilon)\Gamma(\beta+\epsilon-1)\beta}{\Gamma(1+\beta-\epsilon)} z^{\beta+\epsilon}, \quad (40a)$$

$$\theta_2 z^\beta = \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{\Gamma(1-\epsilon)\Gamma(\beta+\epsilon)\beta}{\Gamma(2+\beta-\epsilon)} z^{\beta+\epsilon}, \quad (40b)$$

which completely specify Θ , and consequently Ω^ξ .

B. Consistency requirement as recurrence relations

Based on previous analysis, for a given ansatz for the fermion-photon vertex that ensures the vertex structure $S_F(k)\Gamma^\mu(k, p)S_F(p)$ being linear in $\rho(W)$, the corresponding distributions Ω_{ij} can be calculated. Such an ansatz is consistent with LKFT if and only if Eq. (26) is satisfied. Independent of any ansatz, two terms Ω^m and Ω^ξ are now known exactly.

In this subsection we explore how Eq. (26) is satisfied incorporating Ω^Δ , *i.e.* with Ω^m and Ω^ξ explicitly included. Straightforwardly, one could substitute Eq. (31) with known components into the consistency requirement Eq. (26), and obtain

$$\Omega^\Delta = e^{-\nu\Phi} (\Omega^m + \Omega_0^\Delta) e^{\nu\Phi} - \Omega^m + \nu\Phi, \quad (41)$$

as the consistency requirement on Ω^Δ . Alternatively, with LKFT for fermion propagator spectral functions given by Eq. (13), we have

$$\Omega_\xi = e^{-\nu\Phi} \Omega_0 e^{\nu\Phi}, \quad (42)$$

where the subscript of Ω_ξ highlights the ξ dependent Ω in Eq. (31), therefore $\Omega_0 = \lim_{\xi \rightarrow 0} \Omega_\xi$. To see how infinitesimal

changes in ξ affect Ω^Δ , consider taking the derivative with respect to ν (effectively ξ) of Eq. (42),

$$\partial_\nu \Omega_\xi = -\Phi e^{-\nu\Phi} \Omega_0 e^{\nu\Phi} + e^{-\nu\Phi} \Omega_0 e^{\nu\Phi} \Phi = [\Omega_\xi, \Phi].$$

Substituting in Eq. (31) and Eq. (37) produces

$$\partial_\nu \Omega^\Delta + [\Phi, \Omega^\Delta] = -[\Phi, \Omega^m] + \Theta + \nu[\Phi, \Theta]. \quad (43)$$

In order to recover the corresponding terms using the spectral representation for the fermion propagator, one calculates

$$\int ds ds' \frac{1}{p^2 - s + i\epsilon} \Omega(s, s') \rho(s'). \quad (44)$$

Since the z^β expansion is in fact the p^2/s expansion of the free-particle propagator, commutators of operations on z^β should be calculated with z^β to the left. There exists an alternative convention to Eq. (44) that locates the free-particle propagator to the right of the operation, which subsequently modifies Eq. (26). The net effect of adopting the alternative convention to solutions of Eq. (43) is, however, zero compared with the convention given by Eq. (44) because deriving Eq. (26) using the alternative convention for the location of free-particle propagator leads to exchanging \mathcal{K}_j^ξ with $\mathcal{K}_j^{-\xi}$.

Within this convention of locations, the right-hand side of Eq. (43), operating on z^β can be calculated according to Eqs. (B1, B2).

Since physical Ω^Δ are generated by loop-corrections to the fermion propagator, the following criteria apply:

- while the dependence of Ω^Δ on $\nu = \alpha\xi/(4\pi)$ is allowed to be any order, Ω^Δ cannot depend on the

bare coupling alone because of the renormalizability of fermion propagator SDE, the bare and renormalized forms of $\alpha\xi$ being identical.

- for diagonal elements of Ω^Δ , a trivial solution exists with $\Omega^\Delta = \nu\Theta = -\Omega^\xi$. However, in this case there is no correction to the free-particle propagator.

More generally, we define

$$\Omega^\Delta = \begin{pmatrix} \Omega_{11}^\Delta & -\frac{m_B}{p^2} \Omega_{12}^\Delta \\ -m_B \Omega_{21}^\Delta & \Omega_{22}^\Delta \end{pmatrix} \quad (45)$$

such that Ω_{ij}^Δ correspond to dimensionless transforms. In addition, from Eq. (13), one can easily verify $\partial_\nu \mathcal{K}_\xi + \Phi \mathcal{K}_\xi = 0$. The similarity of Eq. (43) to this differential equation for \mathcal{K}_ξ indicates the following expansions for Ω_{ij}^Δ ,

$$\Omega_{ij}^\Delta z^\beta = \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{m\epsilon} \omega_{ij}(\beta, m) z^{\beta+m\epsilon}, \quad (46a)$$

for $(ij) \neq (12)$, and

$$\Omega_{12}^\Delta z^\beta = \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{m\epsilon} \omega_{12}(\beta, m) z^{\beta+m\epsilon+1}. \quad (46b)$$

where the expansion coefficients $\omega_{ij}(\beta, m)$ are allowed to implicitly depend on ϵ . The ‘12’ component of Ω^Δ is expanded differently from other components to ensure that Ω^Δ given by Eq. (45) translates into operations solely on the spectral variables.

With Eq. (46), the left-hand side of Eq. (43) can be calculated according to Eq. (B3). Then recurrence relations for $\omega_{ij}(\beta, m)$ are obtained by the comparison of $\mathcal{O}(\nu^m)$ terms in Eq. (43). As a result, we have

$$\begin{aligned} & -\omega_{11}(\beta, m+1) + \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} \omega_{11}(\beta+\epsilon, m) - \omega_{11}(\beta, m) \frac{\Gamma(2+\beta+m\epsilon)\Gamma(\beta+(m+1)\epsilon)}{\Gamma(2+\beta+(m-1)\epsilon)\Gamma(\beta+m\epsilon)} \right\} \\ & = \begin{cases} \Gamma(1-\epsilon) \frac{\Gamma(\beta+\epsilon-1)\Gamma(\beta+1)}{\Gamma(1+\beta-\epsilon)\Gamma(\beta)} & \text{for } m=0, \\ -\frac{\Gamma(\epsilon)[\Gamma(1-\epsilon)]^2}{\Gamma(1+\epsilon)} \left(\frac{\beta+1}{\beta+2\epsilon} - \frac{1+\beta+\epsilon}{\beta+1} \right) \frac{\Gamma(\beta+2\epsilon)\Gamma(\beta+\epsilon+1)}{\Gamma(\beta-\epsilon+2)\Gamma(\beta)} & \text{for } m=1, \\ 0 & \text{for } m \geq 2. \end{cases} \quad (47a) \end{aligned}$$

$$\begin{aligned} & -\omega_{12}(\beta, m+1) + \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} \omega_{12}(\beta+\epsilon, m) - \omega_{12}(\beta, m) \frac{\Gamma(2+\beta+m\epsilon)\Gamma(\beta+1+(m+1)\epsilon)}{\Gamma(2+\beta+(m-1)\epsilon)\Gamma(\beta+1+m\epsilon)} \right\} \\ & = \begin{cases} \Gamma(1-\epsilon) \frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta+1)} & \text{for } m=0, \\ 0 & \text{for } m \geq 1. \end{cases} \quad (47b) \end{aligned}$$

$$\begin{aligned}
& -\omega_{21}(\beta, m+1) + \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(1+\beta)\Gamma(\beta+\epsilon)}{\Gamma(1+\beta-\epsilon)\Gamma(\beta)} \omega_{21}(\beta+\epsilon, m) - \omega_{21}(\beta, m) \frac{\Gamma(2+\beta+m\epsilon)\Gamma(\beta+(m+1)\epsilon)}{\Gamma(2+\beta+(m-1)\epsilon)\Gamma(\beta+m\epsilon)} \right\} \\
& = \begin{cases} \Gamma(1-\epsilon) \frac{\Gamma(1+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} & \text{for } m=0, \\ 0 & \text{for } m \geq 1. \end{cases} \quad (47c)
\end{aligned}$$

$$\begin{aligned}
& -\omega_{22}(\beta, m+1) + \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(1+\beta)\Gamma(\beta+\epsilon)}{\Gamma(1+\beta-\epsilon)\Gamma(\beta)} \omega_{22}(\beta+\epsilon, m) - \omega_{22}(\beta, m) \frac{\Gamma(1+\beta+m\epsilon)\Gamma(\beta+(m+1)\epsilon)}{\Gamma(1+\beta+(m-1)\epsilon)\Gamma(\beta+m\epsilon)} \right\} \\
& = \begin{cases} \Gamma(1-\epsilon) \frac{\Gamma(\beta+\epsilon)\Gamma(\beta+1)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} & \text{for } m=0, \\ -\frac{\Gamma(\epsilon)[\Gamma(1-\epsilon)]^2}{\Gamma(1+\epsilon)} \left(\frac{1}{\beta+1} - \frac{1}{1+\beta-\epsilon} \right) \frac{\Gamma(\beta+2\epsilon)\Gamma(\beta+\epsilon+1)}{\Gamma(\beta-\epsilon+1)\Gamma(\beta)} & \text{for } m=1, \\ 0 & \text{for } m \geq 2. \end{cases} \quad (47d)
\end{aligned}$$

These recurrence relations specify how gauge covariance is satisfied when distributions Ω_{ij}^Δ are expanded as Taylor series in $\nu = \alpha\xi/4\pi$ written in Eq. (46). On one hand, when the Ω_{ij}^Δ are only known in the Landau gauge, Eq. (47) can be used to calculate Ω_{ij}^Δ in any other covariant gauge. On the other hand, when an ansatz for $S_F(k)\Gamma^\mu(k, p)S_F(p)$ is known, the operations of Ω_{ij}^Δ on z^β can be calculated. Eq. (47) then works to verify if this ansatz ensures that solutions to fermion propagator SDE are consistent with LKFT.

C. Example: The Gauge Technique in the quenched approximation in 4D

In the quenched approximation with the Gauge Technique ansatz for $S_F(k)\Gamma^\mu(k, p)S_F(p)$ [18], we deduce the Ω_{ij} to be

$$\begin{aligned}
\Omega_{11}(s, s'; \xi) &= -\frac{3\alpha}{4\pi} \left\{ \left(\frac{1}{\epsilon} - \gamma_E + \ln 4\pi + \frac{4}{3} + \ln \frac{\mu^2}{s} \right) \right. \\
&\quad \left. \times \delta(s-s') - \frac{s'}{s^2} \theta(s-s') \right\} - \frac{\alpha\xi}{4\pi} \frac{1}{s} \theta(s-s'), \\
\Omega_{12}(s, s'; \xi) &= -\frac{m_B}{s} \delta(s-s'), \\
\Omega_{21}(s, s'; \xi) &= -m_B \delta(s-s'), \\
\Omega_{22}(s, s'; \xi) &= -\frac{3\alpha}{4\pi} \left\{ \left(\frac{1}{\epsilon} - \gamma_E + \ln 4\pi + \frac{4}{3} + \ln \frac{\mu^2}{s} \right) \right. \\
&\quad \left. \times \delta(s-s') - \frac{1}{s} \theta(s-s') \right\} - \frac{\alpha\xi}{4\pi} \frac{s'}{s^2} \theta(s-s'), \quad (48)
\end{aligned}$$

where m_B is the bare mass and $d = 4 - 2\epsilon$. Equivalently written as operators on z , Ω_{ij} become

$$\begin{aligned}
\Omega_{11}(\xi) &= -\frac{3\alpha}{4\pi} \left[\tilde{C} + \ln(z) - z^{-1}I \right] - \frac{\alpha\xi}{4\pi} I z^{-1}, \\
\Omega_{12} &= -\frac{m_B}{p^2} z, \\
\Omega_{21} &= -m_B, \\
\Omega_{22}(\xi) &= -\frac{3\alpha}{4\pi} \left[\tilde{C} + \ln(z) - I z^{-1} \right] - \frac{\alpha\xi}{4\pi} z^{-1}I, \quad (49)
\end{aligned}$$

where $\tilde{C} = 1/\epsilon - \gamma_E + \ln(4\pi\mu^2/p^2) + 4/3$.

Meanwhile, since in 4D the LKFT for the fermion propagator reduces to Eq. (21), we have,

$$\begin{aligned}
z^\beta \mathcal{K}_1(\xi) &= \left(\frac{\mu^2}{p^2} \right)^{-\nu} \exp \left\{ -\nu \left[\frac{1}{\epsilon} + \gamma_E + \ln(4\pi) \right] \right\} \\
&\quad \times \frac{\Gamma(\beta-\nu)\Gamma(2+\beta-\nu)}{\Gamma(\beta)\Gamma(2+\beta)} z^{\beta-\nu}, \quad (50a)
\end{aligned}$$

$$\begin{aligned}
z^\beta \mathcal{K}_2(\xi) &= \left(\frac{\mu^2}{p^2} \right)^{-\nu} \exp \left\{ -\nu \left[\frac{1}{\epsilon} + \gamma_E + \ln(4\pi) \right] \right\} \\
&\quad \times \frac{\Gamma(\beta-\nu)\Gamma(1+\beta-\nu)}{\Gamma(\beta)\Gamma(1+\beta)} z^{\beta-\nu}. \quad (50b)
\end{aligned}$$

For the consistency requirement, it is more convenient to write Eq. (26) as

$$\begin{aligned}
& \begin{pmatrix} \Omega_{11}(\xi) & \Omega_{12}(\xi) \\ \Omega_{21}(\xi) & \Omega_{22}(\xi) \end{pmatrix} \\
& = \begin{pmatrix} \mathcal{K}_1(\xi)\Omega_{11}(0)\mathcal{K}_1(-\xi) & \mathcal{K}_1(\xi)\Omega_{12}(0)\mathcal{K}_2(-\xi) \\ \mathcal{K}_2(\xi)\Omega_{21}(0)\mathcal{K}_1(-\xi) & \mathcal{K}_2(\xi)\Omega_{22}(0)\mathcal{K}_2(-\xi) \end{pmatrix}. \quad (51)
\end{aligned}$$

With the assistance of the following four identities for

fractional calculus,

$$I^\alpha z^\beta = \frac{\Gamma(\beta+1)}{\Gamma(\alpha+\beta+1)} z^{\alpha+\beta}, \quad (52a)$$

$$D^\alpha z^\beta = \frac{\Gamma(\beta+1)}{\Gamma(-\alpha+\beta+1)} z^{-\alpha+\beta}, \quad (52b)$$

$$I^\alpha z^\beta \ln(z) = \frac{\Gamma(\beta+1)}{\Gamma(\alpha+\beta+1)} \left\{ \psi(\beta+1) - \psi(\alpha+\beta+1) + \ln(z) \right\} z^{\alpha+\beta}, \quad (52c)$$

$$D^\alpha z^\beta \ln(z) = \frac{\Gamma(\beta+1)}{\Gamma(-\alpha+\beta+1)} \left\{ \psi(\beta+1) - \psi(-\alpha+\beta+1) + \ln(z) \right\} z^{-\alpha+\beta}, \quad (52d)$$

where $\psi(\beta)$ is the digamma function, one then obtains

$$\begin{aligned} & z^\beta \mathcal{K}_1(\xi) \Omega_{11}(0) \mathcal{K}_1(-\xi) \\ &= -\frac{3\alpha}{4\pi} \left\{ \tilde{C} - \frac{1}{\beta-\nu+1} + \psi(\beta) - \psi(\beta-\nu) + \right. \\ & \quad \left. \psi(\beta+2) - \psi(\beta+2-\nu) + \ln z \right\} z^\beta, \end{aligned} \quad (53a)$$

$$z^\beta \mathcal{K}_1(\xi) \Omega_{12}(0) \mathcal{K}_2(-\xi) = -\frac{m_B}{p^2} \frac{\beta}{\beta-\nu} z^{\beta+1}, \quad (53b)$$

$$z^\beta \mathcal{K}_2(\xi) \Omega_{21}(0) \mathcal{K}_1(-\xi) = -m_B \frac{\beta+1}{\beta+1-\nu} z^\beta, \quad (53c)$$

$$\begin{aligned} & z^\beta \mathcal{K}_2(\xi) \Omega_{22}(0) \mathcal{K}_2(-\xi) \\ &= -\frac{3\alpha}{4\pi} \left\{ \tilde{C} - \frac{1}{\beta-\nu} + \psi(\beta) - \psi(\beta-\nu) + \right. \\ & \quad \left. \psi(\beta+1) - \psi(\beta+1-\nu) + \ln z \right\} z^\beta. \end{aligned} \quad (53d)$$

While from Eq. (49), we have

$$z^\beta \Omega_{11}(\xi) = \left\{ -\frac{3\alpha}{4\pi} \left[\tilde{C} - \frac{1}{\beta+1} + \ln z \right] - \frac{\nu}{\beta} \right\} z^\beta, \quad (54a)$$

$$z^\beta \Omega_{12}(\xi) = -\frac{m_B}{p^2} z^{\beta+1}, \quad (54b)$$

$$z^\beta \Omega_{21}(\xi) = -m_B z^\beta, \quad (54c)$$

$$z^\beta \Omega_{22}(\xi) = \left\{ -\frac{3\alpha}{4\pi} \left[\tilde{C} - \frac{1}{\beta} + \ln z \right] - \frac{\nu}{\beta+1} \right\} z^\beta \quad (54d)$$

Observe that the digamma functions only occur in Eq. (53), not in Eq. (54). Additionally, the dependence on ν is only linear in Eq. (54), but not in Eq. (53). Therefore the consistency requirement given by Eq. (51) is not satisfied by the Gauge Technique in 4D. The same conclusion has been realized by Delbourgo, Keck and Parker [52] in a completely different approach.

VI. SUMMARY

In this paper we have formulated the fermion propagator SDE in terms of propagator spectral functions. With

the fermion-photon vertex structure $S_F(k)\Gamma^\mu(k,p)S_F(p)$ being linear in the $\rho_j(s;\xi)$ as implied by the equality of renormalization factors $Z_1 = Z_2$, we have derived the necessary and sufficient condition for the solutions of the fermion propagator SDE to be consistent with LKFT in covariant gauges. With known contributions to the fermion propagator SDE calculated, this reduces the consistency requirement to that for the contribution to Ω in Eq. (8) from the Landau gauge photon propagator. Next, an expansion of the operator Ω^Δ (defined in Eq.(31)), similar to that of \mathcal{K}_j in Eq. (19), has been postulated in Eq. (46). The consistency requirements can then be converted into the form of recurrence relations of this expansion, shown in Eq. (47). The requirement on $S_F(k)\Gamma^\mu(k,p)S_F(p)$ to ensure the gauge invariance of $\Pi(q^2)$ was also derived.

We observe that the Gauge Technique [15–18] does not ensure gauge covariance for the fermion propagator in QED. In fact, when fermions are massive, dimension-odd operators are required in $S_F(k)\Gamma^\mu(k,p)S_F(p)$ to ensure gauge covariance. Our formalism for the SDEs using a spectral representation allows propagators to be solved in Minkowski space. Furthermore, our consistency requirements can be used as criteria for truncating the SDEs for QED propagators.

Importantly, our calculations have been performed in arbitrary dimensions. Keeping $\epsilon = 2 - d/2$ explicit to the end turns out to give concise and meaningful results in the case of the σ_j^ξ in Eq. (33), the fermion Schwinger-Dyson equation, as well as the LKFT for the fermion propagator. Results are concise in the sense that one hypergeometric function describes the p^2 dependence for each Dirac component of every loop integral. Meaningful in the sense that the results apply to any number of spacetime dimensions as long as hypergeometric functions converge. Based on these two merits, one might suspect that dimensional regularization evaluated by keeping ϵ explicit to the last step is intrinsic to QED itself.

This work marks a path towards ensuring consistent truncations of the Schwinger-Dyson equations for the fermion and boson propagators yield gauge covariant fermion mass functions like that in Fig. 1: an essential requirement for validating any truncation scheme used.

ACKNOWLEDGMENTS

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract DE-AC05-06OR23177 that funds Jefferson Lab research. The authors would like to thank Professor Keith Ellis and other members of the Institute for Particle Physics Phenomenology (IPPP) of Durham University for kind hospitality during their visit when this article was finalized.

Appendix A: Simplification of $\sigma_j^\xi(p^2; \xi)$

To simplify Eq. (35), we will need contiguous relations for hypergeometric functions from Ref. [51] and the following identity

$$\begin{aligned} (Ap + BW)(p + W) &= s(Az + B) + (A + B)pW \\ &= s(z - 1) \left[\left(A + \frac{A + B}{z - 1} \right) + \frac{A + B}{s(z - 1)} pW \right]. \end{aligned} \quad (\text{A1})$$

Equations referred to by Eq. (15.2.XX) are identities in Ref. [51]. With $a = 3$, $b = 1 + \epsilon$, $c = 4 - \epsilon$, Eq. (15.2.19) becomes

$$\begin{aligned} & \frac{2(3 - 2\epsilon)}{(3 - \epsilon)(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon, 3; 4 - \epsilon; z) \\ &= \frac{2}{(3 - \epsilon)(2 - \epsilon)} {}_2F_1(2, 1 + \epsilon; 4 - \epsilon; z) + \\ & \frac{2(1 - z)}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(3, 1 + \epsilon; 4 - \epsilon; z). \end{aligned} \quad (\text{A2})$$

Explicitly then,

$$\begin{aligned} A &= \frac{-\epsilon}{(3 - \epsilon)(2 - \epsilon)} {}_2F_1(1 + \epsilon, 2; 4 - \epsilon; z) + \\ & \frac{3\epsilon - 4}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon, 2; 3 - \epsilon; z) + \\ & \frac{2(3 - 2\epsilon)}{(3 - \epsilon)(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon, 3; 4 - \epsilon; z) \\ &= \frac{2(1 - z)}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(1 + \epsilon, 3; 4 - \epsilon; z) + \\ & \frac{1}{3 - \epsilon} {}_2F_1(1 + \epsilon, 2; 4 - \epsilon; z) + \\ & \frac{3\epsilon - 4}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon, 2; 3 - \epsilon; z), \end{aligned} \quad (\text{A3})$$

where Eq. (A2) is used to derive Eq. (A3). From Eq. (15.2.17) with $a = 1$, $b = \epsilon$, $c = 3 - \epsilon$ we have

$$\begin{aligned} B &= \frac{1}{1 - \epsilon} {}_2F_1(\epsilon, 1; 2 - \epsilon; z) \\ &= \frac{1}{2 - \epsilon} {}_2F_1(\epsilon, 1; 3 - \epsilon; z) + \\ & \frac{1}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon, 2; 3 - \epsilon; z). \end{aligned} \quad (\text{A4})$$

Next, with $a = \epsilon$, $b = 2$, $c = 3 - \epsilon$, Eq. (15.2.15) becomes

$$\begin{aligned} & \frac{1}{2 - \epsilon} {}_2F_1(\epsilon, 1; 3 - \epsilon; z) \\ &= \frac{1 - 2\epsilon}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(2, \epsilon; 3 - \epsilon; z) + \\ & \frac{\epsilon(1 - z)}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(2, \epsilon + 1; 3 - \epsilon; z). \end{aligned} \quad (\text{A5})$$

With $a = \epsilon$, $b = 3$, $c = 4 - \epsilon$, Eq. (15.2.17) becomes

$$\begin{aligned} & \frac{-1}{1 - \epsilon} {}_2F_1(\epsilon, 2; 3 - \epsilon; z) \\ &= \frac{-(3 - 2\epsilon)}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon, 2; 4 - \epsilon; z) + \\ & \frac{-\epsilon}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon + 1, 2; 4 - \epsilon; z). \end{aligned} \quad (\text{A6})$$

With $a = 2$, $b = 1 + \epsilon$, $c = 4 - \epsilon$, Eq. (15.2.15) becomes

$$\begin{aligned} & -(3 - 2\epsilon) {}_2F_1(\epsilon, 2; 4 - \epsilon; z) + \\ & (1 - 2\epsilon) {}_2F_1(\epsilon + 1, 2; 4 - \epsilon; z) \\ &= -2(1 - z) {}_2F_1(\epsilon + 1, 3; 4 - \epsilon; z). \end{aligned} \quad (\text{A7})$$

Therefore

$$\begin{aligned} & A + B \\ &= \frac{2(1 - z)}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(1 + \epsilon, 3; 4 - \epsilon; z) + \\ & \frac{\epsilon(1 - z)}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon + 1, 2; 3 - \epsilon; z) + \end{aligned} \quad (\text{A8a})$$

$$\begin{aligned} & \frac{-1}{1 - \epsilon} {}_2F_1(\epsilon, 2; 3 - \epsilon; z) + \\ & \frac{1}{3 - \epsilon} {}_2F_1(1 + \epsilon, 2; 4 - \epsilon; z) \\ &= \frac{2(1 - \epsilon)}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(1 + \epsilon, 3; 4 - \epsilon; z) + \end{aligned} \quad (\text{A8b})$$

$$\begin{aligned} & \frac{\epsilon(1 - z)}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon + 1, 2; 3 - \epsilon; z) + \\ & \frac{-(3 - 2\epsilon)}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon, 2; 4 - \epsilon; z) + \\ & \frac{1 - 2\epsilon}{(3 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon + 1, 2; 4 - \epsilon; z) \\ &= \frac{\epsilon(1 - z)}{(2 - \epsilon)(1 - \epsilon)} {}_2F_1(\epsilon + 1, 2; 3 - \epsilon; z), \end{aligned} \quad (\text{A8c})$$

where Eqs. (A5, A6, A7) are used to derive Eqs. (A8a, A8b, A8c), respectively. In addition, with $a = \epsilon$, $b = 2$, $c = 4 - \epsilon$, Eq. (15.2.14) becomes

$$\begin{aligned} & \epsilon {}_2F_1(\epsilon + 1, 2; 4 - \epsilon; z) \\ &= 2 {}_2F_1(\epsilon, 3; 4 - \epsilon; z) - (2 - \epsilon) {}_2F_1(\epsilon, 2; 4 - \epsilon; z). \end{aligned} \quad (\text{A9})$$

With $a = \epsilon$, $b = 2$, $c = 3 - \epsilon$, Eq. (15.2.14) becomes

$$\begin{aligned} & \epsilon {}_2F_1(\epsilon + 1, 2; 3 - \epsilon; z) \\ &= 2 {}_2F_1(\epsilon, 3; 3 - \epsilon; z) - (2 - \epsilon) {}_2F_1(\epsilon, 2; 4 - \epsilon; z). \end{aligned} \quad (\text{A10})$$

With $a = \epsilon$, $b = 2$, $c = 4 - \epsilon$, Eq. (15.2.24) becomes

$$\begin{aligned} & (1 - \epsilon) {}_2F_1(\epsilon, 2; 4 - \epsilon; z) + 2 {}_2F_1(\epsilon, 3; 4 - \epsilon; z) \\ &= (3 - \epsilon) {}_2F_1(\epsilon, 2; 3 - \epsilon; z). \end{aligned} \quad (\text{A11})$$

With $a = \epsilon$, $b = 2$, $c = 3 - \epsilon$, Eq. (15.2.24) becomes

$$\begin{aligned} & -\epsilon {}_2F_1(\epsilon, 2; 3 - \epsilon; z) + 2 {}_2F_1(\epsilon, 2; 3 - \epsilon; z) \\ &= (2 - \epsilon) {}_2F_1(\epsilon, 2; 2 - \epsilon; z). \end{aligned} \quad (\text{A12})$$

Then

$$\begin{aligned}
A + \frac{A+B}{(z-1)} &= \frac{-\epsilon}{(3-\epsilon)(2-\epsilon)} {}_2F_1(1+\epsilon, 2; 4-\epsilon; z) + \frac{3\epsilon-4}{(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon, 2; 3-\epsilon; z) + \\
&\quad \frac{2(3-2\epsilon)}{(3-\epsilon)(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon, 3; 4-\epsilon; z) + \frac{-\epsilon}{(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon+1, 2; 3-\epsilon; z) \\
&= \frac{1}{(3-\epsilon)(2-\epsilon)} \left\{ \frac{2(2-\epsilon)}{1-\epsilon} {}_2F_1(\epsilon, 3; 4-\epsilon; z) + (2-\epsilon) {}_2F_1(\epsilon, 2; 4-\epsilon; z) \right\} + \tag{A13a}
\end{aligned}$$

$$\begin{aligned}
&\quad \frac{1}{(2-\epsilon)(1-\epsilon)} \left\{ -2 {}_2F_1(\epsilon, 3; 3-\epsilon; z) + 2(\epsilon-1) {}_2F_1(\epsilon, 2; 3-\epsilon; z) \right\} \\
&= \frac{2}{(3-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon, 3; 4-\epsilon; z) + \frac{1}{3-\epsilon} {}_2F_1(\epsilon, 2; 4-\epsilon; z) + \tag{A13b}
\end{aligned}$$

$$\begin{aligned}
&\quad \frac{-2}{(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon, 3; 3-\epsilon; z) + \frac{-2}{2-\epsilon} {}_2F_1(\epsilon, 2; 3-\epsilon; z) \\
&= \frac{\epsilon}{(1-\epsilon)(2-\epsilon)} {}_2F_1(\epsilon, 2; 3-\epsilon; z) + \frac{-2}{(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon, 3; 3-\epsilon; z) \tag{A13c}
\end{aligned}$$

$$= \frac{-1}{1-\epsilon} {}_2F_1(\epsilon, 2; 2-\epsilon; z), \tag{A13d}$$

where Eqs. (A9, A10, A11, A12) have been utilized to derive Eqs. (A13a, A13b, A13c, A13d), respectively. Finally we obtain,

$$\begin{aligned}
ie^2\xi \int d\mathbf{k} \not{k} \frac{1}{\not{k}-W} \not{q} \frac{1}{q^4} \frac{1}{\not{p}-W} &= \frac{-\alpha\xi}{4\pi} \Gamma(\epsilon) \left(\frac{4\pi\mu^2}{s} \right)^\epsilon \left\{ \frac{-1}{1-\epsilon} {}_2F_1(\epsilon, 2; 2-\epsilon; z) + \right. \\
&\quad \left. \frac{-\epsilon}{(2-\epsilon)(1-\epsilon)} {}_2F_1(\epsilon+1, 2; 3-\epsilon; z) \frac{\not{p}}{W} \right\}. \tag{A14}
\end{aligned}$$

Appendix B: Operations on z^β from terms in Eq. (43)

For commutators on the right-hand side of Eq. (43), explicit calculation shows that

$$z^\beta \phi_3 \theta_1 = \frac{\Gamma(\epsilon)[\Gamma(1-\epsilon)]^2}{\Gamma(1+\epsilon)} \left(\frac{4\pi\mu^2}{p^2} \right)^{2\epsilon} \frac{\beta+1}{\beta+2\epsilon} \frac{\Gamma(\beta+2\epsilon)\Gamma(\beta+\epsilon+1)}{\Gamma(\beta-\epsilon+2)\Gamma(\beta)} z^{\beta+2\epsilon}, \tag{B1a}$$

$$z^\beta \theta_1 \phi_3 = \frac{\Gamma(\epsilon)[\Gamma(1-\epsilon)]^2}{\Gamma(1+\epsilon)} \left(\frac{4\pi\mu^2}{p^2} \right)^{2\epsilon} \frac{1+\beta+\epsilon}{\beta+1} \frac{\Gamma(\beta+2\epsilon)\Gamma(\beta+\epsilon+1)}{\Gamma(\beta-\epsilon+2)\Gamma(\beta)} z^{\beta+2\epsilon}, \tag{B1b}$$

$$z^\beta \phi_2 \theta_2 = \frac{\Gamma(\epsilon)[\Gamma(1-\epsilon)]^2}{\Gamma(1+\epsilon)} \left(\frac{4\pi\mu^2}{p^2} \right)^{2\epsilon} \frac{1}{\beta+1} \frac{\Gamma(\beta+2\epsilon)\Gamma(\beta+\epsilon+1)}{\Gamma(\beta-\epsilon+1)\Gamma(\beta)} z^{\beta+2\epsilon}, \tag{B1c}$$

$$z^\beta \theta_2 \phi_2 = \frac{\Gamma(\epsilon)[\Gamma(1-\epsilon)]^2}{\Gamma(1+\epsilon)} \left(\frac{4\pi\mu^2}{p^2} \right)^{2\epsilon} \frac{1}{1+\beta-\epsilon} \frac{\Gamma(\beta+2\epsilon)\Gamma(\beta+\epsilon+1)}{\Gamma(\beta-\epsilon+1)\Gamma(\beta)} z^{\beta+2\epsilon}, \tag{B1d}$$

and

$$\begin{aligned}
z^\beta (\phi_3 z - z \phi_3) &= \Gamma(\epsilon) \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left[\frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} - \frac{\Gamma(2+\beta)\Gamma(\beta+1+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta+1)} \right] z^{\beta+\epsilon+1} \\
&= -\Gamma(1-\epsilon) \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta+1)} z^{\beta+\epsilon+1} \tag{B2a}
\end{aligned}$$

$$\begin{aligned}
z^\beta (\phi_2 - \phi_3) &= \Gamma(\epsilon) \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left[\frac{\Gamma(1+\beta)\Gamma(\beta+\epsilon)}{\Gamma(1+\beta-\epsilon)\Gamma(\beta)} - \frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} \right] z^{\beta+\epsilon} \\
&= -\Gamma(1-\epsilon) \left(\frac{4\pi\mu^2}{p^2} \right)^\epsilon \frac{\Gamma(1+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} z^{\beta+\epsilon}. \tag{B2b}
\end{aligned}$$

Up until now all terms on the right-hand side of Eq. (43) are explicit. For the left-hand side, we have

$$z^\beta \partial_\nu \Omega_{ij}^\Delta = \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{(m+1)\epsilon} [-\omega_{ij}(\beta, m+1)] z^{\beta+(m+1)\epsilon} \quad (i, j) \neq (1, 2), \quad (\text{B3a})$$

$$z^\beta \partial_\nu \Omega_{12}^\Delta = \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{(m+1)\epsilon} [-\omega_{12}(\beta, m+1)] z^{1+\beta+(m+1)\epsilon} \quad (\text{B3b})$$

$$\begin{aligned} z^\beta (\phi_3 \Omega_{11}^\Delta - \Omega_{11}^\Delta \phi_3) &= \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{(m+1)\epsilon} \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} \omega_{11}(\beta+\epsilon, m) \right. \\ &\quad \left. - \omega_{11}(\beta, m) \frac{\Gamma(2+\beta+m\epsilon)\Gamma(\beta+(m+1)\epsilon)}{\Gamma(2+\beta+(m-1)\epsilon)\Gamma(\beta+m\epsilon)} \right\} z^{\beta+(m+1)\epsilon} \end{aligned} \quad (\text{B3c})$$

$$\begin{aligned} z^\beta (\phi_3 \Omega_{12}^\Delta - \Omega_{12}^\Delta \phi_2) &= \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{(m+1)\epsilon} \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(2+\beta)\Gamma(\beta+\epsilon)}{\Gamma(2+\beta-\epsilon)\Gamma(\beta)} \omega_{12}(\beta+\epsilon, m) \right. \\ &\quad \left. - \omega_{12}(\beta, m) \frac{\Gamma(2+\beta+m\epsilon)\Gamma(\beta+1+(m+1)\epsilon)}{\Gamma(2+\beta+(m-1)\epsilon)\Gamma(\beta+1+m\epsilon)} \right\} z^{1+\beta+(m+1)\epsilon} \end{aligned} \quad (\text{B3d})$$

$$\begin{aligned} z^\beta (\phi_2 \Omega_{21}^\Delta - \Omega_{21}^\Delta \phi_3) &= \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{(m+1)\epsilon} \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(1+\beta)\Gamma(\beta+\epsilon)}{\Gamma(1+\beta-\epsilon)\Gamma(\beta)} \omega_{21}(\beta+\epsilon, m) \right. \\ &\quad \left. - \omega_{21}(\beta, m) \frac{\Gamma(2+\beta+m\epsilon)\Gamma(\beta+(m+1)\epsilon)}{\Gamma(2+\beta+(m-1)\epsilon)\Gamma(\beta+m\epsilon)} \right\} z^{\beta+(m+1)\epsilon} \end{aligned} \quad (\text{B3e})$$

$$\begin{aligned} z^\beta (\phi_2 \Omega_{22}^\Delta - \Omega_{22}^\Delta \phi_2) &= \sum_{m=0}^{+\infty} \frac{(-\nu)^m}{m!} \left(\frac{4\pi\mu^2}{p^2} \right)^{(m+1)\epsilon} \Gamma(\epsilon) \frac{\Gamma(1-\epsilon)}{\Gamma(1+\epsilon)} \left\{ \frac{\Gamma(1+\beta)\Gamma(\beta+\epsilon)}{\Gamma(1+\beta-\epsilon)\Gamma(\beta)} \omega_{22}(\beta+\epsilon, m) \right. \\ &\quad \left. - \omega_{22}(\beta, m) \frac{\Gamma(1+\beta+m\epsilon)\Gamma(\beta+(m+1)\epsilon)}{\Gamma(1+\beta+(m-1)\epsilon)\Gamma(\beta+m\epsilon)} \right\} z^{\beta+(m+1)\epsilon}. \end{aligned} \quad (\text{B3f})$$

-
- [1] A. Bashir, L. Chang, I. C. Cloet, B. El-Bennich, Y.-X. Liu, C. D. Roberts, and P. C. Tandy, “Collective perspective on advances in Dyson-Schwinger Equation QCD,” *Commun. Theor. Phys.* **58** (2012) 79–134, [arXiv:1201.3366 \[nucl-th\]](#).
- [2] C. D. Roberts and A. G. Williams, “Dyson-Schwinger equations and their application to hadronic physics,” *Prog. Part. Nucl. Phys.* **33** (1994) 477–575, [arXiv:hep-ph/9403224 \[hep-ph\]](#).
- [3] P. Maris and P. C. Tandy, “Bethe-Salpeter study of vector meson masses and decay constants,” *Phys. Rev.* **C60** (1999) 055214, [arXiv:nucl-th/9905056 \[nucl-th\]](#).
- [4] R. Williams, C. S. Fischer, and M. R. Pennington, “Anti-q q condensate for light quarks beyond the chiral limit,” *Phys. Lett.* **B645** (2007) 167–172, [arXiv:hep-ph/0612061 \[hep-ph\]](#).
- [5] M. S. Bhagwat, M. A. Pichowsky, C. D. Roberts, and P. C. Tandy, “Analysis of a quenched lattice QCD dressed quark propagator,” *Phys. Rev.* **C68** (2003) 015203, [arXiv:nucl-th/0304003 \[nucl-th\]](#).
- [6] I. C. Cloet, C. D. Roberts, and A. W. Thomas, “Revealing dressed-quarks via the proton’s charge distribution,” *Phys. Rev. Lett.* **111** (2013) 101803, [arXiv:1304.0855 \[nucl-th\]](#).
- [7] D. J. Wilson, I. C. Cloet, L. Chang, and C. D. Roberts, “Nucleon and Roper electromagnetic elastic and transition form factors,” *Phys. Rev.* **C85** (2012) 025205, [arXiv:1112.2212 \[nucl-th\]](#).
- [8] C. D. Roberts, “Hadron Physics and QCD: Just the Basic Facts,” *J. Phys. Conf. Ser.* **630** no. 1, (2015) 012051, [arXiv:1501.06581 \[nucl-th\]](#).
- [9] C. D. Roberts, “Running Masses in the Nucleon and its Resonances,” *JPS Conf. Proc.* **10** (2016) 010012, [arXiv:1509.08952 \[nucl-th\]](#).
- [10] M. R. Pennington, “Strong Coupling QCD in the Continuum,” *PoS EFT09* (2009) 047, [arXiv:0906.1067 \[hep-ph\]](#).
- [11] A. Kizilersu, T. Sizer, and A. G. Williams, “Strongly-Coupled Unquenched QED4 Propagators

- Using Schwinger-Dyson Equations,” *Phys. Rev.* **D88** (2013) 045008, [arXiv:1310.3160](https://arxiv.org/abs/1310.3160) [hep-ph].
- [12] R. Williams, *Schwinger-Dyson equations in QED and QCD: The Calculation of fermion-antifermion condensates*. PhD thesis, Durham U., IPPP, 2007. <http://www.ippp.dur.ac.uk/export/sites/IPPP/Research/Theses/rwilliams.pdf>.
- [13] B. Zumino, “Gauge properties of propagators in quantum electrodynamics,” *J.Math.Phys.* **1** (1960) 1–7.
- [14] S. Jia and M. R. Pennington, “Gauge covariance of the fermion Schwinger-Dyson equation in QED,” *to appear* (2016), [arXiv:1610.06001](https://arxiv.org/abs/1610.06001) [nucl-th].
- [15] A. Salam, “Renormalizable Electrodynamics of Vector Mesons,” *Phys. Rev.* **130** (May, 1963) 1287–1290. <http://link.aps.org/doi/10.1103/PhysRev.130.1287>.
- [16] A. Salam and R. Delbourgo, “Renormalizable electrodynamics of scalar and vector mesons. ii,” *Phys. Rev.* **135** (Sep, 1964) B1398–B1427. <http://link.aps.org/doi/10.1103/PhysRev.135.B1398>.
- [17] J. Strathdee, “Modified Perturbation Method for Spinor Electrodynamics,” *Phys. Rev.* **135** (Sep, 1964) B1428–B1429. <http://link.aps.org/doi/10.1103/PhysRev.135.B1428>.
- [18] R. Delbourgo and P. C. West, “A Gauge Covariant Approximation to Quantum Electrodynamics,” *J. Phys.* **A10** (1977) 1049.
- [19] D. C. Curtis and M. R. Pennington, “Nonperturbative study of the fermion propagator in quenched QED in covariant gauges using a renormalizable truncation of the Schwinger-Dyson equation,” *Phys. Rev.* **D48** (1993) 4933–4939.
- [20] A. Kizilersü, T. Sizer, M. R. Pennington, A. G. Williams, and R. Williams, “Dynamical mass generation in unquenched QED using the Dyson-Schwinger equations,” *Phys. Rev.* **D91** no. 6, (2015) 065015, [arXiv:1409.5979](https://arxiv.org/abs/1409.5979) [hep-ph].
- [21] R. Fukuda and T. Kugo, “Schwinger-Dyson Equation for Massless Vector Theory and Absence of Fermion Pole,” *Nucl. Phys.* **B117** (1976) 250–264.
- [22] D. Atkinson, J. C. R. Bloch, V. P. Gusynin, M. R. Pennington, and M. Reenders, “Strong QED with weak gauge dependence: Critical coupling and anomalous dimension,” *Phys. Lett.* **B329** (1994) 117–122.
- [23] D. C. Curtis and M. R. Pennington, “Generating fermion mass in quenched QED in four-dimensions,” *Phys. Rev.* **D46** (1992) 2663–2667.
- [24] A. Kizilersu, A. W. Schreiber, and A. G. Williams, “Regularization independent studies of nonperturbative field theory,” *Phys. Lett.* **B499** (2001) 261–269, [arXiv:hep-th/0010161](https://arxiv.org/abs/hep-th/0010161) [hep-th].
- [25] A. Kizilersu, T. Sizer, and A. G. Williams, “Regularization independent study of renormalized nonperturbative quenched QED,” *Phys. Rev.* **D65** (2002) 085020, [arXiv:hep-ph/0101188](https://arxiv.org/abs/hep-ph/0101188) [hep-ph].
- [26] J. C. R. Bloch, *Numerical investigation of fermion mass generation in QED*. PhD thesis, Durham U., 1995. [arXiv:hep-ph/0208074](https://arxiv.org/abs/hep-ph/0208074) [hep-ph]. <http://alice.cern.ch/format/showfull?sysnb=2333793>.
- [27] A. Bashir and M. R. Pennington, “Gauge independent chiral symmetry breaking in quenched QED,” *Phys. Rev.* **D50** (1994) 7679–7689, [arXiv:hep-ph/9407350](https://arxiv.org/abs/hep-ph/9407350) [hep-ph].
- [28] A. Bashir, R. Bermudez, L. Chang, and C. D. Roberts, “Dynamical chiral symmetry breaking and the fermion–gauge-boson vertex,” *Phys. Rev.* **C85** (2012) 045205, [arXiv:1112.4847](https://arxiv.org/abs/1112.4847) [nucl-th].
- [29] K.-I. Kondo, “Phase Structure of Unquenched QED With Four Fermion Interaction,” *Nucl. Phys.* **B351** (1991) 259–276.
- [30] K.-i. Kondo and H. Nakatani, “Strong coupling unquenched QED. 2. Numerical study,” *Prog. Theor. Phys.* **88** (1992) 737–750.
- [31] A. Bashir, C. Calcano-Roldan, L. X. Gutierrez-Guerrero, and M. E. Tejeda-Yeomans, “Critical Number of Flavours in QED,” *Phys. Rev.* **D83** (2011) 033003, [arXiv:1101.5458](https://arxiv.org/abs/1101.5458) [hep-ph].
- [32] P. Maris, “Analytic structure of the full fermion propagator in quenched and unquenched QED,” *Phys. Rev.* **D50** (1994) 4189–4193.
- [33] P. Maris and H. A. Holties, “Determination of the singularities of the Dyson-Schwinger equation for the quark propagator,” *Int. J. Mod. Phys.* **A7** (1992) 5369–5386.
- [34] D. Atkinson, P. W. Johnson, and P. Maris, “Dynamical Mass Generation in QED in Three-dimensions: Improved Vertex Function,” *Phys. Rev.* **D42** (1990) 602–609.
- [35] K. Kusaka, K. M. Simpson, and A. G. Williams, “Solving the Bethe-Salpeter equation for bound states of scalar theories in Minkowski space,” *Phys. Rev.* **D56** (1997) 5071–5085, [arXiv:hep-ph/9705298](https://arxiv.org/abs/hep-ph/9705298) [hep-ph].
- [36] V. Sauli and J. Adam, Jr., “Study of relativistic bound states for scalar theories in the Bethe-Salpeter and Dyson-Schwinger formalism,” *Phys. Rev.* **D67** (2003) 085007, [arXiv:hep-ph/0111433](https://arxiv.org/abs/hep-ph/0111433) [hep-ph].
- [37] C. Gutierrez, V. Gigante, T. Frederico, G. Salmè, M. Viviani, and L. Tomio, “Bethe-Salpeter bound-state structure in Minkowski space,” *Phys. Lett.* **B759** (2016) 131–137, [arXiv:1605.08837](https://arxiv.org/abs/1605.08837) [hep-ph].
- [38] M. Bhagwat, M. A. Pichowsky, and P. C. Tandy, “Confinement phenomenology in the Bethe-Salpeter equation,” *Phys. Rev.* **D67** (2003) 054019, [arXiv:hep-ph/0212276](https://arxiv.org/abs/hep-ph/0212276) [hep-ph].
- [39] K. Raya, L. Chang, A. Bashir, J. J. Cobos-Martinez, L. X. Gutiérrez-Guerrero, C. D. Roberts, and P. C. Tandy, “Structure of the neutral pion and its electromagnetic transition form factor,” *Phys. Rev.* **D93** no. 7, (2016) 074017, [arXiv:1510.02799](https://arxiv.org/abs/1510.02799) [nucl-th].
- [40] L. Chang, I. C. Cloet, J. J. Cobos-Martinez, C. D. Roberts, S. M. Schmidt, and P. C. Tandy, “Imaging dynamical chiral symmetry breaking: pion wave function on the light front,” *Phys. Rev. Lett.* **110** no. 13, (2013) 132001, [arXiv:1301.0324](https://arxiv.org/abs/1301.0324) [nucl-th].
- [41] J. C. Ward, “An identity in quantum electrodynamics,” *Phys. Rev.* **78** (Apr, 1950) 182–182. <http://link.aps.org/doi/10.1103/PhysRev.78.182>.
- [42] D. C. Curtis and M. R. Pennington, “Truncating the Schwinger-Dyson equations: How multiplicative renormalizability and the Ward identity restrict the three point vertex in QED,” *Phys. Rev.* **D42** (1990) 4165–4169.
- [43] A. Kizilersu and M. R. Pennington, “Building the Full Fermion-Photon Vertex of QED by Imposing Multiplicative Renormalizability of the Schwinger-Dyson Equations for the Fermion and Photon Propagators,” *Phys. Rev.* **D79** (2009) 125020, [arXiv:0904.3483](https://arxiv.org/abs/0904.3483) [hep-th].

- [44] A. Bashir and A. Raya, “Gauge symmetry and its implications for the Schwinger-Dyson equations,” *arXiv preprint* (2004) , [arXiv:hep-ph/0411310](https://arxiv.org/abs/hep-ph/0411310) [[hep-ph](#)].
- [45] H.-x. He, “Identical relations among transverse parts of variant Green functions and the full vertices in gauge theories,” *Phys. Rev.* **C63** (2001) 025207, [arXiv:hep-ph/9910373](https://arxiv.org/abs/hep-ph/9910373) [[hep-ph](#)].
- [46] H.-X. He, F. C. Khanna, and Y. Takahashi, “Transverse Ward-Takahashi identity for the fermion boson vertex in gauge theories,” *Phys. Lett.* **B480** (2000) 222–228.
- [47] H.-X. He, “Quantum anomaly of the transverse Ward-Takahashi relation for the axial-vector vertex,” *Phys. Lett.* **B507** (2001) 351–355, [arXiv:hep-th/0201217](https://arxiv.org/abs/hep-th/0201217) [[hep-th](#)].
- [48] H.-X. He, “Transverse vector vertex function and transverse Ward-Takahashi relations in QED,” *Commun. Theor. Phys.* **46** (2006) 109–112.
- [49] S. Jia and M. R. Pennington, “Landau-Khalatnikov-Fradkin transformation for the fermion propagator in QED in arbitrary dimensions,” *to appear* (2016) , [arXiv:1610.06436](https://arxiv.org/abs/1610.06436) [[hep-th](#)].
- [50] P. Lizorkin, *Fractional integration and differentiation*. Encyclopaedia of Mathematics. Springer, 1992.
- [51] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions: With Formulas, Graphs, and Mathematical Tables*. Applied mathematics series. Dover Publications, 1964. <https://books.google.com/books?id=MtU8uP7XMvoC>.
- [52] R. Delbourgo, B. Keck, and C. Parker, “Gauge Covariance and the Gauge Technique,” *J.Phys.* **A14** (1981) 921.