



W&M ScholarWorks

Arts & Sciences Articles

Arts and Sciences

2015

Covariant spectator theory of np scattering: Deuteron quadrupole moment

Franz Gross

Franz Gross

College of William and Mary

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Gross, F. (2014). Covariant spectator theory of np scattering: Deuteron quadrupole moment. arXiv preprint arXiv:1411.7076.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



Covariant spectator theory of np scattering: Deuteron quadrupole moment

Franz Gross

*Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA
and College of William and Mary, Williamsburg, Virginia 23185, USA*

(Received 30 November 2014; published 26 January 2015)

The deuteron quadrupole moment is calculated using two covariant spectator theory model wave functions obtained from the 2007 high-precision fits to np scattering data. Included in the calculation are a new class of isoscalar np interaction currents automatically generated by the nuclear force model used in these fits. The prediction for model WJC-1, with larger relativistic P -state components, is 2.5% smaller than the experimental result, in common with the inability of models prior to 2014 to predict this important quantity. However, model WJC-2, with very small P -state components, gives agreement to better than 1%, similar to the results obtained recently from chiral effective field theory predictions to next-to-next-to-next-to-leading order.

DOI: [10.1103/PhysRevC.91.014005](https://doi.org/10.1103/PhysRevC.91.014005)

PACS number(s): 13.40.Em, 03.65.Pm, 13.75.Cs, 21.45.Bc

I. INTRODUCTION AND BACKGROUND

Until recently, calculations of the deuteron quadrupole moment consistently underpredicted its value by several percent [1–4]. Some of these calculations are summarized in Table I; all of the results shown use realistic nucleon-nucleon (NN) scattering models with kernels or potentials adjusted to fit the low-energy NN data. Because of these fits, predictions of the quadrupole moment are very tightly constrained, with uncertainties coming only from relativistic corrections, including those to the current operator, which have been difficult to determine. The difficulty of avoiding these constraints led Machleidt [3] to identify the underprediction of the quadrupole moment as an “unresolved problem.”

Now, a new chiral effective field theory (χ EFT) calculation, done to next-to-next-to-next-to-leading order (N^3 LO) by the ODU-Pisa group [5], has obtained very good agreement. To this order the three-vector current contains two unknown isoscalar low-energy constants (LECs), but the charge operator from which the quadrupole moment is calculated contains no LEC and is uniquely determined. (At higher order the charge operator contains an LEC that can be used to fix the quadrupole moment [6].) However, all of their results still depend on the cutoff Λ , which is needed to renormalize the calculations, and the dependence of their result for the quadrupole moment on Λ (for the two values of 500 and 600 MeV studied) is shown in Table I. The dependence on Λ is not strong, and both results are much closer to the experimental result than was found previously.

The principal purpose of this paper is to report the new results for the quadrupole moment obtained from the two high-precision models of NN scattering (WJC-1 and WJC-2) that Stadler and I found in 2008 [7] using covariant spectator theory (CST) [8–10]. (The features and differences between these two models will be briefly reviewed in Sec. III.) The predictions obtained for these two models are shown in the last line of Table I. The physics that went into these calculations of the quadrupole moment (which contain *no free parameters*) will be very briefly summarized in Sec. II, with all of the extensive details moved to the Appendix. The implications of these results are discussed in Sec. III.

This paper is the third in a series of four planned papers. These papers grew out of the need for a new treatment of the NN current required by the nature of the kernels used in the high-precision fits of 2008. At that time it was found that a kernel consisting of a sum of covariant one-boson exchange (OBE) diagrams would give an excellent high-precision fit to the NN data (with a $\chi^2/\text{datum} \simeq 1$) provided that the vertex function, Λ^{σ_0} , that describes the coupling of the scalar-isoscalar boson σ_0 to the nucleon, included momentum-dependent terms of the form

$$\Lambda^{\sigma_0}(p, p') = g_{\sigma_0} \mathbf{1} - v_{\sigma_0} [\Theta(p) + \Theta(p')], \quad (1.1)$$

where g_s and v_s are parameters adjusted to fit the data, p and p' are the four-momenta of the outgoing and incoming nucleons, respectively, and the operator Θ is the negative-energy projection operator for a spin 1/2 nucleon,

$$\Theta(p) = \frac{m - \not{p}}{2m}. \quad (1.2)$$

For more discussion of these OBE models, see Ref. [7].

These v -dependent terms, which vanish when the nucleons are on-shell, introduce a new kind of energy dependence into the kernel, generating a new class of isoscalar interaction currents. The first paper in this series, referred to as Ref. I [11], showed how current conservation [12] and the principles of picture independence and simplicity could be used to *uniquely* determine these interaction currents. Then, in the second paper, referred to as Ref. II [13], I calculated the deuteron magnetic moment and showed that both high-precision models gave a nearly identical prediction that is only about 1% larger than the experimental value. The magnetic moment cannot distinguish between the two models. However, the predictions for the quadrupole moment shown in Table I provide a basis for distinguishing between the two models and this will be discussed in Sec. III.

II. SUMMARY OF THE CALCULATION

In CST, the two-body current is given by the five diagrams shown in Fig. 1 (also shown in Ref. II). These include the interaction current contributions derived in Ref. I, expressed

TABLE I. Predictions of the quadrupole moment, expressed as an “error” defined by $\delta Q_{\text{pred}} = (Q_{\text{pred}} - Q_{\text{exp}})/Q_{\text{exp}}$.

Reference	δQ_{pred} (model)
GVOH [1]	-9.0% (IIB), -8.1% (IIB with RC)
Argonne [2]	-3.8% (with MEC)
CD Bonn [3]	-5.6% (no MEC), -2.1% (MEC est.)
Light Front [4]	-5.7% (IM), -3.8% (IM + Ex)
χ EFT (ODU-Pisa) [5]	-0.3% (500), -1.4% (600)
this work	-2.5% (WJC-1), -0.8% (WJC-2)

in terms of the the effective wave functions $\Psi^{(2)}$ and the subtracted vertex functions $\hat{\Gamma}$ (directly related to $\hat{\Psi}$) with two particles off-shell. These contributions are discussed below, but for a complete discussion of the physics, see Refs. I and II.

The quadrupole moment, Q_d , in units of e/m_d^2 , is extracted by taking the $Q^2 \rightarrow 0$ limit of the difference of two matrix elements of the two body current, \mathcal{J}_i ,

$$Q_d = \lim_{Q^2 \rightarrow 0} \frac{m_d}{Q^2} [\mathcal{J}_1 - \mathcal{J}_2], \quad (2.1)$$

where the current matrix elements are

$$\begin{aligned} \mathcal{J}_1 &\equiv G_{00}^0 = 2D_0 (G_C + \frac{4}{3}\eta G_Q), \\ \mathcal{J}_2 &\equiv G_{+-}^0 = 2D_0 (G_C - \frac{2}{3}\eta G_Q), \end{aligned} \quad (2.2)$$

with $G_{\lambda\lambda'}^{\lambda\gamma}$ the matrix element for an incoming (outgoing) deuteron with four-moments $P_-(P_+)$ and helicity $\lambda'(\lambda)$ and a virtual photon with helicity λ_γ ,

$$G_{\lambda\lambda'}^{\lambda\gamma} \equiv \langle P_+ \lambda | J_\mu | P_- \lambda' \rangle \epsilon_{\lambda_\gamma}^\mu. \quad (2.3)$$

TABLE II. Physical origin of the eight different types of terms that contribute to the quadrupole moment.

Term	Physical origin
Q_{NR}	nonrelativistic contribution from the S and D states
Q_{Rc}	relativistic corrections to S and D terms
$Q_{h'}$	dependence on the strong form factor, h
Q_{V_2}	interaction currents: off-shell particle 2
Q_{V_1}	interaction currents: on-shell particle 1
Q_{int}	S -, D -, and P -state interference
Q_P	P -state squared terms
Q_χ	P -state and negative ρ -spin z_ℓ^- interference

Equation (2.2) has been evaluated in the Breit frame, where the photon four-momentum is $q = \{0, \mathbf{q}\}$, and $Q^2 = \mathbf{q}^2$, $P_\mp = (D_0, \mp \frac{1}{2}\mathbf{q})$, and $D_0 = \sqrt{m_d^2 + Q^2}/4$. Details can be found in Ref. II.

The calculation of the quadrupole moment is described in the Appendix. The final result can be arranged into a sum of the eight terms summarized in Tables II and III and given explicitly in Eq. (A79). To understand the origin of these terms, recall that the relativistic deuteron wave function with one particle on-shell (and the other off-shell) can be expanded in terms of four relativistic wave functions: u (S state), w (D state), v_t (a P -state wave function with spin triplet structure), and v_s (a P -state wave function with a spin singlet structure) [14,15]. When both particles are off-shell, an additional four wave functions could contribute, but only one combination, the z_s defined in Eq. (A64), contributes in leading order. The eight terms can now be described.




FIG. 1. (Color online) Diagrammatic representation of the two-body current operator in the covariant spectator theory using vertex functions with particle 2 off-shell. The interaction current contributions are contained in diagrams (A $_{\pm}$) and parts of the (B) diagrams, as discussed in the text. Off-shell nucleon lines are thicker than on-shell lines, which are also labeled with an \times . Diagrams (A) and (A $_{\pm}$) describe the interaction of the photon with particle 2, allowing particle 1 to be on-shell in both the initial and the final states. Diagrams (B $_{\pm}$) describe the interaction of the photon with particle 1, so that both particles must off-shell in either the initial state (diagram B $_{+}$) or in the final state (diagram B $_{-}$).

TABLE III. Contributions to the quadrupole moment from the eight different types of corrections discussed in the text. All terms are normalized by the experimental value of the quadrupole moment ($Q_{\text{exp}} = 0.286$), with $Q_{\text{NR}}^{\Delta} = (Q_{\text{NR}} - Q_{\text{exp}})/Q_{\text{exp}}$, so that all of these terms must sum to zero to get the correct experimental value.

	WJC-1		WJC-2	
	u, w only	all	u, w only	all
Q_{NR}^{Δ}	-0.011	-0.011	-0.018	-0.018
Q_{Rc}	0.010	0.010	0.010	0.010
$Q_{h'}$	0.001	0.001	0.001	0.001
Q_{V_2}	-0.004	-0.004	-0.001	-0.001
Q_{V_1}	-0.004	-0.003	-0.001	-0.002
Q_{int}	-	-0.014	-	0.002
Q_P	-	-0.002	-	0.000
Q_{χ}	-	-0.002	-	-0.000
total	-0.008	-0.025	-0.009	-0.008

The largest contribution, Q_{NR} , is familiar from the first days of nuclear physics [16]:

$$Q_{\text{NR}} = \frac{\sqrt{2}}{10} \int_0^{\infty} r^2 dr \left\{ uw - \frac{w^2}{\sqrt{8}} \right\}. \quad (2.4)$$

However, while the same formula (2.4) arises in both the nonrelativistic theory and (as the leading contribution) in CST the two results are numerically very different because the normalization of the u and w wave functions in the two cases is very different. In the nonrelativistic theory, the normalization is

$$\int_0^{\infty} k^2 dk (u^2 + w^2) = 1, \quad (2.5)$$

while in CST it is

$$\int_0^{\infty} k^2 dk (u^2 + w^2) = 1 + N_{\text{CST}}, \quad (2.6)$$

where

$$N_{\text{CST}} = - \left\langle \frac{\partial V}{\partial m_d} \right\rangle - \int_0^{\infty} k^2 dk (v_t^2 + v_s^2) \quad (2.7)$$

with V the NN kernel, including the strong nucleon form factors h , and the derivative with respect to the deuteron mass (or, alternatively, the total energy in the deuteron rest system) is a consequence of the interaction current, as discussed in Ref. II. The contributions to N_{CST} , discussed in detail in Ref. II, are summarized in Table IV.

Hence the Q_{NR} of Eq. (2.4) is larger than the nonrelativistic result by a factor of N_{CST} but this correction is “hidden” in the sense that it is already included in the leading term Q_{NR}^{Δ} given in Table III. One may infer from Q_{NR}^{Δ} that using the (incorrect) nonrelativistic normalization would give a result for the quadrupole moment about 6% too small for WJC-1 and 4% too small for WJC-2.

While the relativistic normalization (2.6) makes a significant contribution, the calculation is not complete and the result is not believable until all of the other effects that come from the relativistic structure of the interaction current and the deuteron

TABLE IV. Contributions to the normalization factor N_{CST} from the four different types of corrections discussed in the text (extracted from Tables I and II of Ref. II).

	WJC-1		WJC-2	
	u, w only	all	u, w only	all
$N_{h'}$	-0.036	-0.025	-0.018	-0.018
N_{V_2}	0.022	0.023	0.011	0.011
N_{V_1}	0.052	0.057	0.032	0.030
N_P	-	-0.003	-	0.000
total	0.038	0.052	0.025	0.023

wave functions are also calculated. Each of these remaining effects, in the order listed in Table II, will be discussed briefly.

As in Ref. II, only the leading contributions to these corrections (those believed to be larger than 0.001) are retained. A detailed discussion of which terms can be expected to be “leading” was presented in Ref. II, and the same guidelines are followed here.

The Q_{Rc} term includes the corrections of order k^2/m^2 coming from the expansion of the relativistic kinetic energy, $E_k = \sqrt{m^2 + \mathbf{k}^2}$, which appears in many places through the calculation. Only corrections to products involving the largest wave functions (u and w) are leading. This kinematical relativistic correction is one of the largest effects, and it is of comparable size for both models.

The $Q_{h'}$ and $N_{h'}$ terms include corrections to the quadrupole moment that come from the strong nucleon form factor $h(p)$. This form factor is a function of p^2 , the four-momentum of the off-shell nucleon (only), and is normalized to unity when $p^2 = m^2$. As shown in Eq. (2.1), the calculation of the quadrupole moment requires expanding the electromagnetic form factors around $Q^2 = 0$, requiring that the strong form factor be expanded around its mass-shell point, introducing correction terms proportional to $a(p^2) = d \log(h)/dp^2|_{p^2=m^2}$. As shown in Table IV, terms of this type make about a -2% contribution to the relativistic normalization already included in the leading Q_{NR}^{Δ} ; the *additional* corrections to the quadrupole moment contained in $Q_{h'}$ turn out to be negligible.

The Q_{V_2} and N_{V_2} terms include contributions from the isoscalar exchange current generated by the momentum dependence included in the projection operators Θ [defined in Eq. (1.2)] that operate on the off-shell particle 2 [illustrated in diagrams (A $_{\pm}$) shown in Fig. 1]. Terms of this type are present in the vertex functions for the exchange of all mesons (except the axial-vectors present in model WJC-1), but the contributions from the pseudoscalar exchanges (π and η) cancel. The way in which Θ appears in the sNN vertex functions for scalar (s) exchange was already illustrated in Eq. (1.1). The structure of the exchange current implied by the appearance of these operators Θ was uniquely determined in Ref. I, where it was shown how their contributions can be expressed in terms of new deuteron wave functions generically denoted by $z^{(2)}$. As shown in Table IV, terms of this type are already included in Q_{NR}^{Δ} , where they make about a 2% (1%) contribution for models WJC-1 (WJC-2); the *additional* corrections shown in Table III are much smaller.




FIG. 2. (Color online) Running sum of the corrections (in %) to the quadrupole moment, in the order that they are listed in Tables II and III. Panel (a) is model WJC-1 and panel (b) is model WJC-2. The dashed line is the experimental value (zero correction). The error bars are $\pm 0.002 = \pm 0.2\%$, an estimate of the size of the terms missing from the approximation of Eq. (A79).

The Q_{V_1} and N_{V_1} terms include contributions from that part of the isoscalar interaction current that contributes when (the usually on-shell) particle 1 is forced off-shell by the kinematics. Explicitly, in diagram (B₊) of Fig. 1, particle 1 has four-momentum k_- before the interaction, while in diagram (B₋) it has four-momentum k_+ after the interaction, where $k_\pm = k \pm q$ with $k = \{E_k, \mathbf{k}\}$. Therefore, in both cases particle 1 is off-shell unless $q = 0$, so that, as one makes the expansion (2.1) needed to calculate the quadrupole moment, the behavior of the vertex function when both particles are off-shell is probed. However, even if there were no interaction current, there would still be contributions of this type from the vertex function itself. It turns out that the interaction current cancels some of these contributions, and this subtracted vertex function is denoted by $\widehat{\Gamma}_{BS}$. It depends on wave functions generically denoted by \widehat{z} . I have made no attempt to separate the contributions of interaction current from that of the vertex function itself, so these contributions include both effects. The contribution of these terms to the normalization (Table IV) gives a large contribution of almost 6% (3%) to the quadrupole moment from WJC-1 (WJC-2), and the additional contributions from Q_{V_1} is about 1/10 as large.

The Q_{int} interference term includes contributions from the product of the w and v_t wave functions not contained in the other terms. Note that it makes a large contribution of almost -1.5% to the quadrupole moment for WJC-1 and a very small contribution for WJC-2. This term is largely the cause of the small WJC-1 result.

The Q_P and N_P terms include contributions from the square of the P states and are quite small for both models.

Finally, the interesting Q_χ term is the interference between the v_s P state and the combination of negative energy helicity states z_s . It is quite small in both models, but for WJC-1 it is larger than the estimated theoretical error of 0.001, and it is therefore included.

Looking at the cumulative totals shown in Fig. 2, one concludes that the result for model WJC-2 is quite close to the experimental value and well given by the

normalization correction, N_{CST} , alone. The case for model WJC-1 is quite different however; here the additional corrections shown in Table III reduce the quadrupole moment to an unacceptably low value, due largely to the single term Q_{int} . I discuss the significance of these results in the next section.

III. CONCLUSIONS AND OUTLOOK

In this paper I presented an approximate calculation (accurate to about 0.1%) of the deuteron quadrupole moment for two recent models that both give a high-precision fit ($\chi^2/\text{datum} \simeq 1$) to the 2007 np database below 350 MeV laboratory energy. Model WJC-1, designed to give the best fit possible, has 27 parameters, $\chi^2/\text{datum} \simeq 1.06$, and a large $v_{\sigma_0} = -15.2$. Model WJC-2, designed to give an excellent fit with as few parameters as possible, has only 15 parameters, $\chi^2/\text{datum} \simeq 1.12$, and a smaller $v_{\sigma_0} = -2.6$. Both models also predict the correct triton binding energy [7,17] and give the same magnetic moment (with an uncertainty of 0.001) about 1% larger than the experimental value.

Until now, the major distinction between these two models has been their deuteron momentum distributions. Model WJC-1 gives a much harder distribution than WJC-2 [18] and other models [18–20], but since the momentum distribution is not an observable, it may be inappropriate to use this as a means of distinguishing between them. The prediction of the quadrupole moment presented in this paper clearly favors WJC-2. The simplicity of model WJC-2, with only 15 parameters and a pure pseudo vector πNN coupling, might also favor WJC-2, even though the χ^2 of the fit to the np database is very slightly larger than that of WJC-1 (1.12 versus 1.06). Perhaps a calculation of the form factors, planned for the last paper in this series, will be definitive.

How close can one expect the agreement to be between experimental data and CST? Perhaps agreement to about 1% should be expected if the theory is to be taken seriously, and (in agreement with Machleidt [3]) I take the error of -2.5%

in the WJC-1 prediction to be a serious problem. On the other hand, should the error of -0.8% in the WJC-2 prediction be accepted? One answer is that the χ EFT prediction is comparable, and the claim is that this is a theory and not just a model. If an exact prediction is wanted, recall that the deuteron binding energy and the 1S_0 scattering lengths were already constrained when fitting the np database [7], so perhaps the deuteron quadrupole moment could also be constrained at the same time. Since model WJC-2 agrees so closely without this constraint, perhaps it could be included without

seriously degrading the χ^2 . These possibilities await future study.

ACKNOWLEDGMENTS

It is a pleasure to thank Rocco Schiavilla for stimulating conversations about the χ EFT predictions. This material is based upon work supported by the Jefferson Science Associates, LLC, under US DOE Contract No. DE-AC05-06OR23177.

APPENDIX: DETAILS OF THE CALCULATION

For any quantity not defined in the discussion below, refer to Ref. II.

1. Diagrams (A) and (A_{\pm})

a. Exact expressions

The quadrupole form factor, G_Q , is obtained directly from the difference between \mathcal{J}_1 and \mathcal{J}_2 . By using Eq. (2.2) and the results from Ref. II, this is

$$4D_0\eta G_Q(Q^2)|_{A+A_{\pm}} = e_0 F_1(Q^2) \int_k \left\{ f_0(p_+, p_-) \delta \mathcal{A}_1(k, Q) - \frac{h_+}{h_-} \delta \mathcal{A}_1^{(2)}(k, Q) - \frac{h_-}{h_+} \delta \mathcal{A}_1^{(2)}(k, -Q) \right\} \\ + e_0 F_2(Q^2) \int_k \left\{ f_0(p_+, p_-) \delta \mathcal{A}_2(k, Q) - \frac{h_+}{h_-} \delta \mathcal{A}_2^{(2)}(k, Q) + \frac{h_-}{h_+} \delta \mathcal{A}_2^{(2)}(k, -Q) \right\} \\ + e_0 F_3(Q^2) \int_k \frac{g_0(p_+, p_-)}{4m^2} \delta \mathcal{A}_3(k, Q), \quad (\text{A1})$$

where $\delta \mathcal{A}_i$ are differences of the traces $\mathcal{A}_{n,i}$ defined in Ref. II,

$$\delta \mathcal{A}_i(k, Q) \equiv \mathcal{A}_{1,i}(\Psi_+, \Psi_-) - \mathcal{A}_{2,i}(\Psi_+, \Psi_-), \quad \delta \mathcal{A}_i^{(2)}(k, Q) \equiv \mathcal{A}_{1,i}(\Psi_+, \Psi_-^{(2)}) - \mathcal{A}_{2,i}(\Psi_+, \Psi_-^{(2)}). \quad (\text{A2})$$

By introducing the convenient averages

$$\bar{A}_i(k) = \lim_{Q^2 \rightarrow 0} \frac{m_d}{Q^2} \frac{1}{2} \int_{-1}^1 dz \delta \mathcal{A}_i(k, Q), \quad \bar{A}_{i\pm}^{(2)}(k) = \lim_{Q^2 \rightarrow 0} \frac{m_d}{Q^2} \frac{1}{2} \int_{-1}^1 dz \delta \mathcal{A}_i^{(2)}(k, \pm Q), \quad (\text{A3})$$

the contributions of diagrams A and A_{\pm} to the quadrupole moment can be written as

$$Q_A = e_0 \int \frac{k^2 dk}{2\pi^2} \frac{m}{E_k} \{ f_{00} \bar{A}_1(k) - \bar{A}_{1+}^{(2)}(k) - \bar{A}_{1-}^{(2)}(k) \} + e_0 \kappa_s \int \frac{k^2 dk}{2\pi^2} \frac{m}{E_k} \{ f_{00} \bar{A}_2(k) - \bar{A}_{2+}^{(2)}(k) - \bar{A}_{2-}^{(2)}(k) \} \\ + e_0 \int \frac{k^2 dk}{2\pi^2} \frac{m}{E_k} \frac{g_{00}}{4m^2} \bar{A}_3(k), \quad (\text{A4})$$

where f_{00} and g_{00} are coefficients of the off-shell nucleon current defined in Eq. (3.24) of Ref. II.

To work out the limits (A3), expand the differences (A2) to order Q^2 . Making the approximation $m_d \simeq 2m$ gives

$$\delta \mathcal{A}_1(k, Q) = \frac{2k^2}{m^3} P_2(z) \{ B_+ B_- E_k - 4C_+ C_- (2m - E_k) + 4D_+ D_- E_k + 2(A_+ D_- + D_+ A_-) m \\ - 2(B_+ C_- + C_+ B_-)(E_k - m) - 2(B_+ D_- + D_+ B_-) m \} \\ + \frac{k_z Q}{2m^3} \{ [A_+(B_- - 2C_-) - (B_+ - 2C_+)A_-] m + 2(A_+ D_- - D_+ A_-) E_k \\ - 2(B_+ C_- - C_+ B_-)(E_k - m) - 4(C_+ D_- - D_+ C_-)(E_k - m) \} + \frac{Q^2}{2m^3} E_k \left\{ C^2 \left(1 - \frac{2k^2}{3m^2} \right) \right. \\ \left. - \left(\frac{1}{4} B^2 + D^2 \right) \left(1 + \frac{2k^2}{3m^2} \right) + \frac{1}{2} AB - AC + BC \frac{2k^2}{3m^2} + 2CD \left(1 - \frac{m}{E_k} - \frac{2k^2}{3E_k m} \right) \right\},$$

$$\begin{aligned} \delta A_2(k, Q) = & \frac{k_z Q}{m^3} \left\{ (A_+ B_- - B_+ A_-) E_k - 2(A_+ C_- - C_+ A_-)(E_k - 2m) - 2(A_+ D_- - D_+ A_-) m \right. \\ & \left. + 2(B_+ C_- - C_+ B_-)(E_k - m) - 2(B_+ D_- - D_+ B_-) \frac{k^2}{m} P_2(z) - 4(C_+ D_- - D_+ C_-) \left[E_k - m - \frac{k^2}{m} P_2(z) \right] \right\} \\ & - \frac{Q^2}{2m^2} \left\{ A^2 - (4C^2 - 2BC - 4CD) \left(1 - \frac{E_k}{m} + \frac{2k^2}{3m^2} \right) - AB \left(1 + \frac{2k^2}{3m^2} \right) - 2AC \left(1 - \frac{2k^2}{3m^2} \right) + 2AD \frac{E_k}{m} \right\}, \end{aligned} \quad (\text{A5})$$

where $P_2(z)$ is the $\ell = 2$ Legendre polynomial with $z = k_z/k$ the cosine of the polar angle, and $Z_{\pm} = Z(R_{\pm})$ (where Z is a generic name for the $A, B, C,$ or D invariants defined in Ref. II) and R_{\pm} is the covariant generalization of the magnitude of the rest frame three-momentum $|\mathbf{k}|$ for the outgoing (R_+) and incoming (R_-) deuteron states. From Ref. II, these arguments, expanded to order Q^2 , are

$$R_{\pm} \simeq \left[\mathbf{k}^2 \mp \frac{k_z Q E_k}{m_d} + \eta(E_k^2 + k_z^2) \right]^{1/2} \simeq k \mp z Q \frac{E_k}{2m_d} + \frac{\eta}{2k} (E_k^2 - m^2 z^2), \quad (\text{A6})$$

where now $|\mathbf{k}| \rightarrow k$. In calculating the average $\bar{A}_1(k)$, the first term will get contributions of order Q^2 from the expansions of the wave functions, but only terms proportional to $z^2 P_2(z)$ will survive. Hence, for arbitrary $\{X, Y\} = \{A, B, C, D\}$, the expansion needed is

$$X_+ Y_- = X(R_+) Y(R_-) \rightarrow -z^2 \eta \left\{ (X'Y + XY') \frac{m^2}{2k} + X'Y' E_k^2 - \frac{1}{2} (X''Y + XY'') E_k^2 \right\}, \quad (\text{A7})$$

where $X = X(k)$, $X' = dX(k)/dk$, etc. Only derivative terms contribute to the terms proportional to $k_x Q$, and for these one needs

$$X_+ Y_- \rightarrow -\frac{z Q E_k}{2m_d} (X'Y - XY'). \quad (\text{A8})$$

Making these substitutions and continuing to let $m_d \rightarrow 2m$ reduces the averages (A3) to

$$\begin{aligned} \bar{A}_1(k) = & -\frac{k^2 E_k^2}{30m^4} \{ (B^2 + 4D^2 - B''B - 4D''D) E_k - 2(2B'C' - B''C - C''B)(E_k - m) + 4(C'^2 - C''C)(E_k - 2m) \\ & + 2(2A'D' - A''D - D''A + B''D + D''B - 2B'D') m \} \\ & + \frac{k}{30m^3} \{ [5(AB' - A'B) - B'B + 10(A'C - AC') - 4D'D] m E_k + 2(B'D + D'B) m^2 - 4C'C(E_k - 2m) m \\ & + 20(C'D - D'C) E_k (E_k - m) - 2C'B(E_k - m)(5E_k - m) + 2B'C(E_k - m)(5E_k + m) \\ & - 2A'D(5E_k^2 + m^2) + 2D'A(5E_k^2 - m^2) \} \\ & - \frac{1}{12m^4} \{ (B^2 + 4D^2)(2E_k^2 + m^2) E_k - 4BDm(2E_k^2 + m^2) - 6(AB - 2AC) E_k m^2 - 8BCk^2 E_k \\ & + 8CDm(E_k - m)(2E_k - m) + 4C^2 E_k (2E_k^2 - 5m^2) \}, \end{aligned} \quad (\text{A9})$$

$$\begin{aligned} \bar{A}_2(k) = & \frac{4k^3 E_k}{15m^4} \{ B'D - D'B - 2C'D + 2D'C \} - \frac{k E_k}{3m^3} \{ (A'B - B'A) E_k - 2(A'D - D'A) m \\ & + 2(B'C - C'B - 2C'D + 2D'C)(E_k - m) - 2(A'C - C'A)(E_k - 2m) \} - \frac{1}{3m^3} \{ 3A^2 m^2 + 6ADE_k m \\ & - AB(2E_k^2 + m^2) + 2AC(2E_k^2 - 5m^2) + 2(BC - 2C^2 + 2CD)(E_k - m)(2E_k - m) \}, \end{aligned} \quad (\text{A10})$$

where $\bar{A}_3(k)$ is $\bar{A}_1(k)$ with $A \rightarrow F$, etc. As expected, $\bar{A}_2(k)$ includes no terms involving Z'' or Z'^2 because it is already $\mathcal{O}(Q)$ without expansions of the wave functions.

b. Leading terms in momentum space

Equations (A9) and (A10) give the exact results for the quadrupole moment, and they can be easily evaluated numerically. However, our goal here is to obtain some insight into the physical content of the result, and to this end it is sufficient to compute the quadrupole moment to an accuracy of about 0.1% as I did for the magnetic moment in Ref. II. This is done by expanding the exact results in terms of the four deuteron wave functions $z_{\ell} = \{u, w, v_l, v_s\}$ (where z_{ℓ} is the generic name for any of the wave functions, and the expansions were given in Ref. II) and retaining only the leading terms, as defined in Ref. II. These leading

terms are obtained by expanding the coefficients of the leading products of the wave functions u and w to order k^2/m^2 and expanding coefficients of all products involving P -state wave functions to order k/m . In comparing derivative terms, $z'_\ell, z'_\ell/k$, and z_ℓ/k^2 are considered to be of the same order. Pulling out an overall factor of E_k , and integrating by parts to remove all of the double derivatives and to make other simplifications, gives

$$\begin{aligned}\bar{A}_1(k) &\simeq 2\pi^2 \frac{E_k m^2}{m} \frac{1}{10} \left\{ -4\sqrt{2} u' w' - 2w'^2 + 2v_t'^2 - 4v_s'^2 - 12\sqrt{2} \frac{u' w'}{k} - \frac{1}{k^2} [12w^2 - 4v_t'^2 + 8v_s'^2] + \frac{k}{\sqrt{3}m} A_{\text{int}} + \Delta A_1 \right\}, \\ \bar{A}_2(k) &\simeq 2\pi^2 \frac{E_k m^2}{m} \frac{1}{10} \left\{ \frac{2k}{\sqrt{3}m} A_{\text{int}} + \Delta A_2 \right\}, \\ \bar{A}_3(k) &\simeq 2\pi^2 \frac{E_k}{m} \frac{4m^4}{10} \left\{ 2 \left[v_t'^2 + \frac{2v_t'^2}{k^2} \right] - 4 \left[v_s'^2 + \frac{2v_s'^2}{k^2} \right] \right\},\end{aligned}\tag{A11}$$

where the interference terms, multiplied by a factor of k/m , are

$$A_{\text{int}} = -\frac{2}{k} \left\{ 5u'(\sqrt{2}v_t + 2v_s) - 4w \left(v_t' - \frac{v_t}{k} \right) - \sqrt{2}w \left(v_s' - \frac{v_s}{k} \right) \right\}\tag{A12}$$

and ΔA is the k^2/m^2 correction to the leading terms

$$\Delta A_1 = \frac{k^2}{m^2} \left[4\sqrt{2} u' w' + 2w'^2 + \frac{6\sqrt{2} u' w'}{k} + \frac{3w^2}{k^2} \right], \quad \Delta A_2 = \frac{6k}{m^2} \left[2\sqrt{2} u' w + \frac{1}{k} (\sqrt{2} u w + w^2) \right].\tag{A13}$$

Note that the interference terms, smaller by one power of k/m , might be ignored, and, as it turns out, the contributions from the (B) diagrams contribute interference terms that are larger by a factor of m^2/k^2 . The contributions from A_{int} will therefore be ignored when the contributions from the (A) and (B) diagrams are combined.

As discussed in Ref. II, the A_{\pm} contributions to the interaction currents are obtained by substituting $\Psi^{(2)}$ wave functions for Ψ wave functions in the initial state, giving rise to the $A^{(2)}$ terms in Eq. (A1). These can be constructed directly from (A11). Since the $\Psi^{(2)}$ contributions are already small, they will be kept only to leading order, so that any contributions that might have come from A_{int} will be discarded. The δA_2 terms in Eq. (A11) can therefore be ignored. To find the $\Psi^{(2)}$ contributions $\delta A_1^{(2)}$, recall from Ref. II that the helicity traces $\mathcal{A}_{n,1}$ from which δA_1 is calculated satisfy the symmetry relation (for $n = 1, 2$)

$$\mathcal{A}_{n,1}(\Psi_1 \Psi_2) = \mathcal{A}_{n,1}(\Psi_2 \Psi_1)|_{q \rightarrow -q}.\tag{A14}$$

Note that a typical term in the expansion (A5) satisfies this symmetry and is of the form

$$\langle X_+ Y_- \rangle \rightarrow (P_2(z)c_0 + Q^2 c_2)(X_+ Y_- + Y_+ X_-) + k_z Q c_1 (X_+ Y_- - Y_+ X_-),\tag{A15}$$

where the c_i include all of the additional factors present in the expansions. Replacing the initial state by $\Psi^{(2)}$, and exploiting this symmetry, means that the typical XY contribution to the $\delta A_1^{(2)}(k, Q)$ term in (A1) becomes

$$\langle X_+ Y_-^{(2)} \rangle|_Q \rightarrow (P_2(z)c_0 + Q^2 c_2)(X_+ Y_-^{(2)} + Y_+ X_-^{(2)}) + k_z Q C_1 (X_+ Y_-^{(2)} - Y_+ X_-^{(2)}).\tag{A16}$$

Adding the second contribution in Eq. (A1), $\delta A_1^{(2)}(k, -Q)$, gives a combined result

$$\begin{aligned}\langle X_+ Y_-^{(2)} \rangle|_Q + \langle X_+ Y_-^{(2)} \rangle|_{-Q} &\rightarrow (P_2(z)c_0 + Q^2 c_2)(X_+ Y_-^{(2)} + X_+^{(2)} Y_- + Y_+ X_-^{(2)} + Y_+^{(2)} X_-) \\ &\quad + k_z Q C_1 (X_+ Y_-^{(2)} + X_+^{(2)} Y_- - Y_+ X_-^{(2)} - Y_+^{(2)} X_-),\end{aligned}\tag{A17}$$

showing that all terms are obtained by the expected substitution $XY \rightarrow XY^{(2)} + X^{(2)}Y$ where either X or Y may contain one or two derivatives. The contributions from $\Psi^{(2)}$ therefore reduce to

$$\begin{aligned}\bar{A}_{1+}^{(2)}(k) + \bar{A}_{1-}^{(2)}(k) &\simeq 2\pi^2 \frac{E_k m^2}{m} \frac{1}{10} \left\{ -4\sqrt{2} (u' w^{(2)'} + u^{(2)'} w') - 4w' w^{(2)'} + 4v_t' v_t^{(2)'} - 8v_s' v_s^{(2)'} \right. \\ &\quad \left. - \frac{12\sqrt{2}}{k} (u^{(2)'} w + u' w^{(2)}) - \frac{1}{k^2} [24w w^{(2)} - 8v_t v_t^{(2)} + 16v_s v_s^{(2)}] \right\}.\end{aligned}\tag{A18}$$

Finally, the contributions from the derivatives of the strong form factor, h , expressed in terms of $a(p)$ defined in Sec. II [and Eq. (3.25) of Ref. II], are extracted from contributions from f_{00} and g_{00} . These terms will be simplified by integrating by parts as I did for the leading contributions (A11). In doing this integration, I use the fact that $a(p^2)$ is a function of $p^2 = m^2 - \bar{m}_d(2E_k - m_d) \simeq m^2 - 2k^2$, so that $da(p^2)/dk$ is suppressed by one power of k and can be ignored. The contributions from \bar{A}_2 are not of leading order, so that the $a(p^2)$ contributions that might have come from this term can be neglected. The

leading contributions from \bar{A}_1 and \bar{A}_3 combine to give

$$\begin{aligned} Q_A|_{u'} &= e_0 \int_0^\infty \frac{k^2 dk}{2\pi^2} \frac{m}{E_k} 2a(p^2) \{m_d(2E_k - m_d)\bar{A}_1(k) - \bar{A}_3(k)\} \\ &\simeq -e_0 \frac{m^2}{10} \int_0^\infty k^2 dk 2a(p^2) \left\{ 2k^2 \left[4\sqrt{2} u' w' + 2w'^2 + \frac{12\sqrt{2} u' w}{k} + \frac{6w^2}{k^2} \right] + 8m^2 \left[v_t'^2 + \frac{2v_t'^2}{k^2} - 2v_s'^2 - \frac{4v_s'^2}{k^2} \right] \right\}, \end{aligned} \quad (\text{A19})$$

where, when integrating the u, w terms by parts, I use the fact that the volume element is $k^4 dk$ (instead of $k^2 dk$ as it was for \bar{A}_1), giving integrated contributions to (A19) that differ from those shown in (A11).

c. Leading terms in coordinate space

In view of the rich history and importance of this quantity, it is instructive to cast the leading contributions into coordinate space where they have a simple and familiar form.

To aid transforming the terms of $\mathcal{O}(1)$, use the general identities (for arbitrary ℓ and ℓ')

$$\begin{aligned} a \int_0^\infty dk \frac{d}{dk} (k z_\ell z_{\ell'}) &= a \int_0^\infty k^2 dk \left(\frac{z'_\ell z_{\ell'} + z_\ell z'_{\ell'}}{k} + \frac{z_\ell z_{\ell'}}{k^2} \right) = 0, \\ b \int_0^\infty dk \frac{d^2}{dk^2} (k^2 z_\ell z_{\ell'}) &= b \int_0^\infty k^2 dk \left(z''_\ell z_{\ell'} + z_\ell z''_{\ell'} + 2z'_\ell z'_{\ell'} + \frac{4(z'_\ell z_{\ell'} + z_\ell z'_{\ell'})}{k} + \frac{2z_\ell z_{\ell'}}{k^2} \right) = 0, \\ c_\ell \int_0^\infty dk \frac{d}{dk} (k^2 z'_\ell z_{\ell'}) &= c_\ell \int_0^\infty k^2 dk \left(z''_\ell z_{\ell'} + z'_\ell z'_{\ell'} + \frac{2z'_\ell z_{\ell'}}{k} \right) = 0. \end{aligned} \quad (\text{A20})$$

Using these in the calculation of the uw terms gives

$$\begin{aligned} Q_A|_{u,w} &= e_0 \int_0^\infty \frac{k^2 dk}{2\pi^2} \frac{m}{E_k} \bar{A}_1(k) \Big|_{u,w} = \frac{e_0 m^2}{5\sqrt{2}} \int_0^\infty k^2 dk \left\{ -4u' w' - 12 \frac{u' w}{k} \right\} \\ &= \frac{e_0 m^2}{5\sqrt{2}} \int_0^\infty k^2 dk \left\{ (b + c_0) u'' w + (b + c_2) u w'' + (2b + c_0 + c_2 - 4) u' w' \right. \\ &\quad \left. + (2c_0 + 4b + a - 12) \frac{u' w}{k} + (2c_2 + 4b + a) \frac{u w'}{k} + (2b + a) \frac{u w}{k^2} \right\} \\ &= e_0 \frac{4m^2}{5\sqrt{2}} \int_0^\infty k^2 dk \left(u'' - \frac{u'}{k} \right) w, \end{aligned} \quad (\text{A21})$$

where, for any $c_0, a = 2c_0 - 8, b = 4 - c_0$, and $c_2 = c_0 - 4$. To reduce this further, use the fact that the momentum and position space wave functions are related by the spherical Bessel transforms

$$z_\ell(k) = \sqrt{\frac{2}{\pi}} \int_0^\infty r dr j_\ell(kr) z_\ell(r), \quad \frac{z_\ell(r)}{r} = \sqrt{\frac{2}{\pi}} \int_0^\infty k^2 dk j_\ell(kr) z_\ell(k), \quad (\text{A22})$$

where j_ℓ is the spherical Bessel function of order ℓ , satisfying the equation

$$\left(\frac{d^2}{dx^2} + \frac{2}{x} \frac{d}{dx} - \frac{\ell(\ell+1)}{x^2} + 1 \right) j_\ell(x) = 0 \quad (\text{A23})$$

with the convenient recursion relations

$$j_\ell(z) = z^\ell \left(-\frac{1}{z} \frac{d}{dz} \right)^\ell \frac{\sin z}{z} \quad (\text{A24})$$

and the normalization condition

$$\int_0^\infty k^2 dk j_\ell(kr) j_\ell(kr') = \frac{\pi}{2r^2} \delta(r - r'). \quad (\text{A25})$$

Hence, the Bessel transform (A22), and the recursion relation (A24), give

$$u''(k) - \frac{u'(k)}{k} = \sqrt{\frac{2}{\pi}} \int_0^\infty r dr \left(\frac{d^2}{dk^2} - \frac{1}{k} \frac{d}{dk} \right) j_0(kr) u(r) = \sqrt{\frac{2}{\pi}} \int_0^\infty r^3 dr j_2(kr) u(r), \quad (\text{A26})$$

reducing (A21) to

$$Q_A|_{u,w} = e_0 \frac{4m^2}{5\sqrt{2}} \left[\frac{2}{\pi} \right] \int_0^\infty k^2 dk \int_0^\infty r^3 dr j_2(kr) u(r) \int_0^\infty r' dr' j_2(kr') w(r') = e_0 \frac{4m^2}{5\sqrt{2}} \int_0^\infty r^2 dr u(r) w(r). \quad (\text{A27})$$

The leading w^2 term can be similarly reduced. Using the identities (A20) and (A23) gives

$$\begin{aligned} Q_A|_{w^2} &= -e_0 \frac{m^2}{5} \int_0^\infty k^2 dk \left\{ w'^2 + \frac{6w^2}{k^2} \right\} \\ &= -e_0 \frac{m^2}{5} \int_0^\infty k^2 dk \left\{ (2b + c_2) w w'' + (2b + c_2 + 1) w'^2 + (2c_2 + 8b + 2a) \frac{w w'}{k} + (2b + a + 6) \frac{w^2}{k^2} \right\} \\ &= e_0 \frac{m^2}{5} \int_0^\infty k^2 dk \left(w'' + \frac{2w'}{k} - \frac{6w}{k^2} \right) w = -e_0 \frac{m^2}{5} \int_0^\infty r^2 dr w^2(r), \end{aligned} \quad (\text{A28})$$

where $c_2 = -1 - 2b$ and $a = -2b$. Similarly, by using (A23) for $\ell = 1$ the leading P -wave terms become

$$\begin{aligned} Q_A|_{P^2} &= e_0 \frac{m^2}{5} \int_0^\infty k^2 dk \left\{ v_t'^2 - 2v_s'^2 + \frac{2v_t'^2 - 4v_s'^2}{k^2} \right\} \\ &= e_0 \frac{m^2}{5} \int_0^\infty k^2 dk \left\{ c_1 v_t v_t'' + (c_1 + 1) v_t'^2 + 2c_1 \frac{v_t v_t'}{k} + \frac{2v_t'^2}{k} + c_1' v_s v_s'' + (c_1' - 2) v_s'^2 + 2c_1' \frac{v_s v_s'}{k} - \frac{4v_s'^2}{k} \right\} \\ &= e_0 \frac{m^2}{5} \int_0^\infty k^2 dk \left\{ - \left(v_t'' + \frac{2v_t'}{k} - \frac{2v_t'^2}{k^2} \right) v_t + 2 \left(v_s'' + \frac{2v_s'}{k} - \frac{2v_s'^2}{k^2} \right) v_s \right\} = e_0 \frac{m^2}{5} \int_0^\infty r^2 dr (v_t^2 - 2v_s^2), \end{aligned} \quad (\text{A29})$$

where $c_1 = -1$ and $c_1' = 2$.

Summing (A27)–(A29) gives the leading contribution to the quadrupole moment from the (A) diagrams. Multiplying this dimensionless quantity by $1/m_d^2 \simeq 1/(4m^2)$ gives the physical quadrupole moment for a deuteron with unit charge,

$$Q_d|_0 = \frac{1}{4m^2} Q_A|_0 = e_0 \frac{\sqrt{2}}{10} \int_0^\infty r^2 dr \left\{ u(r) w(r) - \frac{1}{\sqrt{8}} [w^2(r) - v_t^2(r) + 2v_s^2(r)] \right\}. \quad (\text{A30})$$

Since $e_0 = 1/2$, this is *one-half* of the relativistic impulse approximation (RIA) result, and it agrees with the leading terms in Eq. (1.16) of Ref. [21]; note that the uw and w^2 terms are identical to 1/2 of the familiar nonrelativistic result (the other 1/2 coming from the B diagrams).

Next I evaluate the terms of order k/m , which arise only from interference between the leading S - and D -state components and the smaller P -state components. These are the A_{int} terms defined in Eq. (A12). Their contribution is

$$Q_A|_1 = e_0(1 + 2\kappa_s) \frac{m}{10\sqrt{3}} \int_0^\infty k^2 dk \left\{ -10\sqrt{2} u' v_t - 20 u' v_s + 8 \left[w v_t' - \frac{w v_t}{k} \right] + 2\sqrt{2} \left[w v_s' - \frac{w v_s}{k} \right] \right\}. \quad (\text{A31})$$

Next, by using

$$\begin{aligned} u'(k) &= \sqrt{\frac{2}{\pi}} \int_0^\infty r^2 dr \left(\frac{1}{r} \frac{d}{dk} \right) j_0(kr) u(r) = -\sqrt{\frac{2}{\pi}} \int_0^\infty r^2 dr j_1(kr) u(r), \\ v'(k) - \frac{v(k)}{k} &= \sqrt{\frac{2}{\pi}} \int_0^\infty r^2 dr \left(\frac{1}{r} \frac{d}{dk} - \frac{1}{kr} \right) j_1(kr) v(r) = -\sqrt{\frac{2}{\pi}} \int_0^\infty r^2 dr j_2(kr) v(r) \end{aligned} \quad (\text{A32})$$

and the normalization condition (A25), these terms give the following contributions to the quadrupole moment:

$$Q_d|_1 = \frac{1}{4m^2} Q_A|_1 = e_0(1 + 2\kappa_s) \frac{1}{2\sqrt{3}} \int_0^\infty dr \frac{r}{m} \left\{ u(r) \left[\frac{1}{\sqrt{2}} v_t(r) + v_s(r) \right] - \frac{2}{5} w(r) \left[v_t(r) + \frac{1}{2\sqrt{2}} v_s(r) \right] \right\}. \quad (\text{A33})$$

Multiplying by 2 gives the RIA result, which also agrees with Ref. [21].

2. Diagrams (B) and (B_±)

By using results from Ref. II, the contributions from diagrams B plus B_± to the quadrupole moment are

$$4D_0\eta G_Q(Q^2)|_{B+B_{\pm}} = e_0 F_1(Q^2) \int_k \left\{ \frac{m}{\kappa_z} \left[\frac{\delta\mathcal{B}_1(k_0, Q)}{k_0} \Big|_{-} - \frac{\delta\mathcal{B}_1(k_0, Q)}{k_0} \Big|_{+} \right] - \frac{1}{m} [\delta\mathcal{C}_1(k, Q) + \delta\mathcal{C}_1(k, -Q)] \right\} \\ + e_0 F_2(Q^2) \int_k \left\{ \frac{m}{\kappa_z} \left[\frac{\delta\mathcal{B}_2(k_0, Q)}{k_0} \Big|_{-} - \frac{\delta\mathcal{B}_1(k_0, Q)}{k_0} \Big|_{+} \right] - \frac{1}{m} [\delta\mathcal{C}_2(k, Q) - \delta\mathcal{C}_2(k, -Q)] \right\}, \quad (\text{A34})$$

where $|_{\pm} \rightarrow |_{k_0=E_{\pm}}$ with $E_{\pm} = \sqrt{m^2 + (\mathbf{k} \pm \mathbf{q}/2)^2}$ and $\kappa_z \equiv \mathbf{k} \cdot \mathbf{q}/E_k = k_z Q/E_k$, and where the $\delta\mathcal{B}_i$ and $\delta\mathcal{C}_i$ differences are

$$\delta\mathcal{B}_i(k_0, Q) = \mathcal{B}_{1,i}(k_0) - \mathcal{B}_{2,i}(k_0), \quad \delta\mathcal{C}_i(k, Q) = \mathcal{C}_{1,i}(\Gamma\Gamma_{\text{off}}) - \mathcal{C}_{2,i}(\Gamma\Gamma_{\text{off}}), \quad (\text{A35})$$

where the traces $\mathcal{B}_{n,i}$ and $\mathcal{C}_{n,i}$ were defined in Ref. II. By introducing the averages

$$\bar{\mathcal{B}}_i(k_0) = \lim_{Q^2 \rightarrow 0} \frac{m_d}{Q^2} \frac{1}{2} \int_{-1}^1 dz \frac{E_k}{k_0} \delta\mathcal{B}_i(k_0, Q), \quad \bar{\mathcal{C}}_{i\pm}(k) = \lim_{Q^2 \rightarrow 0} \frac{m_d}{Q^2} \frac{1}{2} \int_{-1}^1 dz \delta\mathcal{C}_i(k, \pm Q) \quad (\text{A36})$$

and the combinations

$$\bar{B}_i(k) \equiv \frac{m}{k_z Q} (\bar{\mathcal{B}}_i|_{-} - \bar{\mathcal{B}}_i|_{+}), \quad \bar{C}_i(k) \equiv \frac{1}{m} (\bar{\mathcal{C}}_{i+} + \bar{\mathcal{C}}_{i-}) \quad (\text{A37})$$

the contributions of diagrams B plus B_± to the quadrupole moment become

$$Q_B = e_0 \int \frac{k^2 dk}{2\pi^2} \frac{m}{E_k} \{ \bar{B}_1(k) + \kappa_s \bar{B}_2(k) - \bar{C}_1(k) - \kappa_s \bar{C}_2(k) \}. \quad (\text{A38})$$

I will refer to the \mathcal{B} contributions as the ‘‘singular’’ terms, even though the singularity at $k_z = 0$ is canceled by the subtraction of two terms evaluated at $k_0 = E_{\pm}$. The \mathcal{C} contributions are individually finite and depend on the vertex function with *both* nucleons off-shell, Γ_{off} , introduced in Eq. (2.12) of Ref. II.

a. Evaluation of the singular terms

At small Q , the factor $\bar{\mathcal{B}}_i/k_0$ can be expanded in a power series in $(k_0 - E_k)^n$, and the differences $\bar{\mathcal{B}}_i|_{-} - \bar{\mathcal{B}}_i|_{+}$ evaluated. These differences, weighted by the factor κ_z , cannot contribute to the quadrupole moment if they are of higher order than Q^2 . By introducing

$$E_{\pm} - E_k \simeq \pm \frac{k_z Q}{2E_k} + \frac{Q^2}{8E_k^3} (E_k^2 - k_z^2) \mp \frac{k_z Q^3}{16E_k^5} (E_k^2 - k_z^2) \equiv \epsilon_{\pm}, \quad (\text{A39})$$

these differences, up to order Q^2 , are

$$\frac{1}{\kappa_z} (\epsilon_+ - \epsilon_-) \simeq \left[1 - \frac{Q^2}{8E_k^4} (E_k^2 - k_z^2) \right] \rightarrow \left[1 + \frac{z^2 Q^2 k^2}{8E_k^4} \right], \\ \frac{1}{\kappa_z} (\epsilon_+^2 - \epsilon_-^2) \simeq \frac{Q^2}{4E_k^3} (E_k^2 - k_z^2) \rightarrow -\frac{z^2 Q^2 k^2}{4E_k^3}, \quad (\text{A40}) \\ \frac{1}{\kappa_z} (\epsilon_+^3 - \epsilon_-^3) \simeq \frac{k_z^2 Q^2}{4E_k^2} \rightarrow \frac{z^2 Q^2 k^2}{4E_k^2},$$

where contributions from all other powers of $(k_0 - E_k)^n$ are negligible, and, at order Q^2 , only the $z^2 Q^2$ terms will contribute, as explained below. By expanding the coefficients of $\delta\mathcal{B}_i/k_0$ in a power series in Q , Eq. (A40) shows that only the lowest order contribution from the term linear in $k_0 - E_k$ can contribute to the terms of order Q and Q^2 but that all three powers could, in principle, contribute to the term of order Q^0 . However, it turns out that the zeroth order term is accompanied by the Legendre polynomial $P_2(z)$, so that only the contributions proportional to z^2 will survive the integration over z weighted by $P_2(z)$. By recalling the definition of the reduced invariants $X_+ = h\tilde{X}(\tilde{R}_+, R_0^+)$ and $Y_- = h\tilde{Y}(\tilde{R}_-, R_0^-)$ (with X, Y generic names for F, G, H , or I), with $h = h(\tilde{p})$ the strong form factor (which for these contributions is a function of $\tilde{p}^2 = (D_0 - k_0)^2 - k^2$), the contribution from a typical product of invariants $X_+ Y_-$ has the form

$$\frac{\delta\mathcal{B}_i}{k_0} \Big|_{XY} = P_2(z) [B_{00,i}^{XY} + (k_0 - E_k) B_{01,i}^{XY}] h^2 \tilde{X}_+ \tilde{Y}_- + P_2(z) [(k_0 - E_k)^2 B_{02,i}^{XY} + (k_0 - E_k)^3 B_{03,i}^{XY}] XY \\ + k_z Q [B_{10,i}^{XY}(z^2) + (k_0 - E_k) B_{11,i}^{XY}(z^2)] h^2 \tilde{X}_+ \tilde{Y}_- + Q^2 [B_{20,i}^{XY}(z^2) + (k_0 - E_k) B_{21,i}^{XY}(z^2)] h^2 \tilde{X}_+ \tilde{Y}_-, \quad (\text{A41})$$

where the coefficient $B_{nm,i}^{XY}$ multiplies $Q^n(k_0 - E_k)^m$. All of these coefficients are independent of Q and k_0 , but they may be a linear function of z^2 , as indicated. Note the factor of $P_2(z)$ multiplying the terms of $\mathcal{O}(Q^0)$ and that the form of the terms proportional to $(k_0 - E_k)^{2,3}$ anticipates that the arguments of the invariants must be evaluated at $Q = 0$; the differences (A40) ensure that higher order terms will not contribute.

To complete the evaluation of (A41), the vertex functions must also be expanded around the point $Q = 0$ and $k_0 = E_k$. This is done using the arguments of the off-shell vertex functions given in Ref. II. Expanding these arguments to order Q^2 , but at order Q^2 keeping only those terms with a factor of z^2 [because only they will survive the z integration weighted by $P_2(z)$], gives

$$\tilde{R}_\pm = k + \mathbf{R}_\pm + (k_0 - E_k)\mathbf{S}_\pm, \quad R_0^\pm = E_k + \mathcal{E}_\pm + (k_0 - E_k), \quad (\text{A42})$$

where the small quantities are

$$\mathbf{R}_\pm = \pm \frac{zQ}{2m_d}(m_d - E_k) + \frac{z^2 Q^2}{8k m_d^2}[k^2 - (m_d - E_k)^2], \quad \mathbf{S}_\pm = \mp \frac{zQ}{2m_d} + \frac{z^2 Q^2}{4k m_d^2}(m_d - E_k), \quad \mathcal{E}_\pm = \mp \frac{k_z Q}{2m_d}, \quad (\text{A43})$$

and here it is not necessary to retain any higher powers of $(k_0 - E_k)$, because they are multiplied by Q in \tilde{R}_\pm (and hence are negligible) and are altogether absent from R_0^\pm . Note that \tilde{R}_\pm and R_0^\pm reduce to k and k_0 at $Q = 0$, as expected.

Expanding the structure functions to the same order as the expansions (A42) gives

$$h\tilde{X}_\pm \simeq X + \mathbf{R}_\pm X_k + \mathcal{E}_\pm X_{k_0} + \frac{1}{2}[\mathbf{R}_\pm^2 X_{kk} + 2\mathbf{R}_\pm \mathcal{E}_\pm X_{kk_0} + \mathcal{E}_\pm^2 X_{k_0 k_0}] + (k_0 - E_k)\{X_{k_0} + \mathbf{S}_\pm X_k + \mathbf{R}_\pm X_{kk_0} + \mathcal{E}_\pm X_{k_0 k_0}\}, \quad (\text{A44})$$

where

$$X_k = h \frac{\partial}{\partial k} \tilde{X}(k, k_0) \Big|_{Q=0}, \quad X_{k_0} = h \frac{\partial}{\partial k_0} \tilde{X}(k, k_0) \Big|_{Q=0} \quad (\text{A45})$$

and similarly for the other derivatives. The expansion of the strong form factor will also contribute, and these terms will be discussed separately below.

It is convenient to express X_k in terms of X' , where $X' = h \partial \tilde{X}(k, E_k) / (\partial k)$ is the derivative that appears in the calculation of the (A) diagrams. Substituting the relations

$$X_k = X' - \frac{k}{E_k} X_{k_0}, \quad X_{kk_0} = X'_{k_0} - \frac{k}{E_k} X_{k_0 k_0}, \quad X_{kk} = X'' - 2 \frac{k}{E_k} X'_{k_0} - \frac{m^2}{E_k^3} X_{k_0} + \frac{k^2}{E_k^2} X_{k_0 k_0}, \quad (\text{A46})$$

where $X'_{k_0} = h \frac{d}{dk} (\frac{\partial \tilde{X}}{\partial k_0} |_{k_0=E_k})$, into (A44) gives

$$h\tilde{X}_\pm \simeq X \pm \frac{k_z Q}{2k m_d} D_{01}(X) + \frac{z^2 Q^2}{8k m_d^2} D_{02}(X) + (k_0 - E_k) \left\{ X_{k_0} \mp \frac{k_z Q}{2k m_d} D_{11}(X) + \frac{z^2 Q^2}{8k m_d^2} D_{12}(X) \right\}, \quad (\text{A47})$$

where the D_{ij} 's will be given shortly.

Calculation of these contributions is very lengthy, and it is therefore useful to estimate the leading terms at the start. To this end, for the purposes of making estimates only, one recognizes that the leading part of the S -state wave function, u , goes like the inverse of the positive energy propagator, which for $k_0 \neq E_k$ is

$$u(k, k_0) \sim \frac{N_0}{\delta_+} = \frac{N_0}{E_k + k_0 - m_d} \rightarrow N_0 \left[\frac{k^2}{m} + \epsilon + (k_0 - E_k) \right]^{-1}, \quad (\text{A48})$$

where N_0 is an asymptotic normalization constant and $\epsilon > 0$ is the deuteron binding energy. When $k_0 = E_k$ this estimate gives the familiar asymptotic wave function for the deuteron S state. From it the size of various derivatives can be estimated:

$$u \sim k u' \sim k^2 u'' \sim \frac{k^2}{m} u_{k_0} \sim \frac{k^3}{m} u'_{k_0} \sim \frac{k^4}{m^2} u_{k_0 k_0}. \quad (\text{A49})$$

This shows that each k_0 derivative of the ‘‘positive’’ energy wave functions (u , w , and z_ℓ^{-+} , denoted collectively by y_+) is large, of order m/k times larger than each k derivative. However, the expressions for the invariants obtained in Ref. II show that these wave functions are all accompanied by the factor δ_+ , and the k_0 derivatives of the products $(\delta y)_+ \equiv [\delta_+ y_+]_{k_0}$ are small corrections, as was shown in the calculation of the magnetic moment presented in Ref. II. [Similarly, the ‘‘negative’’ energy wave functions (v_r , v_s , and z_ℓ^{--} , denoted collectively by y_-) are all accompanied by the factor δ_- , so for these the corresponding derivatives are $(\delta y)_- \equiv [\delta_- y_-]_{k_0}$, and they are also small.] Since these are small corrections, and the second k_0 derivatives are even smaller, I will neglect the second derivatives $[\delta_\pm y_\pm]_{k_0 k_0}$. With these estimates, the k_0 derivatives of the wave functions are replaced by

$$(y_+)_k \rightarrow \frac{m}{k^2} [(\delta y)_+ - y_+], \quad (y_+)_{k_0 k_0} \sim -\frac{2m}{k^2} (y_+)_{k_0} \rightarrow -\frac{2m^2}{k^4} [(\delta y)_+ - y_+], \quad (\text{A50})$$

where, when $k_0 = E_k$, $\delta_+ = \delta_k \simeq k^2/m$ (neglecting the deuteron binding energy) and $(\delta y)_+$ is a small quantity.

Similar considerations apply to the mixed derivatives, $(y_+)'_{k_0}$. These are large, but the quantity $(\delta'y)_+ \equiv [\delta_+ y_+]'_{k_0}$ is small, leading to the following substitution:

$$(y_+)'_{k_0} \rightarrow \frac{m}{k^2} \left\{ (\delta'y)_+ - y'_+ - \frac{2}{k} [(\delta y)_+ - y] \right\}. \quad (\text{A51})$$

Note that both the second k_0 derivatives and mixed derivatives of y_+ generate large contributions to the leading terms involving y_+ . Ignoring these contributions will give an incorrect result for the nonrelativistic limit.

With this understanding, the D_{ij} 's and their leading terms are

$$\begin{aligned} D_{01}(X) &= (m_d - E_k)X' - \frac{km_d}{E_k}X_{k_0} \rightarrow mX' - 2kX_{k_0}, \\ D_{02}(X) &= [k^2 - (m_d - E_k)^2]X' + k(m_d - E_k)^2X'' + \frac{k^3m_d^2}{E_k^2}X_{k_0k_0} - \frac{2k^2m_d}{E_k}(m_d - E_k)X'_{k_0} - \frac{k^3m_d}{E_k^3}(2E_k - m_d)X_{k_0} \\ &\rightarrow -m^2(X' - kX'') - 4mk^2X'_{k_0} + 4k^3X_{k_0k_0}, \\ D_{11}(X) &= X' - \frac{k}{E_k}X_{k_0} - (m_d - E_k)X'_{k_0} + \frac{k m_d}{E_k}X_{k_0k_0} \rightarrow X' - \frac{k}{m}X_{k_0} - mX'_{k_0} + 2kX_{k_0k_0}, \\ D_{12}(X) &= 2(m_d - E_k)\left(X' - \frac{k}{E_k}X_{k_0}\right) + [k^2 - (m_d - E_k)^2]\left(X'_{k_0} - \frac{k}{E_k}X_{k_0k_0}\right) \\ &\rightarrow 2(mX' - kX_{k_0}) - m(mX'_{k_0} - kX_{k_0k_0}), \end{aligned} \quad (\text{A52})$$

where the double derivative $X_{k_0k_0}$ does not include any of the double k_0 derivatives of the $[\delta_{\pm} y_{\pm}]$ terms listed above.

Using this expansion, the generic product of two invariants picks up some cross terms at $\mathcal{O}(Q^2)$:

$$h^2 X_+ Y_- \simeq XY + \frac{k_z Q}{2km_d} D_{01}(XY) + \frac{z^2 Q^2}{8km_d^2} D_{02}(XY) + (k_0 - E_k) \left\{ X_{k_0} Y + XY_{k_0} + \frac{k_z Q}{2km_d} D_{11}(XY) + \frac{z^2 Q^2}{8km_d^2} D_{12}(XY) \right\}, \quad (\text{A53})$$

where the product coefficients [distinguished from the $D_{ij}(X)$ only by their arguments] are

$$\begin{aligned} D_{01}(XY) &= D_{01}(X)Y - XD_{01}(Y), \\ D_{02}(XY) &= D_{02}(X)Y + XD_{02}(Y) - 2kD_{01}(X)D_{01}(Y), \\ D_{11}(XY) &= -D_{11}(X)Y + XD_{11}(Y) + D_{01}(X)Y_{k_0} - X_{k_0}D_{01}(Y), \\ D_{12}(XY) &= D_{12}(X)Y + XD_{12}(Y) + D_{02}(X)Y_{k_0} + X_{k_0}D_{02}(Y) + 2k[D_{11}(X)D_{01}(Y) + D_{01}(X)D_{11}(Y)], \end{aligned} \quad (\text{A54})$$

with leading contributions obtained from the leading terms given in (A52).

Substituting the expansion (A53) into (A41), taking the differences at $k_0 = E_{\pm}$, and then computing the averages (A36) gives one set of terms coming from the k_0 dependence of the arguments of the invariants proportional to the factors $B_{n0,i}^{XY} D_{1(2-n)}^{XY}$, and another coming from the k_0 dependence of the expansion coefficients proportional to the factors $B_{n1,i} D_{0(2-n)}^{XY}$. The generic term is a sum of these two contributions. Being careful to recall that, through Eq. (A40), the factor of $k_0 - E_k$ gets converted into the factor $-k_z Q/E_k$, and remembering the terms proportional to B_{02} and B_{03} gives

$$\begin{aligned} \bar{B}_i(k)|_{XY} &= - \lim_{Q^2 \rightarrow 0} \frac{m_d m}{Q^2} \frac{1}{2} \int_{-1}^1 dz \left\{ \frac{P_2(z)z^2 Q^2}{8km_d^2} B_{00,i}^{XY} D_{12}(XY) + \frac{(k_z Q)^2}{2km_d} B_{10,i}^{XY} (z^2) D_{11}(XY) + Q^2 B_{20,i}^{XY} (z^2) (XY)_{k_0} \right\} \\ &\quad - \lim_{Q^2 \rightarrow 0} \frac{m_d m}{Q^2} \frac{1}{2} \int_{-1}^1 dz \left\{ \frac{P_2(z)z^2 Q^2}{8km_d^2} B_{01,i}^{XY} D_{02}(XY) + \frac{(k_z Q)^2}{2km_d} B_{11,i}^{XY} (z^2) D_{01}(XY) \right. \\ &\quad \left. + Q^2 \left[B_{21,i}^{XY} (z^2) + \frac{P_2(z)z^2 k^2}{4E_k^4} \left(\frac{1}{2} B_{01,i}^{XY} - E_k B_{02,i}^{XY} + E_k^2 B_{03,i}^{XY} \right) \right] XY \right\} \\ &= - \frac{m}{60km_d} [B_{00,i}^{XY} D_{12}(XY) + B_{01,i}^{XY} D_{02}(XY)] - \frac{km}{6} [\bar{B}_{10,i}^{XY} D_{11}(XY) + \bar{B}_{11,i}^{XY} D_{01}(XY)] \\ &\quad - m_d m [\bar{B}_{20,i}^{XY} (X_{k_0} Y + XY_{k_0}) + \bar{B}_{21,i}^{XY} XY] - \frac{k^2 m_d m}{30E_k^4} \left(\frac{1}{2} B_{01,i}^{XY} - E_k B_{02,i}^{XY} + E_k^2 B_{03,i}^{XY} \right), XY. \end{aligned} \quad (\text{A55})$$

where $(XY)_{k_0} = X_{k_0} Y + XY_{k_0}$, $\bar{B}_{1m,i}^{XY} = B_{1m,i}^{XY} (z^2 = \frac{3}{5})$, and $\bar{B}_{2m,i}^{XY} = B_{2m,i}^{XY} (z^2 = \frac{1}{3})$.

Now consider the contributions from the Q and k_0 dependence of the strong form factors. Expanding the arguments of the form factors to order Q^2 and $(k_0 - E_k)^3$ gives

$$\tilde{p}^2 = (D_0 - k_0)^2 - k^2 \rightarrow p^2 - 2(k_0 - E_k)(m_d - E_k) + (k_0 - E_k)^2 + \frac{Q^2}{4m_d}[m_d - E_k - (k_0 - E_k)], \quad (\text{A56})$$

with $p^2 = m^2 + m_d^2 - 2m_d E_k \simeq m^2 - 2k^2$. Hence, the expansion of the form factors can be written as

$$h^2(\tilde{p}) \simeq h^2 + h^2 \sum_{nm} Q^n (k_0 - E_k)^m B_{nm}^h, \quad (\text{A57})$$

where $B_{1m}^h = B_{00}^h = B_{22}^h = B_{23}^h = 0$ and the exact coefficients, together with their leading values, are

$$\begin{aligned} h^2 B_{01}^h &= -2(h^2)'(m_d - E_k) \rightarrow -h^2 4m a(p^2), \\ h^2 B_{02}^h &= (h^2)' + 2(h^2)''(m_d - E_k)^2 \rightarrow h^2[2a(p^2) + 4a_2(p^2)], \\ h^2 B_{03}^h &= -2(h^2)''(m_d - E_k) - \frac{4}{3}(h^2)'''(m_d - E_k)^3 \rightarrow -h^2 \frac{1}{m} \left[4a_2(p^2) + \frac{8}{3}a_3(p^2) \right], \\ h^2 B_{20}^h &= \frac{(h^2)'}{4m_d}(m_d - E_k) \rightarrow h^2 \frac{1}{4}a(p^2), \\ h^2 B_{21}^h &= -\frac{(h^2)'}{4m_d} \rightarrow -h^2 \frac{1}{4m}a(p^2), \end{aligned} \quad (\text{A58})$$

where the derivatives of h^2 are with respect to \tilde{p}^2 evaluated at $\tilde{p}^2 = p^2$. The first derivative is $(h^2)' = 2h^2 a(p^2)$, where $a(p^2)$ was defined previously, and appeared in the discussion of the (A) diagrams. This definition is generalized to the higher derivatives

$$m^2(h^2)'' \equiv 2h^2 a_2(p^2), \quad m^4(h^2)''' \equiv 2h^2 a_3(p^2), \quad (\text{A59})$$

with $a(p^2) \equiv a_1(p^2)$.

By using the expansion (A57) the dependence of the strong form factors can be included by redefining six of the eight expansion coefficients $B_{nm,i}^{XY}$ as follows:

$$\begin{aligned} B_{01,i}^{XY} &\rightarrow B_{01,i}^{XY} + B_{00,i}^{XY} B_{01}^h, \\ B_{02,i}^{XY} &\rightarrow B_{02,i}^{XY} + B_{01,i}^{XY} B_{01}^h + B_{00,i}^{XY} B_{02}^h, \\ B_{03,i}^{XY} &\rightarrow B_{03,i}^{XY} + B_{02,i}^{XY} B_{01}^h + B_{01,i}^{XY} B_{02}^h + B_{00,i}^{XY} B_{03}^h, \\ B_{11,i}^{XY} &\rightarrow B_{11,i}^{XY} + B_{10,i}^{XY} B_{01}^h, \\ B_{20,i}^{XY} &\rightarrow B_{20,i}^{XY} + P_2(z) B_{00,i}^{XY} B_{20}^h, \\ B_{21,i}^{XY} &\rightarrow B_{21,i}^{XY} + B_{20,i}^{XY} B_{01}^h + P_2(z) B_{00,i}^{XY} B_{21}^h, \end{aligned} \quad (\text{A60})$$

where care has been taken to include the factor of $P_2(z)$ from Eq. (A41), needed in the last two equations. Since neither $B_{00,i}^{XY}$ nor the B^h have any z dependence, these $P_2(z)$ terms integrate to zero, and there are no contributions from the B_{2m}^h , and $B_{20,i}^{XY}$ is not modified.

The exact expansion has been retained to this point, but the derivatives of h^2 are quite small. At small k^2 , $p^2 \sim m^2$, $h = 1$, and by using the form of h given in Ref. [7] (denoted by H in that reference), each successive derivative of h^2 is smaller by a factor of $(\Lambda^2 - m^2)^{-1} \simeq (2m^2)^{-1}$ (near $k^2 \sim 0$ and for the values of Λ_N found to fit the np data). Hence successive derivatives of h^2 are suppressed by factors of k^2/m^2 , and to leading order only the first derivative, proportional to $a(p^2)$, need be retained. With this approximation, and by dropping the $P_2(z)$ terms, the relations (A60) reduce to

$$\begin{aligned} B_{01,i}^{XY} &\rightarrow B_{01,i}^{XY} - 4m a(p^2) B_{00,i}^{XY}, \\ B_{02,i}^{XY} &\rightarrow B_{02,i}^{XY} - 2a(p^2)(2m B_{01,i}^{XY} - B_{00,i}^{XY}), \\ B_{03,i}^{XY} &\rightarrow B_{03,i}^{XY} - 2a(p^2)(2m B_{02,i}^{XY} - B_{01,i}^{XY}), \\ B_{11,i}^{XY} &\rightarrow B_{11,i}^{XY} - 4m a(p^2) B_{10,i}^{XY}, \\ B_{20,i}^{XY} &\rightarrow B_{20,i}^{XY}, \\ B_{21,i}^{XY} &\rightarrow B_{21,i}^{XY} - 4m a(p^2) B_{20,i}^{XY}. \end{aligned} \quad (\text{A61})$$

By using (A55), $\bar{B}_i(k)$ can be expressed in terms of the invariants F , G , H , and I and their first and second derivatives. These in turn can be written in terms of the wave functions u , w , v_t , and v_s and $\chi_\ell = \{z_0^{--}, z_1^{--}, z_0^{++}, z_1^{++}\}$, the negative ρ -spin helicity amplitudes for particle 1, which contribute because the k_0 derivatives of the invariants depend on them. These terms also contributed to the magnetic moment, as discussed in Ref. I. The result of the (B) contributions, as introduced in Eq. (A38), are

$$\begin{aligned} \bar{B}_1(k) &= 2\pi^2 \frac{E_k m^2}{m} \frac{1}{10} \left\{ -4\sqrt{2}u'w' - 2w'^2 - 3v_t'^2 + 6v_s'^2 - \frac{2\sqrt{2}}{k}(6u'w + 5v_t'v_s) - \frac{1}{k^2}[12w^2 + 35v_t'^2 - 10\sqrt{2}v_s v_t - 48v_s'^2] \right. \\ &\quad \left. - \frac{\sqrt{3}m}{k} \left[w'v_t' - 3\frac{w'v_t}{k} - 10\frac{wv_t}{k^2} \right] + \frac{2m}{k} \left[3v_s'z_\delta' + \frac{1}{k}v_s'z_\delta + \frac{14}{k^2}v_s z_\delta \right] + B_{1D} + 2a(p^2)B_1^h + \Delta B_1 \right\}, \quad (\text{A62}) \\ \bar{B}_2(k) &= 2\pi^2 \frac{E_k m^2}{m} \frac{1}{10} \left\{ -20\sqrt{2} \frac{v_t'v_s}{k} - \frac{1}{k^2}[30v_t'^2 - 20\sqrt{2}v_t v_s] + \Delta B_2 \right\}, \end{aligned}$$

where the D-type corrections are

$$\begin{aligned} B_{1D} &= 3 \left\{ \sqrt{2}(u'[\delta_+ \hat{w}]'_{k_0} + w'[\delta_+ \hat{u}]'_{k_0}) + w'[\delta_+ \hat{w}]'_{k_0} + v_t'[\delta_- \hat{v}_t]'_{k_0} - 2v_s'[\delta_- \hat{v}_s]'_{k_0} \right\} \\ &\quad + \frac{2}{k} \left[(9\sqrt{2}u' + 5w')[\delta_+ \hat{w}]_{k_0} - 5\sqrt{2}w'[\delta_+ \hat{u}]_{k_0} + 2v_t'[\delta_- \hat{v}_t]_{k_0} - 4v_s'[\delta_- \hat{v}_s]_{k_0} \right] \\ &\quad + \frac{2}{k^2} \left[(10\sqrt{2}u + 27w - 2[\delta_+ \hat{w}]_{k_0})[\delta_+ \hat{w}]_{k_0} - \sqrt{2}(w + 4[\delta_+ \hat{w}]_{k_0})[\delta_+ \hat{u}]_{k_0} + 7v_t[\delta_- \hat{v}_t]_{k_0} - 14v_s[\delta_- \hat{v}_s]_{k_0} \right], \quad (\text{A63}) \end{aligned}$$

and I introduced the difference

$$z_\delta = \sqrt{2}z_0^{--} - z_1^{--}. \quad (\text{A64})$$

The m/k terms were reduced by using the identities

$$\begin{aligned} \int_0^\infty dk \frac{d}{dk}(kz_1 z_2') &= \int_0^\infty k^2 dk \left[\frac{1}{k}(z_1' z_2' + z_1 z_2'') + \frac{z_1 z_2'}{k^2} \right] = 0, \\ \int_0^\infty dk \frac{d}{dk}(z_1 z_2) &= \int_0^\infty k^2 dk \left(\frac{z_1' z_2 + z_1 z_2'}{k^2} \right) = 0. \end{aligned} \quad (\text{A65})$$

The leading contributions from the derivatives of h^2 are

$$B_1^h = -2k^2 \left\{ 4\sqrt{2}u'w' + 2w'^2 + \frac{12\sqrt{2}}{k}u'w + \frac{1}{k^2}(\sqrt{2}uw + 6w^2) \right\} + \frac{m}{k} 16\sqrt{3}wv_t - 8m^2 \left\{ v_t'^2 - 2v_s'^2 + \frac{2}{k^2}(v_t'^2 - 2v_s'^2) \right\}, \quad (\text{A66})$$

and the k^2/m^2 corrections to the leading terms are

$$\Delta B_1 = -\frac{k^2}{m^2} \left\{ 4\sqrt{2}u'w' + 2w'^2 + \frac{19\sqrt{2}}{k}u'w - \frac{1}{2k^2}(27\sqrt{2}uw - 88w^2) \right\}, \quad \Delta B_2 = \frac{3}{m^2} \{ \sqrt{2}uw - w^2 \}. \quad (\text{A67})$$

b. Evaluation of the regular terms

The contributions from the \mathcal{C} traces are finite, and the generic term from $\delta\mathcal{C}_i$ that contributes to the quadrupole moment has the form

$$\delta\mathcal{C}_i|_{XK} = [P_2(z)C_{0,i}^{XZ} + k_z Q C_{1,i}^{XK} + Q^2 C_{2,i}^{XK}] h^2 \tilde{X}_+ \tilde{K}_-, \quad (\text{A68})$$

where $C_{1,i}$ is linear and $C_{2,i}$ quadratic in k_z^2 . The contributions from the first term come from the expansion of the arguments of the wave functions to order Q^2 [but, because of the presence of $P_2(z)$, only coefficients proportional to z^2 will contribute] and from the second term to order Q . Expanding the arguments given in Ref. II up to order Q^2 gives

$$R_+ \simeq k - \frac{zQ}{2m_d} E_k - \frac{z^2 Q^2 m^2}{8km_d^2}, \quad \hat{R}_- \simeq k - \frac{zQ}{2m_d} m_\Delta - \frac{z^2 Q^2}{8km_d^2} [m_\Delta^2 - k^2], \quad \hat{R}_0^- \simeq E_k + \frac{k_z Q}{2m_d}, \quad (\text{A69})$$

where I have introduced $m_\Delta \equiv 2m_d - E_k$, and R_+ is the argument of the final on-shell vertex function invariants X_+ , and \hat{R}_- and \hat{R}_0^- are the arguments of the initial K_- invariants with both particles off-shell. Hence, expanding a typical product of vertex invariants to order Q^2 gives

$$h^2 \tilde{X}_+ \tilde{K}_- \simeq XK - \frac{zQ}{2m_d} D_1(XK) - \frac{z^2 Q^2}{8km_d^2} D_2(XK), \quad (\text{A70})$$

where

$$\begin{aligned}
D_1(XK) &= E_k X'K + m_\Delta XK_k - kXK_{k_0} = E_k X'K + m_\Delta XK' - \frac{2km_d}{E_k} XK_{k_0} \rightarrow m(X'K + 3XK') - 4kXK_{k_0}, \\
D_2(XK) &= m^2 X'K + [m_\Delta^2 - k^2]XK_k - kE_k^2 X''K - km_\Delta^2 XK_{kk} - k^3 XK_{k_0k_0} + 2k^2 m_\Delta XK_{kk_0} - 2kE_k m_\Delta X'K_k + 2k^2 E_k X'K_{k_0} \\
&= m^2 X'K + [m_\Delta^2 - k^2]XK' - kE_k^2 X''K - km_\Delta^2 XK'' - \frac{4k^3 m_d^2}{E_k^2} XK_{k_0k_0} \\
&\quad + \frac{4k^2 m_d m_\Delta}{E_k} XK'_{k_0} - 2kE_k m_\Delta X'K' + 4k^2 m_d X'K_{k_0} - \frac{4k^3 m_d}{E_k^3} (m_d - E_k)XK_{k_0} \\
&\rightarrow m^2(X'K + 9XK' - kX''K - 9kXK'' - 6kX'K') - 16k^3 XK_{k_0k_0} + 8k^2 m(3XK'_{k_0} + X'K_{k_0}).
\end{aligned} \tag{A71}$$

These were transformed by using (A46) before the leading terms were extracted. Hence the contributions to the quadrupole moment coming from the \bar{C} traces are of the form

$$\begin{aligned}
\bar{C}_i(k) &= \frac{1}{m}(\bar{C}_{i+} + C_{i-}) = \lim_{Q^2 \rightarrow 0} \frac{m_d}{mQ^2} \int_{-1}^1 dz \left\{ -\frac{z^2 P_2(z) Q^2}{8km_d^2} C_{0,i}^{XK} D_2(XK) - \frac{z^2 k Q^2}{2m_d} C_{1i}^{XK} D_1(XK) + Q^2 C_{2,i}^{XK} XK \right\} \\
&= -\frac{1}{30kmm_d} C_{0,i}^{XK} D_2(XK) - \frac{k}{3m} \bar{C}_{1i}^{XK} D_1(XK) + \frac{2m_d}{m} \bar{C}_{2,i}^{XK} XK,
\end{aligned} \tag{A72}$$

where $\bar{C}_{1i}^{XK} = C_{1i}^{XK}(z^2 = \frac{3}{5})$ and $\bar{C}_{2i}^{XK} = C_{2i}^{XK}(z^2 = \frac{1}{3}, z^4 = \frac{1}{5})$ and I used the fact that $\bar{C}_{i+} = \bar{C}_{i-}$.

To include the contributions from the derivatives of the strong form factor, $h_+ = h(p_+)$ (where $p_+^2 = m_d^2 + m^2 - 2D_0 E_k + Qk_z$), expand to order Q^2 , giving

$$h_+^2 \simeq h^2 + 2h^2 a(p^2) \left(Qk_z - \frac{Q^2}{4m_d} E_k \right). \tag{A73}$$

Because the Q^2 term includes no z dependence, it will make no contribution, and the effect of the linear term is to modify the $C_{2,1}^{XK}$ of Eq. (A68) by adding a term

$$C_{2,1}^{XK} \rightarrow C_{2,1}^{XK} + 2a(p^2) k_z^2 C_{1,i}^{XK}. \tag{A74}$$

However, there are no leading contributions from these terms.

The leading contributions to the quadrupole moment coming from the \bar{C} traces are therefore

$$\begin{aligned}
\bar{C}_1(k) &= 2\pi^2 \frac{E_k m^2}{m} \left\{ 12\sqrt{2} v'_t v'_s + \frac{92\sqrt{2}}{k} v'_t v_s - \frac{1}{4k^2} (15v_t^2 + 68\sqrt{2} v_t v_s) + \frac{m}{k} \left[20v'_s z'_\delta - \frac{122}{k} v'_s z_\delta - \frac{164}{k^2} v_s z_\delta \right] + \Delta C_1 \right\}, \\
\bar{C}_2(k) &= 2\pi^2 \frac{E_k m^2}{m} \left\{ -20\sqrt{2} \frac{v'_t v_s}{k} - \frac{1}{k^2} (30v_t^2 - 20\sqrt{2} v_t v_s) + \Delta C_2 \right\},
\end{aligned} \tag{A75}$$

where z_δ was defined in Eq. (A64). Note that these terms all depend the P -state components but that they have very large coefficients. The k^2/m^2 corrections from the large components are

$$\Delta C_1 = \frac{1}{2} \Delta C_2 = -\frac{5k}{m^2} \left[\sqrt{2} u' w + \frac{1}{2k} (3\sqrt{2} u w + w^2) \right]. \tag{A76}$$

Finally, the combined contribution to the quadrupole moment from the (B) + B_\pm terms is the sum of the terms from (A62) and (A75):

$$\begin{aligned}
Q_B &= e_0 \frac{m^2}{10} \int_0^\infty k^2 dk \left\{ -(4\sqrt{2} u' w' + 2w'^2) \left[1 + \frac{k^2}{m^2} \right] + 12\sqrt{2} v'_t v'_s - 3v_t'^2 + 6v_s'^2 - \frac{\sqrt{2}}{k} (12u' w - 82v'_t v_s) \right. \\
&\quad - \frac{1}{k^2} [12w^2 + 50v_t'^2 + 58\sqrt{2} v_s v_t - 48v_s^2] - \frac{\sqrt{3} m}{k} \left[w' v'_t - 3 \frac{w' v_t}{k} - 10 \frac{w v_t}{k^2} \right] \\
&\quad + \frac{m}{k} \left[26v'_s z'_\delta - \frac{120}{k} v'_s z_\delta - \frac{136}{k^2} v_s z_\delta \right] + B_{1D} + 2a(p^2) B_1^h - \kappa_s \left[40\sqrt{2} \frac{v'_t v_s}{k} + \frac{1}{k^2} (60v_t'^2 - 40\sqrt{2} v_t v_s) \right] \\
&\quad \left. - \frac{1}{m^2} \left[\sqrt{2} k u' w (24 + 10\kappa_s) + \sqrt{2} u w (12 - 6\kappa_s) + \frac{1}{2} w^2 (93 + 16\kappa_s) \right] \right\}.
\end{aligned} \tag{A77}$$

3. Total contribution

Adding the contributions from (A11), (A18), (A19), and (A77), and setting $2e_0 = 1$, gives the leading result for the quadrupole moment as the sum of eight terms. Dividing by $m_d^2 \simeq 4m^2$ gives

$$Q_d = Q_{\text{NR}} + Q_{\text{Rc}} + Q_{h'} + Q_{V_2} + Q_{V_1} + Q_{\text{int}} + Q_P + Q_\chi, \quad (\text{A78})$$

where these terms are

$$\begin{aligned} Q_{\text{NR}} &= -\frac{1}{40} \int_0^\infty k^2 dk \left\{ 4\sqrt{2} u' w' + 2w^2 + \frac{12\sqrt{2}}{k} u' w + \frac{12}{k^2} w^2 \right\} = \frac{\sqrt{2}}{10} \int_0^\infty r^2 dr \left\{ u w - \frac{w^2}{\sqrt{8}} \right\}, \\ Q_{\text{Rc}} &= \frac{1}{80} \int_0^\infty \frac{k^4}{m^2} dk \left\{ \frac{\sqrt{2}}{k} u' w (2\kappa_s - 18) + \frac{6\sqrt{2}}{k^2} u w (1 - \kappa_s) - \frac{1}{2k^2} w^2 (87 + 4\kappa_s) \right\}, \\ Q_{h'} &= \frac{1}{80} \int_0^\infty k^2 dk 2a(p^2) \left\{ -2k^2 \left[8\sqrt{2} u' w' + 4w^2 + \frac{24\sqrt{2}}{k} u' w + \frac{1}{k^2} (\sqrt{2} u w + 12w^2) \right] + 16\sqrt{3} \frac{m}{k} w v_t \right. \\ &\quad \left. - 16m^2 [v_t'^2 - 2v_s'^2 + \frac{2}{k^2} (v_t^2 - 2v_s^2)] \right\}, \\ Q_{V_2} &= \frac{1}{20} \int_0^\infty k^2 dk \left\{ \sqrt{2} (u' w^{(2)'} + u^{(2)'} w') + w' w^{(2)'} - v_t' v_t^{(2)'} + 2v_s' v_s^{(2)'} + \frac{3\sqrt{2}}{k} (u^{(2)'} w + u' w^{(2)}) \right. \\ &\quad \left. + \frac{1}{k^2} [6w w^{(2)} - 2v_t v_t^{(2)} + 4v_s v_s^{(2)}] \right\}, \\ Q_{V_1} &= \frac{1}{80} \int_0^\infty k^2 dk B_{1D}, \\ Q_{\text{int}} &= -\frac{\sqrt{3}}{80} \int_0^\infty k^2 dk \frac{m}{k} \left[w' v_t' - 3 \frac{w' v_t}{k} - 10 \frac{w v_t}{k^2} \right], \\ Q_P &= \frac{1}{80} \int_0^\infty k^2 dk \left\{ 2v_s'^2 - v_t'^2 + 12\sqrt{2} v_t' v_s' + (82 - 40\kappa_s) \frac{\sqrt{2}}{k} v_t' v_s - \frac{2}{k^2} [(23 + 30\kappa_s) v_t^2 + (29 - 20\kappa_s) \sqrt{2} v_t v_s - 20v_s^2] \right\}, \\ Q_\chi &= \frac{1}{40} \int_0^\infty k^2 dk \frac{m}{k} \left[13v_s' z_\delta' - \frac{60}{k} v_s' z_\delta - \frac{68}{k^2} v_s z_\delta \right], \end{aligned} \quad (\text{A79})$$

where B_{1D} was given in Eq. (A63).

-
- [1] F. Gross, J. W. Van Orden, and K. Holinde, *Phys. Rev. C* **45**, 2094 (1992).
[2] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, *Phys. Rev. C* **51**, 38 (1995).
[3] R. Machleidt, *Phys. Rev. C* **63**, 024001 (2001).
[4] Y. Huang and W. N. Polyzou, *Phys. Rev. C* **80**, 025503 (2009).
[5] M. Piarulli, L. Girlanda, L. E. Marcucci, S. Pastore, R. Schiavilla, and M. Viviani, *Phys. Rev. C* **87**, 014006 (2013).
[6] D. R. Phillips, *J. Phys. G* **31**, S1263 (2005).
[7] F. Gross and A. Stadler, *Phys. Rev. C* **78**, 014005 (2008).
[8] F. Gross, *Phys. Rev.* **186**, 1448 (1969).
[9] F. Gross, *Phys. Rev. D* **10**, 223 (1974).
[10] F. Gross, *Phys. Rev. C* **26**, 2203 (1982).
[11] F. Gross, *Phys. Rev. C* **89**, 064001 (2014) (referred to as Ref. I).
[12] F. Gross and D. O. Riska, *Phys. Rev. C* **36**, 1928 (1987).
[13] F. Gross, *Phys. Rev. C* **89**, 064002 (2014) (referred to as Ref. II).
[14] W. W. Buck and F. Gross, *Phys. Rev. D* **20**, 2361 (1979).
[15] F. Gross and A. Stadler, *Phys. Rev. C* **82**, 034004 (2010).
[16] V. Z. Jankus, *Phys. Rev.* **102**, 1586 (1956).
[17] A. Stadler and F. Gross, *Phys. Rev. Lett.* **78**, 26 (1997).
[18] W. P. Ford, S. Jeschonnek, and J. W. Van Orden, *Phys. Rev. C* **90**, 064006 (2014).
[19] P. J. Ehlers, A. Accardi, L. T. Brady, and W. Melnitchouk, *Phys. Rev. D* **90**, 014010 (2014).
[20] J. J. Ethier, N. Doshi, S. Malace, and W. Melnitchouk, *Phys. Rev. C* **89**, 065203 (2014).
[21] R. G. Arnold, C. E. Carlson, and F. Gross, *Phys. Rev. C* **21**, 1426 (1980).