



W&M ScholarWorks

Arts & Sciences Articles

Arts and Sciences

2009

Preservers of spectral radius, numerical radius, or spectral norm of the sum on nonnegative matrices

Chi-Kwong Li

William & Mary, ckli@wm.edu

Leiba Rodman

William & Mary

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Li, C. K., & Rodman, L. (2009). Preservers of spectral radius, numerical radius, or spectral norm of the sum on nonnegative matrices. *Linear Algebra and its Applications*, 430(7), 1739-1761.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



Preservers of spectral radius, numerical radius, or spectral norm of the sum on nonnegative matrices

Chi-Kwong Li¹, Leiba Rodman^{*,2}

Department of Mathematics, College of William and Mary, Williamsburg, VA 23187-8795, USA

Received 22 January 2008; accepted 14 April 2008

Available online 29 May 2008

Submitted by R. Loewy

Dedicated to Tom Laffey on the occasion of his 65th birthday.

Abstract

Let M_n^+ be the set of entrywise nonnegative $n \times n$ matrices. Denote by $r(A)$ the spectral radius (Perron root) of $A \in M_n^+$. Characterization is obtained for maps $f : M_n^+ \rightarrow M_n^+$ such that $r(f(A) + f(B)) = r(A + B)$ for all $A, B \in M_n^+$. In particular, it is shown that such a map has the form

$$A \mapsto S^{-1}AS \quad \text{or} \quad A \mapsto S^{-1}A^t S$$

for some $S \in M_n^+$ with exactly one positive entry in each row and each column. Moreover, the same conclusion holds if the spectral radius is replaced by the spectrum or the peripheral spectrum. Similar results are obtained for maps on the set of $n \times n$ nonnegative symmetric matrices. Furthermore, the proofs are extended to obtain analogous results when spectral radius is replaced by the numerical range, or the spectral norm. In the case of the numerical radius, a full description of preservers of the sum is also obtained, but in this case it turns out that the standard forms do not describe all such preservers.

© 2008 Elsevier Inc. All rights reserved.

AMS classification: 15A04; 15A18; 15A60

Keywords: Nonnegative matrices; Spectral radius; Numerical radius; Numerical range; Spectral norm

* Corresponding author.

E-mail addresses: ckli@math.wm.edu (C.-K. Li), lxrod@math.wm.edu (L. Rodman).

¹ Partially supported by a USA NSF Grant DMS 0600859 and a HK RCG Grant.

² Research of the second author was supported in part by NSF Grant DMS-0456625.

1. Introduction

Preserver problems concern the characterization of maps on matrices or operators leaving invariant a certain function, a certain subset of a certain relation. Earlier studies focused on *linear* maps with these properties. The literature on this subject is extensive; see, for example, [12,23] and monographs [19,20,21]. Recently, researchers have studied preserver problems under mild assumptions. In particular, for a given function ν on a matrix set \mathbf{M} with a binary operator $A \circ B$, maps $f : \mathbf{M} \rightarrow \mathbf{M}$ have been studied, that satisfy

$$\nu(f(A) \circ f(B)) = \nu(A \circ B) \quad \forall A, B \in \mathbf{M} \quad (1.1)$$

but not a priori assumed linear or continuous; [5,6,13,14,26] is a small selection of recent works on the topic. There has been interest in studying such problems when $\nu(A)$ is the spectrum, the peripheral spectrum, the numerical radius, the spectral norm, etc. (see the definitions below). See for example the papers [7,15,22], where preserver problems have been studied for ν the peripheral spectrum in the context of uniform algebras; in fact, these works served as motivation for the present study of preservers on nonnegative matrices, as for nonnegative matrices the peripheral spectrum always contains the spectral radius. Moreover, the problems have also been considered for in more general contexts such as function or operator algebras [19]. It is worth noting that even without the linearity assumption, the preservers often end up to be linear and have certain “standard” or “expected” form. Although the statements of results in many cases look similar to those of linear preservers, researchers often have to develop new techniques to solve the preserver problems under mild assumptions; sometimes these assumptions involve nothing more than validity of (1.1). In some cases, one may get unexpected forms for preservers, which lead to deeper understanding and insight to the structures under consideration.

The purpose of this paper is to characterize preservers of the spectral radius, numerical radius, or spectral norm of the sum of nonnegative matrices. There are not many works in the literature on preservers in the context of real entrywise nonnegative matrices: we mention [18], where spectrum preservers are described, [1,10,24,25] that deal with column rank preservers; [2] is concerned with primitivity preservers, and in [4] preserver problems that have to do with irreducibility are considered. In all these works, the linearity of the map f is assumed. In contrast, in the present work we do not assume a priori any additional hypotheses on f except for (1.1) for $A \circ B = A + B$ and a suitable choice of ν .

Let M_n^+ be the set of real entrywise nonnegative matrices, and let $r(A)$ be the spectral radius of a square matrix A . In Section 2, we characterize maps $f : M_n^+ \rightarrow M_n^+$ such that

$$r(f(A) + f(B)) = r(A + B) \quad \forall A, B \in M_n^+.$$

In particular, it is shown that such a map has the form

$$A \mapsto S^{-1}AS \quad \text{or} \quad A \mapsto S^{-1}A^t S, \quad (1.2)$$

for some $S \in M_n^+$ with exactly one positive entry in each row and each column. Moreover, as byproducts, we show that the same conclusion holds if the spectral radius is replaced by the spectrum or the peripheral spectrum. Similar results are obtained for maps on the set of $n \times n$ nonnegative symmetric matrices in Section 3. Furthermore, the proofs are extended to obtain analogous results when spectral radius is replaced by the numerical range, radius, or the spectral norm in Sections 4 and 5. In the case of the numerical radius, a characterization of preservers of the sum is also obtained, but in this case it turns out that the standard forms (1.2) do not describe all such preservers.

The following notation will be used throughout the paper:

M_n the set of all $n \times n$ real matrices.

K_n the set of all $n \times n$ real skew-symmetric matrices.

M_n^+ the set of $n \times n$ real matrices with nonnegative entries.

S_n^+ the set of symmetric matrices in M_n^+ .

To avoid trivialities, we assume $n \geq 2$ throughout our discussion.

$i = \sqrt{-1}$ complex unit.

\mathbb{C} and \mathbb{R} stand for the complex field and the real field, respectively.

$\|x\|$ Euclidean length of a vector x .

e_i is the i th coordinate vector: 1 in the i th position and zeros elsewhere.

$E_{ij} \in M_n^+$ the matrix unit: 1 in the (i, j) th position and zeros everywhere else.

$r(A)$ the spectral radius of a matrix A .

$\sigma(A)$ the spectrum (the set of eigenvalues) of a matrix A .

$\sigma_p(A) = \sigma(A) \cap \{\lambda \in \mathbb{C} : |\lambda| = r(A)\}$ the peripheral spectrum of A .

A^{tr} the transpose of A .

A^* the conjugate transpose of A .

$W(A) = \{x^*Ax : x \in \mathbb{C}^n, x^*x = 1\}$ the numerical range of A .

$w(A) = \max\{|\mu| : \mu \in W(A)\}$ the numerical radius of A .

$\|A\| = \max\{|x^*Ay| : x, y \in \mathbb{C}^n, x^*x = y^*y = 1\}$ the spectral norm of A .

$X \oplus Y := \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}$.

$0_{p \times q}$ the $p \times q$ zero matrix.

$\mathcal{P} \subset M_n^+$ the group of permutation matrices.

$\mathcal{D} \subset M_n^+$ the group of diagonal matrices with positive entries on the diagonal.

$\mathcal{PD} \subset M_n^+$ the group of matrices of the form PD where $P \in \mathcal{P}$ and $D \in \mathcal{D}$.

The role of \mathcal{PD} is exemplified by the following well-known fact:

Fact. A matrix $A \in M_n^+$ has the property that A is invertible and $A^{-1} \in M_n^+$ if and only if $A \in \mathcal{PD}$.

To see the fact, suppose A has columns x_1, \dots, x_n and A^{-1} has rows $y_1^{\text{tr}}, \dots, y_n^{\text{tr}}$. Suppose x_1 has k positive entries. Then for $j = 2, \dots, n$, y_j will have zero entries in the corresponding nonzero positions of x_1 because y_j is nonnegative and $y_j^{\text{tr}}x_1 = 0$. So, all the nonzero entries of the linearly independent vectors y_2, \dots, y_n will lie in fewer than $n - k$ positions. As a result, $k \leq 1$ so that x_1 has only one positive entry. Similar arguments apply to the other columns. Clearly, the nonzero entries of A must lie in different rows because A is invertible.

2. Spectral radius preservers on M_n^+

Here is our main theorem of this section.

Theorem 2.1. *The following statements (1)–(4) are equivalent for a function $f : M_n^+ \rightarrow M_n^+$.*

$$(i) \ r(A + B) = r(f(A) + f(B)) \quad \forall A, B \in M_n^+. \tag{2.1}$$

$$(ii) \ \sigma_p(A + B) = \sigma_p(f(A) + f(B)) \quad \forall A, B \in M_n^+. \tag{2.2}$$

$$(iii) \ \sigma(A + B) = \sigma(f(A) + f(B)) \quad \forall A, B \in M_n^+. \tag{2.3}$$

(iv) There exists a matrix $Q \in \mathcal{PD}$ such that either

$$f(A) = Q^{-1}AQ \quad \forall A \in M_n^+,$$

or

$$f(A) = Q^{-1}A^tQ \quad \forall A \in M_n^+. \tag{2.4}$$

Since for $A \in M_n^+$ we always have $r(A) \in \sigma_p(A)$, the implications (3) \Rightarrow (2) \Rightarrow (1) are clear. Also, (4) \Rightarrow (3) is not difficult to see. It remains to prove (1) \Rightarrow (4).

First, we present some general results and easy observations that will be often used, sometimes without explicit reference, throughout the paper. We will use the directed graph $\Gamma(A)$ associated with $A \in M_n^+$. Recall that $\{1, 2, \dots, n\}$ is the set of vertices of $\Gamma(A)$, and (i, j) is a directed edge in $\Gamma(A)$ if and only if the (i, j) th entry of A is positive.

A matrix $A \in M_n^+$ is said to be irreducible if there is no permutation matrix P such that $PAP^t = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}$ such that A_{11} and A_{22} are non-trivial square matrices. A useful well-known criterion for irreducibility is given in terms $\Gamma(A)$:

Lemma 2.2. $A \in M_n^+$ is irreducible if and only if $\Gamma(A)$ is strongly connected.

Next, we list several well-known properties of nonnegative matrices and their spectral radii (see, for example [8, Theorem 8.4.5] or [3]).

Lemma 2.3. Let $A \in M_n^+$. Then:

- (a) $r(A) \geq r(A')$ for any principal submatrix A' of A . In particular, $r(A) \geq \max\{d : d \text{ is a diagonal entry of } A\}$.
- (b) If $A \in M_n^+$ is nilpotent, i.e., $r(A) = 0$, then all diagonal entries of A are zeros.
- (c) If $A \in M_n^+$ is irreducible and $B \in M_n^+$ is nonzero, then $r(A + B) > r(A)$.
- (d) If $A \in M_n^+$ is reducible, then there is a permutation matrix P such that PAP^t is upper triangular block form $[A_{ij}]_{1 \leq i, j \leq k}$ such that A_{11}, \dots, A_{kk} are irreducible square matrices and $r(A) = \max\{r(A_{jj}) : 1 \leq j \leq k\}$.
- (e) If $A, B \in M_n^+$ and $A \geq B$ entrywise, then $r(A) \geq r(B)$.

Notice that (b) is an immediate consequence of (a).

Lemma 2.4. Let $A_1, A_2 \in M_n^+$ have irreducible principal submatrices B_1 and B_2 , respectively, such that $r(A_1) = r(B_1), r(A_2) = r(B_2)$. If the row and column indices of B_1 and B_2 have non-empty intersection, then

$$r(A_1 + A_2) > \max\{r(A_1), r(A_2)\}. \tag{2.5}$$

Proof. For $t_1, t_2 \in (0, 1]$ consider $t_1A_1 + t_2A_2$ and its irreducible principal submatrix $B(t_1, t_2)$ whose set of row and column indices is the union of the set of row and column indices of B_1 and that of B_2 . Since row and column indices of B_1 and B_2 have non-empty intersection, the matrix $B(t_1, t_2)$ is irreducible in view of Lemma 2.2, for all $t_1, t_2 \in (0, 1]$. Now

$$\begin{aligned} r(A_1 + A_2) &\geq r(B(1, 1)) > \max\{r(B(1, 1/2)), r(B(1/2, 1))\} \\ &\geq \max\{r(B_1), r(B_2)\} = \max\{r(A_1), r(A_2)\}, \end{aligned}$$

where the strict inequality holds by Lemma 2.3 (c), and the non-strict inequalities hold in view of Lemma 2.3 (e). \square

Proof of Theorem 2.1. We focus on the implication (1) \Rightarrow (4). Assume that the function f satisfies the condition (1) of Theorem 2.1. We divide the proof into several assertions.

Assertion 2.5. (a) For any $A \in M_n^+$ we have $r(A) = r(f(A))$.

(b) $A \in M_n^+$ is nilpotent if and only if $f(A)$ is nilpotent.

(c) If A is nonzero, then $f(A)$ is nonzero.

Proof. Condition (a) follows from setting $A = B$ in (2.1).

Condition (b) follows readily from (a).

Suppose A is nonzero and the (i, j) entry of A is nonzero. If $i = j$ then A is not nilpotent and neither is $f(A)$. Thus, $f(A)$ is nonzero. If $i \neq j$, then for $B = E_{ji}$, the submatrix of $A + B$ with row and column indices $\{i, j\}$ has positive spectral radius. If $f(A) = 0$ then $r(f(A) + f(B)) = r(f(B)) = r(B) = 0$, which is a contradiction. \square

Assertion 2.6. There is a permutation P such that for any $\mu > 0$ the diagonal of the matrix $Pf(\mu E_{ii})P^{\text{tr}}$ is the same as that of μE_{ii} for $i = 1, \dots, n$.

Proof. In what follows we let $F_{ij} = f(E_{ij})$. First, consider $\mu = 1$. For each $j = 1, \dots, n$, let G_{jj} be an irreducible principal submatrix of F_{jj} such that

$$r(G_{jj}) = r(F_{jj}) = 1$$

(The existence of principal submatrices G_{jj} is guaranteed by Lemma 2.3 (d).) We will show that $G_{jj} = [1]$. Note that the row (column) indices of G_{11}, \dots, G_{nn} cannot overlap. If it is not true and the row indices of G_{ii} and G_{jj} overlap, then by Lemma 2.4,

$$r(F_{ii} + F_{jj}) > r(F_{ii}) = 1 = r(E_{ii} + E_{jj}),$$

which is a contradiction. Thus, G_{11}, \dots, G_{nn} are one-by-one with non-overlapping row (column) indices. Since $r(G_{jj}) = 1$, we see that $G_{jj} = [1]$ for all $j = 1, \dots, n$. Thus, there exists $P \in \mathcal{P}$ such that $PF_{jj}P^{\text{tr}}$ has one in the (j, j) position. Suppose $i \neq j$, the (i, i) entry $PF_{jj}P^{\text{tr}}$ is zero. Otherwise, the (i, i) entry of $P(F_{ii} + F_{jj})P^{\text{tr}}$ is larger than 1 so that by Lemma 2.3(a),

$$r(F_{ii} + F_{jj}) = r(P(F_{ii} + F_{jj})P^{\text{tr}}) > 1 = r(E_{ii} + E_{jj}).$$

For any $\mu > 0$, we can apply the preceding proof to show that there is a permutation matrix P_μ such that $P_\mu f(\mu E_{ii})P_\mu^{\text{tr}}$ has μ at the (i, i) position and all other diagonal entries equal to zero. If $P_\mu \neq P$, then there will be indices $i \neq j$ and k so that $f(\mu E_{ii})$ has μ in the (k, k) position, and $f(E_{jj})$ has one in the (k, k) position. But then by Lemma 2.3 (a),

$$r(f(\mu E_{ii}) + f(E_{jj})) \geq 1 + \mu > r(\mu E_{ii} + E_{jj}),$$

which is a contradiction. \square

Assertion 2.7. Let P be the permutation satisfying the conclusion of Assertion 2.6. Then for any $i \neq j$, the 2×2 submatrix of $P(E_{ij} + E_{ji})P^{\text{tr}}$ lying at rows and columns with indices $\{i, j\}$ has the form $\begin{bmatrix} 0 & g_{12} \\ g_{21} & 0 \end{bmatrix}$ with $g_{12}g_{21} = 1$.

Proof. For simplicity, we assume that P is the identity matrix. Otherwise, consider the map $X \mapsto Pf(X)P^{\text{tr}}$.

For each $i \neq j$, let $X = E_{ij} + E_{ji}$ and let G_{ij} be an irreducible principal submatrix of $f(X)$ such that

$$r(G_{ij}) = r(f(X)) = r(X) = 1. \tag{2.6}$$

We claim that G_{ij} must lie in a submatrix of $f(X)$ with row and column indices in the set $\{i, j\}$. Indeed, suppose this is not true, and let k be a row and column index of G_{ij} different from i and from j . Denote by $[f(E_{kk})]$ the principal submatrix of $f(E_{kk})$ having the same row and column indices as G_{ij} does. Then:

$$\begin{aligned} r(f(X) + f(E_{kk})) &\geq r(G_{ij} + [f(E_{kk})]) > r(G_{ij}) \\ &= 1 = r(X + E_{kk}) = r(f(X) + f(E_{kk})), \end{aligned}$$

where the strict inequality follows from Lemma 2.3(c). A contradiction is obtained.

Suppose $G_{ij} = \begin{bmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{bmatrix}$. If at least one of g_{12} and g_{21} is zero, then, in view of (2.6) we must have $g_{11} = 1$ or $g_{22} = 1$. But then for $Y = E_{ii}$ or E_{jj} , $f(X) + f(Y)$ a diagonal entry larger than or equal to 2. By Lemma 2.3 (a), we have

$$r(f(X) + f(Y)) \geq 2 > \frac{1 + \sqrt{5}}{2} = r(X + Y),$$

which is a contradiction. Thus, $g_{12}g_{21} \neq 0$.

Next, we claim that $g_{11} = 0$. If it is not true, then for sufficiently large $\mu > 0$, the matrix $f(\mu E_{ii}) + f(X)$ has $\mu + g_{11}$ at the (i, i) position so that

$$r(f(\mu E_{ii}) + f(X)) \geq \mu + g_{11} > \left[\mu + \sqrt{\mu^2 + 4} \right] / 2 = r(\mu E_{ii} + X),$$

which is a contradiction. Similarly, we can show that $g_{22} = 0$. Since $r(G_{ij}) = 1$, we see that $g_{12}g_{21} = 1$. \square

In the rest of the proof, we assume that Assertions 2.6 and 2.7 hold with $P = I$ for simplicity.

Assertion 2.8. For every $\mu > 0$ and every pair of indices $i \neq j$, $f(\mu E_{ij})$ is a nonzero multiple of E_{ij} or of E_{ji} .

Proof. By Assertion 2.7, for any $i \neq j$, the matrix $f(E_{ij} + E_{ji})$ has a submatrix $G_{ij} = \begin{bmatrix} 0 & g_{ij} \\ g_{ij} & 0 \end{bmatrix}$ with row and column indices in $\{i, j\}$ such that $g_{ij}g_{ij} = 1$.

Now, suppose $\mu > 0$, $i \neq j$, and let $f(\mu E_{ij}) = [z_{rs}]_{r,s=1}^n$. Then $z_{kk} = 0$ for all k . Otherwise, the (k, k) entry of $f(E_{kk}) + f(\mu E_{ij})$ is larger than 1 so that

$$r(f(E_{kk}) + f(\mu E_{ij})) > 1 = r(E_{kk} + \mu E_{ij}),$$

which is a contradiction. We also have $z_{pq} = 0$ if at least one of the indices p and q ($p \neq q$) does not belong to the two-element set $\{i, j\}$. Otherwise, the submatrix of $f(E_{pq} + E_{qp}) + f(\mu E_{ij})$ with row and column indices in $\{p, q\}$ has the form

$$C = \begin{bmatrix} 0 & g_{pq} + z_{pq} \\ g_{qp} + z_{qp} & 0 \end{bmatrix}$$

with $g_{qp}g_{pq} = 1$, so that

$$r(f(E_{pq} + E_{qp}) + f(\mu E_{ij})) \geq r(C) > 1 = r(E_{pq} + E_{qp} + \mu E_{ij}),$$

which is a contradiction.

Since $0 = r(\mu E_{ij}) = r(f(\mu E_{ij}))$, we see that $z_{ij}z_{ji} = 0$. Hence $f(\mu E_{ij})$ is a multiple of E_{ij} or E_{ji} . Similarly, $f(\mu E_{ji})$ is a multiple of E_{ij} or of E_{ji} . \square

Assertion 2.9. Let $X = X_0 \oplus 0_{n-3}$, where $X_0 \in M_3^+$ is nilpotent. Then $f(X) = Z_0 \oplus 0_{n-3}$ such that $Z_0 \in M_3^+$ is nilpotent with at most 3 nonzero entries. Moreover, let

$$S = \{E_{ij} : 1 \leq i, j \leq 3, i \neq j\} \tag{2.7}$$

and

$$f(S) = \{\mu_{ij} E_{ij} : 1 \leq i, j \leq 3, i \neq j\} \text{ for some } \mu_{ij} > 0$$

(The form of $f(S)$ follows from Assertion 2.8.) One of the following is true:

- (1) If $f(X)$ has only one nonzero entry, then $r(f(X) + Z) > 0$ for only one matrix Z in $f(S)$.
- (2) If $f(X)$ has exactly two nonzero entries, and they lie in the same row or the same column, then $r(f(X) + Z) > 0$ for exactly two matrices Z in $f(S)$.
- (3) If $f(X)$ has two or three nonzero entries such that two of them are not in the same row or column, then $r(f(X) + Z) > 0$ for at least three matrices Z in $f(S)$.

Proof. Let $f(X) = [y_{pq}]_{p,q=1}^n$. Then $f(X)$ is nilpotent so that $y_{jj} = 0$ for all $j = 1, \dots, n$. Also, if $i \neq j$ and at least one of i and j is larger than 3, then $y_{ij} = 0$. Otherwise,

$$r(f(X) + f(E_{ij})) > 0 = r(X + E_{ij}) \quad \text{or} \quad r(f(X) + f(E_{ji})) > 0 = r(X + E_{ji})$$

by the fact that $f(E_{ij})$ or $f(E_{ji})$ is a multiple of E_{ij} . So, if $y_{ij} \neq 0$, then $i \neq j$ and $1 \leq i, j \leq 3$. Moreover, since $0 = r(X) = r(f(X))$, we see that $y_{ij}y_{ji} = 0$ for $i \neq j$ (otherwise, the 2×2 principal submatrix of $f(X)$ with row and column indices $\{i, j\}$ would have a positive spectral radius, a contradiction with Lemma 2.3(a)). Thus, there are at most three nonzero entries in $f(X)$, and they all lie in the leading 3×3 principal submatrix of $f(X)$. Using the condition that $f(X) = [y_{ij}]_{i,j=1}^n$ with $y_{ij}y_{ji} = 0$, we see that one of the condition (1)–(3) is true. \square

Assertion 2.10. There is $D \in \mathcal{D}$ such that either

- (1) $f(\mu E_{ij}) = \mu D E_{ij} D^{-1}$ for all $\mu > 0$ and all pairs (i, j) , or
- (2) $f(\mu E_{ij}) = \mu D E_{ji} D^{-1}$ for all $\mu > 0$ and all pairs (i, j) .

Proof. First consider the case when $\mu = 1$, for all pairs of indices (i, j) such that $i \neq j$.

By Assertion 2.8, $f(E_{12}) = \mu_2 E_{12}$ or $f(E_{12}) = \mu_2 E_{21}$ for some $\mu_2 > 0$. Assume $f(E_{12}) = \mu_2 E_{12}$. Otherwise, replace f by the map $X \mapsto f(X)^u$. Since

$$1 = r(E_{12} + E_{21}) = r(f(E_{12}) + f(E_{21})) = r(\mu_2 E_{12} + f(E_{21})),$$

using the result of Assertion 2.8 again, we see that $f(E_{21}) = E_{21}/\mu_2$. We get the desired conclusion for $f(E_{ij})$ with $i \neq j$ if $n = 2$.

Assume $n \geq 3$. For any $j \geq 2$, we claim that $f(E_{1j}) = \mu_j E_{1j}$ for some $\mu_j > 0$. For simplicity, suppose that $j = 3$. Let $X = E_{12} + E_{13}$ and let $f(X) = [z_{ij}]_{i,j=1}^n$. By Assertion 2.9, z_{ij} can be nonzero only if $i \neq j$ and $1 \leq i, j \leq 3$; also, $z_{ij}z_{ji} = 0$ for all i, j . Since $r(X + Y) > 0$ for exactly two matrices $Y \in S$ (the set S is defined in (2.7)), we conclude that $r(f(X) + Z) > 0$ for exactly two matrices Z in $f(S)$, and therefore condition (2) of Assertion 2.9 holds. Note that

$$1 = r(X + E_{21}) = r(f(X) + f(E_{21})) = r(f(X) + E_{21}/\mu_2). \tag{2.8}$$

As a result, z_{12} must be one of the two nonzero entries of $f(X)$ in the same row or same column. Thus, either

$$(a) \quad z_{12}z_{13} \neq 0, \quad \text{or} \quad (b) \quad z_{12}z_{32} \neq 0.$$

If (a) holds, then

$$1 = r(X + E_{31}) = r(f(X) + f(E_{31})).$$

Applying Assertion 2.8 for $f(E_{31})$ we see that $f(E_{31})$ is a multiple of E_{31} , and $f(E_{13}) = \mu_3 E_{13}$ as asserted. Suppose (b) holds. Then

$$f(E_{23} + E_{32}) - (kE_{23} + k^{-1}E_{32})$$

is nonnegative for some $k > 0$ by Assertion 2.7. It follows that

$$r(X + E_{23} + E_{32}) = 1 < r(f(X) + kE_{23} + k^{-1}E_{32}) \leq r(f(X) + f(E_{23} + E_{32}))$$

(the inequality \leq holds by Lemma 2.3 (e)), which is a contradiction.

Now, we have $f(E_{1j}) = \mu_j E_{1j}$ with $\mu_j > 0$ for $j = 2, \dots, n$. Let

$$D = \text{diag}(1, \mu_2, \dots, \mu_n).$$

We may replace f by the map $X \mapsto Df(X)D^{-1}$ so that $f(E_{1j}) = E_{1j}$ for $j = 2, \dots, n$. Since

$$1 = r(E_{1j} + E_{j1}) = r(f(E_{1j}) + f(E_{j1})) = r(E_{1j} + f(E_{j1}))$$

and since by Assertion 2.8 $f(E_{j1})$ is a multiple of either E_{1j} or E_{j1} , we have in fact $f(E_{j1}) = E_{j1}$ for all $j = 2, \dots, n$.

Next, we show that $f(E_{ij}) = E_{ij}$ if $i \neq j$ and $i, j \geq 2$. Assume that $(i, j) = (2, 3)$ for simplicity. Let $X = E_{12} + E_{31}$ and $f(X) = [z_{ij}]_{i,j=1}^n$. We claim that $f(X) = X$. Note that $X = X_0 \oplus 0_{n-3}$ with $X_0 \in M_3^+$ is nilpotent. Since $r(X + Y) > 0$ for at least three matrices Y in S , it follows that $r(f(X) + \widehat{Y}) > 0$ for at least three matrices \widehat{Y} in $f(S)$. Hence, $f(X)$ satisfies condition (3) of Assertion 2.9. Since

$$0 = r(X + Y) = r(f(X) + f(Y))$$

for $Y = E_{12}, E_{31}$ and E_{32} , we see that $z_{21} = 0, z_{13} = 0$ and $z_{23} = 0$, i.e.,

$$f(X) = \begin{bmatrix} 0 & z_{12} & 0 \\ 0 & 0 & 0 \\ z_{31} & z_{32} & 0 \end{bmatrix}.$$

Since

$$1 = r(X + E_{21}) = r(f(X) + E_{21}),$$

we see that $z_{12} = 1$; since

$$1 = r(X + E_{13}) = r(f(X) + E_{13}),$$

we see that $z_{31} = 1$. If $Y = E_{23} + E_{32}$, then by Assertion 2.7 there is $\nu > 0$ such that

$$f(Y) - \nu E_{23} - E_{32}/\nu \tag{2.9}$$

is nonnegative. Let

$$\widehat{Z} = E_{12} + E_{31} + z_{32}E_{32} + \nu E_{23} + E_{32}.$$

Assuming for the moment that $\nu \geq 1$, we have

$$r(f(X) + f(Y)) \geq r(\widehat{Z}) \geq r(X + Y) = r(f(X) + f(Y)), \tag{2.10}$$

where the second inequality follows by comparison between the largest roots of the characteristic polynomials $-\lambda^3 + \lambda + 1$ and $-\lambda^3 + (\nu z_{32} + 1)\lambda + \nu$ of $X + Y$ and of \widehat{Z} , respectively. Since the second inequality in (2.10) is an equality, we see that in fact

$$\nu = 1 \quad \text{and} \quad z_{32} = 0. \tag{2.11}$$

Hence $f(X) = X$. Now, by Assertion 2.8, $f(E_{23})$ is a multiple of E_{23} or E_{32} . Since

$$1 = r(X + E_{23}) = r(X + f(E_{23})),$$

we see that $f(E_{23}) = E_{23}$ as asserted.

If ν of (2.9) is smaller than 1, we apply the arguments in the preceding paragraph to $\widehat{X} := E_{21} + E_{13}$ rather than to X , replacing ν by ν^{-1} and interchanging everywhere the subscripts. Then a contradiction with (2.11) will be obtained, thus $\nu < 1$ is not possible.

At this point, we may assume that $f(E_{ij}) = E_{ij}$ if $i \neq j$. Now consider $f(\mu E_{ij}) = [z_{pq}]_{p,q=1}^n$ for $\mu \geq 0$ and $i \neq j$. Then $z_{pp} = 0$ for all $p \in \{1, \dots, n\}$. Otherwise, we obtain a contradiction (in the next formula W stands for a matrix with zero diagonal):

$$\begin{aligned} r(f(\mu E_{ij}) + f(E_{pp})) &= r(f(\mu E_{ij}) + E_{pp} + W) \geq r((1 + z_{pp})E_{pp}) > 1 \\ &= r(\mu E_{ij} + E_{pp}), \end{aligned}$$

where the first equality follows from Assertion 2.6, and the non-strict inequality follows from Lemma 2.3 (e). Also, $z_{pq} = 0$ for $p \neq q$ if $(p, q) \neq (i, j)$. Otherwise, a contradiction again:

$$\begin{aligned} r(f(\mu E_{ij}) + f(E_{qp})) &\geq r(z_{pq}E_{pq} + f(E_{qp})) = r(z_{pq}E_{pq} + E_{qp}) > 0 \\ &= r(\mu E_{ij} + E_{qp}). \end{aligned}$$

Finally,

$$\sqrt{\mu} = r(\mu E_{ij} + E_{ji}) = r(f(\mu E_{ij}) + f(E_{ji})) = r(f(\mu E_{ij}) + E_{ji})$$

implies that

$$f(\mu E_{ij}) = \mu E_{ij}. \tag{2.12}$$

Next, consider $f(\mu E_{ii}) = [z_{rs}]_{r,s=1}^n$ for fixed $\mu > 0$ and fixed $i \in \{1, \dots, n\}$. Then $z_{ii} = \mu$ and $z_{jj} = 0$ for $j \neq i$ by Assertion 2.6. Also, $z_{pq} = 0$ for any $p \neq q$. Otherwise,

$$\begin{aligned} r(f(\mu E_{ii}) + f(\nu E_{pq})) &\geq r(z_{pq}E_{pq} + f(\nu E_{pq})) \\ &= r(z_{pq}E_{pq} + \nu E_{pq}) \quad (\text{using (2.12)}) \\ &> \mu = r(\mu E_{ii} + \nu E_{pq}) \end{aligned}$$

for a sufficiently large ν . Hence $f(\mu E_{ii}) = \mu E_{ii}$. \square

Assertion 2.11. The function f has the form as in (4) of Theorem 2.1.

Proof. Let $D \in \mathcal{D}$ satisfy the conclusion of Assertion 2.10. We may replace f by the map $X \mapsto D^{-1}f(X)D$ and assume that $D = I$. We may further assume that $f(\mu E_{ij}) = \mu E_{ij}$ for all $\mu > 0$ and (i, j) pairs. Otherwise, replace f by the map $X \mapsto f(X)^{\text{tr}}$.

Suppose $A = [a_{ij}]_{i,j=1}^n \in M_n^+$ and $f(A) = [z_{ij}]_{i,j=1}^n$. First, we show that $z_{jj} = a_{jj}$ for each j . For simplicity, we consider z_{11} . Let

$$A = \begin{bmatrix} a_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad \text{and} \quad f(A) = \begin{bmatrix} z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}.$$

Suppose $t > r(A) = r(f(A)) \geq \max\{r(A_{22}), r(Z_{22})\}$,

$$B_t := A_{12}(tI_{n-1} - A_{22})^{-1}A_{21} \quad \text{and} \quad \widehat{B}_t := Z_{12}(tI_{n-1} - Z_{22})^{-1}Z_{21}.$$

Since $\det(tI_n - (A + \mu E_{11}))$ is equal (as a function of μ) to

$$-\mu(\det(tI_{n-1} - A_{22})) + \det(tI_n - A),$$

it follows that there is (unique) $\mu_t > 0$ such that

$$\det(tI_n - (A + \mu_t E_{11})) = 0.$$

Using Schur complements, we see that

$$\det(tI_n - (A + \mu_t E_{11})) = (t - a_{11} - \mu_t - B_t) \det(tI_{n-1} - A_{22}) = 0,$$

i.e.,

$$t - a_{11} - \mu_t - \sum_{k=0}^{\infty} t^{-k-1} A_{12} A_{22}^k A_{21} = 0.$$

Obviously,

$$s - a_{11} - \mu_t - \sum_{k=0}^{\infty} s^{-k-1} A_{12} A_{22}^k A_{21} > 0$$

for every $s > t$; thus

$$t = r(A + \mu_t E_{11}) = r(f(A) + \mu_t E_{11}).$$

Now

$$0 = \det(tI_n - (f(A) + \mu_t E_{11})) = (t - z_{11} - \mu_t - \widehat{B}_t) \det(tI_{n-1} - Z_{22}),$$

i.e.,

$$0 = t - z_{11} - \mu_t - \sum_{k=0}^{\infty} t^{-k-1} Z_{12} Z_{22}^k Z_{21}.$$

As a result,

$$a_{11} + \sum_{k=0}^{\infty} t^{-k-1} A_{12} A_{22}^k A_{21} = \mu_t - t = z_{11} + \sum_{k=0}^{\infty} t^{-k-1} Z_{12} Z_{22}^k Z_{21}$$

for all sufficiently large t , and hence $a_{11} = z_{11}$ as asserted.

Next, we show that $a_{ij} = z_{ij}$ for $i \neq j$. For simplicity, we consider z_{12} . First, suppose $n = 2$. Since

$$r(A + tE_{21}) = \left[(a_{11} + a_{22}) + \sqrt{(a_{11} - a_{22})^2 + 4a_{12}(a_{21} + t)} \right] / 2$$

and

$$r(f(A) + tE_{21}) = \left[(z_{11} + z_{22}) + \sqrt{(z_{11} - z_{22})^2 + 4z_{12}(z_{21} + t)} \right] / 2$$

are equal for all $t > 0$, and using $a_{11} = z_{11}$, $a_{22} = z_{22}$, we see that $a_{12} = b_{12}$.

Next, suppose $n > 2$. Let

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad \text{and} \quad f(A) = \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix},$$

with $A_{11}, Z_{11} \in M_2^+$. Arguing by contradiction, assume that

$$\varepsilon := a_{12} - z_{12} > 0 \tag{2.13}$$

[If the opposite inequality holds, interchange the roles of A and $f(A)$ in the following argument.]
Suppose $t > r(A) = r(f(A))$ and

$$B_t := A_{12}(tI - A_{22})^{-1}A_{21} = \sum_{k=0}^{\infty} t^{-1}A_{12}(t^{-1}A_{22})^k A_{21}.$$

There is $T > 0$ such that each entry of B_t lies in $[0, \varepsilon/3)$ whenever $t \geq T$. If

$$B_t = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} \quad \text{and} \quad C_{t,\mu} = A_{11} + \mu E_{21} + B_t \quad (\mu > 0),$$

then $C_{t,\mu}$ has eigenvalues

$$\begin{aligned} & \left[(a_{11} + a_{22} + b_{11} + b_{22}) \right. \\ & \left. \pm \sqrt{(a_{11} + b_{11} - a_{22} - b_{22})^2 + 4(a_{12} + b_{12})(a_{21} + b_{21} + \mu)} \right] / 2. \end{aligned} \tag{2.14}$$

Note that

$$\det(tI_n - (A + \mu E_{21})) = \det(tI_2 - C_{t,\mu}) \det(tI_{n-2} - A_{22}) \tag{2.15}$$

so that $\det(tI_2 - C_{t,\mu}) > 0$ if $\mu = 0$. Inequality (2.13) implies that $a_{12} > 0$, which, together with formula (2.14), shows that there is (unique) $v_t > 0$ such that the larger eigenvalue of C_{t,v_t} equals t . Moreover, for any $\lambda > t$, we have

$$\det(\lambda I - A - v_t E_{21}) \neq 0 = \det(tI - A - v_t E_{21}).$$

Hence,

$$t = r(A + v_t E_{21}) = r(f(A) + v_t E_{21}).$$

Similarly, if

$$\tilde{B}_t := Z_{12}(tI_{n-2} - Z_{22})^{-1}Z_{21} = [\tilde{b}_{ij}]_{i,j=1}^2 \in M_2^+ \quad \text{and} \quad \tilde{C}_t := Z_{11} + v_t E_{21} + \tilde{B}_t,$$

then

$$\det(tI_n - f(A) - v_t E_{21}) = \det(tI_2 - \tilde{C}_t) \det(tI_{n-2} - Z_{22}),$$

and there exists $\tilde{T} > 0$ such that every entry of \tilde{B}_t is smaller than $\varepsilon/3$ whenever $t > \tilde{T}$. Observe that \tilde{C}_t has eigenvalues

$$\left[(z_{11} + z_{22} + \tilde{b}_{11} + \tilde{b}_{22}) \pm \sqrt{(z_{11} + \tilde{b}_{11} - z_{22} - \tilde{b}_{22})^2 + 4(z_{12} + \tilde{b}_{12})(z_{21} + \tilde{b}_{21} + v_t)} \right] / 2.$$

So, $2r(A + v_t E_{21})$ and $2r(f(A) + v_t E_{21})$ are equal to the following quantities, respectively:

$$a_{11} + a_{22} + b_{11} + b_{22} + \sqrt{(a_{11} + b_{11} - a_{22} - b_{22})^2 + 4(a_{12} + b_{12})(a_{21} + b_{21} + v_t)} \tag{2.16}$$

and

$$z_{11} + z_{22} + \tilde{b}_{11} + \tilde{b}_{22} + \sqrt{(z_{11} + \tilde{b}_{11} - z_{22} - \tilde{b}_{22})^2 + 4(z_{12} + \tilde{b}_{12})(z_{21} + \tilde{b}_{21} + v_t)}. \tag{2.17}$$

Evidently, $v_t \rightarrow \infty$ as $t \rightarrow \infty$. Since

$$a_{12} + b_{12} - (z_{12} + \tilde{b}_{12}) > \varepsilon - 2(\varepsilon/3) > 0 \quad \text{for } t > \max\{T, \tilde{T}\},$$

we have

$$r(A + v_t E_{21}) > r(f(A) + v_t E_{21})$$

for sufficiently large t , which is the desired contradiction. \square

3. Spectral radius preservers on S_n^+

An adaptation of the proof of Theorem 2.1 yields the following preserver result on the set S_n^+ of $n \times n$ symmetric nonnegative matrices.

Theorem 3.1. *The following statements (1)–(4) are equivalent for a function $f : S_n^+ \rightarrow S_n^+$.*

$$(1) \quad r(A + B) = r(f(A) + f(B)) \quad \forall A, B \in S_n^+. \tag{3.1}$$

$$(2) \quad \sigma_p(A + B) = \sigma_p(f(A) + f(B)) \quad \forall A, B \in S_n^+. \tag{3.2}$$

$$(3) \quad \sigma(A + B) = \sigma(f(A) + f(B)) \quad \forall A, B \in S_n^+. \tag{3.3}$$

$$(4) \text{ There exists a matrix } Q \in \mathcal{P} \text{ such that } f(A) = Q^{-1}AQ \quad \forall A \in S_n^+.$$

Proof. We only need to deal with the non-trivial implication (1) \Rightarrow (4). So assume that f satisfies (3.1). Then $r(A) = r(f(A))$ for every $A \in S_n^+$, and in particular $f(A) = 0$ if and only if $A = 0$.

We divide the rest of the proof into several steps.

Step 1. Assertion 2.6, together with its proof, remains valid. Thus, there exists a permutation Q such that for any $\mu > 0$ the diagonal of the matrix $Qf(\mu E_{ii})Q^t$ is the same as that of μE_{ii} for $i = 1, \dots, n$.

Step 2. Let Q be the matrix in Step 1. We show, by following the proof of Assertion 2.7 that for $i \neq j$, the 2×2 submatrix in $Qf(E_{ij} + E_{ji})Q^{-1}$ with row and column indices i, j has the form $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.

Step 3. Assuming the matrix Q in Step 1 equal I_n , we prove that $f(X)$ is a nonzero multiple of X , for $X = \mu(E_{ij} + E_{ji})$ with $\mu > 0$ and $i \neq j$.

Proof of Step 3. By Step 2, for any $i \neq j$, the matrix $f(E_{ij} + E_{ji})$ has a submatrix $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ with row and column indices in $\{i, j\}$. Let $f(X) = [z_{rs}]_{r,s=1}^n$. As in the proof of Assertion 2.8 (but using $E_{ij} + E_{ji}$ in place of E_{ij}) we show that $z_{kk} = 0$ for all k , and that $z_{pq} = 0$ for all pairs $\{p, q\}$, $p \neq q$ such that at least one of p and q does not belong to $\{i, j\}$. Since $f(X)$ is symmetric and $f(X) \neq 0$, the result of Step 3 follows. \square

Step 4. Again assuming $Q = I$, we prove the symmetric analog of Assertion 2.10: The equality

$$f(\mu(E_{ij} + E_{ji})) = \mu(E_{ij} + E_{ji}) \tag{3.4}$$

holds for all $\mu > 0$ and all pairs (i, j) .

Proof of Step 4. For $i \neq j$, the result follows easily from Step 3: $f(\mu(E_{ij} + E_{ji})) = \mu'(E_{ij} + E_{ji})$ for some $\mu' > 0$, but the equality

$$r(f(\mu(E_{ij} + E_{ji}))) = r(\mu(E_{ij} + E_{ji}))$$

yields $\mu' = \mu$, as claimed.

Next, consider $f(\mu E_{ii}) = [z_{rs}]_{r,s=1}^n \in S_n^+$ for fixed $\mu > 0$ and fixed $i \in \{1, \dots, n\}$. Then $z_{ii} = \mu$ and $z_{jj} = 0$ for $j \neq i$ by Step 1. Also, $z_{pq} = 0$ for any $p \neq q$. Suppose it is not true and $z_{pq} = z_{qp} \neq 0$ for some $p \neq q$. Then using the already proved part of (3.4), we can choose $v > \mu$ so that

$$\begin{aligned} & r(f(\mu E_{ii}) + f(v(E_{qp} + E_{pq}))) \\ & \geq r(\mu E_{ii} + z_{pq}(E_{pq} + E_{qp}) + f(v(E_{qp} + E_{pq}))) \\ & = r(\mu E_{ii} + z_{pq}(E_{pq} + E_{qp}) + v(E_{qp} + E_{pq})) \\ & = \begin{cases} z_{pq} + v & \text{if } p \neq i, q \neq i, \\ \frac{1}{2} \left(\mu + \sqrt{\mu^2 + 4(z_{pq} + v)^2} \right) & \text{if } p = i \text{ or } q = i \end{cases} \end{aligned} \tag{3.5}$$

and

$$r(\mu E_{ii} + v(E_{pq} + E_{qp})) = \begin{cases} v & \text{if } p \neq i, q \neq i, \\ \frac{1}{2} \left(\mu + \sqrt{\mu^2 + 4v^2} \right) & \text{if } p = i \text{ or } q = i. \end{cases} \tag{3.6}$$

But then the right hand side of (3.6) is smaller than that of (3.5), a contradiction with (3.1). Hence $f(\mu E_{ii}) = \mu E_{ii}$. \square

Step 5. Conclusion of the proof that (assuming $Q = I$) $f(A) = A$ for all $A \in S_n^+$.

Proof of Step 5. Suppose $A = [a_{ij}]_{i,j=1}^n \in S_n^+$ and $f(A) = [z_{ij}]_{i,j=1}^n$. As in the proof of Assertion 2.11, we show that $z_{jj} = a_{jj}$ for each j .

Next, we will prove the equalities $a_{ij} = z_{ij}$ for $i \neq j$, following the (suitably modified) arguments of the proof of Assertion 2.11. We use the notation introduced in the proof of Assertion 2.11, with obvious additional properties that follow from symmetry; thus $A_{12}^r = A_{21}$, $A_{22}^r = A_{22}$, etc. For simplicity, consider z_{12} . First, suppose $n = 2$. Since

$$r(A + t(E_{21} + E_{12})) = \left[(a_{11} + a_{22}) + \sqrt{(a_{11} - a_{22})^2 + 4(a_{12} + t)(a_{21} + t)} \right] / 2$$

and

$$r(f(A) + t(E_{21} + E_{12})) = \left[(z_{11} + z_{22}) + \sqrt{(z_{11} - z_{22})^2 + 4(z_{12} + t)(z_{21} + t)} \right] / 2$$

are equal for all $t > 0$, and using $a_{11} = z_{11}, a_{22} = z_{22}, a_{12} = a_{21}, z_{12} = z_{21}$, we see that $a_{12} = z_{12}$.

Now suppose $n > 2$. We argue as in the proof of Assertion 2.11, replacing everywhere E_{21} with $E_{21} + E_{12}$, and using the partitions

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}, \quad f(A) = \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix},$$

where $A_{21} = A_{12}^{\text{tr}}$, $Z_{21} = Z_{12}^{\text{tr}}$, and A_{jj}, Z_{jj} are symmetric for $j = 1, 2$,

$$B_t := A_{12}(tI_{n-2} - A_{22})^{-1}A_{21} = [b_{ij}]_{i,j=1}^2 \in S_2^+,$$

$$\tilde{B}_t := Z_{12}(tI_{n-2} - Z_{22})^{-1}Z_{21} = [\tilde{b}_{ij}]_{i,j=1}^2 \in S_2^+.$$

Then

$$r(A + v_t(E_{21} + E_{12})) = r(f(A) + v_t(E_{21} + E_{12})), \tag{3.7}$$

on the other hand, $r(A + v_t(E_{21} + E_{12}))$ and $r(f(A) + v_t(E_{21} + E_{12}))$ are equal to the quantities (2.16) and (2.17), respectively, with b_{12} replaced by $b_{12} + v_t$, and with \tilde{b}_{12} replaced by $\tilde{b}_{12} + v_t$. Let $\varepsilon = a_{12} - z_{12} = a_{21} - z_{21} > 0$. Then

$$a_{12} + b_{12} + a_{21} + b_{21} - (z_{12} + \tilde{b}_{12} + z_{21} + \tilde{b}_{21}) > \frac{1}{3}\varepsilon > 0$$

for large t , a contradiction with (3.7). \square

4. Numerical radius and numerical range preservers

It turns out that preservers of the numerical radius of the sum of nonnegative matrices have more complicated form than the “standard” maps as in other results of this paper.

To state and prove the result, we need to work with the set K_n of $n \times n$ real skew-symmetric matrices.

Theorem 4.1. *Let $f : M_n^+ \rightarrow M_n^+$. Then*

$$w(A + B) = w(f(A) + f(B)) \quad \forall A, B \in M_n^+ \tag{4.1}$$

if and only if there is a permutation matrix P and a function $g : M_n^+ \rightarrow K_n$ satisfying $A + A^{\text{tr}} - g(A) \in M_n^+$ for each A such that

$$f(A) = P(A + A^{\text{tr}} + g(A))P^{\text{tr}}/2 \quad \text{for all } A \in M_n^+. \tag{4.2}$$

Proof. Observe that we have

$$w(A) = r(A + A^{\text{tr}})/2, \quad A \in M_n, \tag{4.3}$$

because for any unit length vector x we can let $|x|$ be obtained from x by replacing all its entries by their absolute values so that

$$|x^*Ax| \leq |x|^{\text{tr}}A|x| = |x|^{\text{tr}}(A + A^{\text{tr}})|x|/2 \leq r(A + A^{\text{tr}})/2 \tag{4.4}$$

and for x a nonnegative eigenvector corresponding to the largest eigenvalue of the symmetric matrix $A + A^{\text{tr}}$ the equality prevails in (4.4). Thus, (4.1) reads

$$r(A + A^{\text{tr}} + B + B^{\text{tr}}) = r(f(A) + f(A)^{\text{tr}} + f(B) + f(B)^{\text{tr}}) \quad \forall A, B \in M_n^+. \tag{4.5}$$

With this observation, the “if” part of Theorem 4.1 is clear.

We focus on the “only if” part. First, note that $w(f(A) + f(A)) = w(A + A)$ implies that $w(A) = w(f(A))$ for all $A \in M_n^+$. Also, $w(A) = r(A)$ for all $A \in S_n^+$. \square

Assertion 4.2. There is a permutation matrix P such that for any $A \in S_n^+$ we have

$$f(A) = P(A + A_K)P^t$$

with $A_K \in K_n$ such that $A + A_K \in M_n^+$.

Proof. Consider the map $f_0 : S_n^+ \rightarrow S_n^+$ defined by $f_0(A) = [f(A) + f(A)^t]/2$. Then

$$r(f_0(A) + f_0(B)) = w(f(A) + f(B)) = w(A + B) = r(A + B) \quad \forall A, B \in S_n^+.$$

By Theorem 3.1, we see that f_0 has the form $A \mapsto PAP^t$ for some permutation matrix P , and Assertion 4.2 follows. \square

Assertion 4.3. Let P be as in Assertion 4.2. For any $A \in M_n^+$, we have $f(A) = P(A + A^t + A_K)P^t/2$ with $A_K \in K_n$ such that $A + A^t + A_K \in M_n^+$.

Proof. For simplicity, we may assume that $P = I$. Suppose $A = A_1 + A_2$ and $f(A) = Z_1 + Z_2$ with $(A_1, A_2), (Z_1, Z_2) \in S_n^+ \times K_n$. Then for any $B \in S_n^+$, we have

$$\begin{aligned} r(A_1 + B) &= w(A + B) = w(f(A) + f(B)) = r\left(Z_1 + \frac{1}{2}(f(B) + f(B)^t)\right) \\ &= r(Z_1 + B), \end{aligned} \tag{4.6}$$

where the last but one equality follows from (4.3), and the last equality holds by Assertion 4.2.

We now prove that

$$A_1 = [a_{ij}]_{i,j=1}^n = [z_{ij}]_{i,j=1}^n = Z_1.$$

First we prove $a_{ii} = z_{ii}$, and for simplicity assume $i = 1$. Then we argue as in the proof of Assertion 2.11, using the partitions

$$A_1 = A = \begin{bmatrix} a_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad \text{and} \quad Z_1 = \begin{bmatrix} z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}$$

and the property (which follows from (4.6)) that

$$r(A_1 + \mu E_{11}) = r(Z_1 + \mu E_{11}) \quad \forall \mu > 0.$$

For the proof that $a_{ij} = z_{ij}, i \neq j$, and assume for simplicity $(i, j) = (1, 2)$, proceed in the same way as in Step 5 of the proof of Theorem 3.1; here, we use the partitions

$$A_1 = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}, \quad Z_1 = \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}, \quad A_{11}, Z_{11} \in S_2^+$$

and the property that

$$r(A_1 + \nu(E_{12} + E_{21})) = r(Z_1 + \nu(E_{12} + E_{21})) \quad \forall \nu > 0.$$

Now, define $g : M_n^+ \rightarrow K_n$ by $g(A) = 2f(A) - P(A + A^t)P^t$. In view of Assertion 4.3 we see that f has the desired form (4.2). \square

Theorem 4.4. Let $f : M_n^+ \rightarrow M_n^+$. Then

$$W(A + B) = W(f(A) + f(B)) \quad \forall A, B \in M_n^+ \tag{4.7}$$

if and only if there is a permutation matrix Q such that either

$$f(A) = Q^{-1}AQ \quad \forall A \in M_n^+,$$

or

$$f(A) = Q^{-1}A^tQ \quad \forall A \in M_n^+.$$

In the proof the following well-known facts will be used. See, for example, Theorem 1.3.6 and Theorem 1.5.2 in [9].

Lemma 4.5. (a) For $n \times n$ complex matrices X and Y , the equality $W(X) = W(Y)$ holds if and only if the largest eigenvalues of the two matrices $e^{it}X + e^{-it}X^*$ and $e^{it}Y + e^{-it}Y^*$ are always the same for every $t \in [0, 2\pi)$.

(b) For a complex 2×2 matrix X , $W(X)$ is an elliptical disk with foci at the eigenvalues of X .

Proof of Theorem 4.4. The implication “if” is clear. (Note that $W(X) = W(X^t)$ for any $n \times n$ complex matrix X .) We focus on the converse. Thus, suppose f satisfies (4.7). Note that $W(f(A) + f(A)) = W(A + A)$ implies that $W(f(A)) = W(A)$ for all $A \in M_n^+$.

Clearly, since (4.7) holds, then (4.1) holds as well. By Theorem 4.1, $f(A)$ has symmetric part $P(A + A^t)P^t/2$ for each $A \in M_n^+$. For simplicity, we may assume that $P = I_n$. If $A \in S_n^+$, then $W(f(A)) = W(A) \subseteq \mathbb{R}$ and hence $f(A) = f(A)^t \in S_n^+$. It follows that

$$f(A) = A \quad \forall A \in S_n^+. \tag{4.8}$$

We divide the rest of the proof into two steps.

Step 1. One of the following holds:

- (a) $f(\mu E_{ij}) = \mu E_{ij}$ for all $i \neq j$ and $\mu > 0$, or
- (b) $f(\mu E_{ij}) = \mu E_{ji}$ for all $i \neq j$ and $\mu > 0$.

Proof of Step 1. Let $f(E_{12}) = X + Y$ with $(X, Y) \in S_n^+ \times K_n$. Then $X = (E_{12} + E_{21})/2$ and $X + Y \in M_n^+$, only the (1, 2) and (2, 1) entries of Y can be nonzero. Since $W(E_{12}) = W(X + Y)$, by Lemma 4.5(b) we see that $X + Y$ is nilpotent. Thus, $Y = (E_{12} - E_{21})/2$ or $(E_{21} - E_{12})/2$. Hence $f(E_{12}) = E_{12}$ or E_{21} . We may assume that the former case holds. Otherwise, replace f by the map $X \mapsto f(X)^t$.

Now, we will show that (a) holds. First, we can use the argument in the preceding paragraph to show that for $\mu > 0$, either $f(\mu E_{21}) = \mu E_{21}$ or $f(\mu E_{21}) = \mu E_{12}$ holds. Since

$$W(\mu E_{21} + E_{12}) = W(f(\mu E_{21}) + f(E_{12})) = W(f(\mu E_{21}) + E_{12}),$$

Lemma 4.5(b) yields $f(\mu E_{21}) = \mu E_{21}$. Now, change the roles of E_{12} and E_{21} in the above argument. We see that $f(\mu E_{12}) = \mu E_{12}$ for any $\mu > 0$. We are done if $n = 2$.

Suppose $n \geq 3$. We can show (as in the preceding paragraph) that for $\mu > 0$ and $j > 2$, either $f(\mu E_{1j}) = \mu E_{1j}$ or $f(\mu E_{1j}) = \mu E_{j1}$. For simplicity, assume that $j = 3$. Suppose $f(\mu E_{13}) = \mu E_{31}$. Let $A = \mu(E_{23} + E_{32})$ and $B = \mu(E_{12} + E_{13})$. Then

$$f(A) = A, \quad f(B) + f(B)^t = B + B^t.$$

Since

$$B + B^{\text{tr}} + f(B) - f(B)^{\text{tr}} = f(B) + f(B)^{\text{tr}} + f(B) - f(B)^{\text{tr}} = 2f(B) \in M_n^+,$$

the skew-symmetric matrix $f(B) - f(B)^{\text{tr}}$ can have nonzero entries only in (1,2), (1,3), (2,1), (3,1) positions, and the absolute value of these entries cannot exceed μ . On the other hand, $W(f(B)) = W(B)$, which is known to be the circular disk centered at zero with radius $\mu/\sqrt{2}$ (see [16] or [11, Theorem 4.1], for example), and therefore $\pm i\mu\sqrt{2}$ are eigenvalues of the matrix $f(B) - f(B)^{\text{tr}}$. It follows that the (1,2), (2,1), (1,3), (3,1) entries of $f(B) - f(B)^{\text{tr}}$ have absolute values equal to μ , and $f(B)$ must be one of the following four matrices:

$$\mu(E_{12} + E_{13}), \quad \mu(E_{21} + E_{31}), \quad \mu(E_{12} + E_{31}), \quad \mu(E_{21} + E_{13}).$$

Suppose the third or the fourth case holds. Then for $X = A + B$ and $Y = f(A) + f(B)$, the largest eigenvalues of $e^{i\pi/3}X + e^{-i\pi/3}X^{\text{tr}}$ and $e^{i\pi/3}Y + e^{-i\pi/3}Y^{\text{tr}}$ are 1.6861μ and 1.6007μ , respectively, by a Matlab computation. Thus, $W(X) \neq W(Y)$, which is a contradiction with (4.7). Now, if $f(B) = \mu(E_{21} + E_{31})$, then for $X = \mu E_{12} + B$ and $Y = \mu E_{12} + f(B)$, the largest eigenvalues of $i(X - X^{\text{tr}})$ and $i(Y - Y^{\text{tr}})$ are μ and $\sqrt{5}\mu$, respectively. Thus, $W(\mu E_{12} + B) \neq W(\mu E_{12} + f(B))$, a contradiction again. So, we must have $f(B) = \mu(E_{12} + E_{13}) = B$. Now, consider $X = \mu E_{13} + B$ and $Y = f(\mu E_{13}) + B = \mu E_{31} + B$. But then $W(X) \neq W(Y)$ (indeed, $W(X)$ is a circular disk but $W(Y)$ is not because Y has 3 distinct eigenvalues [11, Corollary 2.5]), a contradiction. So $f(\mu E_{13}) = \mu E_{31}$ is impossible, and we see that $f(\mu E_{13}) = \mu E_{13}$ holds. Analogously we show that $f(\mu E_{1j}) = \mu E_{1j}$ and $f(\mu E_{j1}) = \mu E_{j1}$ for all $j > 2$.

Now, consider $i, j \geq 2$ and $i \neq j$. Repeat the arguments of the preceding paragraph with $E_{12}, E_{21}, E_{13}, E_{31}$ replaced by $E_{i1}, E_{1i}, E_{ij}, E_{ji}$, respectively, thereby proving the equalities $f(\mu E_{ij}) = \mu E_{ij}, \mu > 0$. \square

Step 2. Assume that condition (a) of Step 1 holds. Then $f(A) = A$ for all $A \in M_n^+$.

Proof of Step 2. Suppose $f(A) = [z_{ij}]_{i,j=1}^n$. Since $f(A) + f(A)^{\text{tr}} = A + A^{\text{tr}}$, we see that $z_{jj} = a_{jj}$ for all $j = 1, \dots, n$. Suppose

$$A - A^{\text{tr}} = [x_{ij}]_{i,j=1}^n \quad \text{and} \quad f(A) - f(A)^{\text{tr}} = [y_{ij}]_{i,j=1}^n.$$

Suppose there is $x_{ij} > y_{ij} \geq 0$ for some $i \neq j$. (We may interchange the roles of A and $f(A)$ in the following if $0 \leq x_{ij} < y_{ij}$.) Say,

$$x_{12} > y_{12} \geq 0. \tag{4.9}$$

Then for sufficiently large $\mu > 0$, one can use a similar argument in the proof of Assertion 2.11 to show that

$$r((A + \mu E_{12}) - (A + \mu E_{12})^{\text{tr}}) \neq r(f(A) + \mu E_{12}) - (f(A) + \mu E_{12})^{\text{tr}}). \tag{4.10}$$

For the reader’s benefit, we provide details.

By Step 1 and (4.8) we know that $f(\mu E_{ij}) = \mu E_{ij}$ for all pairs (i, j) and all $\mu > 0$. If $n = 2$, inequality (4.10) is immediate. So assume $n \geq 3$. Partition:

$$A - A^{\text{tr}} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad \text{and} \quad f(A) - f(A)^{\text{tr}} = \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}$$

with

$$A_{11} = -A_{11}^{\text{tr}} \in M_2, \quad Z_{11} = -Z_{11}^{\text{tr}} \in M_2, \quad A_{21} = -A_{12}^{\text{tr}}, \tag{4.11}$$

$$Z_{21} = -Z_{12}^{\text{tr}}, \quad A_{22} = -A_{22}^{\text{tr}} \in M_{n-2}, \quad Z_{22} = -Z_{22}^{\text{tr}} \in M_{n-2}. \tag{4.12}$$

For sufficiently large $t \in \mathbb{R}$ and for $\mu > 0$, consider

$$B_t := A_{12}(itI - A_{22})^{-1}A_{21} = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} \quad \text{and} \quad C_{t,\mu} := A_{11} + (\mu E_{12} - \mu E_{21}) + B_t.$$

Note that B_t and $C_{t,\mu}$ are complex skew-Hermitian matrices. Then $C_{t,\mu}$ has eigenvalues (note that $x_{11} = x_{22} = 0$)

$$\left[b_{11} + b_{22} \pm \sqrt{(b_{11} - b_{22})^2 + 4(x_{12} + b_{12} + \mu)(x_{21} + b_{21} - \mu)} \right] / 2 \tag{4.13}$$

(Here and in the rest of the proof, for a negative number w , we denote $\sqrt{w} = i|\sqrt{-w}|$.) Also, since $\text{Trace}(B_t)$ is obviously an analytic function of t in a neighborhood of infinity, we have

$$\text{Trace}(B_t) = iq(t),$$

where $q(t) \in \mathbb{R}$ has a fixed sign for all sufficiently large values of t ; say $q(t) \geq 0$. We note also the formula

$$B_t = \sum_{k=0}^{\infty} (it)^{-1} A_{12}((it)^{-1} A_{22})^k A_{21}. \tag{4.14}$$

Formula (4.13) shows that there is (unique) $v_t > 0$ such that the eigenvalue of C_{t,v_t} with the larger absolute value equals it ; here we use the inequality $q(t) \geq 0$. Moreover, for any $|\lambda| > t$, $\lambda \in \mathbb{C}$, we have

$$\det(\lambda I - (A - A^{\text{tr}}) - v_t(E_{12} - E_{21})) \neq 0 = \det(itI - (A - A^{\text{tr}}) - v_t(E_{12} - E_{21}))$$

(this follows from a Schur complement equality analogous to (2.15)). Hence,

$$t = r((A - A^{\text{tr}}) + v_t(E_{12} - E_{21}))$$

and

$$\begin{aligned} & -i2r(A - A^{\text{tr}} + v_t(E_{12} - E_{21})) \\ & = b_{11} + b_{22} + \sqrt{(b_{11} - b_{22})^2 + 4(x_{12} + b_{12} + v_t)(x_{21} + b_{21} - v_t)}. \end{aligned} \tag{4.15}$$

Similarly, if

$$\begin{aligned} \tilde{B}_t & := Z_{12}(itI_{n-2} - Z_{22})^{-1}Z_{21} = [\tilde{b}_{ij}]_{i,j=1}^2 \in M_2 \quad \text{and} \\ \tilde{C}_t & := Z_{11} + v_t(E_{12} - E_{21}) + \tilde{B}_t, \end{aligned}$$

then \tilde{C}_t has eigenvalues

$$\left[(\tilde{b}_{11} + \tilde{b}_{22}) \pm \sqrt{(\tilde{b}_{11} - \tilde{b}_{22})^2 + 4(y_{12} + \tilde{b}_{12} + v_t)(y_{21} + \tilde{b}_{21} - v_t)} \right] / 2.$$

So, for sufficiently large t , and hence for sufficiently large v_t , we have

$$\begin{aligned} & -i2r(f(A) - f(A)^{\text{tr}} + v_t(E_{12} - E_{21})) \\ & = \tilde{b}_{11} + \tilde{b}_{22} \pm \sqrt{(\tilde{b}_{11} - \tilde{b}_{22})^2 + 4(y_{12} + \tilde{b}_{12} + v_t)(y_{21} + \tilde{b}_{21} - v_t)}. \end{aligned} \tag{4.16}$$

Since $x_{12} > y_{12}, x_{21} = -x_{12}, y_{21} = -y_{12}$, and since the absolute values of b_{ij} and $\tilde{b}_{ij}, i, j = 1, 2$, are small in view of (4.14) and an analogous formula for \tilde{B}_t , we see that the right hand sides of (4.15) and (4.16) are not equal for sufficiently large v_t . This proves (4.10).

Now, in view of (4.10), we have

$$W(A + \mu E_{12}) \neq W(f(A) + \mu E_{12}) = W(f(A) + f(\mu E_{12})),$$

where the equality follows from Step 1. This is a contradiction with (4.7). So, $A - A^{\text{tr}} = f(A) - f(A)^{\text{tr}}$, and we conclude that $f(A) = A$. \square

5. Spectral norm preservers

In this section, we consider spectral norm preservers on nonnegative matrices. In contrast with other sections in the paper, here it is natural to prove the result in the framework of the set $M_{m,n}^+$ of $m \times n$ entrywise nonnegative matrices.

Theorem 5.1. *Let $f : M_{m,n}^+ \rightarrow M_{m,n}^+$. Then*

$$\|A + B\| = \|f(A) + f(B)\| \quad \forall A, B \in M_{m,n}^+ \tag{5.1}$$

if and only if there exist permutation matrices $P \in M_m^+$ and $Q \in M_n^+$ such that one of the following holds.

- (a) $f(A) = PAQ$ for all $A \in M_{m,n}^+$.
- (b) $m = n$ and $f(A) = PA^{\text{tr}}Q$ for all $A \in M_{m,n}^+$.

Proof. We focus on the non-trivial “only if” part. Thus, assume (5.1) holds. We may assume that $m \leq n$ and $n \geq 2$ (The case $m > n$ can be treated similarly, and the case $m = n = 1$ is trivial.) The following easy observation will be used repeatedly:

Observation 5.2. (a) For $\mu > 0$, we have $1 + \mu = \|E_{ij} + \mu E_{pq}\|$ if and only if $(i, j) = (p, q)$.
 (b) The equality $\sqrt{2} = \|E_{ij} + E_{pq}\|$ holds if and only if either $i = p, j \neq q$, or $i \neq p, j = q$.

We divide the proof into several steps.

Step 1. For every $\mu > 0$, there exist permutation matrices $P \in M_m^+$ and $Q \in M_n^+$ (which a priori may depend on μ) such that

- (a) $Pf(\mu E_{ii})Q = \mu E_{ii}$ for $i = 1, \dots, m$, and
- (b) for $j = 1, \dots, n - m$, the equalities $Pf(\mu E_{1,m+j})Q = \mu E_{1,m+j}$ hold, in case $n > m$.

Proof of Step 1. Fix $\mu > 0$, and let $f(\mu E_{ii}) = F_{ii}$ for $1 \leq i \leq m$. The condition (5.1) implies that $\|f(A)\| = \|A\|$ for every $A \in M_{m,n}^+$; in particular,

$$\|F_{ii}\| = \mu. \tag{5.2}$$

Since F_{ii} is entrywise nonnegative, there exist entrywise nonnegative vectors of unit length $x_i \in \mathbb{R}^m$ such that

$$\|x_i^{\text{tr}} F_{ii}\| = \mu, \quad i = 1, \dots, m.$$

For any $i \neq j$, since

$$\mu^2 = \|\mu E_{ii} + \mu E_{jj}\|^2 = \|F_{ii} + F_{jj}\|^2$$

$$\begin{aligned} &\geq x_i^{\text{tr}}(F_{ii} + F_{jj})(F_{ii} + F_{jj})^{\text{tr}}x_i \\ &\geq x_i^{\text{tr}}F_{ii}F_{ii}^{\text{tr}}x_i = \mu^2, \end{aligned}$$

we see that $x_i^{\text{tr}}(F_{jj}F_{jj}^{\text{tr}})x_i = 0$. So, x_i is an eigenvector of $F_{jj}F_{jj}^{\text{tr}}$ corresponding to the (smallest) eigenvalue 0. Recall that x_j is the eigenvector of $F_{jj}F_{jj}^{\text{tr}}$ corresponding to the (largest) eigenvalue μ^2 . So, x_i and x_j are orthogonal. As a result, $\{x_1, \dots, x_m\}$ is an orthonormal basis of \mathbb{R}^m . Since x_1, \dots, x_m are nonnegative, we can conclude that x_1, \dots, x_m is a permutation of e_1, \dots, e_m . We may replace f by a map of the form $A \mapsto P(f(A))$ for a suitable permutation matrix P and assume that $x_j = e_j$. Then $e_i^{\text{tr}}F_{jj}F_{jj}^{\text{tr}}e_i = 0$ ($i \neq j$). It follows that the (i, i) entry of $F_{jj}F_{jj}^{\text{tr}}$ is zero for all $i \neq j$. Since $F_{jj}F_{jj}^{\text{tr}}$ is positive semidefinite of norm μ^2 , we see that $F_{jj}F_{jj}^{\text{tr}} = \mu^2 E_{jj}$. As a result, $F_{jj} = \mu e_j v_j^{\text{tr}}$ for some nonnegative vector $v_j \in \mathbb{R}^n$ of unit length, for $j = 1, 2, \dots, m$. Moreover, the equation $\|F_{ii} + F_{jj}\| = \mu$ for $i \neq j$ implies that $v_i^{\text{tr}}v_j = 0$ for $i \neq j$, i.e., the vectors v_1, \dots, v_m have positive entries at different positions.

If $m = n$, then the vectors v_1, \dots, v_m are a permutation of e_1, \dots, e_m , and the proof of Step 1 is complete. Suppose $n > m$. Consider $F(\mu E_{1j}) = F_{1j}$ for $j > m$. Applying the preceding argument to $F_{1j}, F_{22}, \dots, F_{mm}$, we see that $F_{1j} = \mu e_1 w_j^{\text{tr}}$ for some nonnegative unit length vector $w_j \in \mathbb{R}^n$ such that w_j and v_k has positive entries at different positions for any $k = 2, \dots, m$. Note that (for $j > m$)

$$\begin{aligned} 2\mu^2 &= \|\mu E_{11} + \mu E_{1j}\|^2 = \|F_{11} + F_{1j}\|^2 = \|e_1(\mu v_1^{\text{tr}} + \mu w_j^{\text{tr}})\|^2 \\ &= \|\mu v_1^{\text{tr}} + \mu w_j^{\text{tr}}\|^2 = (\mu v_1 + \mu w_j)^{\text{tr}}(\mu v_1 + \mu w_j) = 2\mu^2 + 2w_j^{\text{tr}}v_1, \end{aligned}$$

hence w_j and v_1 also have positive entries at different positions. Applying the same reasoning to $\mu E_{1,j_1}$ and $\mu E_{1,j_2}$, ($j_1, j_2 > m$), we see that also w_{j_1} and w_{j_2} have positive entries at different positions. Now it follows that each of the vectors

$$v_1, \dots, v_m, w_1, \dots, w_{n-m} \tag{5.3}$$

has exactly one positive entry and the positions of these positive entries are different for different vectors in the set (5.3). Thus, the set (5.3) is a permutation of e_1, \dots, e_n , and the results of Step 1 follows. \square

Step 2. There exist permutation matrices $P \in M_m^+$ and $Q \in M_n^+$ such that

- (a) $Pf(\mu E_{ii})Q = \mu E_{ii}$ for $i = 1, \dots, m$ and all $\mu > 0$, and
- (b) for $j = 1, \dots, n - m$, the equalities $Pf(\mu E_{1,m+j})Q = \mu E_{1,m+j}$ hold for all $\mu > 0$, in case $n > m$.

Proof of Step 2. By Step 1, there exist permutations $P(\mu)$ and $Q(\mu)$ such that

$$P(\mu)f(\mu E_{ii})Q(\mu) = \mu E_{ii}, \quad i = 1, \dots, m,$$

and

$$P(\mu)f(\mu E_{1,m+j})Q(\mu) = \mu E_{1,m+j}, \quad j = 1, \dots, n - m$$

(if $m < n$).

We may assume that $P(1) = I_m$ and $Q(1) = I_n$. Otherwise, replace f by the map of the form $X \mapsto P(1)^{-1}f(X)Q(1)^{-1}$. Hence, if

$$\mathcal{S} = \{E_{jj} : 1 \leq j \leq n\} \cup \{E_{1j} : m < j \leq n\},$$

then $f(X) = X$ for any $X \in \mathcal{S}$. Moreover, for any $\mu > 0$ and $X \in \mathcal{S}$, $f(\mu X) = P(\mu)\mu X Q(\mu) = \mu E_{pq}$ for some (p, q) pair. Since

$$1 + \mu = \|X + \mu X\| = \|f(X) + f(\mu X)\| = \|X + f(\mu X)\|,$$

we see (using Observation 5.2) that $f(\mu X) = \mu X$. \square

Step 3. Assume that $P = I$ and $Q = I$ in Step 2. Then one of the two following possibilities holds:

- (a) $f(\mu E_{ij}) = \mu E_{ij}$ for all $\mu > 0$ and (i, j) pairs,
- (b) $m = n$ and $f(\mu E_{ij}) = \mu E_{ji}$ for all $\mu > 0$ and (i, j) pairs.

Proof of Step 3. We may suppose $i \neq j$ (the cases when $i = j$ are taken care of in Step 2). Here $1 \leq i \leq m; 1 \leq j \leq n$.

First, we prove Step 3 for the case $m = 1$. By Step 2, we have $f(\mu E_{11}) = \mu E_{11}$ for all $\mu > 0$. If $f([x_1, x_2, \dots, x_n]) = [z_1, \dots, z_n]$, then

$$\begin{aligned} (\mu + x_1)^2 + \sum_{j=2}^n x_j^2 &= \|\mu E_{11} + [x_1, x_2, \dots, x_n]\|^2 = \|\mu E_{11} + [z_1, \dots, z_n]\|^2 \\ &= (\mu + z_1)^2 + \sum_{j=2}^n z_j^2 \end{aligned}$$

for all $\mu > 0$ which implies $x_1 = z_1$. In particular, f maps the set $\{[a_1, \dots, a_n] \in M_{1,n}^+ : a_1 = 0\}$ to itself, and using the induction on n , we obtain the equalities $f(\mu E_{1j}) = \mu E_{1j}$ for all $\mu > 0$ and $j = 2, 3, \dots, n$. From now on in the proof of Step 3 we assume $m \geq 2$.

Next, for any pair (i, j) , $1 \leq i \leq m, 1 \leq j \leq n$, we can find permutation matrices R (of size $m \times m$) and S (of size $n \times n$) such that $E_{ij} = RE_{11}S$. Then, applying the result of Step 2 to the map $\widehat{f}(X) = f(RXS)$, $X \in M_{m,n}^+$, we see that $f(\mu E_{ij}) = \mu E_{pq}$ for some index pair (p, q) which is independent of μ . It remains to show that

- (a) (p, q) always equals (i, j) , or
- (b) $m = n$ and (p, q) always equals (j, i) .

To this end, consider $f(E_{ij})$ with $i \neq j$. By Step 1,

$$\|f(E_{ij}) + E_{kk}\| = \|f(E_{ij}) + f(E_{kk})\| = \|E_{ij} + E_{kk}\| = \sqrt{2}$$

for $k \in \{i, j\}, j \leq m$. By Observation 5.2 (b), we see that

$$f(E_{ij}) = E_{ij} \quad \text{or} \quad f(E_{ij}) = E_{ji}. \tag{5.4}$$

Consider $f(E_{12})$. If $n > m$, then

$$\sqrt{2} = \|E_{12} + E_{1,m+1}\| = \|f(E_{12}) + E_{1,m+1}\|.$$

Thus, $f(E_{12}) = E_{12}$. Suppose $m = n$ and $f(E_{12}) = E_{21}$. We may replace f by the map $A \mapsto f(A)^{\text{tr}}$ and assume that $f(E_{12}) = E_{12}$.

Assuming that $f(E_{12}) = E_{12}$, we can easily shows that $f(E_{1j}) = E_{1j}$ for all $j = 3, \dots, m$, because

$$\sqrt{2} = \|E_{12} + E_{1j}\| = \|f(E_{12}) + f(E_{1j})\| = \|E_{12} + f(E_{1j})\|,$$

where (5.4) was used. Note that $f(E_{1j}) = E_{1j}$ for $j > m$ by Step 1. Recall that for $i = 3, \dots, m$, we have $f(E_{i1}) = E_{i1}$ or $f(E_{i1}) = E_{1i}$. Since

$$1 = \|E_{1i} + E_{i1}\| = \|f(E_{1i}) + f(E_{i1})\| = \|E_{1i} + f(E_{i1})\|,$$

we see that $f(E_{i1}) = E_{i1}$ for $i = 2, \dots, m$.

Now, for any E_{ij} for $2 \leq i, j \leq m$, since

$$\sqrt{2} = \|E_{1j} + E_{ij}\| = \|f(E_{1j}) + f(E_{ij})\| = \|E_{1j} + f(E_{ij})\|,$$

and using Observation 5.2, we see that $f(E_{ij}) = E_{ij}$. For E_{ij} with $i \geq 2$ and $j > m$, we have

$$\sqrt{2} = \|E_{1j} + E_{ij}\| = \|f(E_{1j}) + f(E_{ij})\| = \|E_{1j} + E_{pq}\|,$$

where the pair (p, q) is such that $f(E_{ij}) = E_{pq}$, and

$$\sqrt{2} = \|E_{ir} + E_{ij}\| = \|E_{ir} + E_{pq}\|, \quad r = 1, 2, \dots, m,$$

so by Observation 5.2 we must have $(p, q) = (i, j)$. So, for any (r, s) pair, we have $f(E_{rs}) = E_{rs}$. \square

Step 4. Assume that (a) in Step 3 holds, and assume also $P = I, Q = I$. Then $f(A) = A$ for all $A \in M_{m,n}^+$.

Proof of Step 4. Let $A = [a_{ij}]$ and $f(A) = [z_{ij}]$ in $M_{m,n}^+$. We show that $a_{ij} = z_{ij}$ for each (i, j) pair. Arguing by contradiction, assume $a_{ij} \neq z_{ij}$ for some pair (i, j) . Say, $(i, j) = (1, 1)$ (for other pairs (i, j) the proof is exactly the same). Suppose $a_{11} > z_{11}$ (if the opposite inequality holds, interchange the roles of A and $f(A)$ in the subsequent argument). Then for $\mu > 0$,

$$(A + \mu E_{11})^*(A + \mu E_{11}) = \mu^2 E_{11} + \mu(A^* E_{11} + E_{11} A) + A^* A.$$

Note that the largest eigenvalue of $\tilde{A} = \mu^2 E_{11} + \mu(A^* E_{11} + E_{11} A)$ equals

$$\left[\mu(\mu + 2a_{11}) + \sqrt{\mu^2(\mu + 2a_{11})^2 + \mu^2\alpha} \right] / 2 \quad \text{with } \alpha = 4 \sum_{j=2}^n a_{1j}^2.$$

Similarly, we have

$$(f(A) + \mu E_{11})^*(f(A) + \mu E_{11}) = \mu^2 E_{11} + \mu(f(A)^* E_{11} + E_{11} f(A)) + f(A)^* f(A)$$

and the largest eigenvalue of $\tilde{Z} = \mu^2 E_{11} + \mu(f(A)^* E_{11} + E_{11} f(A))$ equals

$$\left[\mu(\mu + 2z_{11}) + \sqrt{\mu^2(\mu + 2z_{11})^2 + \mu^2\beta} \right] / 2 \quad \text{with } \beta = 4 \sum_{j=2}^n z_{1j}^2.$$

Denote by $\lambda_1(X)$ the largest eigenvalue of $X \in S_n^+$. Since $a_{11} > z_{11}$, there is a sufficiently large $\mu > 0$ such that

$$\mu(\mu + 2a_{11}) > \mu(\mu + 2z_{11}) + 2\lambda_1(f(A)^* f(A))$$

and

$$(\mu + 2a_{11})^2 + \alpha \geq (\mu + 2z_{11})^2 + \beta.$$

Consequently,

$$\begin{aligned}\lambda_1((\mu E_{11} + A)^*(\mu E_{11} + A)) &\geq \lambda_1(\tilde{A}) > \lambda_1(\tilde{Z}) + \lambda_1(f(A)^* f(A)) \\ &\geq \lambda_1((\mu E_{11} + f(A))^*(\mu E_{11} + f(A))).\end{aligned}$$

It follows that $\|A + \mu E_{11}\| > \|f(A) + \mu E_{11}\|$, which is the desired contradiction, because by Step 3 we have $f(\mu E_{11}) = \mu E_{11}$. \square

References

- [1] L.B. Beasley, S.-Z. Song, K.-T. Kang, B.K. Sarma, Column ranks and their preservers over nonnegative real matrices, *Linear Algebra Appl.*, 399 (2005) 3–16.
- [2] L.B. Beasley, N.J. Pullman, Linear operators that strongly preserve primitivity, *Linear and Multilinear Algebra* 25 (1989) 205–213.
- [3] A. Berman, P. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, SIAM Press, Philadelphia, 1994.
- [4] R.A. Brualdi, Q. Li, Linear transformations on nonnegative matrices preserving properties of irreducibility and full indecomposability, *SIAM J. Algebraic Discrete Methods* 4 (1983) 116–131.
- [5] J.T. Chan, C.K. Li, N.S. Sze, Mappings on matrices: invariance of functional values of matrix products, *J. Aust. Math. Soc.* 81 (2006) 165–184.
- [6] G. Dolinar, P. Šemrl, Determinant preserving maps on matrix algebras, *Linear Algebra Appl.* 348 (2002) 189–192.
- [7] S.A. Grigoryan, T.V. Tonev, *Shift-invariant uniform algebras on groups*, Birkhäuser Verlag, Basel, 2006.
- [8] R.A. Horn, C.R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.
- [9] R.A. Horn, C.R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.
- [10] S.-G. Hwang, S.-J. Kim, S.-Z. Song, Linear operators that preserve spanning column ranks of nonnegative matrices, *J. Korean Math. Soc.* 31 (1994) 645–657.
- [11] D.S. Keeler, L. Rodman, I.M. Spitkovsky, The numerical range of 3×3 matrices, *Linear Algebra Appl.* 252 (1997) 115–139.
- [12] C.K. Li, S. Pierce, Linear preserver problems, *Amer. Math. Monthly* 108 (2001) 591–605.
- [13] C.K. Li, E. Poon, Schur product of matrices and numerical radius (range) preserving maps, *Linear Algebra Appl.* 424 (2007) 8–24.
- [14] C.K. Li, E. Poon, A. Rastogi, Maps preserving spectral radius, numerical radius, spectral norm, *Electron. Linear Algebra* 16 (2007) 347–359.
- [15] A. Luttmann, T.V. Tonev, Uniform algebra isomorphisms and peripheral multiplicativity, *Proc. Amer. Math. Soc.* 135 (2007) 3589–3598., electronic.
- [16] M. Marcus, C. Pesce, Computer generated numerical ranges and some resulting theorems, *Linear and Multilinear Algebra* 20 (1987) 121–157.
- [17] I. Marek, Frobenius theory of positive operators: Comparison theorems and applications, *SIAM J. Applied Math.* 19 (1970) 607–628.
- [18] H. Minc, Linear transformations on nonnegative matrices, *Linear Algebra Appl.* 9 (1974) 149–153.
- [19] L. Molnár, *Selected Preserver Problems on Algebraic Structures of Linear Operators and on Function Spaces*, Springer-Verlag, Berlin, 2007.
- [20] G. de Oliveira, M.A. Duffner, O. Azenhas (Eds.), *Linear preserver problems*, *Linear and Multilinear Algebra* 48 (2001) 281–408.
- [21] S. Pierce, et al., A survey of linear preserver problems, *Linear and Multilinear Algebra* 33 (1992) 1–129.
- [22] N.V. Rao, T.V. Tonev, E.T. Toneva, Uniform algebra isomorphisms and peripheral spectra, topological algebras and applications, *Contemp. Math.* 427 (2007) 401–416., Amer. Math. Soc., Providence, RI.
- [23] P. Šemrl, Maps on matrix spaces, *Linear Algebra Appl.* 413 (2006) 364–393.
- [24] S.-Z. Song, L.B. Beasley, Linear operators that preserve commuting pairs of nonnegative real matrices, *Linear and Multilinear Algebra* 51 (2003) 279–283.
- [25] S.-Z. Song, S.-G. Hwang, Spanning column ranks and their preservers of nonnegative matrices, *Linear Algebra Appl.* 254 (1997) 485–495.
- [26] V. Tan, F. Wang, On determinant preserver problems, *Linear Algebra Appl.* 369 (2003) 311–317.