

# Kenyon College

## Digital Kenyon: Research, Scholarship, and Creative Exchange

---

Kenyon Summer Science Scholars Program

Summer Student Research Scholarship

---

Summer 2015

## Fluctuating Asymmetry as an Indicator of Reproductive Effort in the Leach's Storm-Petrel (*Oceanodroma leucorhoa*)

Lauren Elisabeth Michael  
*Kenyon College*, [michaell@kenyon.edu](mailto:michaell@kenyon.edu)

Robert A. Mauck  
[mauckr@kenyon.edu](mailto:mauckr@kenyon.edu)

Follow this and additional works at: <https://digital.kenyon.edu/summerscienceprogram>



Part of the [Biology Commons](#)

---

### Recommended Citation

Michael, Lauren Elisabeth and Mauck, Robert A., "Fluctuating Asymmetry as an Indicator of Reproductive Effort in the Leach's Storm-Petrel (*Oceanodroma leucorhoa*)" (2015). *Kenyon Summer Science Scholars Program*. Paper 4.  
<https://digital.kenyon.edu/summerscienceprogram/4>

This Article is brought to you for free and open access by the Summer Student Research Scholarship at Digital Kenyon: Research, Scholarship, and Creative Exchange. It has been accepted for inclusion in Kenyon Summer Science Scholars Program by an authorized administrator of Digital Kenyon: Research, Scholarship, and Creative Exchange. For more information, please contact [noltj@kenyon.edu](mailto:noltj@kenyon.edu).

# Fluctuating Asymmetry as an Indicator of Reproductive Effort in the Leach's Storm-Petrel (*Oceanodroma leucorhoa*)

L. E. Michael '17, R. A. Mauck, Department of Biology, Kenyon College

## Background

- Fluctuating asymmetry (FA) is defined as an organism's deviation from bilateral symmetry in a morphological trait, usually as a consequence of an environmental stressor (Galban 2011, De Coster et al. 2013).
- Fluctuating Asymmetry has been used as a phenotypic measurement of developmental instability (Swaddle 2003).
- Parents balance the energetic investment in young against their own condition. because significant health deterioration can decrease their chances of survival and future reproduction (Erikstad et al. 1997).
- The Leach's Storm-Petrel (LHSP) is a pelagic seabird that forms long-term pair bonds and lays one egg per year for up to 38 years.
- We measured FA in a population of Leach's storm-petrels at the Bowdoin Scientific Station that has been studied continuously since 1953.

## Do storm-petrels show FA and can we measure it?


### Inducing Feather Growth

- Plucking a feather induces growth of a replacement feather
- Rate of growth shown reflects nutritional condition while grown
- We plucked right and left rectrices from 35 breeding individuals.
- This allowed us to:
  - compare our measurements of feathers *in situ* against the same feather *ex situ*.
  - assess our ability to accurately measure *in situ* morphology.

### Measuring Feathers *in situ*

- With a wing ruler, we measured both original and induced right and left 5<sup>th</sup> rectrices.

### Measuring Feathers *ex situ*



### Feathers measured *in situ* accurately reflect actual feather length




Figure 1. *Ex situ* measurements were correlated with *in situ* measurements. The red line represents measurements for the right side of the bird; the blue line represents measurements for the left side of the bird. Correlation test,  $r(\text{red}) = 0.89$ ,  $df = 33$ ,  $p < 0.01$ ,  $r(\text{blue}) = 0.89$ ,  $df = 33$ ,  $p < 0.01$ .

### Storm-Petrels show asymmetry in both original and induced feathers




Figure 2. Distribution of asymmetry values of (A) raw induced tail feather measurements and (B) the absolute value of original tail feather.

**Acknowledgements:** First and foremost, I would like to thank Professor Robert Mauck for giving me this incredible opportunity and supporting me every step of the way. I would also like to thank Liam Taylor for helping me keep a sense of humor during those long days in the pouring rain, and Sarah Adrianowycz for all of her help with data collection. Finally, I would like to thank the Kenyon Biology Department and Kenyon Summer Science for funding my research and supporting me throughout this process.

## Does FA in storm-petrels reflect reproductive effort?

### Measuring FA in the field

- Asymmetry measurements were taken on 91 storm-petrels.
- The induced fifth rectrix (tail feather) was measured on the birds from which an original rectrix was plucked.
- Measurements were taken on left and right wing length, tarsus length, and fifth rectrix length *in situ*.
- Assumption: FA of original feathers reflect stress during molt (immediately following previous breeding season).
- Assumption: FA of induced feathers reflect stress since original feather was plucked.

### Measures of Reproductive Effort

- Assumption: egg size positively correlated with energy allocated to the egg
- Assumption: egg lay date negatively correlated with nutritional condition entering breeding season.
- Assumption: Age and pair bond length may affect energetic demands of incubation due to benefits of experience.

### Predictions

- If FA reflects Reproductive Effort, then FA will be:
  - Correlated (+) with egg size and lay date.
  - Correlated (-) with pair bond length and years of site presence.

### Weak evidence that FA reflects reproductive effort in storm-petrels




Figure 3. The relationship between *in situ* tail feather asymmetry values and (A) egg lay date (Pearson's correlation test,  $r=0.093$ ,  $df = 86$ ,  $p=0.39$ ), (B) pair bond length (Pearson's correlation test,  $r=0.0069$ ,  $df = 40$ ,  $p=0.96$ ), (C) years of petrel site presence (Pearson's correlation test,  $r=0.19$ ,  $df = 41$ ,  $p=0.21$ ) and (D) egg volume (Pearson's correlation test,  $r=0.26$ ,  $df = 85$ ,  $p=0.012$ ).

### Literature Cited:

- Galban, I. 2011. *Journal of Ornithology*. 152:223-230.
- De Coster, G., Van Dongen, S., Malaki, P., Muchane, M., Alcantara-Exposito, A., et al. 2013. *Plos One*. 8:3.
- Swaddle, J. P. 2003. *Advances in the Study of Behavior*. 32:169-205.
- Erikstad, K. E. et al. 1997. *Behavioral Ecology and Sociobiology*. 40:95-100.