

Henry Ford Health System

Henry Ford Health System Scholarly Commons

Center for Health Policy and Health Services
Research Articles

Center for Health Policy and Health Services
Research

3-28-2016

Pharmacogenetic Risk Scores for Perindopril Clinical and Cost Effectiveness in Stable Coronary Artery Disease: When Are We Ready to Implement?

Jasmine A. Luzum

David E. Lanfear

Henry Ford Health System, dlanfea1@hfhs.org

Follow this and additional works at: https://scholarlycommons.henryford.com/chphsr_articles

Recommended Citation

Luzum JA, Lanfear DE. Pharmacogenetic Risk Scores for Perindopril Clinical and Cost Effectiveness in Stable Coronary Artery Disease: When Are We Ready to Implement?. *J Am Heart Assoc* 2016; 5(3):003440.

This Article is brought to you for free and open access by the Center for Health Policy and Health Services Research at Henry Ford Health System Scholarly Commons. It has been accepted for inclusion in Center for Health Policy and Health Services Research Articles by an authorized administrator of Henry Ford Health System Scholarly Commons.

Pharmacogenetic Risk Scores for Perindopril Clinical and Cost Effectiveness in Stable Coronary Artery Disease: When Are We Ready to Implement?

Jasmine A. Luzum, PharmD, PhD, BCPS; David E. Lanfear, MD, MS, FAHA

In this issue of the *Journal of the American Heart Association*, Oemrawsingh et al present a very interesting and well-executed analysis of clinical and pharmacogenetic risk scores for perindopril effectiveness in 8726 patients with stable coronary artery disease.¹ Their results demonstrate that the clinical risk score can predict patient risk but not the relative risk reduction of the therapy, which was homogeneous across clinical risk groups. In contrast, the pharmacogenetic risk scores could differentiate patients in terms of the effectiveness of therapy; in the 26.5% of patients with a risk score >2 , perindopril was *not* associated with clinical benefit (primary end point: cardiovascular mortality, nonfatal myocardial infarction, or resuscitated cardiac arrest), while an enhanced benefit was seen in the remaining 73.5%. The authors then added to this by quantifying the clinical and cost implications of various clinical and pharmacogenetic risk score strategies. Genetically testing all patients and only treating those with a pharmacogenetic risk score ≤ 2 was equally clinically effective and more cost-effective compared to standard care (ie, no genetic testing, all patients are treated with perindopril).

By combining the pharmacogenetic risk score with a published clinical risk score² and analyzing the cost-effectiveness of the combination, this study by Oemrawsingh et al is an important extension of their initial publication of the pharmacogenetic risk score in 2010.³ The current study

has several strengths: large sample size, randomized controlled data, assessment of pharmacogenetic risk factors in the context of clinical risk strata, inclusion of a cost-effectiveness analysis, and the relative simplicity of the pharmacogenetic score. However, like many excellent publications, this one prompts as many questions as it answers. Are we ready to accept the likelihood that 1 in 4 stable coronary artery disease patients are treated ineffectively (or perhaps even harmfully?) with perindopril, wasting precious resources and effort? Should we implement pharmacogenetic tailored therapy now, and if not, when will we have sufficiently convincing data to do so? While the United States Food and Drug Administration (FDA) has incorporated genetics into the prescribing information for >130 drugs⁴ (Table), and the Clinical Pharmacogenetics Implementation Consortium has published 33 guidelines on pharmacogenetic information (Table),⁵ pharmacogenetics has not achieved significant use in cardiology. There are several important barriers including logistical and educational concerns,⁶ but equally important and relevant to the work by Oemrawsingh et al are the cost implications and unclear standards for levels of evidence required to justify pharmacogenetic testing.

Regarding costs, the current work is very helpful. Oemrawsingh et al estimated the cost to genotype all 3 variants used in their pharmacogenetic risk score to be only €15 euros (approximately \$17 USD), and currently, 2.5 million genetic variants in an individual patient can be simultaneously tested in a single sample for around \$300 USD. For comparison, a single comprehensive metabolic panel, which is often repeated annually, costs around \$600. Unlike many other types of laboratory tests, the patient's DNA is static, and thus genetic testing only needs to be performed once to provide lifetime results. Despite the declining costs of genetic testing and examples of cost-effectiveness as reported by Oemrawsingh et al, the barrier that patients and providers most often face is lack of reimbursement for pharmacogenetic testing (not only in the United States but in Europe as well).^{7,8} Pharmacogenetic tests for most cardiovascular drugs are not reimbursed, except for clopidogrel and warfarin. For example, 7 different third-party payers, including Medicare, have

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

From the Ohio State University, College of Medicine, Center for Pharmacogenomics, Columbus, OH (J.A.L.); Heart and Vascular Institute, Henry Ford Hospital, Detroit, MI (D.E.L.); Center for Health Policy and Health Services Research, Henry Ford Hospital, Detroit, MI (D.E.L.).

Correspondence to: David E. Lanfear, MD, MS, Henry Ford Hospital, 2799 W. Grand Blvd. K14, Detroit, MI 48202. E-mail: dlanfea1@hfhs.org
J Am Heart Assoc. 2016;5:e003440 doi: 10.1161/JAHA.116.003440.

© 2016 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Table. Cardiovascular Drugs With Pharmacogenetic Information Included in the US FDA Prescribing Information and CPIC guidelines.

Drug	Source	Summary of Pharmacogenetic Information
Carvedilol	US FDA	“Retrospective analysis of side effects in clinical trials showed that poor 2D6 metabolizers had a higher rate of dizziness during up-titration”
Clopidogrel	US FDA	“Consider alternative treatment or treatment strategies in patients identified as CYP2C19 poor metabolizers.” (Black Box Warning)
Clopidogrel	CPIC	“The CPIC Dosing Guideline for clopidogrel recommends an alternative antiplatelet therapy (eg, prasugrel, ticagrelor) for CYP2C19 poor or intermediate metabolizers if there is no contraindication.”
Isosorbide and hydralazine	US FDA	“Hydralazine is metabolized by acetylation . . . About 50% of patients are fast acetylators and have lower exposure” “In patients with heart failure, mean absolute bioavailability of a single oral dose of hydralazine 75 mg varies from 10% to 26%, with the higher percentages in slow acetylators.”
Metoprolol	US FDA	“Metoprolol is metabolized predominantly by CYP2D6, an enzyme that is absent in about 8% of Caucasians (poor metabolizers) and about 2% of most other populations.” “Poor metabolizers and extensive metabolizers who concomitantly use CYP2D6 inhibiting drugs will have increased (several-fold) metoprolol blood levels, decreasing metoprolol’s cardioselectivity.”
Prasugrel	US FDA	“There is no relevant effect of genetic variation in CYP2B6, CYP2C9, CYP2C19, or CYP3A5 on the pharmacokinetics of prasugrel’s active metabolite or its inhibition of platelet aggregation.”
Propafenone	US FDA	“Simultaneous use with both a CYP3A4 and CYP2D6 inhibitor (or in patients with CYP2D6 deficiency) should be avoided.”
Propranolol	US FDA	“In healthy subjects, no difference was observed between CYP2D6 extensive metabolizers (EMs) and poor metabolizers (PMs) with respect to oral clearance or elimination half-life.”
Quinidine	US FDA	“Quinidine is not metabolized by cytochrome P450IID6, but therapeutic serum levels of quinidine inhibit the action of cytochrome P450IID6, effectively converting extensive metabolizers into poor metabolizers. Caution must be exercised whenever quinidine is prescribed together with drugs metabolized by cytochrome P450IID6.”
Ticagrelor	US FDA	“In a genetic substudy cohort of PLATO, the rate of thrombotic CV events in the BRILINTA arm did not depend on CYP2C19 loss of function status.”
Simvastatin	CPIC	“The FDA recommends against 80 mg daily simvastatin dosage. In patients with the C allele at SLC01B1 rs4149056, there are modest increases in myopathy risk even at lower simvastatin doses (40 mg daily); if optimal efficacy is not achieved with a lower dose, alternate agents should be considered.”
Warfarin	US FDA	“CYP2C9 and VKORC1 genotype information, when available, can assist in selection of the initial dose of warfarin.” (A table of expected warfarin maintenance doses based on CYP2C9 and VKORC1 genotypes is provided.)
Warfarin	CPIC	“The best way to estimate the anticipated stable dose of warfarin is to use the algorithms available on http://www.warfarindosing.org .” (This recommendation is from 2011. CPIC guideline authors are aware of more recently published warfarin pharmacogenetic studies and will incorporate them into an updated guideline.)

Data from <http://www.fda.gov/Drugs/ScienceResearch/ResearchAreas/Pharmacogenetics> and with guidelines published by CPIC (www.PharmGKB.org). 2D6 indicates cytochrome P450 family 2 subfamily D member 6; CPIC, Clinical Pharmacogenetics Implementation Consortium of the U.S. National Institutes of Health’s Pharmacogenomics Research Network; CYP2B6, cytochrome P450 family 2 subfamily B member 6; CYP2C9, cytochrome P450 family 2 subfamily C member 9; CYP2C19, cytochrome P450 family 2 subfamily C member 19; CYP2D6, cytochrome P450 family 2 subfamily D member 6; CYP3A4, cytochrome P450 family 3 subfamily A member 4; CYP3A5, cytochrome P450 family 3 subfamily A member 5; P450IID6, cytochrome P450 family 2 subfamily D member 6; PLATO, trial of ticagrelor vs clopidogrel in patients with acute coronary syndromes; SLC01B1, solute carrier organic anion transporter family member 1B1; US FDA, United States Food and Drug Administration; VKORC1, vitamin K epoxide reductase complex subunit 1.

reimbursed for a clopidogrel pharmacogenetic test, with an 85% reimbursement rate.⁹ Although Oemrawsingh et al demonstrated cost-effectiveness for their perindopril pharmacogenetic strategy, cost-effectiveness is often not enough to gain reimbursement. According to case studies published in 2010, the strongest predictor of pharmacogenetic test reimbursement is the strength of available evidence.⁷

In terms of level of evidence, the PERGENE study and idea of pharmacogenetic-guided treatment for angiotensin-converting enzyme (ACE) inhibitors in stable coronary artery disease still needs some additional steps. We feel most observers would agree that the pharmacogenetic risk score presented by Oemrawsingh et al must be, at a minimum,

replicated in an independent data set before it can be taken forward. However, this should be quite achievable, and should be a high-priority research topic. We also do not know that the same pharmacogenetic effect is at play for other ACE inhibitors, though a class effect seems likely. Ideally a validating study could answer both of these questions (ie, testing a distinct patient cohort and a different ACE inhibitor). Another issue is that the mechanisms of the genetic variants used in the pharmacogenetic risk score are unknown (Figure); the authors selected genetic variants based on linkage disequilibrium.³ Moreover, linkage disequilibrium differs depending on ancestry, and thus this pharmacogenetic risk score may not be applicable to patients who are not of




Figure. Agents acting on the renin–angiotensin system pathway ©PharmGKB. Reproduced with permission from the Pharmacogenomics Knowledge Base (PharmGKB) and Stanford University (<https://www.pharmgkb.org/pathway/PA165110622#PGG>).⁵ ACE indicates angiotensin-converting enzyme; AGT, angiotensinogen; AGTR1, angiotensin II receptor type 1; AGTR2, angiotensin II receptor type 2; Ang, angiotensin; ARB, angiotensin receptor blocker; ATP6AP2, ATPase H⁺ transporting accessory protein 2; BDKRB1, bradykinin receptor B1; BDKRB2, bradykinin receptor B2; CMA1, chymase 1; CTSG, cathepsin G; CYP11B2, cytochrome P450 family 11 subfamily B member 2; KNG, kininogen; MAPK1, mitogen-activated protein kinase 1; MAPK3, mitogen-activated protein kinase 3; MAS1, MAS1 proto-oncogene; MME, membrane metallo-endopeptidase; NOS3, nitric oxide synthase 3; NR3C2, nuclear receptor subfamily 3 group C member 2; PK, pharmacokinetic; REN, renin; TGFB1, transforming growth factor β1.

European ancestry. Given this, and the well-recognized racial variation in effectiveness of many cardiovascular medications, additional validation studies of this pharmacogenetic test in patients of non-European ancestry are necessary and would be required prior to contemplating using it for clinical purposes in other racial groups.

Assuming that the Oemrawsingh et al findings can be replicated, should this strategy then be clinically implemented? First consider the myriad of nongenetic patient factors that we constantly use to guide cardiovascular drug therapy without prospective trial data. For example, co-treatment of omeprazole with clopidogrel is often avoided due

to a drug–drug interaction *without prospective trial data*. However, the pharmacogenetic effects on clopidogrel are just as strong as the drug–drug interaction with omeprazole.¹⁰ Warfarin dose is empirically decreased in patients starting amiodarone due to a drug–drug interaction *without prospective trial data*.¹¹ However, the pharmacogenetic effects on warfarin dose are stronger than the drug–drug interaction with amiodarone.¹² Patients are advised to avoid grapefruit juice while taking simvastatin due to a drug–food interaction *without prospective trial data*.¹³ However, the pharmacogenetic effects on simvastatin blood concentrations are just as large as the effects of the drug–food interaction with grapefruit juice.¹⁴ Similar to nongenetic factors used to guide drug therapy, the decision to use pharmacogenetic testing should consider the total risks and benefits, including, for example, the severity of potential side effects averted, or the difference in efficacy, and even the cost/potential savings.

An argument often used against comparing genetic to nongenetic factors to guide drug therapy is that nongenetic factors are usually readily available, but genetic test results are not. However, keep in mind that we also delay drug therapy based on nongenetic factors. For example, the initiation of ACE inhibitor therapy is delayed until the return of laboratory results ruling out renal insufficiency. Moreover, the lack of readily available genetic test results may no longer be an issue since many institutions and laboratories (<http://www.ncbi.nlm.nih.gov/gtr/>) now offer pharmacogenetic testing. More importantly, with the continually decreasing cost and increasing scale of genetic testing, it is anticipated that genetics will ultimately become a routine part of the electronic medical record for all patients. Thus, there are many current situations where genetic information could impact clinical decisions in terms of “best available evidence,” even if it does not meet the commonly held evidence-based medicine standards of having multiple clinical trials on a single intervention.

A second consideration is whether we are holding pharmacogenetics to an unequal standard. In a 2015 Scientific Statement by the American Heart Association,¹⁵ the authors recommended against pharmacogenetic testing for clopidogrel because “. . . no clinical trials assessing the utility of a CYP2C19 genotype test to guide and tailor therapy in a way that leads to improved patient outcomes have been published . . .” However, in the same 2015 Scientific Statement, the authors endorse genetic testing for long QT syndrome, stating it “can potentially help guide patient management,” and they provide recommendations for potential interventions *without prospective trial data*. Moreover, in cancer treatment guidelines, pharmacogenetic recommendations for thiopurine drugs are widely accepted but not informed by a randomized controlled trial.^{16,17}

The standard of evidence required for the clinical implementation of pharmacogenetic testing is widely debated.¹⁸ The “gold standard” of evidence for pharmacogenetic appli-

cations, as stated in a 2012 Special Report on Cardiovascular Pharmacogenomics in the *Journal of the American Heart Association* is “a prospective trial with treatment determined by genotype and with a clinical end point as the primary outcome . . .”¹⁹ The study by Oemrawsingh et al falls short of this because it is a retrospective analysis of a prospective clinical trial. However, the authors of the 2012 Special Report do state “. . . it also recognized that it would be very difficult to sustain and fund a large number of such trials.” This is an understatement; it would be nearly impossible to perform a genetically guided randomized trial for every possible drug and indication, particularly drugs that are already FDA approved. Indeed, a single prospective trial takes several years to complete and costs tens of millions of dollars (or more). The authors of the 2012 Special Report go on to say “. . . in some cases the weight of evidence from observational studies may be so compelling as to render prospective trials unnecessary.”¹⁹ We agree with this assertion and posit that what is needed is more clarity regarding what this level of evidence should look like so that widely acceptable and achievable standards can exist. We would propose that retrospective analyses of multiple prospective clinical trials validating a specified pharmacogenetic approach should be adequate evidence. As a further step, if there is evidence from 1 clinical trial (as in the case of the current study), validation in an independent, adequately sized, and well-phenotyped observational cohort (or vice versa) is likely enough to justify clinical pharmacogenetic implementation. This is of course a debatable point, but with a strong preponderance of evidence, and the best interest of all patients in mind, we feel it may be most satisfactory. The alternative of ignoring congruent data from multiple large studies and simply continuing a standard of care that is wasteful (and even potentially harmful) in an easily identifiable subgroup of patients seems unappealing.

In conclusion, the study by Oemrawsingh et al is an important contribution despite the fact that the pharmacogenetic association is not new. It spotlights a practical path forward to precision medicine for ACE inhibitors in stable coronary artery disease, and it quantifies the motivation for doing so in terms of clinical and cost-effectiveness. The major limitation of the study is the lack of a sufficient replication data set. If validation testing can be achieved, then clinical care can be fundamentally altered. While there are other scientific questions remaining regarding the mechanisms of the genetic variants, the alternative therapy in patients with a pharmacogenetic risk score >2, and application in non-European ancestry patients, these can be addressed in additional future studies. More important is to hear this as a wake-up call so that the clinical and scientific communities can reach a better consensus regarding when and how to put findings like these into action clinically. Improved care for our patients is waiting.

Disclosures

Luzum is funded by a Post-Doctoral Fellowship from the American Heart Association (14POST20100054) and the National Institutes of Health student loan repayment program (L30 HL110279). Luzum is a consultant for Gnome Diagnostics, LLC. Lanfear is supported by the National Heart, Lung, and Blood Institute (R01 HL103871) and has other genetic research grants funded by Janssen and Amgen.

References

- Oemrawsingh RM, Akkerhuis KM, Van Vark LC, Redekop WK, Rudez G, Remme WJ, Bertrand ME, Fox KM, Ferrari R, Jan Danser AH, de Maat M, Simoons ML, Brugts JJ, Boersma E; on behalf of the PERGENE Investigators. Individualized angiotensin-converting enzyme (ACE)-inhibitor therapy in stable coronary artery disease based on clinical and pharmacogenetic determinants: the PERindopril GENETic (PERGENE) risk model. *J Am Heart Assoc*. 2016;5:e002688 doi: 10.1161/JAHA.115.002688.
- Deckers JW, Goedhart DM, Boersma E, Briggs A, Bertrand M, Ferrari R, Remme WJ, Fox K, Simoons ML. Treatment benefit by perindopril in patients with stable coronary artery disease at different levels of risk. *Eur Heart J*. 2006;27:796–801.
- Brugts JJ, Isaacs A, Boersma E, van Duijn CM, Uitterlinden AG, Remme W, Bertrand M, Ninomiya T, Ceconi C, Chalmers J, MacMahon S, Fox K, Ferrari R, Wittman JC, Danser AH, Simoons ML, de Maat MP. Genetic determinants of treatment benefit of the angiotensin-converting enzyme-inhibitor perindopril in patients with stable coronary artery disease. *Eur Heart J*. 2010;31:1854–1864.
- United States Food and Drug Administration. Table of Pharmacogenomic Biomarkers in Drug Labeling. 2015. Available at: <http://www.fda.gov/20drugs/scienceresearch/researchareas/pharmacogenetics/ucm083378.htm>. Accessed March 14, 2016.
- Whirl-Carrillo M, McDonagh EM, Hebert JM, Gong L, Sangkuhl K, Thorn CF, Altman RB, Klein TE. Pharmacogenomics knowledge for personalized medicine. *Clin Pharmacol Ther*. 2012;92:414–417.
- Shuldiner AR, Relling MV, Peterson JF, Hicks JK, Freimuth RR, Sadee W, Pereira NL, Roden DM, Johnson JA, Klein TE; Pharmacogenomics Research Network Translational Pharmacogenetics Program Group. The Pharmacogenomics Research Network Translational Pharmacogenetics Program: overcoming challenges of real-world implementation. *Clin Pharmacol Ther*. 2013;94:207–210.
- Meckley LM, Neumann PJ. Personalized medicine: factors influencing reimbursement. *Health Policy*. 2010;94:91–100.
- Horgan D, Jansen M, Leyens L, Lal JA, Sudbrak R, Hackenitz E, Bußhoff U, Ballensiefen W, Brand A. An index of barriers for the implementation of personalised medicine and pharmacogenomics in Europe. *Public Health Genomics*. 2014;17:287–298.

- Weitzel KW, Elsei AR, Langae TY, Burkley B, Nessler DR, Obeng AO, Staley BJ, Dong HJ, Allan RW, Liu JF, Cooper-Dehoff RM, Anderson RD, Conlon M, Clare-Salzler MJ, Nelson DR, Johnson JA. Clinical pharmacogenetics implementation: approaches, successes, and challenges. *Am J Med Genet C Semin Med Genet*. 2014;166C:56–67.
- Mega JL, Close SL, Wiviott SD, Shen L, Hockett RD, Brandt JT, Walker JR, Antman EM, Macias W, Braunwald E, Sabatine MS. Cytochrome p-450 polymorphisms and response to clopidogrel. *N Engl J Med*. 2009;360:354–362.
- Bungard TJ, Yakiwchuk E, Foisy M, Brocklebank C. Drug interactions involving warfarin: practice tool and practical management tips. *Can Pharm J*. 2011;144:21–25.e9.
- Gage BF, Eby C, Johnson JA, Deych E, Rieder MJ, Ridker PM, Milligan PE, Grice G, Lenzini P, Rettie AE, Aquilante CL, Grosso L, Marsh S, Langae T, Farnett LE, Voora D, Veenstra DL, Glynn RJ, Barrett A, McLeod HL. Use of pharmacogenetic and clinical factors to predict the therapeutic dose of warfarin. *Clin Pharmacol Ther*. 2008;84:326–331.
- Merck Sharp & Dohme Corp. Simvastatin prescribing information. Copyright©1999–2014.
- Pasanen MK, Neuvonen M, Neuvonen PJ, Niemi M. SLCO1B1 polymorphism markedly affects the pharmacokinetics of simvastatin acid. *Pharmacogenet Genomics*. 2006;16:873–879.
- Musunuru K, Hickey KT, Al-Khatib SM, Delles C, Fornage M, Fox CS, Frazier L, Gelb BD, Herrington DM, Lanfear DE, Rosand J. Basic concepts and potential applications of genetics and genomics for cardiovascular and stroke clinicians: a scientific statement from the American Heart Association. *Circ Cardiovasc Genet*. 2015;8:216–242.
- Relling MV, Gardner EE, Sandborn WJ, Schmiegelow K, Pui CH, Yee SW, Stein CM, Carrillo M, Evans WE, Klein TE; Clinical Pharmacogenetics Implementation Consortium. Clinical pharmacogenetics implementation consortium guidelines for thiopurine methyltransferase genotype and thiopurine dosing. *Clin Pharmacol Ther*. 2011;89:387–391.
- Alvarnas JC, Brown PA, Aoun P, Ballen KK, Barta SK, Borate U, Boyer MW, Burke PW, Cassaday R, Castro JE, Coccia PF, Coutre SE, Damon LE, DeAngelo DJ, Douer D, Frankfurt O, Greer JP, Johnson RA, Kantarjian HM, Klisovic RB, Kupfer G, Litzow M, Liu A, Rao AV, Shah B, Uy GL, Wang ES, Zelenetz AD, Gregory K, Smith C. Acute lymphoblastic leukemia, version 2.2015. *J Natl Compr Canc Netw*. 2015;13:1240–1279.
- Ratain MJ, Johnson JA. Meaningful use of pharmacogenetics. *Clin Pharmacol Ther*. 2014;96:650–652.
- Musunuru K, Roden DM, Boineau R, Bristow MR, McCaffrey TA, Newton-Cheh C, Palttoo DN, Rosenberg Y, Wohlgenuth JG, Zineh I, Hasan AA. Cardiovascular pharmacogenomics: current status and future directions—report of a National Heart, Lung, and Blood Institute Working Group. *J Am Heart Assoc*. 2012;1:e000554 doi: 10.1161/JAHA.111.000554.

Key Words: Editorials • angiotensin-converting enzyme inhibitor • cardiovascular disease • pharmacogenetics