

Henry Ford Health System

## Henry Ford Health System Scholarly Commons

---

Clinical Research

Medical Education Research Forum 2019

---

5-2019

### Feasibility of Omitting Outer Renorrhaphy During Robotic Partial Nephrectomy

Sohrab Arora

Henry Ford Health System, sarora3@hfhs.org

Chandler Bronkema

Henry Ford Health System

James R. Porter

Alexander Mottrie

Mani Menon

Henry Ford Health System, MMENON1@hfhs.org

*See next page for additional authors*

Follow this and additional works at: <https://scholarlycommons.henryford.com/merf2019clinres>

---

#### Recommended Citation

Arora, Sohrab; Bronkema, Chandler; Porter, James R.; Mottrie, Alexander; Menon, Mani; Rogers, Craig G.; Jeong, Wooju; Dasgupta, Prokar; Bhandari, Mahendra; and Abdollah, Firas, "Feasibility of Omitting Outer Renorrhaphy During Robotic Partial Nephrectomy" (2019). *Clinical Research*. 15.

<https://scholarlycommons.henryford.com/merf2019clinres/15>

This Poster is brought to you for free and open access by the Medical Education Research Forum 2019 at Henry Ford Health System Scholarly Commons. It has been accepted for inclusion in Clinical Research by an authorized administrator of Henry Ford Health System Scholarly Commons.

---

**Authors**

Sohrab Arora, Chandler Bronkema, James R. Porter, Alexander Mottrie, Mani Menon, Craig G. Rogers, Wooju Jeong, Prokar Dasgupta, Mahendra Bhandari, and Firas Abdollah



## Feasibility of omitting outer (cortical) renorrhaphy during robotic partial nephrectomy - A multi-institutional analysis

Sohrab Arora<sup>a</sup>, Chandler Bronkema<sup>a,b</sup>, James R. Porter<sup>c</sup>, Alexander Mottrie<sup>d</sup>, Mani Menon<sup>a</sup>, Craig Rogers<sup>a</sup>, Wooju Jeong<sup>a</sup>, Prokar Dasgupta<sup>e</sup>, Koon H Rhaf<sup>f</sup>, Rajesh K Ahlawat<sup>g</sup>, Umberto Capitanio<sup>h</sup>, TB Yuvaraja<sup>i</sup>, Sudhir Rawal<sup>j</sup>, Daniel A. Moon<sup>k</sup>, Ananthkrishnan Sivaraman<sup>l</sup>, Kris K. Maes<sup>m</sup>, Fansesco Porgiglia<sup>n</sup>, Gagan Gautam<sup>o</sup>, Levent Turkeri<sup>p</sup>, Mahendra Bhandari<sup>a</sup>, and Firas Abdollah<sup>a</sup>

<sup>a</sup> Vattikuti Urology Institute, Henry Ford Hospital, Detroit, MI, USA, <sup>b</sup> Wayne State University School of Medicine, Detroit, MI, USA, <sup>c</sup> Swedish Medical Center, Seattle, WA, USA, <sup>d</sup> OLV Vattikuti Robotic Surgery Institute, Melle, Belgium, <sup>e</sup> MRC Centre for Transplantation, King's College London, London, UK, <sup>f</sup> Yonsei University Health system, Seoul, South Korea, <sup>g</sup> Medanta Kidney and Urology Institute, Gurgaon, India, <sup>h</sup> Urology Clinic, San Raffaele Hospital, Milan, Italy, <sup>i</sup> Kokilaben Dhirubhai Ambani Hospital, Mumbai, India, <sup>j</sup> Rajiv Gandhi Cancer Institute and Research Centre, New Delhi, India, <sup>k</sup> Peter MacCallum Cancer Centre, Melbourne, Victoria, Australia, <sup>l</sup> Apollo hospitals, Chennai, <sup>m</sup> Center for Robotic and Minimally Invasive Surgery, Hospital Da Luz, Luz Sáude, Portugal, <sup>n</sup> San Luigi Gonzaga University Hospital, Torino, Italy, <sup>o</sup> Max Institute of Cancer Care, New Delhi, India, <sup>p</sup> Acibadem Hospitals Group, Istanbul, Turkey,

# Feasibility of omitting outer (cortical) renorrhaphy during robotic partial nephrectomy - A multi-institutional analysis


## Introduction

- Vattikuti Collective Quality Initiative (VCQI):
  - 41 surgeons
  - 14 centers
  - 9 countries
- Reconstruction technique after robotic partial nephrectomy (RPN) has recently been shown to be a modifiable factor with possible impacts on ischemia time, postoperative bleeding, renal function, and incidence of pseudoaneurysms after surgery.



## Feasibility of omitting outer (cortical) renorrhaphy during robotic partial nephrectomy - A multi-institutional analysis


- Recent literature comparing single versus double layer (cortical) renorrhaphy stems primarily from single institution studies.
- There are currently no randomized trials.
- Objective: to evaluate the feasibility of omitting cortical renorrhaphy in a multi-institutional setting.



\*Standard double layer renorrhaphy technique  
Credit: Laydner H, Kaouk JH. Robotic partial nephrectomy:  
The new horizon. *Arab J Urol.* 2012;10(1):2-9

## Methods

- Inverse probability of treatment weighting (IPTW) was performed to minimize selection bias by adjusting for several preoperative factors.
- Firth correction was applied to the data model to account for center-specific practices.
- Perioperative outcomes were compared between matched cohorts.



## Feasibility of omitting outer (cortical) renorrhaphy during robotic partial nephrectomy - A multi-institutional analysis



Variable	Double Layer	Single Layer	p value
Operative time, minutes, median (IQR)	168 (140-203)	162 (140-202)	0.2
Ischemia time, minutes, median (IQR)	18 (14-22)	17 (13-20)	0.7
Estimated blood loss, mL, median (IQR)	100 (50-200)	100 (50-200)	0.6
% drop in eGFR, ml/min/1.73m <sup>2</sup> , median (IQR)	7.3 (16.9-1.4)	10.4 (17.5-3.6)	0.9
Intraoperative complications, %	7.4	8.9	0.6
Postoperative Clavien grade ≥ 3 complications, %	1.0	2.8	0.2
Positive surgical margin, %	2.2	3.2	0.6
Hospital stay, days, median (IQR)	3 (2-4)	2 (2-4)	0.4
Need for angioembolization within 1 year, %	0.7	1.4	0.4

### Conclusion:

Omission of cortical renorrhaphy did not significantly improve operative or ischemia time; however, it also had no adverse effect on perioperative outcomes after RPN in a multi-institutional setting.

- Won best poster prize at EAU Barcelona
- Being presented at AUA annual meeting, Chicago as we speak.