

21st International Conference on Virus and other Graft Transmissible Diseases of Fruit Crops

## Assessment of susceptibility to European stone fruit yellows phytoplasma of new plum variety and five rootstock/plum variety combinations

Landi, F.<sup>1</sup>, Prandini, A.<sup>1</sup>, Paltrinieri, S.<sup>1</sup>, Missere, D.<sup>2</sup>, Bertaccini, A.<sup>1</sup><sup>1</sup> DiSTA, Patologia vegetale, *Alma Mater Studiorum*, Università di Bologna, viale Fanin 42, 40127 Bologna. Italy<sup>2</sup> Centro Ricerche Produzioni Vegetali, Via dell'Arrigoni 120, 47522 Cesena (FC) - Italy

### Abstract

Two separate experiments were carried out to assess the plum susceptibility to infection by European stone fruit yellows phytoplasmas during a five years period. Commercial varieties/cultivars and new selections grafted on Myrabolan 29C were evaluated in at least two plots of four plants each. Visual inspection and PCR/RFLP identification of phytoplasmas detected an increasing phytoplasma presence in both symptomatic and asymptomatic plants. Eight Japanese plum selections showed ESFY symptoms or pathogen presence in the 50% of the plants and nine selections showed ESFY infection in 20% of the plants. Only nine selections showed absence of both symptoms and pathogen. Although the European selections/cultivars were not symptomatic, plants belonging to six of these cultivars were positive for phytoplasma infection. The evaluation of cultivar/rootstock combinations indicate phytoplasma presence from the first year after plantation on. Two of the rootstocks seem to induce a delay in symptoms appearance and cultivar T.C. Sun resulted to be the most susceptible to the disease independently from the rootstock employed.

Keywords: Japanese plum, European plum, European stone fruit yellows phytoplasmas, resistance, diseases

### Introduction

In the last thirty years an increasing presence of European stone fruit yellows phytoplasma (ESFY, '*Candidatus* Phytoplasma prunorum') (Seemüller and Schneider, 2004) on Japanese plum (*Prunus salicina*) was observed in commercial orchards in several European regions (Giunchedi et al., 1978; Desvignes and Cornaggia, 1982; Dosba et al., 1991; Torres et al., 2004). *Prunus* rootstocks (Jarausch et al., 1998) as well as wild *Prunus* species, e.g. *Prunus spinosa* and *P. cerasifera* (Carraro et al., 2002) and cherry (*Prunus avium*) (Paltrinieri et al., 2001) were also reported as infected by these phytoplasmas. In recent years ESFY phytoplasma has been detected in other wild plants such as *Rosa canina*, *Celtis australis*, *Fraxinus excelsior* (Jarausch et al., 2001), and grapevine in Hungary (Varga et al., 2000) and in Serbia (Duduk et al., 2004). The production losses can reach 40% in Japanese plum (Poggi Pollini et al., 1995; Pastore et al., 1999) due to the lack of sanitary control of propagation materials together with the presence of the ESFY-specific vector, *Cacopsylla pruni* (Carraro et al., 1998b) in the orchards. The most evident symptoms on Japanese plum consist of black phloem and leaf malformations such as excessive elongation and upward rolling. European plum cultivars are usually infected only in a latent way (Carraro et al., 1998a). Diseased plants start producing flowers during mild winters, and show then undersized leaves with very little fruit production. The disease usually starts from a branch, affecting within 2-3 years all the plant; infected plants also show abnormal rootstock proliferation. The correct choice of cultivar/rootstock combination that must be tested for resistance/tolerance to the disease is also very important before distributing new cultivars (Desvignes, 1999). During 2003-2008 a research was carried out to assess the ESFY plum susceptibility under natural conditions of disease spreading by visual inspection coupled with molecular analyses.

### Material and methods

Two experiments were carried out to verify susceptibility of about 60 plum varieties/cultivars and new selections (A), and of 5 routinely used rootstock/scion combinations (B). The orchards were located in an ESFY severely naturally infected area of Northern Italy (Paltrinieri et al., 2004). Monitoring by visual inspection and PCR/RFLP assays was carried every year out from August to October to verify phytoplasma presence/identity in all plants.

**Experiment A:** Japanese and European plum varieties and selections grafted into Myrabolan 29C (Table 1 A and B) were evaluated in at least two plots with four plants each.

**Experiment B:** Three plum commercial cultivars TC Sun, Fortune and Angeleno were grafted on the five rootstocks 'Adesoto 101', 'Ishtara-Ferciana', 'GF 677', 'Montclair-Chanturgue' and 'Myrabolan 29C'; four plants per each combination per at least two plots were employed.

**Tab. 1** A. Results of ESFY phytoplasma identification in Japanese plum cultivars/varieties and selections. B. Results of ESFY phytoplasma identification in European plum varieties/cultivars and selections. In both cases plants were grown in a field located in a severely ESFY naturally infected area.

Japanese plum cultivars and selections			European plum cultivars and selections		
		n. of ESFY infected plants/n. tested			n. of ESFY infected plants/n. tested
Table A	Symptoms	plants	Table B	Symptoms	plants
Anne Gold	yes	2/4	Bellamira	no	0/4
Aphrodite	yes	1/4	Capitana	no	0/4
Black Glow	yes	1/3	Elena	no	0/4
Black Sunrise	yes	3/4	Felsina	no	0/4
Black Top	no	2/4	Grossa Di Felisio	no	0/4
Bragialla	no	0/4	Jojo	no	0/4
Brarossa	no	0/4	Liablu	no	0/4
Carmen Blu	yes	2/4	Maria Novella	no	0/4
Dofi Sandra	yes	0/3	Presenta	no	0/4
Early Fortune	yes	1/4	President	no	0/4
Fortune	no	0/4	Rheingold	yes	1/4
Gaia	no	1/4	Stanley	no	0/4
Golden Plumza	yes	1/4	Tegera	no	0/4
Obilnaja	no	1/4	Tipala	no	0/4
Red Noble	yes	4/4	Top 2000	no	0/4
Ruby Crunch	no	0/3	Topend Plus	no	0/4
Shiro	yes	1/4	Topfive	no	0/4
Dofi selections (Florence University)			Topgigant Plus	no	0/4
89.024.004	yes	2/4	Tophit	no	0/4
89.024.029	no	1/4	Tophit Plus	no	0/4
89.028.047	no	1/4	Topking	no	0/4
89.030.010	no	1/4	Topstar Plus	no	0/4
89.030.020	no	0/4	Valcean	yes	3/4
89.030.030	yes	1/4	Valerie	yes	1/4
89.030.031	no	0/4	Victory	no	0/4
89.036.131	no	0/4	Agri 2000 selections		
CRA Forli' selections			N. 8	no	0/4
IFF on 219	yes	4/4	N. 10	no	0/4
IFF on 221	yes	4/4	Hohenheim University selections		
IFF on 260	no	0/4	N. 1218	no	0/4
IFF on 268	no	3/3	N. 1446	no	0/4
IFF on 271	no	0/4	N. 1462	no	0/4
			N. 1464	no	0/4
			N. 1468	no	0/4
			N. 1474	no	1/4
			N. 1632	no	0/4
			N. 3018	yes	3/4
			N. 3217	no	0/4
			N. 4913	no	0/4

**Molecular analyses:** DNA was extracted from fresh leaf midribs and phloem by a chloroform/phenol procedure (Prince et al., 1993). PCR assays were carried out on the nucleic acid samples diluted in TE buffer [10 mM Tris-HCl, 1 mM EDTA (pH 8.0)] to give a final concentration of 20 ng per  $\mu$ l, in total 25  $\mu$ l reaction mixtures under the conditions described by Schaff et al. (1992). Nested PCR reactions were performed under the same conditions, using as template the products of the previous amplification diluted 1: 30 with sterile water. Positive control samples were DNAs extracted from periwinkle plants infected by phytoplasma strains from the micropropagated collection of DiSTA (University of Bologna), in particular GSFY1, GSFY2 (subgroup 16SrX-B), PD (subgroup 16SrX-C) and AP (subgroup 16SrX-A) were employed.


Phytoplasma primers P1/P7 (Deng and Hiruki, 1991; Schneider et al., 1995) and R16F2/R2 (Lee et al., 1995) amplifying fragments internal to each others in the 16S ribosomal gene were used, respectively in direct and nested PCR. Samples were further tested in nested PCR with primers R16(X)F1/R1 specific for 16SrX (apple proliferation) group (Lee et al., 1994; 1995). Samples with the reaction mixture devoid of DNA template were included in each experiment as negative controls. PCR products were subjected to electrophoresis in a 1% agarose gel and visualised by staining with ethidium bromide and UV illumination.

Three to six µl of PCR products were digested using *MseI*, *SspI* and *RsaI* restriction enzymes at 37°C for at least 16 hours following the instructions of the manufacturer (Fermentas, Vilnius, Lithuania). The restriction patterns were then compared with those of reference strains after electrophoresis through a 5% polyacrylamide gel in 1X TBE buffer followed by staining with ethidium bromide and visualization under an UV transilluminator.


## Results and discussion

**Experiment A:** After five years 8 Japanese plum selections showed ESFY symptoms or pathogen presence in 50% of the plants and nine selections showed ESFY infection in 20% of the plants. Only 9 selections showed absence of both symptoms and pathogen (Table 1A). None of the European selections/cultivars employed was symptomatic, however plants belonging to 6 cultivars were positive for phytoplasma presence (Table 1B). The comparison of symptom expression and results from PCR/RFLP analyses indicates that selections IFF on 221 and IFF on 219, are very susceptible to ESFY infection; while selection IFF on 268 resulted to be phytoplasma infected even if asymptomatic. A medium level of susceptibility i.e. pathogen or symptom presence starting from the second year after plantation in 50% of the plants, was shown by Anne Gold, Black Sunrise, Black Top, Carmen Blue, Dofi Sandra, Gaia, Obilnaja and Dofi 89.024.004 among the Japanese cultivar and varieties. Less susceptible were Shiro, Black Glow, Dofi 89.030.030, 89.024.029, 89.028.047, 89.030.010, Aphrodite, Early Fortune, and Golden Plumza (one infected plant per cultivar). Varieties Bragialla, Brarossa, Fortune, Ruby Crunch and selections Dofi 80.030.020, 89.030.031, 89.030.131 and IFF on 260 and on 271 were always negative to both symptom presence and molecular analyses. Some European plum cultivars or selections such as Rheingold, Presenta, Valerie, Valcean, 3018 and 1474 were positive to the ESFY presence, however only Rheingold, Valerie, Valcean and selection 3018 showed symptoms possibly related to phytoplasma presence.

**Experiment B:** The majority of the cultivar/rootstocks combinations showed phytoplasma symptoms and were positive to the PCR/RFLP analyses from the first year after plantation. Two of the rootstocks induced a delay in symptoms appearance indicating some resistance to ESFY in Japanese plum could be present, but only for one-two years after plantation. Intermediate/high susceptibility was detected in varieties grafted on 'GF 677', 'Adesoto 101' and 'Montclair-Chanturgue' showing plants with severe leaf symptoms and reduced growth; while scions grafted on 'Myrabolan 29/C' were efficiently vegetating and growing in spite of the presence of symptoms in the leaves. TC Sun was the most susceptible cultivar to ESFY phytoplasmas in all tested rootstocks; cultivars grafted on 'Ishtara-Ferciana' showed symptoms only after four years from plantation (Figs. 1 and 2) indicating less susceptibility to ESFY infection. The molecular analyses carried out were very important to identify the latently infected combinations that are the most dangerous allowing the pathogen to survive long time in orchards causing recurrent epidemic outbreaks.



**Fig. 1** Number of infected plants for the different cultivar/rootstock combinations



**Fig. 2** Number of symptomatic plum plants per year grouped by rootstock.

## Literature

- Carraro, L.; Loi, N.; Emarcora, P.; Osler, R.; 1998a: High tolerance of European plum varieties to plum leptonecrosis. *European Journal of Plant Pathology* **104**, 141-145.
- Carraro, L.; Osler, R.; Loi, N.; Emarcora, P.; Refatti, E.; 1998b: Transmission of European stone fruit yellows phytoplasma by *Cacopsylla pruni*. *Journal of Phytopathology* **80**, 233-239.
- Carraro, L.; Ferrini, F.; Emarcora, P.; Loi, N.; 2002: Role of wild *Prunus* species in the epidemiology of European stone fruit yellows. *Plant Pathology* **51**, 513-517.
- Deng, S.; Hiruki, C.; 1991: Genetic relatedness between two nonculturable mycoplasma-like organisms revealed by nucleic acid hybridization and polymerase chain reaction. *Phytopathology* **81**, 1475-1479.
- Desvignes, J.C.; Cornaggia, D.; 1982: Observations on apricot chlorotic leaf roll: sensitiveness of different *Prunus* species, detection, spread in plum orchards. *Acta Horticulturae* **130**, 249-256.
- Desvignes, J.C.; 1999: *Virus Diseases of Fruit Trees*. CTIFL editor, 113-143.
- Dosba, F.; Lansac, M.; Mazy, K.; Garnier, M.; Eyquard, P.J.; 1991: Incidence of different diseases associated with mycoplasma-like organisms in different species of *Prunus*. *Acta Horticulturae* **283**, 311-320.
- Duduk, B.; Botti, S.; Ivanović, M.; Krstić, B.; Dukić, N.; Bertaccini, A.; 2004: Identification of phytoplasmas associated with grapevine yellows in Serbia. *Journal of Phytopathology* **152**, 575-579.
- Giunchedi, L.; Marani, F.; Credi, R.; 1978: Mycoplasma-like bodies associated with plum decline (leptonecrosis). *Phytopathologia Mediterranea* **17**, 205-209.
- Jarausch, W.; Lansac, M.; Saillard, C.; Broquaire, J.M.; Dosba, F.; 1998: PCR assays for specific detection of European stone fruit yellows phytoplasmas and its use for epidemiological studies in France. *European Journal of Plant Pathology* **104**, 17-27.
- Jarausch, W.; Jarausch-Wehrheim, B.; Danet, J.L.; Broquaire, J.M.; Dosba, F.; Saillard, C.; Garnier, M.; 2001: Detection and identification of European stone fruit yellows and other phytoplasmas in wild plants in the surroundings of apricot chlorotic leafroll-affected orchards in southern France. *European Journal of Plant Pathology* **107**, 209-217.
- Lee, I.-M.; Gundersen, D.E.; Hammond, R.W.; Davis, R.E.; 1994: Use of mycoplasma-like organism (MLO) group-specific oligonucleotide primers for nested-PCR assays to detect mixed-MLO infections in a single host plant. *Phytopathology* **84**, 559-566.
- Lee, I.-M.; Bertaccini, A.; Vibio, M.; Gundersen, D.E.; 1995: Detection of multiple phytoplasmas in perennial fruit trees with decline symptoms in Italy. *Phytopathology* **85**, 728-735.
- Paltrinieri, S.; Martini, M.; Stefani, E.; Pondrelli, M.; Fideghelli, C.; Bertaccini, A.; 2001: Phytoplasma infection in peach and cherry in Italy. *Acta Horticulturae* **550**, 365-370.
- Paltrinieri, S.; Lugli, A.; Monari, W.; Bertaccini, A.; 2004: Three years of molecular monitoring of phytoplasma spreading in a plum growing area in Italy. *Acta Horticulturae* **657**, 501-506.
- Pastore, M.; Tian, J.B.; Paltrinieri, S.; Martini, M.; Simeone, A.M.; Santonastaso, M.; Bertaccini, A.; 1999: Incidence of European stone fruit yellows phytoplasma natural infection and of its experimental transmission in different Plum varieties. *Journal of Plant Pathology* **82**, 72.

- Poggi Pollini, C.; Bissani, R.; Giunchedi, L.; Vindimian, E.; 1995: Occurrence of phytoplasma infection in European Plums (*Prunus domestica*). *Journal of Phytopathology* **143**, 701-703.
- Prince, J.P.; Davis, R.E.; Wolf, T.K.; Lee, I.-M.; Mogen, B.D.; Dally, E.L.; Bertaccini, A.; Credi, R.; Barba, M.; 1993: Molecular detection of diverse mycoplasma-like organisms (MLOs) associated with grapevine yellows and their classification with aster yellows, X-disease, and elm yellows MLOs. *Phytopathology* **83**, 1130-1137.
- Schaff, D.A.; Lee, I.-M.; Davis, R.E.; 1992. Sensitive detection and identification of mycoplasma-like organisms by polymerase chain reactions. *Biochemical Biophysics Research Communications* **186**, 1503-1509.
- Schneider, B.; Seemüller, E.; Smart, C.D., Kirkpatrick, B.C.; 1995: Phylogenetic classification of plant pathogenic mycoplasma-like organisms or phytoplasmas. In: Razin R and Tully JG (eds), *Molecular and Diagnostic Procedures in Mycoplasmaology*, Vol. I, pp. 369-380, San Diego: Academic Press.
- Seemüller, E.; Schneider, B.; 2004: '*Candidatus* Phytoplasma mali', '*Candidatus* Phytoplasma pyri', '*Candidatus* Phytoplasma prunorum', the causal agents of apple proliferation, pear decline and European stone fruit yellows, respectively. *International Journal of Systematic and Evolutionary Microbiology* **54**, 1217-1226.
- Torres, E.; Martín, M.P.; Paltrinieri, S.; Vila, A.; Masalles, R.; Bertaccini, A.; 2004: Spreading of ESFY phytoplasmas in stone fruit in Catalonia (Spain). *Journal of Phytopathology* **152**, 432-437.
- Varga, K.; Kölber, M.; Martini, M.; Pondrelli, M.; Ember, I.; Tökés, G.; Lázár, J.; Mikulás, J.; Papp, E.; Szendrey, G.; Schweigert, A.; Bertaccini, A.; 2000: Phytoplasma identification in Hungarian grapevines by two nested-PCR systems. In: Extended abstracts of XIIIth meeting of the International Council for the Study of viruses and virus-like diseases of the grapevine (ICVG). Adelaide, Australia 12-17 March, 113-115.