

12th International Working Conference on Stored Product Protection (IWCSPP) in Berlin, Germany, October 7-11, 2018

- OPIT, G. P. AND J. E. THRONE, 2009. Population growth and development of the psocid *Liposcelis brunnea* (Psocoptera: Liposcelidae) at constant temperatures and relative humidities. *Journal of Economic Entomology* **102**, 1360–1368.
- OPIT, G., ARTHUR, F. H., BONJOUR, E. L., JONES, C. L., AND PHILIPS, T. W., 2011. Efficacy of heat treatment for disinfection of concrete grain silos. *Journal of Economic Entomology* **104**, 1415–1422.
- REES, D. P. AND A. J. WALKER, 1990. The effect of temperature and relative humidity on population growth of three *Liposcelis* species (Psocoptera: Liposcelidae) infesting stored products in tropical countries. *Bulletin of Entomological Research* **80**, 353–358.
- SAS INSTITUTE, 2014. The SAS system for Windows, version 9.2. SAS Institute, Cary, NC.
- SUMMERS, C. G., COVIELLO, R. L., AND GUTIERREZ, A. P., 1984. Influence of constant temperature on the development and reproduction of *Acyrtosiphon kondoi* (Homoptera: Aphididae). *Environmental Entomology* **13**, 236–242.
- SYSTAT SOFTWARE, INC., 2002A. TableCurve 3D, version 4.0. Systat Software Inc., San Jose, CA.
- SYSTAT SOFTWARE, INC., 2002B. TableCurve 2D, version 5.01. Systat Software Inc., San Jose, CA.
- SYSTAT SOFTWARE, INC., 2006. Sigma Plot, version 10.0. Systat Software Inc., San Jose, CA.
- TANG, P. A., WANG, J. J., HE, Y., JIANG, H. B., AND WANG, Z. Y., 2008. Development, survival, and reproduction of the psocid *Liposcelis decolor* (Psocoptera: Liposcelidae) at constant temperatures. *Annals of the Entomological Society of America* **101**, 1017–1025.
- WANG, J. J., ZHAO, Z. M., AND LI, L. S., 1999. Induced tolerance of the psocid *Liposcelis bostrychophila* (Psocoptera: Liposcelidae) to controlled atmosphere. *International Journal of Pest Management* **45**, 75–79.
- WANG, J. J., TSAI, J. H., ZHAO, Z. M., AND LI, L. S., 2000. Development and reproduction of the psocid *Liposcelis bostrychophila* (Psocoptera: Liposcelidae) as a function of temperature. *Annals of the Entomological Society of America* **93**, 261–270.
- WANG, J. J., DONG, P., XIAO, L. S., AND DOU, W., 2008. Effects of removal of *Cardinium* infection on fitness of the stored-product pest *Liposcelis bostrychophila* (Psocoptera: Liposcelidae). *Journal of Economic Entomology* **101**, 1711–1717.
- WANG, J. J., REN, Y., WEI, X. Q., AND DOU, W., 2009. Development, survival, and reproduction of the psocid *Liposcelis paeta* (Psocoptera: Liposcelidae) as a function of temperature. *Journal of Economic Entomology* **102**, 1705–1713.
- ZILAH-BALOGH, G. AND D. PFEIFFER. 1998. Understanding degree-days and using them in pest management decision making. Project for ENT 4987.

Circadian Rhythm of *Liposcelis entomophila* and *Liposcelis paeta* in Paddy Warehouse

Zhenjun Zhang¹, Yanyu Li¹, Zhongming Wang¹, Yang Cao^{1*}, Yanmei Qi², Derong Pan³, Rui He⁴

¹Academy of State Administration of Grain, Beijing 100073;

²Beilun Grain General Corporation, Ningbo, 315800;

³Nanning Grain Reserve Management Corporation, Nanning, 530031;

⁴Zhanjiang North Station National Grain Reserve Depot, Zhanjiang, 524043

Corresponding author: cy@chinagrains.org

DOI 10.5073/jka.2018.463.038

Abstract

Booklice is a small but serious stored grain pest, and understanding the circadian rhythm of booklice help to control. In this study, circadian activity of booklice were monitored with sticky traps in the grain bulk surfaces of two warehouses stored paddy rice in two different provinces in China. The results showed that the species of booklice were different and were *Liposcelis entomophila*, and *Liposcelis paeta* for Nanning's and Zhanjiang's warehouses respectively. In term of *L.entomophila*, its activity intensity gradually decreased from 0 am to 12 pm and reached the lowest level of daily activity at 12pm. After this, there was a steady and straight upward trend, and the peak of its activity intensity is reached at 8 pm. Its circadian activity trend can be represented as: $y = -0.971x^3 + 21.88x^2 - 139.5x + 353.4$ (x: time; y: quantity of booklice). Over the same period, the activity intensity of *L.paeta* varied greatly. It gradually increased, reached a peak at 8 am, dropped dramatically at 12 pm and then climbed the second peak at 6 pm.

Keyword: sticky trap, monitor, *L.entomophila*, *L.paeta*, circadian rhythm

1. Introduction

In control of stored grain pests, insect population dynamics monitoring and density inspection are important. The species, density, distribution, and damage status data of grain stored pests in the grain bulk can be timely detected, predicting the development trend of insects, avoiding unnecessary prevention cost and the economic losses, and providing scientific strategy for insect control (Bai Xuguang, 2002).

At present, the pest detection technology is traditional screening method. There are many disadvantages, such as, labor intensive, low efficiency, and imprecise. Base on these, trapping detections are developed. They are convenient, fast, environmentally friendly, and highly automated (LI Zhishen, 2014) . Sticky board trapping is one of these technologies and has been widely used in the monitoring and control of agricultural and forestry pests. However, there are few reports in stored grain insect pests. In this study, the sticky board trapping technology was applied to monitor the population dynamics of the booklice in two paddy warehouses in Guangdong and Guangxi, southern China. The aim in work reported in this publication was to assess sticky board trapping as an alternative to detection.

2. Materials and Methods

2.1 Test Warehouse

No. 32 large warehouses in Nanning Shajing Grain Warehouse examined was a length of 26.69 m and a width of 19.73 m, storage ability 1250t. The stored grain is indica rice and has been stored since September 2013. The moisture content and impurity ratio of this grain is 12.8%, and 1%, respectively.

No.4 large warehouses (50 m ×20 m) in Guangdong Zhanjiang North Station Grain Warehouse was selected for trails. The indica rice in this warehouse had 10.5% grain moisture content and been stored since August 2013. .

Both warehouses had been fumigated with phosphine.

2.2 Application of sticky board trap

Sticky board traps (20 cm × 25 cm, Beijing Ikoman Bio-Technology Co., Ltd.) were applied different positions of the surface of indica rice in each of two warehouses, including corner, under the fan, window, and check door. The numbers applied were four. The position of the trap location in the real warehouse is shown in Fig. 1.




Fig. 1 sampling points in the warehouse

2.3 Observing the circadian rhythm of the booklice in warehouse

General procedures: the methods of assessment were based on procedures developed to measure booklice population on grain. The methods were:

- Counting of booklice in marked areas.
- Leaving sticky board traps for defined periods (2 hs), shaking out the booklices into trays and counting.
- Sampling grain with a bottom-opening probe, sieving ($\Phi 1.5 \times 2.5$ mm) and count insects.
- The whole test lasted for 24 hours.

The temperature and relative humidity of the pf the warehouses were monitored by Vaisala VAISALAHM34 High-precision Temperature and Humidity Table (VAISALA, Finland).

3. Results

3.1 Circadian rhythm of *L. entomophila* in paddy warehouse in Nanning Shajing Grain Warehouse

The species of booklice in Nanning Shajing Grain Warehouse was *L. entomophila*. Within 24 hours, the temperature ranged over time from 27.48 °C to 29.28°C and the humidity ranged from 68.95% to 73.05% (Fig. 2). The warehouse temperature showed an overall downward trend during the period from 12 am to 10 pm and a rise after 10 pm. During this period, the humidity showed an overall downward trend, except that it was an abnormally high point at 10 am. The trend continued until 2 pm. The quantity of booklice trapped generally decreased during the period from 12 am to 10 am, and reached a minimum at 12 pm. After 12 pm, the temperature in the warehouse experienced an increase, while this phenomenon appeared 4 hours later and 2 hours later in the humidity and the quantity of booklice trapped respectively. Therefore, the Circadian Rhythm can be inferred from the quantity of booklice attracted by the sticky trap at different time periods. The activity frequency of *L. entomophila* in the warehouse is correlated with the temperature and humidity in the warehouse. From early morning, the frequency of booklice gradually decreased, and booklice activities entered a relatively quiet period at 12 pm. After this, with the overall recovery of temperature and humidity, pest activity gradually became active again, and reached relative activity peak period from 6 pm to 2 am. The daily activity trend of *L. entomophila* can be expressed as $y = -0.971x^3 + 21.88x^2 - 139.5x + 353.4$ (x is the 24-hour time, and y is the quantity of booklice)




Fig. 2 Changes in Temperature, Humidity, and quantity of *L. entomophila* trapped with time in the warehouse of No. 32 Shajing Grain Depot

3.2 Circadian Rhythm of *L. paeta* in Zhanjiang Warehouse

The species of booklice in Zhanjiang Warehouse is *L. paeta*. In Zhanjiang North Station State Grain Storage No. 4 warehouse, the temperature varied significantly with time, ranging from 27.30°C to 29.27°C. However, the humidity vary slightly with time, ranging from 74.40% to 79.27% and reached an unusually high point around 2 am. There was a certain correlation between the quantities of booklice and humidity after 10 am. The trend in Fig. 3 showed that the quantity of trapped *L. paeta* was relate to temperature. Interestingly, 12 pm was not the lowest temperature of the day, but the quantity of trapped *L. paeta* reached a minimum value as same as that of *L. entomophila* in Nanning Shajing. This might be caused by other factors expect temperature and humidity, such as insect daily activity rhythm. The peak activity of the *L. paeta* appeared at 2 pm.




Fig. 3 Changes in temperature, humidity, and quantity of *L.paeta* trapped with time in Zhanjiang No. 4 warehouse

4. Discussion

In recent years, booklice has also become a new threat to global grain security (Zhang Shengfang, 1998), which is the problem that needs to be solved urgently (Muhammad Shoaibet al, 2010). The sticky trap in the study can be considered as a physical control method. It provides a new green and effective means for the prevention and treatment of booklice.

Since both the *L.entomophila* and *L. paeta* have obvious light-shielding properties (Yan Xiaoping et al, 2008), the principle of sticky trap remains to be further studied. It may be related to the fact that the *L.entomophila* prefers high humidity environment because of the stickiness. The glue on the glue sheet causes an increase in humidity.

References

- Bai Xuguang et al, 2002, Pest Control in Stores, Science Press, Beijing.
 Li Zhishen, 2014, Application research of Prediction and Rediction and Control of Pests by the Insect Different Light, (master) Henan University of Technology, Zhengzhou, Henan, China.
 Zhang Shengfang, 1998, Stored beetles in China, China Agricultural Science and Technology Press, Beijing.
 Muhammad Shoaibet al, 2010, Psocid: A new risk for global food security and safety, Appl.Entomol **45**, 89-100.
 Yan Xiaoping, Zhou Hao, Shen Zhaopeng, 2008, Summary and Analysis of Previous Surveys of Grain Insects in China, Grain Storage, 3-6.

Development of a suitable rearing media for *Tribolium castaneum*

Kariyawasam Bovithanthri Thanushi Thamodhi Wijerathne, Edirimunhie Vishwa Udani Perera Karunarathne, Dissanayaka Mudiyanseelage Saman Kumara Dissanayaka, Leanage Kanaka Wolly Wijayaratne*

Department of Plant Sciences, Faculty of Agriculture, Rajarata University of Sri Lanka, Puliyankulama, Anuradhapura, Sri Lanka.

*Corresponding author: wollylk@yahoo.com

DOI 10.5073/jka.2018.463.039

Abstract

Tribolium castaneum is a serious pest of cereal flour and flour-based products, and thus a test insect in stored-product research. The composition of the rearing medium affects the progeny production, their performance and handling efficacy. The objective of this research was to develop a suitable rearing media for *T. castaneum*. The research tested wheat flour, crushed broiler feed, crushed dog feed and corn flour alone and in different combinations. Twenty adults of *T. castaneum* were introduced to each medium separately, and removed after 2 weeks. The progeny adults emerged in each rearing medium was determined. The progeny produced differed with the food medium. In general, the rearing media having a combination of ingredients produced more