

Vitis 42 (1), 43–51 (2003)

## Genetic structure in populations of the fungus *Fomitiporia punctata* associated with the esca syndrome in grapevine

I. JAMAUX-DESPRÉAUX and J. P. PÉROS

UMR DGPC, Equipe vigne, INRA, Montpellier, France

### Summary

Six populations of *Fomitiporia punctata* (Fr.) Murrill were sampled in France and Italy from vines showing symptoms of esca syndrome. Genetic variation within and among populations was studied by using 34 random amplified polymorphic DNA (RAPD) markers. All the 192 isolates analyzed were distinguished, and the haplotypic diversity was similar in each population. The hypothesis that markers were randomly associated was not rejected for 5 populations and for the total sample. Comparison of marker frequency showed significant differences among populations for only 4 markers, indicating a low level of genetic differentiation. The analysis of molecular variance (AMOVA) confirmed that most of the variance in RAPD banding patterns was present within populations (98.5 %). These data strongly suggest that the fungus spreads by means of airborne basidiospores and regularly outcrosses in nature. The prospected regions seem to form an epidemiological unit with a panmictic population of *F. punctata*. However, a very low but significant differentiation was detected between the populations in western France and those in Mediterranean locations.

**Key words:** *Fomitiporia punctata*, *Vitis vinifera* L., esca syndrome, genetic diversity, RAPD, population biology.

### Introduction

The esca syndrome is a major threat for grapevine (*Vitis vinifera* L.) longevity worldwide (DUBOS and LARIGNON 1988; MUGNAI *et al.* 1999). Two basidiomycetes, *Stereum hirsutum* (Willd.) Pers. and *Phellinus igniarius* (L.:Fr.) Quél., were formerly suspected of being involved in the wood decay process (VIALA 1926; CHIARAPPA 1959). However, recent investigations indicated that *Fomitiporia punctata* (Fr.) Murrill [= *Phellinus punctatus* (P. Karst.) Pilát] is the basidiomycete responsible for the white rot lesion (MUGNAI *et al.* 1996; LARIGNON and DUBOS 1997; CORTESI *et al.* 2000). *F. punctata* is able to colonize grapevine alone (SPARAPANO *et al.* 2000) but it is accepted that pioneer decay fungi, like *Eutypa lata* (Pers.:Fr.) Tul. *et* C. Tull., *Phaeoconiella chlamydospora* (W. Gams *et al.*) Crous *et* W. Gams (formerly *Phaeoacremonium chlamydosporum* W. Gams *et al.*) and *Phaeoacremonium aleophilum* W. Gams *et al.*, play an important role in the development of the syndrome in vineyards (MUGNAI

*et al.* 1996; JAMAUX-DESPRÉAUX *et al.* 1997; LARIGNON and DUBOS 1997). The banning of sodium arsenite, which was the only mode of chemical control in vineyards, means that now new strategies against esca have to be developed. This requires a better knowledge of the population biology of the fungal species that degrade the wood, and, in particular, of the white-rot pathogen *F. punctata*.

The way *F. punctata* populations are established in vineyards is not known. According to VIALA (1926) and many others, the white rot lesion develops first in the pith of pruning wounds and the vine is killed before the lesion reaches the rootstock. This indicates that the *F. punctata* infection does not start from an inoculum in the soil. The infection of wounded aerial parts may be due to the vegetative spread of the fungus from vine to vine, for instance by contaminated pruning tools. Another possibility is that airborne basidiospores spread the fungus. To address the question of how the fungus spreads, the spatial distribution of vines showing foliar symptoms has been extensively studied. Evidence for aggregation (SURICO *et al.* 1999; POLLASTRO *et al.* 2000 b) as well as for random distribution (CORTESI *et al.* 2000; SURICO *et al.* 2000) has been presented. Spreading by airborne basidiospores could lead to random distribution but aggregation could have other reasons than propagation from vine to vine. For instance, soil heterogeneity may have a considerable effect on the expression of symptoms. Moreover, as several fungi are associated with esca, considering the spatial patterns of the syndrome may be of limited value to establish the spreading mode of one of these fungi.

Assessment of the genetic diversity of *F. punctata* in vineyards could provide more information about the spreading mode of the fungus than the spatial patterns of symptomatic vines. CORTESI *et al.* (2000) described a wide diversity of somatic incompatibility types within Italian vineyards and POLLASTRO *et al.* (2000 a) found that isolates of *F. punctata* taken from adjacent vines were different Random Amplified Polymorphic DNA (RAPD) phenotypes. This high level of fine-scale genetic diversity suggests that airborne basidiospores could be important sources of inoculum and that *F. punctata* is able to outcross in nature.

Whether *F. punctata* occurs in European vineyards as a single population or as genetically distinct subpopulations has not been documented. Addressing this question could also provide information about the spreading mode. For instance, lack of genetic differentiation among geographically distant regions may be explained by a gene flow that occurs through the movement of basidiospores. On the other hand,

a high level of genetic differentiation may reflect the existence of intersterile populations, *i.e.* sibling species, which have been described in several basidiomycetes that cause wood decay (McKEEN 1952; RAYNER and BODDY 1988; FISHER and BRESINSKY 1992; FISCHER, 1994; STENLID *et al.* 1994). *F. punctata* is widespread in Europe where different grapevine cultivars are cultivated in very diverse climates. *F. punctata* is also found on many living trees as well as on logs of hardwoods and conifers (WALLA 1984; RYVARDEN and GILBERTSON 1994; IPPOLITO *et al.* 1998). These different ecological niches could be occupied by different subpopulations.

The objective of this study was to determine the genetic structure of *F. punctata* populations obtained from esca-diseased vines in France and Italy. We used Random Amplified Polymorphic DNA (RAPD) markers to assess the level of diversity within populations and to determine if distinct subpopulations of *F. punctata* exist.

### Material and Methods

**Fungal isolates:** The 192 *F. punctata* isolates studied were collected in different geographical areas in France and Italy (Tab. 1, Fig. 1). Isolates from Languedoc-Roussillon (region F1) were separated in two groups: F1-A ( $N = 46$ ) corresponded to vineyards in the Pyrénées-Orientales and Aude districts (eastern part of the region) whereas F1-B ( $N = 63$ ) corresponded to those in the Hérault and Gard districts (western part of the region). Isolates from southwestern France (region F2) were also divided in two zones: F2-A ( $N = 23$ ) from the Charentes region and F2-B ( $N = 28$ ) from the Aquitaine and Midi-Pyrénées regions. The F2-A population was sampled on a single grapevine cultivar (cv. Ugni blanc) in a rather homogenous area, whereas the F2-B population corresponded to several cultivars cultivated in diverse environmental conditions (Tab. 1). P. LARIGNON




Fig. 1: Location of the *Fomitiporia punctata* populations sampled in this study from southern France (F1), south-western France (F2) and northern Italy (IT). Each region was separated in two areas, A and B.

(INRA, Bordeaux, France) provided isolates from F2-A and F2-B populations. Italian isolates (IT) originated from two different zones in northern Italy separated by approximately 150 km: IT-A ( $N = 13$ ) from near lake of Garda and IT-B ( $N = 19$ ) from Tuscany. Italian isolates were provided by M. MINERVINI (University of Milan, Italy).

Isolates were generally sampled in different vineyards but isolates from different plants were also taken within the same vineyard (Tab. 1). To isolate the fungus in the F1 region, we chose vines with external symptoms of infection, *i.e.* either leaf discoloration or dieback. Trunks were then cut into pieces to look for white-colored soft lesions in the wood. Pieces of wood (10 x 5 x 5 mm) were taken at the lesion margin and disinfected in a 3% active chlorine solution. The pieces were washed twice in sterile water and put in Petri plates containing Potato Dextrose Agar (PDA, Difco) medium. The plates were incubated at  $23 \pm 2$  °C and observed every 2-3 d for two months. Isolated fungi were subcultured on PDA medium, and identification of *F. punctata* was based on its cultural characteristics. Isolates were maintained at 4 °C as culture disks placed in plastic tubes containing 1 ml of distilled water.

**DNA extraction:** PDA plates were inoculated with two plugs taken at the margin of fresh PDA cultures and incubated for two weeks at 23 °C. Mycelium was scraped using a tip and immediately ground with a chilled mortar and pestle in liquid nitrogen. The mycelium powder was put in a microtube. After addition of 0.5 ml of extraction buffer: 2% cetyltrimethylammonium bromide (CTAB), 1.4 M NaCl, 100 mM Tris-HCl, pH 8.0, 20 mM EDTA, the mixture was incubated for 1-2 h at 65 °C. One volume of chloroform was added to the tube, which was then centrifuged at 12,000 g for 15 min at 16 °C. The supernatant was transferred into another microtube containing 0.1 ml of 10% CTAB and 0.4 ml of chloroform. After homogenization and centrifugation, the supernatant was transferred and nucleic acids were precipitated using one volume of cold isopropanol. The tube was kept at -20 °C overnight, then gently agitated and centrifuged. Isopropanol was discarded and the pellet washed with 70% ethanol. The pellet was dried and dissolved in 0.5 ml of 10 mM Tris-HCl, pH 8.0, 0.1 mM EDTA. The DNA was quantified on 0.8% agarose gel stained with ethidium bromide by visual comparison with different quantities of lambda DNA (Life Technologies, Illkirch, France).

**RAPD amplifications:** PCR conditions described by WILLIAMS *et al.* (1990) were used with some modifications. Each reaction volume of 25 µl included 1.5 units of Taq DNA polymerase (Appligene, France), 1x buffer (10 mM Tris-HCl, pH 9, 50 mM KCl, MgCl<sub>2</sub> 1.5 mM, 0.1% tritonX100, 0.2 mg.ml<sup>-1</sup> BSA) provided with the enzyme, 120 µM each of dATP, dCTP, dGTP and dTTP (Boehringer Mannheim, Germany), 30 ng of primer (Life Technologies, Illkirch, France) and approximately 25 ng of template DNA. The reaction was overlaid with a drop of mineral oil. A thermocycler (Biometra, Göttingen, Germany) was programmed for one step of 4 min at 94 °C, followed by 36 cycles of 1 min at 93 °C, 1 min at 38 °C, 1 min at 72 °C, and a final step of 6 min at 72 °C. Amplified products were analyzed by electrophoresis in 1.6% agarose gel at 5 V per cm for 4 h, along a molecular size marker (1-kb ladder, Life Technolo-

Table 1

List of isolates of *Fomitiporia punctata* used in this study with their geographical origin, vine cultivar origin, and date of sampling in three European regions in France (F1, F2) and Italy (IT), each region being subdivided into two areas (A and B)

| Identification <sup>a</sup> | Pop. | Locality <sup>b</sup>         | Vine cultivar          | Sampling date |
|-----------------------------|------|-------------------------------|------------------------|---------------|
| LR6 to LR8                  | F1-A | Antugnac (11)                 | Mauzac                 | Sept. 1996    |
| LR9 to LR13                 | F1-A | Couiza (11)                   | Mauzac                 | Sept. 1996    |
| LR14 to LR18                | F1-A | Montréal (11)                 | Cinsault               | Sept. 1996    |
| LR96 and LR98               | F1-A | St Laurent Cabrerisse (11)    | Grenache               | Sept. 1996    |
| LR104 to LR106              | F1-A | St Laurent Cabrerisse (11)    | Carignan               | Sept. 1996    |
| LR112                       | F1-A | Ribaute (11)                  | Carignan               | Sept. 1996    |
| LR114                       | F1-A | Laure-Minervois (11)          | Cinsault               | Sept. 1996    |
| LR118                       | F1-A | Durban (11)                   | Grenache/Carignan      | Sept. 1996    |
| LR140                       | F1-A | Narbonne (11)                 | Cinsault               | Oct. 1996     |
| LR143                       | F1-A | Treilles (11)                 | Muscat à petits grains | Oct. 1996     |
| LR80 and LR81               | F1-A | Montesquieu (66)              | Cinsault               | Sept. 1996    |
| LR82 to LR84                | F1-A | Montesquieu (66)              | Grenache               | Sept. 1996    |
| LR85 to LR89                | F1-A | Ortoffa (66)                  | Grenache               | Sept. 1996    |
| LR90 to LR94                | F1-A | Tour de France (66)           | Mourvèdre              | Sept. 1996    |
| LR99 to LR103               | F1-A | Banyuls del Aspres (66)       | Muscat à petits grains | Sept. 1996    |
| LR131                       | F1-A | Maury (66)                    | Grenache               | Oct. 1996     |
| LR144                       | F1-A | Le Boulou (66)                | Macabeu                | Nov. 1996     |
| LR146                       | F1-A | Le Boulou (66)                | Macabeu                | Nov. 1996     |
| LR1 to LR5                  | F1-B | Beauvoisin (30)               | Carignan               | Sept. 1996    |
| LR34 to LR38                | F1-B | Les Tavernes (30)             | Cinsault               | Sept. 1996    |
| LR39                        | F1-B | Monteils (30)                 | Cinsault               | Sept. 1996    |
| LR43 to LR47                | F1-B | Cannes et Clairan (30)        | Carignan               | Sept. 1996    |
| LR76                        | F1-B | St Nazaire (30)               | Syrah                  | Sept. 1996    |
| LR126                       | F1-B | Montaren-St Médières (30)     | Cinsault               | Oct. 1996     |
| LR123                       | F1-B | Villeneuve les Maguelone (34) | Carignan               | Sept. 1996    |
| LR19 to LR23                | F1-B | Assignan (34)                 | Grenache               | Sept. 1996    |
| LR24 to LR27                | F1-B | Assignan (34)                 | Sauvignon              | Sept. 1996    |
| LR69                        | F1-B | Adissan (34)                  | Bourret                | Sept. 1996    |
| LR48 to LR 51               | F1-B | Puissalicon (34)              | Carignan               | Sept. 1996    |
| LR52 to LR55                | F1-B | Puissalicon (34)              | Cinsault               | Sept. 1996    |
| LR57, L58 and LR60          | F1-B | Pinet (34)                    | Cinsault               | Sept. 1996    |
| LR61                        | F1-B | Pinet (34)                    | Alicante Bouschet      | Sept. 1996    |
| LR63 to LR65                | F1-B | St Christol (34)              | Grenache               | Sept. 1996    |
| LR66 to LR68                | F1-B | St Christol (34)              | Grenache blanc         | Sept. 1996    |
| LR124                       | F1-B | Villeneuve les Maguelone (34) | Carignan               | Sept. 1996    |
| LR70 to LR72                | F1-B | Vendargues (34)               | Alphonse Lavallée      | Sept. 1996    |
| LR107 to LR111              | F1-B | Villeveyrac (34)              | Cabernet-Sauvignon     | Sept. 1996    |
| LR30                        | F1-B | Leyrac (34)                   | Cinsault               | Sept. 1996    |
| LR79                        | F1-B | Octon (34)                    | Cinsault               | Sept. 1996    |
| LR73 to LR75                | F1-B | Vendargues (34)               | Cinsault               | Sept. 1996    |
| LR134                       | F1-B | St Mathieu de Trévières (34)  | Mourvèdre              | Oct. 1996     |
| LR137                       | F1-B | Montpellier (34)              | Unknown                | Oct. 1996     |
| PC8                         | F2-A | Pérignac (16)                 | Ugni blanc             | Dec. 1996     |
| PC9                         | F2-A | St Laurent de Cognac (16)     | Ugni blanc             | Dec. 1996     |
| PC11                        | F2-A | Montguyon (16)                | Ugni blanc             | Dec. 1996     |
| PC14                        | F2-A | La Couronne (16)              | Ugni blanc             | Dec. 1996     |
| PC15 and PC17               | F2-A | Vaux-Rouillac (16)            | Ugni blanc             | Dec. 1996     |
| PC18                        | F2-A | Monchaude (16)                | Ugni blanc             | Dec. 1996     |
| PC20 to PC22                | F2-A | St Preuil (16)                | Ugni blanc             | Dec. 1996     |
| PC23                        | F2-A | Angeac-Charente (16)          | Ugni blanc             | Dec. 1996     |
| PC25 to PC27                | F2-A | Graves (16)                   | Ugni blanc             | Dec. 1996     |
| PC28                        | F2-A | Julienne (16)                 | Ugni blanc             | Dec. 1996     |

(Table 1, continued)

| Identification <sup>a</sup>    | Pop. | Locality <sup>b</sup>          | Vine cultivar       | Sampling date |
|--------------------------------|------|--------------------------------|---------------------|---------------|
| PC1 to PC 3                    | F2-A | Les Touches de Périgny (17)    | Ugni blanc          | Dec. 1996     |
| PC4 and PC5                    | F2-A | Les Eglises d'Argenteuil (17)  | Ugni blanc          | Dec. 1996     |
| PC6                            | F2-A | Saintes (17)                   | Ugni blanc          | Dec. 1996     |
| PC7                            | F2-A | Le Chay (17)                   | Ugni blanc          | Dec. 1996     |
| PC12                           | F2-A | Pouillac (17)                  | Ugni blanc          | Dec. 1996     |
| AQ15                           | F2-B | Montbazillac (24)              | Sauvignon           | Sept. 1996    |
| AQ18                           | F2-B | St Avit St Nazaire (24)        | Sauvignon           | Sept. 1996    |
| AQ19                           | F2-B | St Méard de Gurçon (24)        | Sauvignon           | Sept. 1996    |
| AQ21                           | F2-B | Fougueyrolles (24)             | Cabernet-Sauvignon  | Sept. 1996    |
| AQ4                            | F2-B | Léognan (33)                   | Cabernet-Sauvignon  | Oct. 1996     |
| AQ5                            | F2-B | St Emilion (33)                | Cabernet-Franc      | Sept. 1996    |
| AQ6                            | F2-B | St Emilion (33)                | Merlot              | Sept. 1996    |
| AQ7                            | F2-B | Grézillac (33)                 | Sauvignon           | Sept. 1996    |
| AQ8                            | F2-B | Tauriac (33)                   | Cabernet-Sauvignon  | Sept. 1996    |
| AQ9                            | F2-B | Labarde (33)                   | Cabernet-Sauvignon  | Sept. 1996    |
| AQ10                           | F2-B | Pessac (33)                    | Cabernet-Sauvignon  | Sept. 1996    |
| AQ12                           | F2-B | Bouqueyrans (33)               | Cabernet-Sauvignon  | Sept. 1996    |
| AQ13                           | F2-B | Bouqueyrans (33)               | Cabernet-Franc      | Sept. 1996    |
| AQ14                           | F2-B | St Julien Beychevelle (33)     | Cabernet-Sauvignon  | Sept. 1996    |
| AQ23                           | F2-B | St Christophe des Bardes (33)  | Cabernet-Sauvignon  | Sept. 1996    |
| AQ33 to AQ37                   | F2-B | Naujan et Postiac (33)         | Cabernet-Sauvignon  | Sept. 1996    |
| AQ27                           | F2-B | La Bastide d'Armagnac (44)     | Baco 22A            | Oct. 1996     |
| AQ29                           | F2-B | Arthez d'Armagnac (44)         | Folle Blanche       | Oct. 1996     |
| AQ28                           | F2-B | Aydie (64)                     | Tannat              | Oct. 1996     |
| AQ30                           | F2-B | Moncaup (64)                   | Cabernet-Franc      | Oct. 1996     |
| AQ31                           | F2-B | Monein (64)                    | Gros Manseng        | Oct. 1996     |
| AQ32                           | F2-B | Jurançon (64)                  | Gros Manseng        | Oct. 1996     |
| AQ43                           | F2-B | Moncaup (64)                   | Cabernet-Franc      | Oct. 1996     |
| MP1                            | F2-B | Soublecause (65)               | Cabernet-Franc      | Oct. 1996     |
| VE3                            | IT-A | Lazise, Custoza                | Corvina             | May 1996      |
| VE6                            | IT-A | Lazise, Custoza                | Trebbiano Toscano   | May 1996      |
| VE7                            | IT-A | St Giorgio in Salici, Custoza  | Trebbiano Toscano   | May 1996      |
| VE10                           | IT-A | St Giorgio in Salici, Custoza  | Corvina             | Mar. 1996     |
| VE11                           | IT-A | St Giorgio in Salici, Custoza  | Trebbiano di Lugana | Sept. 1995    |
| VE12 and VE13                  | IT-A | St Giorgio in Salici, Custoza  | Trebbiano di Lugana | Aug. 1995     |
| VE17                           | IT-A | Lazise, Custoza                | Corvina             | May 1995      |
| VE23 and VE31                  | IT-A | St Benedetto di Lugana, Lugana | Trebbiano di Lugana | May 1996      |
| VE26                           | IT-A | St Benedetto di Lugana, Lugana | Trebbiano di Lugana | Oct. 1996     |
| EM1                            | IT-A | Imola, Emilia                  | Sangiovese          | Jun. 1995     |
| LO1                            | IT-A | Broni, Oltrepo Pavese          | Barbera             | Oct. 1995     |
| TO1                            | IT-B | St Andrea, Chianti             | Sangiovese          | Nov. 1996     |
| TO3                            | IT-B | St Andrea, Chianti             | Sangiovese          | Oct. 1996     |
| TO4                            | IT-B | Villa Branca, Chianti          | Sangiovese          | Oct. 1996     |
| TO5                            | IT-B | Villa Branca, Chianti          | Sangiovese          | Oct. 1996     |
| TO6                            | IT-B | St Andrea, Chianti             | Sangiovese          | Oct. 1996     |
| TO8                            | IT-B | St Cristina, Chianti           | Sangiovese          | Oct. 1996     |
| TO9                            | IT-B | St Cristina, Chianti           | Sangiovese          | Oct. 1996     |
| TO11                           | IT-B | St Teresa, Chianti             | Sangiovese          | Sept. 1996    |
| TO17, TO19<br>and TO21 to TO28 | IT-B | Valledoro, Chianti             | Sangiovese          | Mar. 1997     |

<sup>a</sup> Isolate code includes the geographical origin (LR, Languedoc-Roussillon; PC, Poitou-Charentes; AQ, Aquitaine; MP, Midi-Pyrénées; VE, Venetia; EM, Emilia Romagna; TO, Tuscany) and the order of isolation for each region.

<sup>b</sup> Numbers in parentheses indicate official French district codes, as follows: 11, Aude; 16, Charente; 17, Charente-Maritime; 24, Dordogne; 30, Gard; 33, Gironde; 34, Hérault; 44, Landes; 64, Pyrénées-Atlantiques; 65, Hautes-Pyrénées and 66, Pyrénées-Orientales.

gies, Illkirch, France). Fragments were detected by staining with ethidium bromide and gels were photographed under UV light. A total of 120 decamer primers were initially screened using one DNA sample to identify primers that gave scorable patterns. Twenty primers were then evaluated to find polymorphism among 18 isolates from Languedoc-Roussillon (region F1). Finally, 9 primers that produced strongly amplified, polymorphic and reproducible bands were selected for full analysis. These primers were A08 (5'GTGACGTAGG), A11 (5'CAATCGCCGT), B18 (5'CCACAGCAGT), E11 (5'GAGTCTCAGG), E15 (5'ACGCACAACC), E16 (5'GGTGACTGTG), E18 (5'GGACTGCAGA), P15 (5'GGAAGCCAAC) and P17 (5'TGACCCGCCT). Analyses were repeated twice from extraction to electrophoresis of amplification products. A negative control reaction with no DNA template and a positive control reaction were included in each run.

**Data analysis:** To analyze RAPD patterns, it is commonly assumed that each band represents the phenotype at a single *locus* with two alleles. Without a segregation analysis to validate this hypothesis for each marker, allele frequencies cannot be estimated with precision. For instance, if the absence of the band is due to several genetic events, the diversity will be underestimated. The dominant nature of RAPDs introduces another bias since the marker is present both in homozygotes and heterozygotes of the positive allele. According to FISCHER (1996), hyphal segments of *F. punctata* contain 2-8 nuclei that were presumed to be diploid. Therefore, if *F. punctata* is heterothallic, the dominance of RAPDs would also lead to a false estimation of the gene diversity. Another important bias concerns the analysis of the differentiation between populations. Studies in conifers have shown that estimates of population differentiation derived from dominant RAPD fingerprints were inflated compared to those obtained from haploid tissues (ISABEL *et al.* 1995; SZMIDT *et al.* 1996; ISABEL *et al.* 1999). Although statistical methods have been implemented to reduce the bias (LYNCH and MILLIGAN 1994), we analyzed data in terms of differences in RAPD patterns without attempting to estimate allele frequencies.

Only polymorphic bands were considered and scored as 1 present, or 0 absent. To describe the diversity in populations, we first calculated the number of marker differences ( $d_i$ ) for each pair of isolates within the population  $i$  using a program written in C++ language. The values were averaged over all comparisons to obtain  $D_i$ . For each RAPD marker and each population, the haplotypic diversity (NEI and TAJIMA 1981) corrected for small population sizes was also calculated as  $h_i = N_i(1 - (p_i^2 + q_i^2)) / (N_i - 1)$  where, for the population  $i$ ,  $N_i$  was the number of isolates,  $p_i$  the frequency of isolates showing the band and  $q_i$  the frequency of isolates not showing the band. The values were averaged over all markers to represent the diversity ( $H_i$ ) within each population.  $H$  corresponded to the probability of observing a difference in markers between two isolates taken at random. This quantity might thus also be obtained by dividing the mean number of differences ( $D_i$ ) by the number of markers. The relationships among isolates from each population were studied by cluster analyses applying the unweighed pair group method with arithmetic averages (UPGMA) based on

pair-wise differences. For this purpose, we used the procedures Neighbour of PHYLIP Version 3.5c (J. FELSENSTEIN, Department of Genetics, University of Washington).

The hypothesis that recombination might occur in each population was tested using the method developed by BROWN *et al.* (1980). In the original method, the observed variance in the number of *loci* at which two individuals from a population of  $N$  individuals have different alleles was calculated over  $N(N-1)/2$  pairs and compared to the variance expected under the assumption of random association. Our markers were not genetically characterized, but we applied the same reasoning to the presence/absence of markers. In each population, the observed variance in the number of marker differences ( $V_D$ ) was compared to the variance expected ( $V_E$ ) for random association of the markers. The null hypothesis  $H_0: V_D = V_E$  was tested by simulation using the Monte Carlo procedure (SOUZA *et al.* 1992). Variances, critical value for  $V_D$  and the probability of rejecting by chance only the null hypothesis were obtained using LIAN (HAUBOLD and HUDSON 2000).

To study the genetic differentiation, we compared the observed numbers of isolates showing or not each RAPD marker with the numbers expected under the null hypothesis that no difference occurred between populations. An unbiased estimate of the P-value of a log-likelihood ratio ( $G^2$ )-based exact test was performed for each marker and combined over markers using Fisher's method. These calculations were carried out using GENEPOP Version 3.2 (RAYMOND and ROUSSET 1995).

The analysis of molecular variance (AMOVA) developed by EXCOFFIER *et al.* (1992) was implemented to study the genetic differentiation between the 6 populations in more detail. The analysis was performed using ARLEQUIN Version 2.000 (SCHNEIDER *et al.* 2000) based on the number of differences between all pairs of RAPD phenotypes. The total variation for these differences was partitioned in hierarchical components (among regions IT, F1 and F2, among populations areas A and B within regions, and within populations). ARLEQUIN also gave the pair-wise distance between populations ( $\phi_{st}$ ) and computed by permutation procedures the significance levels for variance component estimates and for  $\phi_{st}$ .

## Results

From the multiple-banding patterns produced with the 9 primers, we selected a total of 34 reproducible and clearly scorable bands that were polymorphic across the whole sample (Tab. 2). These markers allowed us to distinguish each isolate as a distinct RAPD phenotype, the number of marker differences between isolates ranging from 1 to 22 over all populations. The diversity appeared similar within the 6 populations; the haplotypic diversity ranged from 0.292 to 0.322 whereas the mean number of marker differences ranged from 9.9 to 11.2 (Tab. 3). The pattern of clustering of isolates to be expected if geographical proximity structured each population was not observed. In particular, isolates derived from the same vineyard were not grouped in the cluster analysis performed for each population (example in Fig. 2 for F2-A

Table 2

Frequency of 34 Random Amplified Polymorphic DNA (RAPD) markers in 6 populations of *Fomitiporia punctata*.  
For details: Tab. 1

| RAPD     | Population   |              |              |              |              |              | All populations <sup>a</sup><br>N=192 | Pr <sup>b</sup> |
|----------|--------------|--------------|--------------|--------------|--------------|--------------|---------------------------------------|-----------------|
|          | F1-A<br>N=46 | F1-B<br>N=63 | F2-A<br>N=23 | F2-B<br>N=28 | IT-A<br>N=13 | IT-B<br>N=19 |                                       |                 |
| A8-1910  | 0.565        | 0.460        | 0.391        | 0.357        | 0.384        | 0.473        | 0.458                                 | 0.557           |
| A8-1370  | 0.717        | 0.825        | 0.696        | 0.536        | 0.615        | 0.895        | 0.734                                 | 0.032*          |
| A8-870   | 0.348        | 0.175        | 0.261        | 0.357        | 0.154        | 0.211        | 0.255                                 | 0.273           |
| A8-670   | 0.217        | 0.349        | 0.261        | 0.321        | 0.462        | 0.263        | 0.302                                 | 0.552           |
| A11-2300 | 0.152        | 0.222        | 0.087        | 0.143        | 0.154        | 0.105        | 0.161                                 | 0.689           |
| A11-2130 | 0.348        | 0.270        | 0.217        | 0.214        | 0.385        | 0.211        | 0.276                                 | 0.680           |
| A11-1110 | 0.326        | 0.270        | 0.087        | 0.107        | 0.385        | 0.316        | 0.250                                 | 0.063           |
| A11-910  | 0.848        | 0.794        | 0.783        | 0.964        | 0.077        | 0.947        | 0.844                                 | 0.212           |
| A11-740  | 0.652        | 0.714        | 0.870        | 0.679        | 0.769        | 0.421        | 0.688                                 | 0.056           |
| B18-1230 | 0.152        | 0.111        | 0.087        | 0.107        | 0.154        | 0.158        | 0.125                                 | 0.959           |
| B18-470  | 0.696        | 0.698        | 0.913        | 0.964        | 0.462        | 0.473        | 0.724                                 | 0.001*          |
| E11-1030 | 0.261        | 0.079        | 0.348        | 0.250        | 0.154        | 0.105        | 0.188                                 | 0.039*          |
| E15-1270 | 0.217        | 0.317        | 0.348        | 0.393        | 0.385        | 0.263        | 0.307                                 | 0.630           |
| E15-930  | 0.000        | 0.016        | 0.043        | 0.038        | 0.077        | 0.053        | 0.052                                 | 0.288           |
| E15-730  | 0.022        | 0.127        | 0.174        | 0.071        | 0.000        | 0.053        | 0.083                                 | 0.159           |
| E15-560  | 0.565        | 0.556        | 0.391        | 0.536        | 0.615        | 0.421        | 0.526                                 | 0.627           |
| E15-330  | 0.804        | 0.825        | 0.739        | 0.750        | 1.000        | 0.842        | 0.813                                 | 0.465           |
| E16-1580 | 0.761        | 0.667        | 0.435        | 0.643        | 0.692        | 0.632        | 0.656                                 | 0.185           |
| E16-1060 | 0.043        | 0.063        | 0.130        | 0.036        | 0.231        | 0.053        | 0.073                                 | 0.452           |
| E16-840  | 0.609        | 0.778        | 0.696        | 0.750        | 0.692        | 0.684        | 0.708                                 | 0.564           |
| E18-1780 | 0.196        | 0.222        | 0.087        | 0.143        | 0.538        | 0.263        | 0.214                                 | 0.096           |
| E18-1470 | 0.717        | 0.746        | 0.739        | 0.536        | 0.615        | 0.421        | 0.667                                 | 0.061           |
| E18-670  | 0.239        | 0.127        | 0.174        | 0.214        | 0.154        | 0.053        | 0.167                                 | 0.440           |
| E18-590  | 0.065        | 0.095        | 0.043        | 0.107        | 0.077        | 0.053        | 0.078                                 | 0.946           |
| E18-490  | 0.087        | 0.159        | 0.087        | 0.143        | 0.077        | 0.105        | 0.120                                 | 0.863           |
| E18-410  | 0.848        | 0.730        | 1.000        | 0.964        | 1.000        | 1.000        | 0.869                                 | 0.001*          |
| P15-1590 | 0.109        | 0.095        | 0.087        | 0.036        | 0.000        | 0.105        | 0.083                                 | 0.663           |
| P15-1160 | 0.826        | 0.746        | 0.783        | 0.821        | 0.846        | 0.684        | 0.781                                 | 0.772           |
| P15-670  | 0.174        | 0.222        | 0.304        | 0.321        | 0.154        | 0.211        | 0.229                                 | 0.675           |
| P15-460  | 0.174        | 0.095        | 0.174        | 0.036        | 0.154        | 0.263        | 0.135                                 | 0.262           |
| P15-290  | 0.239        | 0.317        | 0.130        | 0.214        | 0.308        | 0.263        | 0.255                                 | 0.593           |
| P17-1590 | 0.217        | 0.175        | 0.174        | 0.143        | 0.077        | 0.105        | 0.167                                 | 0.804           |
| P17-1160 | 0.087        | 0.127        | 0.130        | 0.071        | 0.154        | 0.052        | 0.104                                 | 0.871           |
| P17-970  | 0.152        | 0.127        | 0.261        | 0.107        | 0.154        | 0.158        | 0.151                                 | 0.791           |

<sup>a</sup> Mean allele frequencies for each RAPD locus calculated for the total sample.

<sup>b</sup> Probability to observe the likelihood ratio ( $G^2$ ) under the null hypothesis that frequency did not differ among populations, asterisk shows probability below the 0.05 level of significance.

population). The null hypothesis that markers were randomly associated was tested in each population and for the total population. A significant value that indicated deviation from random association was detected only in the F2-B population (Tab. 3).

The majority of markers were detected in all the populations and their frequency varied from 0.052 to 0.869 over all populations (Tab. 2). The null hypothesis that the number of isolates having or not having the marker was the same in each population was rejected for only 4 markers indicating a low level of genetic differentiation. The P value obtained with Fisher's method over all markers was significant ( $P=0.0088$ ). AMOVA analysis clearly confirmed that most

of the genetic variation was present within populations (Tab. 4). A significant level of genetic differentiation was detected between the three regions, but there was no difference between the two populations within each region (Tab. 4). Pair-wise  $\phi_{st}$  revealed that this differentiation was due to some differences between populations sampled in western France and those taken in Mediterranean areas (Tab. 5).

## Discussion

Our study revealed that *F. punctata* had a high degree of genetic variation. All isolates within populations were

Table 3

Haplotypic diversity, number of marker differences between pairs of isolates and marker association for 34 Random Amplified Polymorphic DNA (RAPD) markers in 6 populations of *Fomitiporia punctata*. For details: Tab. 1

| Population | N   | $H_i^a$ | Number of differences <sup>b</sup> |       | Marker association <sup>c</sup> |       |          |       |
|------------|-----|---------|------------------------------------|-------|---------------------------------|-------|----------|-------|
|            |     |         | $D_i$                              | Range | $V_D$                           | $V_E$ | $L_{MC}$ | P     |
| F1-A       | 48  | 0.317   | 10.8                               | 4-20  | 7.48                            | 6.78  | 7.93     | 0.148 |
| F1-B       | 63  | 0.315   | 10.9                               | 4-20  | 6.40                            | 7.00  | 8.01     | 0.828 |
| F2-A       | 23  | 0.304   | 10.4                               | 4-20  | 7.59                            | 6.57  | 8.47     | 0.175 |
| F2-B       | 28  | 0.292   | 9.9                                | 3-18  | 7.98                            | 6.19  | 7.78     | 0.032 |
| IT-A       | 13  | 0.322   | 11.2                               | 4-19  | 8.28                            | 6.50  | 9.27     | 0.117 |
| IT-B       | 19  | 0.304   | 10.3                               | 5-16  | 5.56                            | 6.36  | 8.33     | 0.779 |
| All        | 192 | 0.315   | 10.7                               | 1-22  | 7.24                            | 6.93  | 7.53     | 0.194 |

<sup>a</sup> Haplotypic diversity.

<sup>b</sup> The number of differences between pairs of isolates (out of 34 markers) was averaged ( $D_i$ ) in each population and was also calculated for the total population.

<sup>c</sup>  $V_D$  = observed variance in number of differences;  $V_E$  = expected variance assuming random association of markers;  $L_{MC}$  = simulated 5 % critical value for  $V_D$  (obtained using a Monte Carlo procedure with 1000 resamplings); P = probability of rejecting by chance alone the null hypothesis that  $V_D = V_E$ .




Fig. 2: Relationships among isolates from a *Fomitiporia punctata* population sampled in south-western France (population F2-A) revealed by UPGMA cluster analysis of the matrix of pair-wise differences for presence of 34 RAPD markers. In 10 vineyards only one isolate was taken whereas in 5 other vineyards (V1 to V5), 2-3 isolates were collected from different vines.

distinguished and no RAPD phenotype was shared between populations. The same high level of diversity was observed within a given vineyard, thus confirming at this spatial scale the findings of CORTESI *et al.* (2000) and POLLASTRO *et al.*

Table 4  
Analysis of molecular variance (AMOVA) of Random Amplified Polymorphic DNA (RAPD) phenotypes for 6 populations of *Fomitiporia punctata*. For details: Tabs 1 and 2

| Source of variation              | df  | Variance component | % total | Pr    |
|----------------------------------|-----|--------------------|---------|-------|
| Among regions                    | 2   | 0.084              | 1.56    | 0.000 |
| Among populations within regions | 3   | -0.012             | -0.04   | 0.462 |
| Within populations               | 186 | 5.307              | 98.48   | 0.002 |

Table 5  
Pair-wise  $\phi_{st}$  between 6 populations of *Fomitiporia punctata*. For details: Tab. 1

| Population | F1-A    | F1-B     | F2-A    | F2-B     | IT-A    |
|------------|---------|----------|---------|----------|---------|
| F1-B       | 0.0046  | -        |         |          |         |
| F2-A       | 0.0155* | 0.0173*  | -       |          |         |
| F2-B       | 0.0116  | 0.0179** | -0.0073 | -        |         |
| IT-A       | 0.0017  | -0.0017  | 0.0301  | 0.0248   | -       |
| IT-B       | 0.0064  | 0.0095   | 0.0356* | 0.0284** | -0.0059 |

\*, \*\*: significant at P=0.05, P=0.01

(2000 a). Furthermore, independent markers appeared to be randomly assorted. This suggests that spreading of *F. punctata* is achieved by airborne basidiospores within outcrossing populations. A low level of fine-scale genetic

diversity would indeed be expected if the fungus was propagated either by mycelium or by basidiospores produced by a homothallic mating system.

In contrast to the evidence we obtained for a high level of recombination, *F. punctata* has been described as a homothallic species after the pairing of mycelia from single spores from the same basidiocarp (FISCHER 1996). Six basidiocarps were analyzed that were not collected from grapevine; two were from *Salix hindsinia* in the United States, two from *Salix caprea* and *Salix fragilis* in Germany, one from *Corylus avellana* in Germany, and one from *Sorbus aucuparia* in Estonia (FISCHER, pers. comm.). The possibility that the mating system in *F. punctata* populations differs depending on the geographical origin or on the origin of the host could explain the conflicting information on the mating system. Such a phenomenon is not uncommon in wood-decaying basidiomycetes where the mating system may differ in various subpopulations. For instance, species of the genus *Stereum* contain both outcrossing and non-outcrossing subpopulations (RAYNER and BODDY 1988). The sexual system of *F. punctata* could be investigated by analyzing the segregation of molecular markers within natural progenies from different locations and hosts. However, a limiting factor for this approach is the poor germination of basidiospores in the laboratory (FISCHER, pers. comm.).

In outcrossing basidiomycetes, a germinating basidiospore produces a homokaryotic, haploid mycelium that grows until another sexually compatible mycelium is encountered to form a heterokaryotic mycelium. Infection of grapevine by *F. punctata* may be initiated by basidiospores with fusion of different homokaryotic mycelia taking place either externally or within the wood. It is generally assumed that homokaryons of *basidiomycetes* have limited development in the wood substrate. However, homokaryons of *Phellinus weirii* performed as well as heterokaryons in degrading inoculated wood strips although not all homokaryons developed successfully (HANSEN 1979). Since *F. punctata* enters through pruning wounds (VIALA 1926), basidiospore infections are expected to take place at scattered positions in the same vine. Using pairing tests, CORTESI *et al.* (2000) obtained some evidence that two different mycelia could be recovered from the same vine but more frequently only one somatic incompatibility type was found. The infrequent occurrence of basidiocarps in vineyards (FISCHER 2000; CORTESI *et al.* 2000) may be partly explained by the rare contact between sexually compatible homokaryons. It would be of great interest to analyze in more detail the composition of *F. punctata* mycelia established within the same vine and to determine how long the homokaryotic condition lasts. However, in hymenochaetales, the order to which *Phellinus* and *Fomitiporia* species belong, it was difficult to distinguish between homokaryons and heterokaryons because both can have oligonucleate cells, and heterokaryons lack clamp connections (HENNON and HANSEN 1987; RIZZO *et al.* 1995). Thus, molecular markers may provide a convenient way to compare the different mycelia of *F. punctata* recovered within and among vines in relation with the presence or absence of basidiocarps.

The low level of genetic differentiation among the populations suggests that *F. punctata* basidiospores can spread over large distances. In other wood-inhabiting, vast numbers of spores were shown to have been transported by air and spores were detected as far as 320 km from extensive sources (RISHBETH 1959). Gene flow may also be due to transport of infected material or to insects carrying spores and mycelial fragments. Only a few immigrants are sufficient to prevent the genetic differentiation of populations (SLATKIN 1987). Another possible explanation for the low level of genetic differentiation is that the populations recently derived from the same population and had no time to diverge by mutation or genetic drift. The low level of differentiation appears to be due to differences between the Mediterranean populations and those sampled in western France. This is surprising since the Alps and the Mediterranean Sea may constitute abrupt geographic barriers to the transport of basidiospores, whereas there is no such barrier between the two French regions. Although RAPD markers are supposed to be neutral, hitch-hiking selection of a few markers might cause some differentiation. Selection may occur, for instance, because of the marked difference between the oceanic climate (western France) and the Mediterranean climate (southern France and Italy).

Isolates of *F. punctata* analyzed in this study originated all from a few European vineyards whereas the fungus is present worldwide (DUBOS and LARIGNON 1988) on several other hosts (WALLA 1984; RYVARDEN and GILBERTSON 1994; IPPOLITO *et al.* 1998). It is therefore possible that subdivision in *F. punctata* occurs at another geographical scale or through a host effect. This possibility is supported by two observations, first, as discussed above, the existence of both outcrossing and non-outcrossing populations would explain the conflicting observations regarding the mating system, and, second, the RFLP (Restriction Fragment Length Polymorphism) analysis of an amplified ITS (Internal Transcribed Spacers) DNA region separated 12 isolates in three different groups: isolates from grapevine in Italy, isolates from *Salix hindsiana* and *Salix lucida* in the United States and isolates from *Salix caprea*, *Rhamnus cathartica* and *Sorbus aucuparia* in northern Europe (FISCHER 2000). Additional locations and other hosts therefore need to be sampled to determine 1) whether the same, highly diverse population of *F. punctata* is widely present in vineyards and 2) whether or not alternative hosts contribute to the epidemiology of Esca in grapevine. The markers identified in our study could help to address these questions.

#### Acknowledgements

The authors thank J. F. BALLESTER and G. BERGER for their technical contribution. They also thank P. LARIGNON and MARIA MINERVINI for providing some of the *F. punctata* isolates and M. FISCHER for helpful comments on the manuscript. This work was supported by the European Union (program FAIR N°95CT654).



## References

- BROWN, A. D. H.; FELDMAN, N. W.; NEVO, E.; 1980: Multilocus structure of natural populations of *Hordeum spontaneum*. *Genetics* **96**, 523-536.
- CHIARAPPA, L.; 1959: Wood decay of the grapevine and its relationship with black measles disease. *Phytopathology* **49**, 510-519.
- CORTESI, P.; FISCHER, M.; MILGROOM, M. G.; 2000: Identification and spread of *Fomitiporia punctata* associated with wood decay of grapevine showing symptoms of esca. *Phytopathology* **90**, 967-972.
- DUBOS, B.; LARIGNON, P.; 1988: Esca and black measles. In: R. C. PEARSON, A. C. GOHEEN (Eds.): *Compendium of Grape Diseases*, 34-35. Am. Phytopathol. Soc., St Paul, USA.
- EXCOFFIER, L.; SMOUSE, P. E.; QUATTRO, J. M.; 1992: Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics* **131**, 479-491.
- FISCHER, M.; 1994: Pairing tests in the *Phellinus pini* group. *Mycologia* **86**, 524-539.
- FISCHER, M.; 1996: On the species complexes within *Phellinus*: *Fomitiporia* revisited. *Mycol. Res.* **100**, 1459-1467.
- FISCHER, M.; 2000: Grapevine wood decay and lignicolous basidiomycetes. *Phytopathol. Mediterr.* **39**, 100-106.
- FISCHER, M.; BRESINSKY, A.; 1992: *Phellinus torulosus*: Sexuality and evidence of intersterility groups. *Mycologia* **84**, 823-833.
- HANSEN, E. M.; 1979: Nuclear condition and vegetative characteristics of homokaryotic and heterokaryotic isolates of *Phellinus weirri*. *Can. J. Bot.* **57**, 1579-1582.
- HAUBOLD, B.; HUDSON, R. R.; 2000: LIAN 3.0: Detecting linkage disequilibrium in multilocus data. *Bioinformatics* **16**, 847-848.
- HENNON, P. E.; HANSEN, E. M.; 1987: Nuclear behavior of *Phellinus arctostaphyli*, *P. ignarius*, and *P. tremulae*. *Mycologia* **79**, 501-507.
- IPPOLITO, A.; NIGRO, F.; DECOCK, C.; 1998: *Phellinus punctatus*, agente di carie in piante di agrumi. *Inform. Fitopatol.* **48**, 36-40.
- ISABEL N.; BEAULIEU, J.; BOUSQUET, J.; 1995: Complete congruence between gene diversity estimates derived from genotypic data at enzyme and random amplified polymorphic DNA loci in black spruce. *Proc. Nat. Acad. Sci. USA* **92**, 6369-6373.
- ISABEL, N.; BEAULIEU, J.; THÉRIAULT, P.; BOUSQUET, J.; 1999: Direct evidence for biased gene diversity estimates from dominant random amplified polymorphic DNA (RAPD) fingerprints. *Mol. Ecol.* **8**, 477-483.
- JAMAUX-DESPRÉAUX, I.; BERGER, G.; PÉROS, J-P.; 1997: Characterization of microflora putatively involved in Esca syndrome on grapevine. In: *Proceedings of the 10th Congress of the Mediterranean Phytopathological Union*, June 1997, 39-41. Montpellier, France.
- LARIGNON, P.; DUBOS, B.; 1997: Fungi associated with esca disease in grapevine. *Eur. J. Plant Pathol.* **103**, 147-157.
- LYNCH, M.; MILLIGAN, B.G.; 1994: Analysis of population genetic structure with RAPD markers. *Mol. Ecol.* **3**, 991-99.
- McKEEN, C.; 1952: A cultural and taxonomic study of three species of *Peniophora*. *Can. J. Bot.* **30**, 764-787.
- MUGNAI, L.; GRANITI, A.; SURICO, G.; 1999: Esca (Black Measles) and brown wood-streaking: Two old and elusive diseases of grapevine. *Plant Dis.* **83**, 404-418.
- MUGNAI, L.; SURICO, G.; ESPOSITO, A.; 1996: Microflora associata al mal dell'esca della vite in Toscana. *Inform. Fitopatol.* **46**, 49-55.
- NEI, M.; TAJIMA, F.; 1981: DNA polymorphism detectable by restriction endonucleases. *Genetics* **97**, 145-163.
- POLLASTRO, S.; ABBATECOLA, A.; DONGIOVANNI, C.; FARETRA, F.; 2000 a: Usage of molecular markers (PCR-RAPD) for studying genetic variability in *Phellinus (Fomitiporia)* sp. *Phytopathol. Mediterr.* **39**, 107-111.
- POLLASTRO, S.; DONGIOVANNI, C.; ABBATECOLA, A.; FARETRA, F.; 2000 b: Observations on the fungi associated with esca and on spatial distribution of esca-symptomatic plants in Apulian (Italy) vineyards. *Phytopathol. Mediterr.* **39**, 206-210.
- RAYMOND, M.; ROUSSET, F.; 1995: GENEPOP (Version 1.2): Population genetics software for exact tests and ecumenism. *J. Hered.* **86**, 248-249.
- RAYNER, A.D.M.; BODDY, L.; 1988: *Fungal decomposition of wood. Its biology and ecology*. John Wiley and Sons, Chichester, UK.
- RISHBETH, J.; 1959: Dispersal of *Fomes annosus* Fr. and *Peniophora gigantea* (Fr.) Masee. *Trans. Brit. Mycol. Soc.* **42**, 243-260.
- RIZZO, D.M.; RENTMEESTER, R.M.; BURDSALL, H.H.; 1995: Sexuality and somatic incompatibility in *Phellinus gilvus*. *Mycologia* **87**, 805-820.
- RYVARDEN, L.; GILBERTSON, R.L.; 1994: *European Polypores. Part 2. Fungiflora*, Oslo, Norway.
- SOUZA, V.; NGUYEN, T.T.; HUDSON, R.R.; PINERO, D.; LENSKI, R.E.; 1992: Hierarchical analysis of linkage disequilibrium in *Rhizobium* populations: Evidence for sex? *Proc. Nat. Acad. Sci. USA* **89**, 8389-8393.
- SCHNEIDER, S.; ROESSLI, D.; EXCOFFIER, L.; 2000: Arlequin ver. 2.000: A Software for Population Genetics Data Analysis. Genetics and Biometry Laboratory, University of Geneva, Switzerland.
- SLATKIN, M.; 1987: Gene flow and the geographic structure of natural populations. *Science* **236**, 787-792.
- SPARAPANO, L.; BRUNO, G.; CICCARONE, C.; GRANITI, A.; 2000: Infection of grapevines by some fungi associated with esca. I. *Fomitiporia punctata* as a wood-rot inducer. *Phytopathol. Mediterr.* **39**, 46-52.
- SURICO, G.; MARCHI, G.; BRACCINI, P.; MUGNAI, L.; 1999: Spatial distribution of esca-diseased grapevine plants in five vineyards in Tuscany (Italy). *J. Plant Pathol.* **81**, 238 (abstract).
- SURICO, G.; MARCHI, G.; FERRANDINO, F. J.; BRACCINI, P.; MUGNAI, L.; 2000: Analysis of the spatial spread of esca in some Tuscan vineyards (Italy). *Phytopathol. Mediterr.* **39**, 211-224.
- STENLID, J.; KARLSSON, J. O.; HÖGGERG, N.; 1994: Intraspecific genetic variation in *Heterobasidion annosum* revealed by amplification of minisatellite DNA. *Mycol. Res.* **98**, 57-63.
- SZMIDT, A. E.; WANG, X. R.; LU, M. Z.; 1996: Empirical assessment of allozyme and RAPD variation in *Pinus sylvestris* (L.) using haploid tissue analysis. *Heredity* **76**, 412-420.
- VIALA, P.; 1926: Recherches sur les maladies de la vigne. *Ann. Epiphyt.* **12**, 5-108.
- WALLA, J. A.; 1984: Incidence of *Phellinus punctatus* on living woody plants in North Dakota. *Plant Dis.* **68**, 252-253.
- WILLIAMS, J. G. K.; KUBELIK, A. R.; LIVAK, K. J.; RAFALSKI, J. A.; TINGEY, S. V.; 1990: DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucl. Acids Res.* **18**, 6531-6535.

Received November 25, 2002