

A new experimental system to study the virus entry into plant growing points

Mayada Abdelgalil, Lilya Kopertekh, Joachim Schiemann
Julius Kühn-Institut, Institute for Biosafety of Genetically Modified Plants
mayada.abdelgalil@jki.bund.de

Understanding the mechanisms of plant-virus interaction is an essential prerequisite to develop appropriate defence approaches against pathogens. Plant growing points containing the shoot apical meristem have been thought to be protected from virus infection, particularly PVX. We developed a new experimental system, consisting of PVX and recombinant PVX-Cre (P1 recombinase), to study virus entry into growing points. In comparison to PVX infected plants, PVX-Cre infected plants were more severely diseased, stunted and later displayed a “recovery” phenotype. It is known that the recovery-inducing RNA viruses have the unusual ability to infect meristems (RATCLIFF *et al.*, 1999). Based on phenotypic observations of PVX and PVX-Cre infected *N. benthamiana* plants we proposed a more efficient apical colonization by PVX-Cre. To test this hypothesis, Cre-transgenic *N. benthamiana* plants were inoculated with PVX-GFP. Most of the apical shoots investigated exhibited GFP fluorescence. Results reported from studies with other RNA viruses indicated that RNA silencing is implicated in virus meristem exclusion (Qu *et al.*, 2005; Schwach *et al.*, 2005). To investigate if PVX-Cre is able to suppress RNA silencing, we performed a transient expression assay using *N.*

benthamiana plants and a *gfp*-expressing construct. In the absence of PVX-Cre the *gfp* expression was weak most probably due to silencing. Co-expression of PVX-Cre with the *gfp*-containing construct leads to enhanced *gfp* expression. These findings indicate that PVX-Cre might provide a silencing suppressor activity, which allows overcoming the meristem exclusion. Detailed analysis of this phenomenon is in progress.

QU, F., X. H. YE, G. C. HOU, S. SATO T. E., CLEMENTE and T. J. MORRIS, 2005: RDR6 has a broad-spectrum but temperature-dependent antiviral defence role in *Nicotiana benthamiana*. *J. Virol.* **79**, 15209-15217.

RATCLIFF, F. G., S. A. MACFARLANE and D. C. BAULCOMBE, 1999: Gene silencing without DNA:RNA-mediated cross-protection between viruses. *Plant Cell.* **11**, 1207-1215.

SCHWACH, F., F. E. VAISTIJ, L. JONES and D. C. BAULCOMBE, 2005. An RNA-dependent RNA polymerase prevents meristem invasion by potato virus X and is required for the activity but not the production of a systemic silencing signal. *Plant Physiol.* **138**, 1842-1852.