

LETTER TO THE EDITOR

Discovery Phytomedicine 2019, Volume 6, Number 4: 172-173

The proliferation and differentiation of pre-osteoblastic MC3T3-E1 cells from Vietnamese drug formulations

Pham Thi Thuy Hang,¹ Chu Thi Bich Viet,¹ and Ninh The Son^{2*}

*Correspondence to:

Ninh The Son Ph.D, Institute of Natural Products Chemistry, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam
yamantson@gmail.com

Cite This Article: Hang, P.T.T., Viet, C.T.B., Son, N.T. 2019. The proliferation and differentiation of pre-osteoblastic MC3T3-E1 cells from Vietnamese drug formulations. *Discovery Phytomedicine* 6(4): 172-173. DOI: [10.15562/phytomedicine.2019.96](https://doi.org/10.15562/phytomedicine.2019.96)

DEAR EDITORS

A public health problem, osteoporosis, is recognized as the prevalent disease, and mainly causes for impairment and loss mass of bone. It closely related the balance between bone formation by osteoblasts and resorption by osteoclasts during the remodeling cycle of bone.^{1,2} Hence, pharmaceutical therapies are looking for the potential agents to stimulate osteoblastic bone formation, as well as inhibit osteoclastic processes.³

In Vietnamese traditional medicine, the combination of the three species, including *Dillenia Indica* (local name “So”), *Schefflera racemosa* (local name “May tang”), *Adiantum raddianum* (local name “Cut nich”), were widely used as a herbal drug of osteoarthritis by several ethnic minorities in the mountainous areas of Laocai-Vietnam. However, up to now, no studies improve the roles of these crucial medicinal herbal plants. In current paper, we set out to investigate the effects of Vietnamese drug formulations of three plants on the proliferation and differentiation of pre-osteoblastic MC3T3-E1 cells through golden criteria, including cell viability, alkaline phosphatase activity (ALP), collagen content, and mineralization (calcium deposition). The dried leaves of these herbal plants were used as materials, whereas utilizing water-ethanol (1:1, v/v) as a solvent for extraction. According to traditional uses, we herein proposed the preparation of three drug formulations CT1-CT3, followed by the material rates, comprising formulation CT1 (3 : 1 : 1, *D. Indica* : *S. racemosa* : *A. raddianum*), formulation CT2 (2 : 1 : 1, respectively), formulation CT3 (1 : 1 : 1, respectively). The osteoblastic protocols were performed as described by.^{4,5}

The fluctuation of viability of pre-osteoblastic MC3T3-E1 cells cultured with or without the presence of three formulations were displayed in [Figure 1A](#). Data clearly revealed that, at the concentrations 50 and 100 µg/mL and 3 days incubation, only CT2 increased while the remainders failed to do so. Therefore, The formulation CT2 can be claimed responsible for next steps. A biochemical method, alkaline phosphatase activity, can be seen as a osteoblast phenotype marker, which provided quantitative information for the initial differentiation.^{4,6} At the concentrations 50 and 100 µg/mL, the drug formulation CT2 significantly promoted the increases of ALP activity up to 7.5 and 20% after 7 days, respectively, as compared to that of the control ([Table 1](#)). Collagen acted as a predominant product in bone cell during the osteoblastic differentiation.² The effects of CT2 on the synthetic collagen were investigated, using Sirius red based colorimetric experiment. As shown in [Table 1](#), accounting for 6 and 25% in the accumulation of collagen was induced by this drug formulation at the corresponding concentrations of 50 and 100 µg/mL and after 10 days treatment. Histochemical analysis also confirmed that, under two concentration conditions, pre-osteoblastic MC3T3-E1 cells took higher colors than that of untreated group ([Figure 1B](#)). During the osteoblastic differentiation, calcium accumulated in the matrix and cell layer of pre-osteoblastic MC3T3-E1 cells, the observation of mineralization was normally assessed by Alirazin red staining.^{2,4,5} After 15 days treatment, the formulation CT2 (50 and 100 µg/mL) notably increased the mineralization content of respective 14 and

Table 1 Effects of drug formulation CT2 (50 and 100 µg/mL) on ALP activity, collagen content, and mineralization of MC3T3-E1; Data are expressed as mean ± SEM (n = 3), *P < 0.05, **P < 0.01 versus control (only MC3T3-E1)

Concentration (mg/µL)	ALP activity (%)	Collagen content (%)	Mineralization (%)
100	119.70 ± 3.94**	124,55 ± 6.74**	128.65 ± 5.02**
50	107.49 ± 0.61*	105,83 ± 5.56	114.38 ± 2.74*
0	100.00 ± 0.94	100.00 ± 0.85	100.00 ± 2.89

¹Lao Cai School, Vietnam;²Institute of Natural Products Chemistry, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam




Figure 1A Effects of drug formulation CT1-CT3 (50, 100 and 200 µg/mL) on viability of MC3T3-E1; Data are expressed as mean ± SEM (n = 3), *P < 0.05 versus control (only MC3T3-E1)




Figure 1B Sirius red staining: a) only MC3T3-E1, b) MC3T3-E1 + CT2 (50 µg/mL), c) MC3T3-E1 + CT2 (100 µg/mL)




Figure 1C Alizarin red staining: a) only MC3T3-E1; b) MC3T3-E1 + CT2 (50 µg/mL); c) MC3T3-E1 + CT2 (100 µg/mL)

29%, especially the bright red color was observed at 100 µg/mL (Table 1 and Figure 1C).

Taken together, at concentration 100 µg/mL, Vietnamese combination drug of three plants

(2 : 1 : 1, *D. Indica* : *S. racemosa* : *A. raddianum*) evidently generated the prospective values in the treatment of bone related diseases. Therefore, extensive researches in either further biological experiments or phytochemical investigations are expected.

ACKNOWLEDGEMENTS

This work was supported by a grant from Laocai Education Department-2017.

REFERENCE

- Kumagai M, Mishima T, Wanatabe A, Harada T, Yoshida I, Fujita K, Watai M, Tawata S, Nishikawa K, Morimoto Y. 5,6-Dehydrokawain from *Alpinia zerumbet* promotes osteoblastic MC3T3-E1 cell differentiation. *Biosci Biotechnol Biochem.* 2016; 80:1425–1432 <https://doi.org/10.1080/09168451.2016.1153959>.
- Thu HE, Mohamed IN, Hussain Z, Mohamed N, Shuid AN. *Eurycoma longifolia*, a Malaysian medicinal herb, significantly upregulates proliferation and differentiation in pre-osteoblasts (MC3T3-E1): An *in vitro* model. *Int J Pharm Pharm Sci.* 2016;8:199–204 <https://doi.org/10.22159/ijpps.2016v8i11.14518>.
- Son NT, A review on the medicinal plant *Dalbergia odorifera* species: Phytochemistry and biological activity. *Evid Based Complement Alternat Med.* 2017;2017:ID 7142370 <https://doi.org/10.1155/2017/7142370>.
- Tai BH, Cuong NM, Huong TT, Choi EM, Kim JA, Kim YH. Chrysoeriol isolated from the leaves of *Eurya ciliata* stimulates proliferation and differentiation of osteoblastic MC3T3-E1 cells. *J Asian Nat Prod Res.* 2009;11:817–823 <https://doi.org/10.1080/10286020903117317>.
- Tai BH, Huyen VT, Huong TT, Nhiem NX, Choi EM, Kim JA, Long PQ, Cuong NM, Kim YH. New pyrano-pyrone from *Goniolamus tamirensis* enhances the proliferation and differentiation of osteoblastic MC3T3-E1 cells. *Chem Pharm Bull.* 2010;58:521–525 <https://doi.org/10.1248/cpb.58.521>.
- Lee YK, Song J, Lee SB, Kim KM, Choi SH, Kim CK, Legeros RZ, Kim KN. Proliferation, differentiation, calcification of preosteoblast-like MC3T3-E1 cells cultured onto noncrystalline calcium phosphate glass. *J. Biomed Mater Res A* 2004;69:188-195 <https://doi.org/10.1002/jbm.a.20137>.



This work is licensed under a Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>