

Final-state interactions in $e^+e^- \rightarrow t \bar{t} \rightarrow b l^+ \nu_b W^-$ near the top quark threshold

| | |
|------------------------------|---|
| 著者 | 隅野 行成 |
| journal or publication title | Physical review. D |
| volume | 57 |
| number | 11 |
| page range | 6912-6927 |
| year | 1998 |
| URL | http://hdl.handle.net/10097/35284 |

doi: 10.1103/PhysRevD.57.6912

Final-state interactions in $e^+e^- \rightarrow t\bar{t} \rightarrow bl^+\nu\bar{b}W^-$ near the top quark threshold

M. Peter* and Y. Sumino†

Institut für Theoretische Teilchenphysik, Universität Karlsruhe, D-76128 Karlsruhe, Germany

(Received 4 August 1997; published 4 May 1998)

We calculate final-state interaction corrections to the energy-angular distribution of l^+ in semileptonic top quark decay, where the parent top quark is produced via $e^+e^- \rightarrow t\bar{t}$ near threshold. These are the corrections due to gluon exchange between t and \bar{b} (\bar{t} and b) and between b and \bar{b} . Combining with other previously known corrections, we explicitly write down the l^+ energy-angular distribution including the full $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections near the $t\bar{t}$ threshold. Numerical analyses of the final-state interaction corrections are given. We find that they deform the l^+ distribution typically at the 10% level. We also find that all qualitative features of the numerical results can be understood from intuitive pictures. The mechanisms of various effects of the final-state interactions are elucidated. Finally we define an observable which is proper to the decay process of the top quark (dependent only on $d\Gamma_{t \rightarrow bl^+\nu}/dE_l d\Omega_l$ of a free polarized top quark) near the $t\bar{t}$ threshold. Such a quantity will be useful in extracting the decay properties of the top quark using the highly polarized top quark samples. [S0556-2821(98)04411-7]

PACS number(s): 13.65.+i, 14.65.Ha

I. INTRODUCTION

A future e^+e^- linear collider operating at energies around the $t\bar{t}$ threshold will be one of the ideal testing grounds for unraveling the properties of the top quark. So far there have been a number of studies of the cross section for top quark pair production near the $t\bar{t}$ threshold, both theoretical and experimental [1–19], in which it has been recognized that this kinematical region is rich in physics and is also apt for extracting various physical parameters efficiently, e.g., m_t , α_s , Γ_t , m_H , g_{tH} , etc.

While most of the previous analyses were solely concerned with the production process of the top quark, one may also analyze the decay process in detail and extract some important physics information. Especially the fact that t and \bar{t} are produced highly polarized in the threshold region is potentially quite advantageous for studying the electroweak properties of the top quark through its decay. Detailed investigations of the decay of free polarized top quarks have already been available including the full $\mathcal{O}(\alpha_s)$ corrections [16,20]. Close to threshold, however, these precise analyses do not apply directly because of the existence of corrections unique to this region, which connect the production and decay processes of the top quark. Specifically, these are the corrections due to gluon exchange between t and \bar{b} (\bar{t} and b) or between b and \bar{b} .

This type of corrections arises when the particles produced decay quickly into many particles, and are referred to as final-state interactions, rescattering corrections, or nonfactorizable corrections in the literature. They generally vanish in inclusive cross sections [9–11], but modify the shape of

differential distributions in a nontrivial manner [10,15,19]. The size of the corrections is at the 10% level in the $t\bar{t}$ threshold region, hence it is inevitable to incorporate their effects in precision studies of top quark production and decay near threshold. The same kind of effects has recently been studied in W pair production [21–24].

The first analysis of the top quark decay in the threshold region was given as a part of the results in Ref. [19]. In that paper, the mean value $\langle n \rangle$ of the charged lepton four-momentum projection on an arbitrarily chosen four-vector n in semileptonic top-quark decays was proposed as an experimentally observable quantity sensitive to top quark polarization, and this quantity was calculated including the final-state interactions. Clearly, and also admittedly in that work, the calculation of the differential distribution of l^+ including the final-state interactions has been demanded.

In this paper, we calculate these final-state interaction corrections to the differential energy-angular distribution of the charged leptons. We find that the corrections deform the l^+ distribution nontrivially at the expected level. Combining with other, previously known results, we write down the explicit formula for the l^+ energy-angular distribution including all $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections near the $t\bar{t}$ threshold.

Another aim of this paper is to present physical descriptions of the final-state interactions which enable us to qualitatively understand the features of our numerical results. Such descriptions would be useful since the systematic calculation of final-state interactions based on quantum field theory is rather complicated, involving box- and pentagon-type diagrams, and it is not easy to make physical sense out of the obtained final expressions. To our knowledge such qualitative explanations have never been put forth, although corresponding theoretical calculations and numerical studies have been partly available.

Finally we propose an observable which is proper to the decay process of the top quark near threshold. Since the final-state interaction connects the production and decay processes of the top quark, it destroys the factorization property

*Present address: Institut für Theoretische Physik, Universität Heidelberg, Philosophenweg 16, D-69120 Heidelberg, Germany.

†On leave of absence from Department of Physics, Tohoku University, Sendai 980-77, Japan.

of the corresponding production and decay cross sections. In order to study the decay of the top quark in a clean environment, it would be useful if we could find an observable which depends only on this process ($d\Gamma_{t \rightarrow bl^+v}/dE_l d\Omega_l$ of a free polarized top quark). In fact, such an observable can be constructed, which at the same time preserves most of the differential information of the l^+ energy-angular distribution.

In our numerical analysis we solve the Schrödinger equation numerically in order to include the QCD binding effects near threshold. We follow both the coordinate-space approach developed in Refs. [2,3] and the momentum-space approach developed in Refs. [4,5] in solving the equation, and compare the results. There are small differences in the numerical results obtained from the two approaches, reflecting the difference in the construction of the potentials at short distance. [The difference is formally $\mathcal{O}(\alpha_s^2)$, of the order beyond our present scope.] This issue is also discussed.

In Sec. II we introduce some notations to be used in later sections. Section III contains the physical descriptions of the effects of final-state interactions. The results of the systematic calculation of final-state interaction corrections to the l^+ energy-angular distribution, as well as the complete formula for the distribution including all $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections, are given in Sec. IV. Section V shows various numerical results and a comparison with the qualitative picture. We define an observable proper to the top-quark decay process in Sec. VI. Discussion and conclusion are presented in Secs. VII and VIII, respectively.

II. DEFINITIONS AND CONVENTIONS

We consider longitudinally polarized e^+e^- beams throughout our analyses. P_{e^\pm} denotes the longitudinal polarization of e^\pm , and we set

$$\chi = \frac{P_{e^+} - P_{e^-}}{1 - P_{e^+} P_{e^-}}. \quad (1)$$

We choose a reference coordinate system in the $t\bar{t}$ c.m. frame. Three orthonormal basis vectors are defined as

$$\hat{\mathbf{n}}_{\parallel} = \frac{\mathbf{p}_{e^-}}{|\mathbf{p}_{e^-}|}, \quad \hat{\mathbf{n}}_N = \frac{\mathbf{p}_{e^-} \times \mathbf{p}_t}{|\mathbf{p}_{e^-} \times \mathbf{p}_t|}, \quad \hat{\mathbf{n}}_{\perp} = \hat{\mathbf{n}}_N \times \hat{\mathbf{n}}_{\parallel}, \quad (2)$$

where \mathbf{p}_{e^-} and \mathbf{p}_t represent the e^- and t momentum, respectively.

Our conventions for the fermion vector and axial-vector couplings to the Z boson are

$$v_f = 2I_f^3 - 4q_f \sin^2 \theta_W, \quad a_f = 2I_f^3, \quad (3)$$

respectively. Certain combinations of these couplings will be useful below:

$$a_1 = q_e^2 q_t^2 + (v_e^2 + a_e^2) v_t^2 d^2 + 2q_e q_t v_e v_t d,$$

$$a_2 = 2v_e a_e v_t^2 d^2 + 2q_e q_t a_e v_t d,$$

$$a_3 = 4v_e a_e v_t a_t d^2 + 2q_e q_t a_e a_t d,$$

$$a_4 = 2(v_e^2 + a_e^2) v_t a_t d^2 + 2q_e q_t v_e a_t d,$$

$$d = \frac{1}{16 \sin^2 \theta_W \cos^2 \theta_W} \frac{s}{s - M_Z^2}, \quad (4)$$

and

$$C_{\parallel}^0(\chi) = -\frac{a_2 + \chi a_1}{a_1 + \chi a_2},$$

$$C_{\parallel}^1(\chi) = (1 - \chi^2) \frac{a_2 a_3 - a_1 a_4}{(a_1 + \chi a_2)^2},$$

$$C_{\perp}(\chi) = -\frac{1}{2} \frac{a_4 + \chi a_3}{a_1 + \chi a_2},$$

$$C_N(\chi) = -\frac{1}{2} \frac{a_3 + \chi a_4}{a_1 + \chi a_2} = -C_{\text{FB}}(\chi). \quad (5)$$

Finally, we define

$$y = \frac{M_W^2}{m_t^2}, \quad \kappa = \frac{1 - 2y}{1 + 2y}. \quad (6)$$

The QCD enhancement of the top-quark production cross section near threshold is incorporated through the S -wave and P -wave Green's functions $\bar{G}(p, E)$ and $\bar{F}(p, E)$ of the nonrelativistic Schrödinger equations in the presence of the QCD potential. These functions are defined by

$$\left[-\frac{\nabla^2}{m_t} + V_{\text{QCD}}(r) - (E + i\Gamma_t) \right] G(\mathbf{x}, E) = \delta^3(\mathbf{x}), \quad (7)$$

$$\left[-\frac{\nabla^2}{m_t} + V_{\text{QCD}}(r) - (E + i\Gamma_t) \right] F^k(\mathbf{x}, E) = -i \partial^k \delta^3(\mathbf{x}), \quad (8)$$

and

$$\bar{G}(p, E) = \int d^3 \mathbf{x} e^{-i\mathbf{p} \cdot \mathbf{x}} G(\mathbf{x}, E), \quad (9)$$

$$p^k \bar{F}(p, E) = \int d^3 \mathbf{x} e^{-i\mathbf{p} \cdot \mathbf{x}} F^k(\mathbf{x}, E). \quad (10)$$

One may obtain the Green's functions either by first solving the Schrödinger equations in coordinate space and taking the Fourier transforms of the solutions [3], or by solving the Schrödinger equations directly in momentum space [4].

Various known $\mathcal{O}(\alpha_s)$ corrections to threshold cross sections can also be expressed in terms of the above Green's functions. In Ref. [19] the following functions have been defined, which are solely determined from QCD, to represent various independent corrections. We will pursue the same conventions in this paper:

$$\varphi(p, E) = \frac{[1 - C_F \alpha_s(m_t)/\pi] p \tilde{F}^*(p, E)}{[1 - 2C_F \alpha_s(m_t)/\pi] m_t \tilde{G}^*(p, E)}, \quad (11)$$

$$\varphi_R = \text{Re}\varphi, \quad \varphi_I = \text{Im}\varphi, \quad (12)$$

$$\begin{aligned} \psi_1(p, E) &= -C_F \cdot 4\pi\alpha_s(\mu_B) \\ &\times \int \frac{d^3\mathbf{q}}{(2\pi)^3} \frac{1}{|\mathbf{q} - \mathbf{p}_t|^3} 2\text{Im} \left[\frac{\tilde{G}(q, E)}{\tilde{G}(p, E)} \right] \frac{\pi}{2}, \end{aligned} \quad (13)$$

$$\begin{aligned} \psi_R(p, E) &= -C_F \cdot 4\pi\alpha_s(\mu_B) \\ &\times \text{Pr} \int \frac{d^3\mathbf{q}}{(2\pi)^3} \frac{1}{|\mathbf{q} - \mathbf{p}_t|^3} \frac{\mathbf{p}_t \cdot (\mathbf{q} - \mathbf{p}_t)}{|\mathbf{p}_t| |\mathbf{q} - \mathbf{p}_t|} 2\text{Re} \left[\frac{\tilde{G}(q, E)}{\tilde{G}(p, E)} \right]. \end{aligned} \quad (14)$$

Here $C_F=4/3$ is a color factor; $p=|\mathbf{p}_t|$ and $q=|\mathbf{q}|$, μ_B is taken to be 15 GeV in our analyses. Equations (13) and (14) differ slightly in their integrands from those defined in Ref. [19].¹ The differences, however, can be regarded as higher order corrections which are beyond the scope of our analysis.

III. QUALITATIVE DESCRIPTION OF FINAL-STATE INTERACTION EFFECTS

In this section a qualitative understanding of the effects of final-state interactions is explained. It is based on the classical picture that t and \bar{t} (\bar{t} and b) attract each other due to their color charges. We will see that all the following qualitative features match well with our numerical studies presented in Sec. V. Moreover, the following argument will help interpreting the formulas derived in the next section.

A. Top momentum distribution

Let us first consider the effect of final-state interactions on the top quark momentum ($|\mathbf{p}_t|$) distribution, where \mathbf{p}_t is reconstructed from the bW^+ momenta at time $\tau \rightarrow \infty$. The energy of a $t\bar{t}$ system before its decay is given by

$$\mathcal{E} = E_t + E_{\bar{t}} + V(|\mathbf{r}_t - \mathbf{r}_{\bar{t}}|), \quad (15)$$

where

$$E_t = m_t + \frac{\mathbf{p}_t^2}{2m_t}, \quad E_{\bar{t}} = m_t + \frac{\mathbf{p}_{\bar{t}}^2}{2m_t}, \quad V(r) = -C_F \frac{\alpha_s}{r}. \quad (16)$$

¹In Ref. [19] the Coulomb propagator between b and \bar{t} (\bar{b} and t) together with the color charges are replaced by the QCD potential between two heavy quarks by hand $-C_F \cdot 4\pi\alpha_s/|\mathbf{q} - \mathbf{p}_t|^2 \rightarrow \tilde{V}_{\text{QCD}}(|\mathbf{q} - \mathbf{p}_t|)$. One advantage of Eqs. (13) and (14) is that one may convert them into one-parameter integral forms by explicitly integrating over $d\Omega_{\mathbf{q}}$ [10,13].




FIG. 1. Attractive Coulomb force between \bar{t} and b (from t decay). The momentum transfer $\delta\mathbf{p}_b = -\delta\mathbf{p}_{\bar{t}}$ due to the attraction is indicated by thick arrows.

Suppose t decays first and let $a \sim (\alpha_s m_t)^{-1} \sim (m_t \Gamma_t)^{-1/2}$ be the typical distance between t and \bar{t} at the time of t decay. Then, just before the decay, the momenta of t and \bar{t} are given by

$$|\mathbf{p}_t| = |\mathbf{p}_{\bar{t}}| \sim \sqrt{m_t [\mathcal{E} - 2m_t + |V(a)]}. \quad (17)$$

Their order of magnitude is $\alpha_s m_t \sim (\text{Bohr radius})^{-1}$. If it were not for the final-state interactions between \bar{t} and b , and if \bar{t} , b , and W^+ traveled as free particles, the above momentum would be transferred to the bW^+ system at time $\tau \rightarrow \infty$:

$$|\mathbf{p}_b + \mathbf{p}_{W^+}|_{\tau \rightarrow \infty} \sim \sqrt{m_t [\mathcal{E} - 2m_t + |V(a)]}. \quad (18)$$

Taking into account the final-state interaction (Coulomb interaction) between \bar{t} and b , the energy of the $\bar{t}bW^+$ system is given by²

$$\mathcal{E} = E_b + E_{W^+} + E_{\bar{t}} + V(|\mathbf{r}_b - \mathbf{r}_{\bar{t}}|), \quad (19)$$

$$E_b = |\mathbf{p}_b|, \quad E_{W^+} = \sqrt{\mathbf{p}_{W^+}^2 + M_{W^+}^2}. \quad (20)$$

As depicted in Fig. 1, the tracks of b and \bar{t} are deflected due to the attraction between the two particles, which lose kinetic energy as b flies off to infinity at the speed of light. The classical equation of motion is given by

$$\frac{d\mathbf{p}_b}{d\tau} = -\frac{d\mathbf{p}_{\bar{t}}}{d\tau} = -\frac{\partial}{\partial \mathbf{r}_b} V(|\mathbf{r}_b - \mathbf{r}_{\bar{t}}|). \quad (21)$$

Substituting the free particle solution $\mathbf{r}_b = \mathbf{v}\tau + \mathbf{r}_0$ on the right-hand side and noting $|\mathbf{r}_b - \mathbf{r}_{\bar{t}}| \simeq \tau$, we may estimate the size of the momentum transfer due to the attractive force as³

$$|\delta\mathbf{p}_b| = |\delta\mathbf{p}_{\bar{t}}| \sim |V(r_{\min})|, \quad (22)$$

where the minimum distance between b and \bar{t} is denoted as r_{\min} . Typically $r_{\min} \sim a$, and we find from Eq. (22) that the

²Here we neglect the interaction of \bar{t} (b) and the magnetic field generated by b (\bar{t}). This approximation is justified in the case of our interest; see Sec. IV.

³It corresponds to solving the equation of motion by a series expansion of α_s .




FIG. 2. Typical configurations in the decay of \bar{t} with definite spin orientation. Transverse $W^- (W_T^-)$ tend to be emitted in the direction of the \bar{t} spin orientation, while longitudinal $W^- (W_L^-)$ are emitted in the opposite direction due to helicity conservation. For $m_t \approx 175$ GeV, \bar{t} decays mainly to W_L^- , hence \bar{b} is emitted more in the \bar{t} spin direction.

effect of final-state interactions on the $|\mathbf{p}_t|$ distribution is of the order of $|\delta\mathbf{p}_t| \sim \alpha_s^2 m_t$. Obviously the effect is to reduce $|\mathbf{p}_t|$.

B. Forward-backward asymmetric distribution

Next we consider the $\cos \theta_{te}$ distribution of the top quark. (θ_{te} denotes the angle between t and e^- in the $t\bar{t}$ c.m. frame.) It is known that a forward-backward asymmetric distribution of the top quark is generated by the final-state interactions [10,19]. We describe its mechanism here.

We consider the case where \bar{t} decays first and examine the interaction between t and \bar{b} . The t and \bar{t} pair produced near threshold in e^+e^- collisions have their spins approximately parallel or antiparallel to the e^- beam direction and the spins are always oriented parallel to each other. In fact in leading order the polarization vector of the top quark is given by $\mathcal{P} = C_{\parallel}^0(\chi) \hat{\mathbf{n}}_{\parallel}$ [25,17]. On the other hand, the decay of \bar{t} occurs via a $V-A$ coupling, and \bar{b} is emitted preferably in the spin direction of the parent \bar{t} , see Fig. 2. More precisely, the excess of the \bar{b} 's emitted in the \bar{t} spin direction over those emitted in the opposite direction is given by κ defined in Eq. (6). Now suppose t and \bar{t} have their spins in the $\hat{\mathbf{n}}_{\parallel}$ direction. Then \bar{b} will be emitted dominantly in the $\hat{\mathbf{n}}_{\parallel}$ direction. One can see from Fig. 3(a) that in this case t is always attracted to the forward direction due to the attractive force between t and \bar{b} . The direction of the attractive force will be opposite if t and \bar{t} have their spins in the $-\hat{\mathbf{n}}_{\parallel}$ direction [Fig. 3(b)].




FIG. 3. Attractive force between t and \bar{b} when the t and \bar{t} spins are oriented in (a) the $\hat{\mathbf{n}}_{\parallel}$ direction and in (b) the $-\hat{\mathbf{n}}_{\parallel}$ direction. The momentum transfer $\delta\mathbf{p}_b = -\delta\mathbf{p}_t$ due to the attraction is indicated by thick black arrows.




FIG. 4. Typical configurations of the particles in semileptonic decay of t when the b quark is emitted in the \bar{t} direction. Due to the boost by W^+ , the energy-angle correlation of l^+ will be either “ E_l is small and l^+ emitted in $-\mathbf{p}_l$ direction” or “ E_l is large and l^+ emitted in \mathbf{p}_l direction.”

Thus, polarized top quarks will be pulled in a definite (forward or backward) direction, and we may expect that a forward-backward asymmetric distribution of the top quark $\sim \kappa C_{\parallel}^0(\chi) \cos \theta_{te}$ is generated by the final-state interaction.

Incidentally, a forward-backward asymmetric distribution of the top quarks is also generated by the interference between the S -wave and P -wave $t\bar{t}$ pair-production amplitudes [6]. It is formally a quantity of the same order as the final-state interaction near threshold, since it arises as an $\mathcal{O}(\beta) = \mathcal{O}(\alpha_s)$ correction to the leading spherically symmetric distribution. Interestingly, we find that each $\cos \theta_{te}$ distribution has quite a different physical explanation for its generation mechanism, although it is a common feature that both originate from the interplay between QCD and electroweak interactions.

C. Top quark polarization vector

The Coulomb attraction between t and \bar{b} also modifies the top quark polarization vector [19]. As we have seen previously, in the \bar{t} decay, \bar{b} tends to be emitted in the direction of the parent \bar{t} 's spin direction (Fig. 2). We then find from Fig. 3 that if the t and \bar{t} spins are oriented in the $\hat{\mathbf{n}}_{\parallel}$ direction, t will be attracted to the forward direction due to the attraction by \bar{b} , and oppositely attracted to the backward direction if the t and \bar{t} spins are in the $-\hat{\mathbf{n}}_{\parallel}$ direction. This means that in the forward region ($\cos \theta_{te} \approx 1$) the number of t 's with spin in the $\hat{\mathbf{n}}_{\parallel}$ direction increases whereas in the backward region the number of those with spin in the opposite direction increases. Or equivalently, the $\hat{\mathbf{n}}_{\parallel}$ component of the top quark polarization vector increases in the forward region and decreases in the backward region. We may thus conjecture that the top quark polarization vector is modified as $\delta\mathcal{P} \sim \kappa \cos \theta_{te} \hat{\mathbf{n}}_{\parallel}$ due to the interaction between t and \bar{b} .

D. l^+ energy-angular distribution

Finally let us examine the effect of the Coulomb attraction between b and \bar{t} on the l^+ energy-angular distribution in the semileptonic decay of t . The b quark from t decay will be attracted in the direction of \bar{t} due to the Coulomb interaction between these two particles. We show schematically typical configurations of the particles in the top quark semileptonic decay in Fig. 4. It can be seen that if the probability for b being emitted in the \bar{t} direction increases, correspondingly the probability for particular l^+ energy-angular con-




FIG. 5. Born-type diagram for the process $e^+e^- \rightarrow t\bar{t} \rightarrow bl^+\nu\bar{b}W^-$.

figurations increases. These configurations are either “ E_l is small and l^+ emitted in $-\mathbf{p}_t$ direction” or “ E_l is large and l^+ emitted in \mathbf{p}_t direction.”

IV. l^+ ENERGY-ANGULAR DISTRIBUTION

We present the formulas for the charged lepton energy-angular distribution in the decay of a top quark that is produced via $e^+e^- \rightarrow t\bar{t}$ near threshold. In the following, $x_l = 2E_l/m_t$ and Ω_l , respectively, denote the normalized energy and the solid angle of the charged lepton as defined in the rest frame of the parent top quark. For simplicity we neglect the decay of W^- in our calculations.

A. Factorizable part

It is well known that the contribution of the Born-type (reducible) diagram (Fig. 5) to the differential distribution of t and l^+ has a form where the production and decay processes of the top quark are factorized:

$$\begin{aligned} \frac{d\sigma_{\text{Born}}(e^+e^- \rightarrow t\bar{t} \rightarrow bl^+\nu\bar{b}W^-)}{d^3\mathbf{p}_t dx_l d\Omega_l} &= \frac{d\sigma_{\text{Born}}(e^+e^- \rightarrow t\bar{t})}{d^3\mathbf{p}_t} \\ &\times \frac{1}{\Gamma_t} \frac{d\Gamma_{t \rightarrow bl^+\nu}(\mathcal{P}_{\text{Born}})}{dx_l d\Omega_l}. \end{aligned} \quad (23)$$

The above form holds true even including $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections to each vertex and propagator in the Born-type diagram. Here, Γ_t [26,27] and $d\Gamma_{t \rightarrow bl^+\nu}(\mathcal{P}_{\text{Born}})/dx_l d\Omega_l$ [28,16], respectively, are the width of a free top quark and the charged lepton (l^+) energy-angular distribution in the decay of a free polarized top quark, both including the corresponding full $\mathcal{O}(\alpha_s)$ corrections. Near threshold, the top quark production cross section for longitudinally polarized e^+e^- beams is given by [17,19]

$$\frac{d\sigma_{\text{Born}}(e^+e^- \rightarrow t\bar{t})}{d^3\mathbf{p}_t} = \frac{d\sigma_{t\bar{t}}^0}{d^3\mathbf{p}_t} [1 + 2C_{FB}(\chi)\varphi_R(p, E)\cos\theta_{te}], \quad (24)$$

$$\begin{aligned} \frac{d\sigma_{t\bar{t}}^0}{d^3\mathbf{p}_t} &= \frac{N_c\alpha^2\Gamma_t}{4\pi m_t^4} (1 - P_{e^+}P_{e^-})(a_1 + \chi a_2) \\ &\times \left(1 - \frac{4C_F\alpha_s(m_t)}{\pi}\right) |\bar{G}(p, E)|^2. \end{aligned} \quad (25)$$

Here, α is the fine structure constant and $N_c = 3$. $\mathcal{P}_{\text{Born}} = \mathcal{P}_{\parallel}\hat{\mathbf{n}}_{\parallel} + \mathcal{P}_{\perp}\hat{\mathbf{n}}_{\perp} + \mathcal{P}_N\hat{\mathbf{n}}_N$ represents the polarization vector of a top quark produced via the Born-type diagram (Fig. 5) near threshold. The components are given by

$$\mathcal{P}_{\parallel}(\mathbf{p}_t, E, \chi) = C_{\parallel}^0(\chi) + C_{\parallel}^1(\chi)\varphi_R(p, E)\cos\theta_{te}, \quad (26)$$

$$\mathcal{P}_{\perp}(\mathbf{p}_t, E, \chi) = C_{\perp}(\chi)\varphi_R(p, E)\sin\theta_{te}, \quad (27)$$

$$\mathcal{P}_N(\mathbf{p}_t, E, \chi) = C_N(\chi)\varphi_l(p, E)\sin\theta_{te}. \quad (28)$$

Let us review briefly how to derive the factorized form of the differential distribution, Eq. (23). First, in calculating the fully differential cross section $d\sigma/d\Phi_5(bl^+\nu\bar{b}W^-)$, one may replace the top quark momentum by an on-shell four-vector as

$$p_t^\mu = (p_t^0, \mathbf{p}_t) \rightarrow \tilde{p}_t^\mu = (\sqrt{\mathbf{p}_t^2 + m_t^2}, \mathbf{p}_t) \quad (29)$$

in vertices and propagator numerators (but not in propagator denominators). The replacement is justified because near threshold relevant kinematical configurations are determined by

$$p_t^0 - m_t \sim \alpha_s^2 m_t, \quad |\mathbf{p}_t| \sim \alpha_s m_t \quad (30)$$

so that the replacement induces differences only at $\mathcal{O}(\alpha_s^2)$, and also because we will not be concerned with the p_t^0 dependence of the cross section. Then, one may use the following identity to factorize the spinor traces that appear in the fully differential cross section into their production and decay parts: For an arbitrary 4×4 spinor matrix G and for a four-vector \tilde{p}_t^μ satisfying $\tilde{p}_t^2 = m_t^2$,

$$\frac{\tilde{p}_t + m_t}{2m_t} G \frac{\tilde{p}_t + m_t}{2m_t} = \frac{\tilde{p}_t + m_t}{2m_t} \frac{1 - \mathcal{P}\gamma_5}{2} C, \quad (31)$$

where the four-vector \mathcal{P}^μ and the constant C are determined from G and \tilde{p}_t^μ via the relation

$$\frac{1 - \mathcal{P} \cdot s}{2} C = \text{Tr} \left[\frac{\tilde{p}_t + m_t}{2m_t} G \frac{\tilde{p}_t + m_t}{2m_t} \frac{1 - \not{s}\gamma_5}{2} \right], \quad (32)$$

provided s^μ and \mathcal{P}^μ satisfy $s \cdot \tilde{p}_t = \mathcal{P} \cdot \tilde{p}_t = 0$.

One may also factorize the phase space as

$$\begin{aligned} d\Phi_5(\gamma^* \rightarrow bl^+\nu\bar{b}W^-) &= \frac{d^4p_t}{(2\pi)^4} d\Phi_3(t^* \rightarrow bl^+\nu) d\Phi_2(\bar{t}^* \rightarrow \bar{b}W^-), \end{aligned} \quad (33)$$

$$d\Phi_3(t^* \rightarrow bl^+\nu) = \left(\frac{1}{4\pi}\right)^5 dx_l d\Omega_l dp_W^2 d\phi_{bl}, \quad (34)$$




FIG. 6. Diagrams for final-state interactions for $e^+e^- \rightarrow t\bar{t} \rightarrow bl^+\nu\bar{b}W^-$: (a) Coulomb-gluon exchange between t and \bar{b} , (b) Coulomb-gluon exchange between \bar{t} and b , (c) Coulomb-gluon exchange between b and \bar{b} , and (d) transverse-gluon exchange between b and \bar{b} .

where p_W^2 is the invariant-mass-squared of $l^+\nu$, and ϕ_{bl} denotes the azimuthal angle of b around l^+ in the top quark rest frame. Then one integrates over dp_W^2 , $d\phi_{bl}$, $d\Phi_2(\bar{b}W^-)$, and $dp_t^0/(2\pi)$; the integration over p_W^2 is trivial since we use the narrow-width approximation for W^+ , the integration over $d\phi_{bl}$ is also trivial since the fully differential cross section is independent of ϕ_{bl} , the integration over the $\bar{b}W^-$ phase space merely replaces the $\bar{b}W^-$ wave functions by Γ_t , the $p_t^0/(2\pi)$ integration is straightforward.

B. Final-State Interaction Corrections

Corrections due to the final-state interactions (rescattering corrections) that originate from the irreducible diagrams (a)–(d) in Fig. 6 are important particularly in the threshold region. In fact their contributions are counted as $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections to the leading threshold enhancement [10]. We calculate the effect of each diagram on the l^+ energy-angular distribution. We chose Coulomb gauge for the QCD part in our calculations.

The contribution of diagram (a) (exchange of one Coulomb gluon between t and \bar{b}) can be regarded, after integrat-

ing over the $\bar{b}W^-$ phase space, as a correction to the production process of t . Thus, the production cross section $d\sigma_{\text{Born}}(e^+e^- \rightarrow t\bar{t})/d^3\mathbf{p}_t$ and the polarization vector $\mathcal{P}_{\text{Born}}$ of the top quark receive corrections by this diagram, whereas the decay distribution $d\Gamma_{t \rightarrow bl^+\nu}(\mathcal{P}_{\text{Born}})/dx d\Omega_l$ remains unaffected (except for the modification of $\mathcal{P}_{\text{Born}}$). In fact the contribution of this diagram can be incorporated by the following substitutions in Eq. (23):

$$\frac{d\sigma_{\text{Born}}(e^+e^- \rightarrow t\bar{t})}{d^3\mathbf{p}_t} \rightarrow \frac{d\sigma_{\text{Born}}(e^+e^- \rightarrow t\bar{t})}{d^3\mathbf{p}_t} (1 + \delta_a), \quad (35)$$

$$\mathcal{P}_{\text{Born}} \rightarrow \mathcal{P}_{\text{Born}} + \delta\mathcal{P}_a \quad (36)$$

with

$$\delta_a = \frac{1}{2} [\psi_1(p, E) + \kappa C_{\parallel}^0 \psi_R(p, E) \cos \theta_{te}], \quad (37)$$

$$\delta\mathcal{P}_a = \frac{1}{2} [1 - (C_{\parallel}^0)^2] \kappa \psi_R(p, E) \cos \theta_{te} \cdot \hat{\mathbf{n}}_{\parallel}. \quad (38)$$

The derivation of the formula goes as follows. Since the relevant kinematical configuration lies in the soft-gluon region, we can use soft-gluon approximation and factor out the part that depends on the loop-momentum q^μ (the propagators of the gluon \bar{b} , t , and \bar{t} together with the loop integral) outside the spinor structure, while the remaining part is similar to the fully differential cross section of the Born-type diagram. The latter part is factorized as before. Due to the soft-gluon factor (the factor pulled outside), at this stage we may interpret that the production cross section and polarization vector of top quark get corrections that depend on t and \bar{b} momenta. Integrations over dp_W^2 and $d\phi_{bl}$ are the same as for the Born-type diagram. For integrations over $d\Phi_2(\bar{b}W^-)$, $dp_t^0/(2\pi)$, and $dq^0/(2\pi)$, we follow the method described in Ref. [10], Appendix D. We are thus led to Eqs. (36)–(38).

Diagram (b) (exchange of one Coulomb gluon between \bar{t} and b) in Fig. 6 gives a correction that connects the production and decay processes of the top quark. In fact one may incorporate the contribution of this diagram by multiplying Eq. (23) by a factor $[1 + \xi(p, E, x_l, \cos \theta_{lt})]$, where

$$\xi(p, E, x_l, \cos \theta_{lt}) = C_F \cdot 4\pi\alpha_s(\mu_B) \int \frac{d^3\mathbf{q}}{(2\pi)^3} \frac{1}{|\mathbf{q}-\mathbf{p}_t|^3} \text{Re} \left[\frac{\tilde{G}^*(q, E)}{\tilde{G}^*(p, E)} \int_0^{2\pi} d\phi_{bl} \frac{|\mathbf{q}-\mathbf{p}_t|}{2\pi \hat{\mathbf{n}}_b \cdot (\mathbf{q}-\mathbf{p}_t) + i\epsilon} \right]. \quad (39)$$

$\hat{\mathbf{n}}_b$ denotes the unit vector in the direction of b . After integration over $d\Omega_q d\phi_{bl}$, one may reduce the expression to a one-parameter integral form as

$$\xi(p, E, x_l, \cos \theta_{lt}) = C_F \cdot 4\pi\alpha_s(\mu_B) \int_0^\infty dq \left\{ w_R \text{Re} \left[\frac{\tilde{G}^*(q, E)}{\tilde{G}^*(p, E)} \right] + w_I \text{Im} \left[\frac{\tilde{G}^*(q, E)}{\tilde{G}^*(p, E)} \right] \right\} \quad (40)$$

with

$$w_R = \frac{1}{4\pi^2} \frac{q}{p^2 - q^2} \{ \theta(z_+^2 - 1) \cosh^{-1}|z_+| - \theta(z_-^2 - 1) \cosh^{-1}|z_-| \}, \quad (41)$$

$$w_I = \frac{1}{4\pi^2} \frac{q}{|p^2 - q^2|} \{ \pi \theta[(1+z_+)(1+z_-)] - \theta(1-z_+^2) \cos^{-1}(z_+) - \theta(1-z_-^2) \cos^{-1}(z_-) \}, \quad (42)$$

$$z_{\pm} = \frac{q/p \pm \cos \theta_{Wl} \cos \theta_{lt}}{\sin \theta_{Wl} \sin \theta_{lt}}, \quad \cos \theta_{Wl} = \frac{1+y}{1-y} - \frac{2y}{x_l(1-y)}, \quad \cos \theta_{lt} = \frac{\mathbf{p}_l \cdot \mathbf{p}_l}{|\mathbf{p}_l| |\mathbf{p}_l|}. \quad (43)$$

Here, θ_{Wl} represents the angle between W^+ and l^+ in the t rest frame (given as a function of x_l); θ_{lt} represents the angle between t and l^+ in the $t\bar{t}$ c.m. frame⁴ ($0 \leq \theta_{Wl}, \theta_{lt} \leq \pi$). The inverse functions \cosh^{-1} and \cos^{-1} in the above formulas take their values within $[0, \infty)$ and $[0, \pi]$, respectively. $\theta(x)$ is the unit step function. It is understood that the principal value should be taken in the integration of the w_R term as $p \rightarrow q$.

We derived Eqs. (39) and (40) in the following manner. As in diagram (a), we used the soft-gluon approximation and factored out a soft-gluon factor (loop integral of the propagators of gluon b , t , and \bar{t}). The remaining part is same as the fully differential cross section of the Born-type diagram except for the t and \bar{t} propagators and Green's functions, which is again factorized. This time, however, the correction cannot be interpreted as associated with the top quark production process since the soft-gluon factor depends on the b momentum. The integrations over dp_W^2 and $d\Phi_2(\bar{b}W^-)$ are the same as those for the Born-type diagram. The function ξ given in Eq. (39) is essentially the soft-gluon factor inte-

grated over $dp_t^0/(2\pi)$, $d^4q/(2\pi)^4$, and $d\phi_{bl}$. Finally, to derive Eq. (40) from (39), it is simpler to integrate over $d\Omega_q$ before $d\phi_{bl}$.

Two noteworthy properties of $\xi(p, E, x_l, \cos \theta_{lt})$ are (1) its l^+ -angular dependence enters only through $\cos \theta_{lt}$ and is independent of the angle from the e^- beam direction or from the top quark polarization vector and (2) it is purely determined by the QCD interaction and free of the coupling parameters of electroweak interactions (except y). As a non-trivial cross check of the formula (40), we integrated $\xi \times \Gamma_t^{-1} d\Gamma_{t \rightarrow bl^+ \nu} / dx_l d\Omega_l$ over the lepton energy-angular variables $\int dx_l d\Omega_l$ analytically and reproduced the one-parameter integral formula [10,13] for the final-state interaction correction [from diagram (b)] to the top quark three-momentum distribution.

The contribution of diagram (c) (exchange of one Coulomb gluon between b and \bar{b}) vanishes within our approximation. We show it in steps. Using the soft-gluon approximation, the contribution of this diagram to the cross section can be written as

$$\begin{aligned} \frac{d\sigma_c}{d^3\mathbf{p}_l dx_l d\Omega_l} &= \int \frac{dp_t^0}{(2\pi)} dp_W^2 d\phi_{bl} d\Phi_2(\bar{b}W^-) \frac{1}{(2\pi)^3} \frac{1}{(4\pi)^5} \times T \\ &\times iC_F \cdot 4\pi\alpha_s \int \frac{d^4k}{(2\pi)^4} [D(p_t) + D(p_{\bar{t}})] [D^*(p_t+k) + D^*(p_{\bar{t}}-k)] \\ &\times \tilde{G}(p, E) \tilde{G}^*(|\mathbf{p}_t + \mathbf{k}|, E) \frac{1}{k^0 - \hat{\mathbf{n}}_b \cdot \mathbf{k} - i\epsilon} \frac{1}{-k^0 + \hat{\mathbf{n}}_{\bar{b}} \cdot \mathbf{k} - i\epsilon} \frac{1}{|\mathbf{k}|^2} + \text{c.c.}, \end{aligned} \quad (44)$$

where k^μ is the gluon momentum and

$$D(p_t) = \frac{1}{p_t^0 - m_t - \mathbf{p}_t^2/2m_t + i\Gamma_t/2} \quad (45)$$

denotes the nonrelativistic top-quark propagator. T is the contraction of hadronic and leptonic tensors resulting from the spinor traces after a soft-gluon factor (the second and third lines) is taken out. It coincides with the fully differen-

tial cross section of the Born-type diagram except for the t and \bar{t} propagators and Green's functions; hence T is real. Next we integrate over $dp_t^0/(2\pi)$, $dk^0/(2\pi)$ and $d\Omega_{\mathbf{k}}$. Let us define

$$\begin{aligned} I(\mathbf{k}, \hat{\mathbf{n}}_b, \hat{\mathbf{n}}_{\bar{b}}) &\equiv \int \frac{dp_t^0}{(2\pi)} \frac{dk^0}{(2\pi)} [D(p_t) + D(p_{\bar{t}})] [D^*(p_t+k) \\ &+ D^*(p_{\bar{t}}-k)] \frac{1}{k^0 - \hat{\mathbf{n}}_b \cdot \mathbf{k} - i\epsilon} \frac{1}{-k^0 + \hat{\mathbf{n}}_{\bar{b}} \cdot \mathbf{k} - i\epsilon} \\ &\simeq \frac{1}{(\hat{\mathbf{n}}_b - \hat{\mathbf{n}}_{\bar{b}}) \cdot \mathbf{k} + i\epsilon} \left(\frac{1}{\hat{\mathbf{n}}_b \cdot \mathbf{k} + i\Gamma_t} + \frac{1}{-\hat{\mathbf{n}}_{\bar{b}} \cdot \mathbf{k} + i\Gamma_t} \right). \end{aligned} \quad (46)$$

⁴Within our approximation, there is no distinction between $\cos \theta_{lt}$ and $-\cos \theta_{l\bar{t}}$, where $\theta_{l\bar{t}}$ denotes the angle between \bar{t} and l^+ in the t rest frame.

Noting that the contribution of diagram (c) to the cross section comes solely from the gluon momentum region

$$k^0, \quad |\mathbf{k}| \sim \alpha_s^2 m_t, \quad (47)$$

we keep $|\mathbf{k}|$ within this region. Then we may substitute $\tilde{G}^*(|\mathbf{p}_t + \mathbf{k}|, E) \rightarrow \tilde{G}^*(p, E)$ in Eq. (44) since the difference is higher order, and find

$$\begin{aligned} \frac{d\sigma_c}{d^3\mathbf{p}_t dx_t d\Omega_t} &= \int dp_W^2 d\phi_b d\Phi_2(\bar{b}W^-) \frac{1}{(2\pi)^6} \frac{1}{(4\pi)^5} \times T \\ &\times iC_F \cdot 4\pi\alpha_s |\tilde{G}(p, E)|^2 \int_{|\mathbf{k}| \sim \alpha_s^2 m_t} |\mathbf{k}|^2 d|\mathbf{k}| \\ &\times \frac{1}{|\mathbf{k}|^2} \int d\Omega_{\mathbf{k}} J(\mathbf{k}, \hat{\mathbf{n}}_b, \hat{\mathbf{n}}_{\bar{b}}) + \text{c.c.} \end{aligned} \quad (48)$$

It is easy to see that $\int d\Omega_{\mathbf{k}} J(\mathbf{k}, \hat{\mathbf{n}}_b, \hat{\mathbf{n}}_{\bar{b}})$ is real using the symmetry of $d\Omega_{\mathbf{k}}$ under $\mathbf{k} \rightarrow -\mathbf{k}$. Thus, we conclude $d\sigma_c/d^3\mathbf{p}_t dx_t d\Omega_t = 0$ in our approximation.

In fact the same proof can be applied to show quite generally that the contribution of diagram (c) vanishes at $\mathcal{O}(\alpha_s)$ *provided* one calculates a cross section where the top quark energy is integrated out; for example, the top quark three-momentum distribution. This is no longer the case when one considers a cross section that depends explicitly on the top quark energy; for example, the top quark four-momentum distribution. Then the diagram in question *does* contribute.

One can also show in a similar way that the kinematical regions Eq. (47) in diagrams (a) and (b) do not contribute to the cross section $d\sigma/d^3\mathbf{p}_t dx_t d\Omega_t$ and that only the gluon momentum region where $|\mathbf{k}| \sim \alpha_s m_t \gg \Gamma_t$ is relevant. We took advantage of this fact in deriving Eqs. (36)–(40).

It can be shown using similar techniques that the contribution of diagram (d) (exchange of one transverse gluon between b and \bar{b}) gets canceled when it is added to that of the corresponding real-gluon emission diagram [interference of diagrams (e) and (f) in Fig. 7]. This cancellation is consistent with the same cancellation that was found in the calculation




FIG. 7. Diagrams for real-gluon emission process $e^+e^- \rightarrow t\bar{t} \rightarrow bl^+v\bar{b}W^-g$.

of the top quark momentum distribution [10]. The contribution from each of these diagrams comes from the gluon momentum region $|\mathbf{k}| \lesssim \alpha_s^2 m_t$ and is in fact logarithmically divergent due to a soft-gluon singularity. The cancellations of the final-state interaction corrections at the various levels of inclusive cross sections are summarized in Sec. VII.

Now let us compare our formulas and the argument given in the previous section. In Secs. III B and III C, the parity-violating nature of the electroweak interactions in top production and decay played an essential role, while this was not the case in Secs. III A and III D. We find that correspondingly the $\cos \theta_{te}$ term of δ_a and $\delta\mathcal{P}_a$ contain electroweak coupling parameters (through C_{\parallel}^0), while the symmetric term of δ_a and ξ are independent of these electroweak parameters. More precisely, the $\cos \theta_{te}$ term of δ_a and $\delta\mathcal{P}_a$ have the forms anticipated in Secs. III B and III C, respectively, if the function $\psi_R(p, E)$ is positive. Indeed, the numerical evaluation in Ref. [19] shows that $\psi_R(p, E) \gtrsim 0$ holds in the entire threshold region. Besides, an additional coefficient $[1 - (C_{\parallel}^0)^2]$ in $\delta\mathcal{P}_a$ can be understood within our previous argument in the extreme cases $C_{\parallel}^0 = \pm 1$. Namely, if the top quark is 100% polarized, there will be no contamination from the opposite spin so that the correction should disappear. It may be interesting to note that the final-state interaction corrections to the polarization vector vanishes for the ideally polarized top quarks $C_{\parallel}^0 = \pm 1$.

C. Formula including full $\mathcal{O}(\alpha_s)$ corrections

In summary, the energy-angular distribution of l^+ including full $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections can be cast into the form

$$\frac{d\sigma(e^+e^- \rightarrow t\bar{t} \rightarrow bl^+v\bar{b}W^-)}{d^3\mathbf{p}_t dx_t d\Omega_t} = \frac{d\sigma(e^+e^- \rightarrow t\bar{t})}{d^3\mathbf{p}_t} \frac{1}{\Gamma_t} \frac{d\Gamma_{t \rightarrow bl^+v}(\mathcal{P})}{dx_t d\Omega_t} (1 + \xi) \quad (49)$$

with

$$\frac{d\sigma(e^+e^- \rightarrow t\bar{t})}{d^3\mathbf{p}_t} = \frac{d\sigma_{t\bar{t}}^0}{d^3\mathbf{p}_t} \left[1 + \frac{1}{2} \psi_1 + \left(2C_{FB}\varphi_R + \frac{\kappa}{2} C_{\parallel}^0 \psi_R \right) \cos \theta_{te} \right], \quad (50)$$

$$\mathcal{P} = \mathcal{P}_{\text{Born}} + \delta\mathcal{P}_a. \quad (51)$$

The above distribution is obtained as the sum of the cross sections for $e^+e^- \rightarrow t\bar{t} \rightarrow bl^+\nu\bar{b}W^-$ and $e^+e^- \rightarrow t\bar{t} \rightarrow bl^+\nu\bar{b}W^-g$. An independent emission of a gluon from the t or \bar{t} side has been included in Eq. (23), while the interference of both has been incorporated in conjunction with the final-state interaction diagram (d) in Sec. IV B.

Experimentally the top quark four-momentum p_t^μ will necessarily be reconstructed from the $\bar{b}W^-$ system in the study of the l^+ distribution. For the case with a gluon in the final state, we assign the ‘‘top quark momentum’’ as

$$\text{case (A): } p_t \equiv (p_{e^-} + p_{e^+}) - (p_{\bar{b}} + p_{W^-} + p_g) \\ \text{if } (p_{\bar{b}} + p_{W^-} + p_g)^2 - m_t^2 \lesssim m_t \Gamma_t,$$

$$\text{case (B): } p_t \equiv (p_{e^-} + p_{e^+}) - (p_{\bar{b}} + p_{W^-}) \\ \text{if } (p_{\bar{b}} + p_{W^-})^2 - m_t^2 \lesssim m_t \Gamma_t.$$

(See, however, the discussion in Sec. VII.) In other kinematical configurations the cross section is suppressed. Experimentally there will be a corresponding cut in the $\bar{b}W^-$ invariant mass. If both conditions in cases (A) and (B) are satisfied simultaneously, the gluon should necessarily be soft, and there will be no difference within our approximation between the cross sections corresponding to the above two assignments of the top quark momentum.

One comment is in order here. In defining the ‘‘rest frame’’ of a top-quark in the study of l^+ distribution, one may use either p_t^μ or \tilde{p}_t^μ [defined in Eq. (29)]. The difference of the cross sections based on the two definitions is of $\mathcal{O}(\alpha_s^2)$ which is beyond the scope of our approximations. Thus, the cross sections defined in both definitions should be measured in experiment and compared. It will serve as a cross check for the stability of our prediction.

V. NUMERICAL RESULTS

In this section we examine the effects of the final-state interaction corrections on the l^+ energy-angular distribution numerically, and compare the results with the qualitative argument given in Sec. III. The numerical results are obtained using both the coordinate-space approach developed in Refs. [2,3] and the momentum-space approach developed in Refs. [4,5]. Conventionally these two approaches have been used independently by different groups, and this is the first time to make a direct comparison of the cross sections calculated in both approaches. Some of the produced results are slightly different. We set $m_t = 175$ GeV, $\alpha_s(M_Z) = 0.118$, $P_{e^+} = 0$, and $\alpha = 1/128$ in all our analyses.

We first examine the contribution of diagram (a) in Figs. 6 (Coulomb interaction between t and \bar{b}) as given in Eqs. (36)–(38). Shown in Fig. 8 are the top quark momentum distribution $d\sigma/d|\mathbf{p}_t|$ for various c.m. energies measured from the lowest-lying resonance⁵ $\Delta E = \sqrt{s} - M_{1S}$. This is

calculated from the top quark production cross section Eq. (36). As expected, the top quark momentum is reduced. The effects are half in magnitude as compared to the final-state interaction corrections given in Refs. [10,19] since only the interaction between t and \bar{b} is included here.

We show the angular distribution of the top quarks in Figs. 9. It can be seen that the final-state interaction increases the top quark distribution in the forward direction for $P_{e^-} = 1$. This is consistent with our argument in Sec. III B since in leading-order approximation t and \bar{t} have their spins aligned perfectly in the $\hat{\mathbf{n}}_{\parallel}$ direction for this e^- polarization. Oppositely we see that the final-state interaction decreases the top quark distribution in the forward direction for $P_{e^-} = -1$. We note that the top quark has a natural polarization $\mathcal{P} \approx -0.4\hat{\mathbf{n}}_{\parallel}$ for unpolarized e^+e^- beams. Hence, the sign of the correction is the same as in the $P_{e^-} = -1$ case. Also we show corrections to the top quark polarization vector in Fig. 10. Although the qualitative behavior meets our expectation, the magnitude of the correction is rather small. Note that $\delta\mathcal{P}_a$ vanishes for $P_{e^-} = \pm 1$ since $C_{\parallel}^0 = \pm 1$.

Next we investigate the irreducible (nonfactorizable) correction that stems from the Coulomb interaction between \bar{t} and b . The correction factor $\xi(p, E, x_l, \cos\theta_l)$ given in Eqs. (40)–(43) depends on four parameters, two of which specify the lepton configuration: x_l and $\cos\theta_l$. Therefore, we will examine the dependence of ξ on these two parameters for several $(p, \Delta E)$ combinations. We fix the top quark momentum p to be the peak momentum of the distribution for each ΔE ; hence its values are slightly different for the two numerical approaches, see the top quark momentum distributions in Fig. 8. Shown in Fig. 11 are three-dimensional plots of ξ as a function of x_l and $\cos\theta_l$. One can see that in the figures ξ takes comparatively large positive values for either ‘‘small x_l and $\cos\theta_l \approx -1$ ’’ or ‘‘large x_l and $\cos\theta_l \approx +1$.’’ Oppositely, in the other two corners of the x_l – $\cos\theta_l$ plane ξ becomes small or becomes negative for smaller ΔE . The typical magnitude of ξ is 10–20 %, which would be a reasonable size for an $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ correction. This behavior holds also true for ξ at p off the peak of the top quark momentum distribution. These features of the correction factor ξ are consistent with our qualitative argument in Sec. III.

We made a cross check of our numerical results for ξ by numerically integrating $\xi \times \Gamma_t^{-1} d\Gamma_{t \rightarrow bl^+\nu} / dx_l d\Omega_l$ over the lepton energy-angular variables $\int dx_l d\Omega_l$ and comparing to the final-state interaction corrections to top quark momentum distribution given in Refs. [10,19]. It is seen that in Figs. 8 and 9 the coordinate-space approach and the momentum-space approach produce slightly different results. In particular the normalization of the cross sections differ at lower c.m. energies, $0 \leq \Delta E \leq 2$ GeV, whereas the differences decrease at higher energies. The cause of the differences can be traced back to the different short-distance QCD potentials employed in the two approaches. As we will discuss in Sec. VII, this difference is formally counted as higher order beyond our approximation, and at present it should be taken as an uncertainty of the theoretical prediction.

⁵ M_{1S} is defined as the real part of the position of the lowest-lying resonance pole in the complex energy plane. The energy measured from M_{1S} is more convenient than the energy measured from the

threshold ($E = \sqrt{s} - 2m_t$) when we compare different potentials in the literature.





FIG. 8. Top quark momentum distributions $d\sigma/d|p_t|$ obtained from Eq. (36) for various c.m. energies measured from the lowest-lying resonance, $\Delta E = \sqrt{s} - M_{1S}$. Solid lines and dashed lines represent the distributions with and without the final-state interaction between t and \bar{b} (δ_a), respectively. We set $P_{e^-} = 0$. The left figure corresponds to the coordinate-space calculation, the right one to the momentum-space calculation.


VI. OBSERVABLE PROPER TO TOP DECAY PROCESSES

As we have seen in the previous section, the final-state interactions affect the l^+ distribution in top quark decays.

The correction factor ξ depends on the kinematical variables of both the top quark and l^+ , and destroys the factorization of the cross section Eq. (23). In this section we define an observable which depends only on the top quark decay pro-




(a) Coordinate-space approach results




(b) Momentum-space approach results

FIG. 9. Top-quark angular distribution $d\sigma/d \cos \theta_{l^+}$ obtained from Eq. (36) for various electron polarizations P_{e^-} . Solid lines and dashed lines represent the distributions with and without the final-state interaction between t and \bar{b} (δ_a), respectively.



(a) Coordinate-space approach results



(b) Momentum-space approach results

FIG. 10. Final-state interaction correction to the $\hat{\mathbf{n}}_{\parallel}$ component of the top quark polarization vector $\delta\mathcal{P}_a$ for $P_{e^-} = +0.8$ (solid line), 0 (dash line), -0.8 (dot-dash line).

cess ($d\Gamma_{t \rightarrow b l^+ \nu} / dx_l d\Omega_l$ of a free polarized top quark).⁶ It is a differential quantity dependent on the l^+ energy-angular variables.

From Eq. (40) one sees that the correction factor ξ is invariant under the simultaneous transformations of the angular variables

$$\cos \theta_{Wl} \rightarrow -\cos \theta_{Wl}, \quad \cos \theta_{lt} \rightarrow -\cos \theta_{lt}, \quad (52)$$

since z_{\pm} is invariant. This invariance may be understood as follows. The soft-gluon factor, which represents the final-state interaction in diagram (b) (Fig. 6), does not depend on the l^+ momentum as long as the W^+ momentum is kept fixed. One sees that accordingly the integrand of Eq. (39) is independent of the l^+ energy and angle. The dependence on these variables enters only through the phase-space integra-

tion over ϕ_{bl} for a given l^+ configuration. Therefore, one may reverse the l^+ momentum in the t rest frame without affecting the phase-space integration, thereby keeping the whole function ξ also unchanged. The above transformations Eq. (52) are essentially this reversal of the l^+ momentum. (Due to the form of the integrand, there are extra degrees of freedom for the transformation of the l^+ direction, see below.)

Using Eq. (43), the above transformation can be written as a transformation of the lepton energy and angle as

$$x_l \rightarrow x'_l = \left(\frac{1+y}{y} - \frac{1}{x_l} \right)^{-1}, \quad \hat{\mathbf{n}}_l \rightarrow \hat{\mathbf{n}}'_l. \quad (53)$$

Here, $\hat{\mathbf{n}}_l = \mathbf{p}_l / |\mathbf{p}_l|$ denotes the unit vector in the direction of l^+ in the top quark rest frame. The choice of $\hat{\mathbf{n}}'_l$ for flipping the sign of $\cos \theta_{lt}$ ($\cos \theta_{lt} \rightarrow -\cos \theta_{lt}$) is not unique. We represent by $\hat{\mathbf{n}}'_l$ an arbitrary one of those choices. The produc-

⁶This property is true only up to $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections and may be violated by yet uncalculated $\mathcal{O}(\alpha_s^2)$ corrections.




FIG. 11. Three-dimensional plots of ξ as a function of x_l (x axis) and $\cos \theta_{te}$ (y axis) at the peak momenta of the $|\mathbf{p}_l|$ -distribution in Fig. 8: $\Delta E=0$ GeV and $\Delta E=5$ GeV.

tion cross section of the top quark is not affected by this transformation since the top quark kinematical variables are not involved. The important point is that neither the final-state interaction correction is affected by it.

Now, using this invariance, we first construct a quantity that has the simplest structure from the theoretical point of view, and afterwards we present an improved quantity that will be more useful for practical purposes. Let us define

$$A(x_l, \hat{\mathbf{n}}_l \cdot \mathcal{P}, \hat{\mathbf{n}}'_l \cdot \mathcal{P}) \equiv \left[\frac{d\sigma(e^+e^- \rightarrow t\bar{t} \rightarrow bl^+v\bar{b}W^-)}{d^3\mathbf{p}_l dx_l d\Omega_l} \right] \bigg/ \left[\frac{d\sigma(e^+e^- \rightarrow t\bar{t} \rightarrow bl^+v\bar{b}W^-)}{d^3\mathbf{p}_l dx_l d\Omega_l} \right]_{x_l \rightarrow x'_l, \hat{\mathbf{n}}_l \rightarrow \hat{\mathbf{n}}'_l}. \quad (54)$$

The production cross section and the correction factor ξ cancel in the numerator and denominator. As a result, this quantity is independent of the top quark momentum \mathbf{p}_l , and is determined only from the free polarized top quark decay cross section. In fact, using Eq. (49), we find that

$$A(x_l, \hat{\mathbf{n}}_l \cdot \mathcal{P}, \hat{\mathbf{n}}'_l \cdot \mathcal{P}) = \left[\frac{d\Gamma_{t \rightarrow bl^+v}(\mathcal{P})}{dx_l d\Omega_l} \right] \bigg/ \left[\frac{d\Gamma_{t \rightarrow bl^+v}(\mathcal{P})}{dx_l d\Omega_l} \right]_{x_l \rightarrow x'_l, \hat{\mathbf{n}}_l \rightarrow \hat{\mathbf{n}}'_l} \quad (55)$$

holds up to (and including) $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections. Note that the polarization vector \mathcal{P} , which specifies the decay distribution, includes the correction induced by the final-state interaction between t and \bar{b} ; see Eq. (51).

Since we take a ratio of differential cross sections in the definition of A in Eq. (54), this quantity would suffer from a large statistical error experimentally. Meanwhile, this quantity is predicted to be dependent only on a few variables theoretically.

[See Eq. (55).] This means that we may integrate out the irrelevant kinematical variables before taking the ratio and reduce its statistical uncertainty.⁷ For instance, we may choose $\hat{\mathbf{n}}_l = -\hat{\mathbf{n}}'_l$ and define

$$\bar{A}(x_l, a) \equiv \frac{\int d^3\mathbf{p}_l d\Omega_l \delta(\hat{\mathbf{n}}_l \cdot \mathcal{P} - a) \left[\frac{d\sigma(e^+ e^- \rightarrow t \bar{t} \rightarrow bl^+ \nu \bar{b} W^-)}{d^3\mathbf{p}_l dx_l d\Omega_l} \right]}{\int d^3\mathbf{p}_l d\Omega_l \delta(\hat{\mathbf{n}}_l \cdot \mathcal{P} + a) \left[\frac{d\sigma(e^+ e^- \rightarrow t \bar{t} \rightarrow bl^+ \nu \bar{b} W^-)}{d^3\mathbf{p}_l dx_l d\Omega_l} \right]_{x_l \rightarrow x'_l}}. \quad (56)$$

Here, the top quark polarization vector in the delta functions should be evaluated as a function of \mathbf{p}_l according to Eqs. (26)–(28), (38), and (51). The numerator and denominator, respectively, depend on two external kinematical variables and all other variables are integrated out.

Again using Eq. (49), one finds that \bar{A} is determined solely from the free polarized top decays

$$\bar{A}(x_l, a) = \left[\frac{d\Gamma_{t \rightarrow bl^+ \nu}(\mathcal{P})}{dx_l d\Omega_l} \right]_{\hat{\mathbf{n}}_l \cdot \mathcal{P} = a} \Big/ \left[\frac{d\Gamma_{t \rightarrow bl^+ \nu}(\mathcal{P})}{dx_l d\Omega_l} \right]_{x_l \rightarrow x'_l, \hat{\mathbf{n}}_l \cdot \mathcal{P} = -a}. \quad (57)$$

This is a general formula that is valid even if the decay vertices of the top quark deviate from the standard-model forms.⁸ We see that the quantity $\bar{A}(x_l, a)$ preserves most of the differential information⁹ contained in $d\Gamma_{t \rightarrow bl^+ \nu}/dx_l d\Omega_l$.

VII. DISCUSSION

In this section we discuss three different issues relevant to our work. These are the difference between the coordinate-space and momentum-space potentials, the misassignment of the top quark momentum, and the disappearance of the final-state interaction corrections at the various levels of inclusive cross sections.

As we saw in Sec. V, our numerical results obtained from the coordinate-space calculations and those obtained from the momentum-space calculations differ slightly, although all the qualitative features are common. The difference can be traced back to the difference in the short-distance part of the QCD potentials used in the two approaches.

Let us remind the reader how each potential is constructed

⁷Consider a ratio of certain physical quantities which depend on a set of kinematical variables

$$R = \frac{X(\phi_i)}{Y(\phi'_i)},$$

where ϕ_i denotes a point in the phase space, and ϕ'_i is obtained by a transformation from it, $\phi'_i = \phi'(\phi_i)$. Whenever the ratio R takes the same value in a subspace U of the whole phase space, we may take sum over the subspace before taking the ratio

$$R = \frac{\sum_{i \in U} X(\phi_i)}{\sum_{i \in U} Y(\phi'_i)}.$$

⁸Note that, quite generally, energy-angular distributions of l^+ from free polarized top quarks have the form

$$\frac{d\Gamma_{t \rightarrow bl^+ \nu}(\mathcal{P})}{dx_l d\Omega_l} = F_0(x_l) + (\hat{\mathbf{n}}_l \cdot \mathcal{P}) F_1(x_l).$$

⁹According to its construction, it satisfies the relation

$$\bar{A}(x_l, a) \cdot \bar{A}(x'_l, -a) = 1.$$

(in the short-distance regime). The large-momentum part of the momentum-space potential V_{JKT} [4,5] is determined as follows. First the potential has been calculated up to the next-to-leading order in a fixed-order calculation. The potential is then improved using the two-loop renormalization group equation in momentum space. On the other hand, the short-distance part of the coordinate-space potential V_{SFHMN} [3] is calculated by taking the Fourier transform of the fixed-order potential in momentum space, and then the potential is improved using the two-loop renormalization group equation in coordinate space. Thus, the two potentials are *not* the Fourier transforms of each other. Only the leading and next-to-leading logarithmic terms of the series expansion in a fixed modified minimal subtraction scheme ($\overline{\text{MS}}$)-coupling are the same for the two potentials. The difference begins at the next-to-next-to-leading order terms. (The nonlogarithmic term in the two-loop fixed-order correction.)

To make a clear comparison, the two potentials are Fourier transformed numerically and we examine their difference both in coordinate space and in momentum space. We show the effective charges defined as

$$\alpha_{\text{cor}}(1/r) = (-C_F/r)^{-1} V(r), \quad (58)$$

$$\alpha_{\text{mom}}(q) = (-4\pi C_F/q^2)^{-1} \bar{V}(q) \quad (59)$$

in Fig. 12, which clearly demonstrates that there is a non-negligible difference between the potentials. The oscillatory behavior of V_{SFHMN} in Fig. 12(b) is an artifact due to the discontinuity in the second derivative of the coordinate-space potential, which is located at the continuation point of the perturbative potential to a long-distance potential. As already stated, the coordinate-space potential V_{SFHMN} follows the form required by the two-loop renormalization group equation in the short-distance region, whereas the momentum-




FIG. 12. The effective charges (a) $\alpha_{\text{coor}}(1/r)$ and (b) $\alpha_{\text{mom}}(p)$ defined via the potentials in coordinate space and momentum space, respectively. V_{JKT} [4,5] and V_{SFHMN} [3] represent the potentials used in the momentum-space approach and the coordinate-space approach, respectively.

space potential V_{JKT} follows the form required by the two-loop renormalization group equation in the large momentum region.

In principle we may reduce the difference by including the two-loop finite correction and invoking the three-loop renormalization group improvement [29]. We shall not do so in this paper, because there are a number of other corrections of the same order of magnitude which are not calculated yet. We will study the difference of the two approaches in more detail in a forthcoming paper.

In Sec. IV C we assumed a perfect assignment of the top quark momentum in cases (A) and (B) for defining the distribution formula, Eq. (49). In real experiments, however, a misassignment of the top quark momentum will be inevitable whenever there is real gluon radiation in the final state because of the typical jet-clustering algorithm that will be used. For instance, when a gluon is indistinguishable from a b jet, the top quark momentum reconstructed by clustering may be off shell; rather grouping the gluon on the other side (with \bar{b}) would result in an on-shell momentum.¹⁰ Reference [10] studied how this misassignment alters the top quark three-momentum distribution near threshold and found that the correction is less than a few percent; the clustering algorithm assumed in that paper, however, is somewhat unrealistic. The effect of the misassignment was also studied in Ref. [30] in the open-top-quark region ($\sqrt{s} \gg 2m_t$) using a Monte Carlo generator and with more realistic experimental assumptions. It was shown that the effects on the top quark invariant-mass distribution and on the angular distributions are substantial at $\sqrt{s} = 400\text{GeV}$ (for $m_t = 175\text{GeV}$) and increase in magnitude and in complexity as the c.m. energy is raised. Clearly, in our case, we need more detailed studies to see how a similar effect may influence our results. For this purpose, studies based on a Monte Carlo generator that produces the fully differential distribution including the full $\mathcal{O}(\alpha_s)$ corrections near threshold would be necessary. (See also Ref. [12] for

¹⁰There is no ambiguity in assigning the gluon to the production of $t\bar{t}$ since real gluon radiation in the top quark production process is suppressed near threshold.

analyses on the radiative color flows from t , \bar{t} , b , and \bar{b} close to the $t\bar{t}$ threshold.)

In the course of calculating the final-state interaction effects on the lepton energy-angular distribution, we found that the final-state interaction between b and \bar{b} vanishes [diagrams (c) and (d)] if we add the corresponding real-emission diagram as well as if we integrate over the top quark energy. This fact was already conjectured in Ref. [19] on account of an estimate of the Coulomb energy between b and \bar{b} . Nevertheless the same final-state interaction modifies the top quark energy distribution. As a similar phenomenon one may be reminded of the cancellation of final-state interactions (including also those between t and \bar{b}) in the total $t\bar{t}$ production cross section, despite the modification of the top quark three-momentum distribution. In fact the cancellation of final-state interactions is a general feature known in a wide class of inclusive hard scattering cross sections both in QED and QCD.¹¹

It may be worth summarizing here at which level of inclusiveness the effects of the final-state interactions cancel in the various cross sections in our particular process, top quark pair production near threshold, and its subsequent decay. This is shown in Table I. Note that there are three typical mass scales involved in this process: the top quark mass m_t , the inverse Bohr radius $\alpha_s m_t$, and the Coulomb energy between the $t\bar{t}$ pair $\alpha_s^2 m_t$. We show in the table the typical momentum scale of the gluon in each diagram.

VIII. CONCLUSION

In this paper we considered the differential distribution of l^+ from the semileptonic decay of the top quark, where the parent top quark is produced in $e^+e^- \rightarrow t\bar{t}$ near threshold.

¹¹For example, it is found in Refs. [21–24] that the final-state interaction correction to the invariant-mass distribution of W changes sign above and below the distribution peak and that the correction vanishes upon integration over the invariant mass. For an enlightening discussion on related problems, see also Ref. [31].

TABLE I. Vanishment or nonvanishment of final-state interaction diagrams in the various inclusive cross sections. The plus sign shows there is a noncancelled contribution from the diagram to the corresponding cross section, while “vanish” shows the cancellation of contributions from the diagram to the corresponding inclusive cross section. For details, see Sec. IV B.

| Inclusiveness | Diagram (a)/(b) | Diagram (c) | Diagram (d)+(e)*(f) |
|--------------------------|-----------------|------------------|---------------------|
| fully differential | + | + | + |
| top energy integrated | + | vanish | vanish |
| $\sigma_{\text{tot}}(s)$ | vanish | vanish | vanish |
| typical gluon momentum | $\alpha_s m_t$ | $\alpha_s^2 m_t$ | $\alpha_s^2 m_t$ |

Particularly, we have calculated the final-state interaction corrections (rescattering corrections) to the energy-angular distribution of leptons in the top quark decay. Also we have explicitly written down the l^+ energy-angular distribution $d\sigma(e^+e^- \rightarrow t\bar{t} \rightarrow b l^+ \nu \bar{b} W^-)/d^3\mathbf{p}_l dx_l d\Omega_l$ including the full $\mathcal{O}(\alpha_s) = \mathcal{O}(\beta)$ corrections near threshold.

We presented numerical studies of the various effects of the final-state interaction corrections. All numerical results can be understood qualitatively from intuitive pictures. Attractive forces between t and \bar{b} and between \bar{t} and b modify not only the momentum distribution of bW^+ or $\bar{b}W^-$ system but also the top quark polarization vector and the lepton energy-angular distribution.

(i) The effect of Coulomb-gluon exchange between \bar{b} and t , when integrated over the $\bar{b}W^-$ phase space, can be regarded as a correction to the top quark production process. The effect can be incorporated by modifying the top quark production cross section and the top quark polarization vector. The top quark momentum distribution is shifted to take a smaller average momentum due to the attraction by \bar{b} . Also since \bar{b} is emitted preferably in the t spin direction, the attraction generates a $\cos\theta_{te}$ distribution of the top quark as well as modifies the top quark polarization vector.

(ii) The Coulomb interaction between b and \bar{t} causes a nonfactorizable correction with respect to the production and decay processes of the top quark. It generates an energy-angle-correlated correction to the lepton distribution. Namely, the l^+ distribution is deformed in favor of the kinematical configurations “small E_l and emitted in \bar{t} direction” or “large E_l and emitted in t direction,” which can be understood as originating from the attraction of b in the direction of \bar{t} .

(iii) Corrections from the gluon exchange between b and

\bar{b} turn out to vanish when the top quark energy is integrated out.

Without the nonfactorizable effect ξ , the l^+ angular distribution is dependent only on the polar angle from the polarization vector of the parent top quark in its rest frame. The final-state interaction brings in another direction into the problem, the direction of \bar{t} , which is a completely new feature in comparison to the decays of free polarized top quarks.

In order to study the decay properties of top quarks near the $t\bar{t}$ threshold, it is desirable to extract the part which is specific to the top quark decay process alone. In the case of semileptonic decay, we defined a quantity which depends only on the decay distribution of a free polarized top quark. The part which depends on $\cos\theta_{te}$ and $\cos\theta_{lt}$ is dropped using the transformation of the l^+ energy and angle which leaves the final-state interaction unchanged. Thus, we recover a differential quantity $\bar{A}(x_l, \hat{\mathbf{n}}_l \cdot \mathcal{P})$ dependent only on the lepton energy and the lepton angle from the parent top quark polarization vector.

This quantity will be useful from the theoretical point of view. It can be calculated from the decay distribution of free top quarks without including the bound-state effects or the final-state interaction corrections that are typical to the threshold region. Therefore, a variety of former studies on free top quark decays may also be applicable in the $t\bar{t}$ threshold region, where highly polarized top quarks are available with the largest cross sections.

ACKNOWLEDGMENTS

The authors wish to thank M. Jezabek and J.H. Kühn for enlightening discussion. This work is supported in part by Graduiertenkolleg “Elementarteilchenphysik an Beschleunigern,” by the “Landesgraduiertenförderung” at the University of Karlsruhe, by BMBF under Contract No. 057KA92P, and by the Alexander von Humboldt Foundation.

[1] V. S. Fadin and V. A. Khoze, JETP Lett. **46**, 525 (1987); Sov. J. Nucl. Phys. **48**, 309 (1988).
 [2] M. Strassler and M. Peskin, Phys. Rev. D **43**, 1500 (1991).
 [3] Y. Sumino, K. Fujii, K. Hagiwara, H. Murayama, and C.-K. Ng, Phys. Rev. D **47**, 56 (1993).
 [4] M. Jezabek, J. H. Kühn, and T. Teubner, Z. Phys. C **56**, 653 (1992).
 [5] M. Jezabek and T. Teubner, Z. Phys. C **59**, 669 (1993).
 [6] H. Murayama and Y. Sumino, Phys. Rev. D **47**, 82 (1993).

[7] P. Igo-Kemenes, M. Martinez, R. Miquel, and S. Orteu, in *Proceedings of the Workshop on Physics and Experiments with Linear e^+e^- Colliders* Waikoloa, Hawaii, 1993, edited by F. Harris *et al.* (World Scientific, Singapore, 1993).
 [8] J. H. Kühn, in *Proceedings of the Workshop on Physics and Experiments with Linear e^+e^- Colliders* [7], p. 72.
 [9] K. Melnikov and O. Yakovlev, Phys. Lett. B **324**, 217 (1994).
 [10] Y. Sumino, Ph.D. thesis, University of Tokyo, 1993.
 [11] V. Fadin, V. Khoze, and A. Martin, Phys. Rev. D **49**, 2247

- (1994); Phys. Lett. B **320**, 141 (1994).
- [12] V. Khoze and W. Sjöstrand, Phys. Lett. B **328**, 466 (1994).
- [13] K. Fujii, T. Matsui, and Y. Sumino, Phys. Rev. D **50**, 4341 (1994).
- [14] W. Mödritsch and W. Kummer, Nucl. Phys. **B430**, 3 (1994); W. Kummer and W. Mödritsch, Z. Phys. C **66**, 225 (1995).
- [15] Y. Sumino, Acta Phys. Pol. B **25**, 1837 (1994).
- [16] M. Jeżabek, in *Physics at LEP200 and Beyond*, edited by T. Riemann and J. Blümlein [Nucl. Phys. (Proc. Suppl.) **37**, 197 (1994)].
- [17] R. Harlander, M. Jeżabek, J. H. Kühn and T. Teubner, Phys. Lett. B **346**, 137 (1995).
- [18] W. Mödritsch, Nucl. Phys. **B475**, 507 (1996).
- [19] R. Harlander, M. Jeżabek, J. Kühn, and M. Peter, Z. Phys. C **73**, 477 (1997).
- [20] M. Jeżabek, Acta Phys. Pol. B **26**, 789 (1995).
- [21] K. Melnikov and O. Yakovlev, Nucl. Phys. **B471**, 90 (1996).
- [22] V. Khoze and W. Stirling, Phys. Lett. B **356**, 373 (1995).
- [23] W. Beenakker, A. Chapovsky, and F. Berends, hep-ph/9706339; hep-ph/9707326.
- [24] V. Khoze and W. Sjöstrand, Z. Phys. C **70**, 625 (1996).
- [25] J. H. Kühn, Acta Phys. Pol. B **12**, 347 (1981).
- [26] M. Jeżabek and J. H. Kühn, Nucl. Phys. **B314**, 1 (1989).
- [27] A. Czarnecki, M. Jeżabek and J. H. Kühn, Nucl. Phys. **B351**, 70 (1991); A. Czarnecki and M. Jeżabek, *ibid.* **B427**, 3 (1994).
- [28] M. Jeżabek and J. H. Kühn, Phys. Rev. D **48**, R1910 (1993); **49**, 4970(E) (1994).
- [29] M. Peter, Phys. Rev. Lett. **78**, 602 (1997); Nucl. Phys. **B501**, 471 (1997).
- [30] C. Schmidt, Phys. Rev. D **54**, 3250 (1996).
- [31] V. Fadin, V. Khoze, A. Martin, and A. Chapovskii, Phys. Rev. D **52**, 1377 (1995).