

Radiation Protection and Management

著者	Miyata T., Yamadera A., Nakamura T., Izumi
	Y., Satake Y.
journal or	CYRIC annual report
publication title	
volume	1991
page range	300-302
year	1991
URL	http://hdl.handle.net/10097/49679

V. 4. Radiation Protection and Management

Miyata T., Yamadera A., Nakamura T., Izumi Y.* and Satake Y.*

Cyclotron and Radioisotope Center, Tohoku University Japan Radiation Protection Co., Ltd.*

(1) Unsealed radionuclides used in the center

The kinds and activities of unsealed radionuclides handled in the center in 1991 are shown in Table 1. The table includes the isotopes produced by the cyclotron, purchased from the Japan Isotope Association and took over from another RI institutes.

(2) Individual monitoring

The exposure doses of the workers in the center in 1991 is given in Table 2. They were less than the permissible doses.

(3) Monitoring of the workplace

Radiation dose rates inside and outside of the controlled areas were monitored periodically and as needed. They were below the legal permissible levels. Surface contamination levels of the floors inside the controlled areas were measured by smear method and with survey meters periodically and as needed. They also cleared the legal regulation levels.

(4) Wastes management

The radioactive wastes delivered to the Japan Radioisotope Association in 1991 are shown in Table 3. The concentration of radioisotopes in the air released after filtration from the stack was monitored with stack gas monitors. The levels were less than the legal regulation levels. The radioactive water was storaged at the tanks at least for 3 days and then released to the sewerage after confirming that the concentration was less than permissible levels.

The treated volume of radioactive waste of organic scintillator was 633 l by the incinerator made by Fujikogyo Co., Ltd.

Table 1. Unsealed radionuclides used in the center in 1991.(kBq)

1. ⁹⁰ Sr	4,590,900	3. ¹¹ C	244,563,358,000	4. ³ H	336,776,060
Total	4,590,900	^{13}N	10,582,000,000	¹⁴ C	1,609,406,500
		15 _O	50,140,000,000	18 _F	199,189,654,000
2. ⁴⁵ Ca	76,533,300	²⁸ Mg	3,082,100	⁵¹ Cr	378,533,730
⁵⁴ Mn	10,000	32 _P	886,225,000 _	²⁰¹ Tl	3,700,000
⁵⁷ Co	240,263	35 _S	308,400,000	Total	201,518,070,290
⁶⁸ Ge	408,432,000	⁴⁵ Ti	1,146,051,300		
85 _{Sr}	175,564,000	^{48}V	1,108,617,400		
125 _I	824,303,880	57 _{Ni}	370,901		
¹³⁷ Cs	37,700	⁶⁴ Cu	12,307		
¹⁶⁰ Tb	24,326,300	$80_{\mathbf{Br}}$	100,000		
170_{Tm}	26,199,000	⁹⁹ Mo	407,100,000		
²⁰⁷ Bi	303,540	99mTc	340,400,000		
Total	1,535,949,983	111 _{In}	107,126,000		
		126 _I	185,239		
		131 _I	925,896,120		
		140 _{La}	213,663,000		
		¹⁵³ S m	160,167,800		
		175 _{Yb}	1,932,500		
	_	177 _{Lu}	57,396,339		
	· · · · · · · · · · · · · · · · · · ·	Total	310,952,084,006		

Table 2. Occupational radiation exposures at the center in 1991.

Dose range (mSv)	Number of individuals
No measurable exposure	33
Measurable exposure less than 1.0	4
1.0 to 2.5	2
2.5 to 5.0	3
Total persons monitored	42

Table 3. Radioactive wastes delivered to the Japan Radioisotope Association in 1991.

Wastes	Container	Number
solids		
combustibles	200 l drum	14
incombustibles	50 <i>l</i> drum	34
dried animals	50 <i>l</i> drum	6
liquids		
inorganic substan	ces 25 ! PE bottle	6
filters		
HEPA filters	$610 \times 610 \times 292_{\text{mm}}$	14.5