

Marine Algae from Hirota Bay on the Pacific Coast of Northeastern Honshu, Japan (Applied Aquatic Bio-Sciences Marine Plant Ecology)

著者	LI Jing-Yu, NARITA Michiko, ENDO Hikaru, AGATSUMA Yukio, TANIGUCHI Kazuya
journal or publication title	Tohoku journal of agricultural research
volume	58
number	3/4
page range	89-97
year	2008-03
URL	http://hdl.handle.net/10097/40434

Marine Algae from Hirota Bay on the Pacific Coast of Northeastern Honshu, Japan

Jing-Yu LI¹, Michiko NARITA¹, Hikaru ENDO¹, Yukio AGATSUMA¹ and
Kazuya TANIGUCHI¹

¹Laboratory of Marine Plant Ecology, Graduate School of Agricultural Science,
Tohoku University, Sendai 981-8555, Japan

(Received, December 25, 2007)

Summary

Monthly collection of marine algae from the intertidal to subtidal zones in Hirota Bay (38°56' N, 141°42' E) on the Pacific coast of northeastern Honshu, Japan, was carried out from August 2005 to July 2006. A total of 95 species were listed, 12 of which belong to Chlorophyta, 24 to Phaeophyta, 58 to Rhodophyta, and one to sea grass. In these algae, the 10 species, *Urospora penicilliformis*, *Pachydictyon coriaceum*, *Chordaria flagelliformis*, *Myagropsis myagroides*, *Bangia gloiopeltidicola*, *Lithophyllum okamurae*, *Titanoderma tumidulum*, *Chondracanthus tenellus*, *Polysiphonia abscissa* and *P. sphaerocarpa*, were newly recorded to the marine algae from the coasts adjacent to Hirota Bay. The floral index of I/H value in Hirota Bay was 1.1, suggesting that the marine algal flora in Hirota Bay belongs to the typical subarctic zone.

Key words: marine algae, flora, maturation, life form

The marine algae from the Pacific coast of northeastern Honshu, Japan, have been extensively investigated in Shimokita Peninsula (Fuji and Yamamoto, 1972), the coast of Iwate Prefecture (Kawashima, 1954, 1955), Miyako (Taniguchi *et al.*, 1979), Okirai (Takamatsu, 1974), Iwai (Sato *et al.*, 2005), Kitakami (Nakata *et al.*, 2001), Ogatsu (Taniguchi *et al.*, 1985), Oshika Peninsula (Ogawa and Machida, 1976, 1977), Tomarihama (Agatsuma *et al.*, 2000), Sasunohama (Endo *et al.*, 2005), Matsushima Bay (Takamatsu, 1936; Narita *et al.*, 2006), and Iwaki (Suda, 1987; Sato *et al.*, 2001). For decades, marine flora is having been changed with the variation of hydrographic conditions, which is in turn affected by global warming (Taniguchi *et al.*, 2006). Narita *et al.* (2006) reported that the flora index of I/H value in Matsushima Bay increased from 1.5 in 1936 to 1.7 in 2006, and the value in Iwaki also increased from 1.4 in 1987 to 1.9 in 2001. Detail records of current status of marine algae off a lot of coasts lead to a better

understanding of changing process of marine flora. In the present study, we investigated the marine algae from Hirota Bay ($38^{\circ}56' \text{ N}$, $141^{\circ}42' \text{ E}$), an outstanding fishing ground under mixture of the Oyashio and Kuroshio Currents. The growth and maturation periods of these algae were also observed.

Materials and Methods

Marine algae and sea grasses were collected monthly by skin diving from the intertidal to subtidal zones in Hirota Bay, on the Pacific coast of northeastern Honshu, Japan (Fig. 1), from August 2005 to July 2006. The plants collected were preserved in 5% formaldehyde seawater and kept in dark conditions.

Each alga and sea grass was identified, and the presence of reproductive cells was recorded based on taxonomic literatures (Okamura, 1907–1933; Okamura, 1936; Segawa, 1956; Chihara, 1983; Yoshida, 1998). The algae identified were classified into five life-form groups: crustaceous algae, small annual algae, small perennial algae, large annual algae and large perennial algae, according to Taniguchi (1996). Finally, the marine algae and sea grass were preserved as dry herbarium specimens. The floral index of I/H value, that is the ratio of the number of species with isomorphic alternation of generations and lacking gametophytic generation, generally found in warm current areas (I) to heteromorphic alternation of generations, generally found in cold current areas (H) in both Chlorophyta and Phaeophyta (Nakahara and Masuda 1971), was also evaluated




FIG. 1. The locations of study site in Hirota Bay and Ofunato in Iwate Prefecture, northeastern Honshu, Japan.

for determining the biogeographic region.

The surface water temperature in Hirota Bay during the study period was monitored with a normal thermometer every month. The data of monthly average water temperature at Ofunato (38°59'N, 141°43'E) (Fig. 1), a port near the study site, from 1980 to 2006 measured by Iwate Farming Fisheries Association were referred.

Results and Discussion

During this study, the surface water temperature in Hirota Bay ranged from a maximum of 23°C in August to a minimum of 5°C in March (Fig. 2). The water temperature was 0.3–1.3°C lower from February to May and 0.6–2.1°C higher from August to November than those of the monthly average water temperature for 27 years at Ofunato.

In this study, one sea grass and 94 species of marine algae were identified, 12 of which belong to Chlorophyta, 24 to Phaeophyta and 58 to Rhodophyta. These algae are shown with their growth and maturation periods and life forms in Fig. 3. In these algae, the 10 species, *Urospora penicilliformis*, *Pachydictyon coriaceum*, *Chordaria flagelliformis*, *Myagropsis myagroides*, *Bangia gloiopeltidicola*, *Lithophyllum okamurae*, *Titanoderma tumidulum*, *Chondracanthus tenellus*, *Polysiphonia abscissa* and *P. sphaerocarpa*, were newly recorded to the marine algae from the coasts adjacent to Hirota Bay (Kawashima, 1954, 1955; Takamatsu, 1974; Taniguchi *et al.*, 1979). In addition, six species of fucoids, *Cystoseira hakodatensis*, *Myagropsis myagroides*, *Sargassum hemiphyllum*, *S. horneri*, *S. macrocarpum* and *S. yezoense*, were recognized as drifting rockweeds.

The algae were classified into the five life forms as follows: 48 small annuals, 31 small perennials, six large annuals, six large perennials and three crustaceans




FIG. 2. Surface water temperature in Hirota Bay from August 2005 to July 2006 and monthly average water temperature for 27 years (1980–2006) at Ofunato.




FIG. 3.

	2005						2006					
	Agu.	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.
32 ■ <i>Myagropsis myagroides</i>												
33 ■ <i>Sargassum hemiphyllum</i>											♂	
34 □ <i>Sargassum horneri</i>											♀	
35 ■ <i>Sargassum macrocarpum</i>											♂	
36 ■ <i>Sargassum yezoense</i>												
RHODOPHYTA												
37 △ <i>Bangia gloiopeltidicola</i>												
38 △ <i>Porphyra yezoensis</i>						♀♂	♀♂	♀♂	♀♂	♀♂	♀♂	
39 △ <i>Palmaria palmata</i>						⊕			⊕	⊕		⊕
40 △ <i>Nemalion vermiculare</i>	⊕											
41 ▲ <i>Calliarthron yessoense</i>			♀									
42 ▲ <i>Corallina pilulifera</i>												
43 ● <i>Lithophyllum okamurae</i>												
44 ● <i>Pneophyllum zostericola</i>	⊕	⊕	⊕	⊕	⊕	⊕	⊕				⊕	⊕
45 ● <i>Titanoderma tumidulum</i>												
46 ▲ <i>Gelidium divaricatum</i>												
47 ▲ <i>Gelidium elegans</i>	♀⊕	♀⊕	⊕	♀⊕		♀						♀⊕
48 ▲ <i>Gelidium vagum</i>	♀	♀⊕										⊕
49 ▲ <i>Pterocladia capillacea</i>		⊕	♀⊕		♀							
50 ▲ <i>Caulacanthus ustulatus</i>	⊕											
51 △ <i>Neodilsea yendoana</i>	⊕	⊕	⊕	♀⊕	⊕	⊕	♀⊕					⊕
52 ▲ <i>Gloiopeltis furcata</i>											♀⊕	♀⊕
53 ▲ <i>Chondracanthus intermedius</i>		♀										
54 ▲ <i>Chondracanthus tenellus</i>						♀						
55 ▲ <i>Chondrus giganteus</i>	⊕	⊕	⊕									
56 ▲ <i>Chondrus ocellatus</i>	♀⊕	♀⊕	⊕	⊕		⊕						
57 ▲ <i>Chondrus verrucosus</i>		♀⊕	♀⊕	♀⊕	⊕	⊕	⊕	⊕				⊕
58 ▲ <i>Chondrus yendoi</i>									♀⊕		♀⊕	♀
59 ▲ <i>Mazzaella japonica</i>	⊕			⊕	♀⊕	♀⊕	♀⊕	♀⊕	♀⊕	♀⊕		♀⊕
60 ▲ <i>Carpopeltis affinis</i>	♀	♀	⊕	♀	♀	♀			♀			
61 ▲ <i>Carpopeltis prolifera</i>	⊕	♀⊕					♀					
62 ▲ <i>Grateloupia elliptica</i>												
63 ▲ <i>Grateloupia filicina</i>												

FIG. 3. (continued)




FIG. 3. Growth and maturation periods, and life forms of marine algae and sea grass collected in Hirota Bay. ■; growing period, ○; unilocular sporangia, ⊙; plurilocular sporangia, P; propagule, M; mature thallus, ♂/♀; male/female reproductive organ, ⊕; tetrasporangia, △; small annual algae, ▲; small perennial algae, □; large annual algae, ■; large perennial algae, ●; crustaceous algae.

ones. Most of small annuals such as *Enteromorpha compressa*, *E. intestinalis*, *E. linza*, *Colpomenia sinuosa*, *Neodilsea yendoana*, *Schizymenia dubyi*, *Champia parvula*, *Ceramium japonicum* and *Polysiphonia japonica* matured from spring to summer although the mature species appeared in every month. For small perennials, the maturation periods were mainly divided into three seasons, spring such as *Analphus japonicus*, autumn such as *Dictyota dichotoma*, *Dilophus okamurai* and *Pachydictyon coriaceum*, and year-round such as *Chondrus verrucosus*, *Mazzaella japonica* and *Ahnfeltiopsis paradoxa*. Among large annuals, only *Undaria pinnatifida* matured from spring to summer. Among large perennials, the three fucoids *Cystoseira hakodatensis*, *Sargassum hemiphyllum* and *S. macrocarpum* matured in summer, while the kelp *Eisenia bicyclis* did in autumn. Among crustaceous algae, the reproductive cells were present only in *Pneophyllum zoestericola* from summer to winter.

Along the Pacific coast of northeastern Honshu from Oshika Peninsula, where are biogeographically located in subarctic floral zone (Okamura, 1931), I/H values were 1.4 in Shimokita Peninsula (Fuji and Yamamoto, 1972), 1.4 in Miyako (Taniguchi *et al.*, 1979), 1.4 in Iwai (Sato *et al.*, 2005), 1.4 in Kitakami (Nakata *et al.*, 2001), 1.9 in Ogatsu (Taniguchi *et al.*, 1985), 1.3 in Tomarihama (Agatsuma *et al.*, 2000) and 1.4 in Sasunohama (Endo *et al.*, 2005). The value in Hirota Bay was 1.1, suggesting that the marine algal flora in Hirota Bay belongs to the typical subarctic zone although it is slightly lower than those on other coasts.

Acknowledgements

We are grateful to Mr. U. Shimizu, Counselor of Hirota Fisheries Cooperative Association for his cooperation in sampling of algae. We also thank Mr. U. Nagahora, director-general of Iwate Farming Fisheries Association for providing the data of surface water temperature measured from 1980 to 2006. This study was funded partly by the project entitled "Study on construction of recycling-oriented economic system modeled on Iwate Prefecture" from Japan Science and Technology Agency.

References

- Agatsuma, Y., Ogawa, M., Taniguchi, K., and Yamada, H., Marine algal flora off the coast of Tomari-Hama along the Oshika Peninsula, Japan. *Wildlife Conservation Japan*, **5**, 47-53 (2000) (in Japanese, with English summary)
- Chihara, M., "The seaweeds of Japan" ed. by M. Chihara, Gakken, Tokyo, pp. 1-292 (1983) (in Japanese)
- Endo, H., Agatsuma, Y., and Taniguchi, K., Marine algae from Sasunohama on the southern coast of Oshika Peninsula, Japan. *Biosphere Conserva-*

- tion, 7, 29-38 (2005)
- Fuji, A. and Yamamoto, H., The benthic organism off the western coast of Shimokita Peninsula. *Shimokita Peninsula National Park Shimokita Umi Kouen Chosa Hokoku-syo*, pp. 55-88 (1972) (in Japanese)
- Kawashima, S., A list of the marine algae from the coast of Iwate Prefecture. I. Chlorophyceae and Phaeophyceae. *Bull. Jap. Soc. Phycol.*, 2, 5-10 (1954) (in Japanese)
- Kawashima, S., A list of the marine algae from the coast of Iwate Prefecture. II. Rhodophyceae. *Bull. Jap. Soc. Phycol.*, 3, 1-7 (1955) (in Japanese)
- Nakahara, H. and Masuda, M., Type of life cycle and geographical distribution of marine green and brown algae in Japan. *Marine Science Monthly*, 3, 24-26 (1971) (in Japanese)
- Nakata, Y., Agatsuma, Y., and Taniguchi, K., Marine algal flora off the coast of Jusan-Hama at Kitakami in Miyagi Prefecture, Japan. *Wildlife Conservation Japan*, 6, 33-39 (2001) (in Japanese, with English summary)
- Narita, M., Endo, H., Agatsuma, Y., and Taniguchi, K., Marine algae around the coasts of Oshika Peninsula, northeastern Honshu, Japan. *Kaiyo monthly*, 38, 216-220 (2006) (in Japanese)
- Ogawa, H. and Machida, M., Marine algae of the Oshika Peninsula. I. Chlorophyceae and Phaeophyceae. *Tohoku Journal of Agricultural Research*, 27, 145-154 (1976)
- Ogawa, H. and Machida, M., Marine algae of the Oshika Peninsula. II. Rhodophyceae. *Tohoku Journal of Agricultural Research*, 28, 151-165 (1977)
- Okamura, K., "Icones of Japanese algae" ed. by K. Okamura, Kazamashobo, Tokyo (1907-1933)
- Okamura, K., "Kaisan Syokubutsu no Chiri-teki Bunpu" ed. by K. Okamura, Iwanami Koza Chirigaku, Tokyo, pp. 1-86 (1931) (in Japanese)
- Okamura, K., "Nippon Kaiso Shi" ed. by K. Okamura, Uchida Roukakuho, Tokyo, pp. 1-964 (1936) (in Japanese)
- Sato, T., Nakata, Y., Agatsuma, Y., and Taniguchi, K., Marine algal flora off the coast of Shimo-Kajiro in Iwaki City, Japan. *Wildlife Conservation Japan*, 6, 41-46 (2001) (in Japanese, with English summary)
- Sato, Y., Tsumoto, T., Endo, H., Agatsuma, Y., Sasaki, R., Oshino, A., and Taniguchi, K., Marine algae from the cape of Iwai on the Pacific coast of northeastern Honshu, Japan. *Tohoku Journal of Agricultural Science*, 55, 93-97 (2005)
- Segawa, S., "Colored illustration of the seaweeds of Japan" ed. by S. Segawa, Hoikusha, Tokyo, pp. 1-196 (1956) (in Japanese)
- Suda, M., Marine algae from the coast of Iwaki City, Fukushima Prefecture. *Jap. J. Phycol.*, 35, 22-33 (1987) (in Japanese, with English summary)
- Takamatsu, M., The marine algae from Matsushima Bay, Miyagi Prefecture, northeastern Honshu, Japan. *Saito Ho-on Kai Museum Research Bulletin*, 8, 1-43 (1936)
- Takamatsu, M., "An atlas of marine algae from Okirai Bay and its vicinities, Iwate prefecture" ed. by M. Takamatsu, Faculty of Fisheries, Kitazato University, Kesenuma, pp. 1-42 (1974) (in Japanese)
- Taniguchi, K., Primary succession of marine algal communities in the sublittoral zone off Oshika Peninsula, Japan. *Nippon Suisan Gakkaishi*, 62,

- 765-771 (1996) (in Japanese, with English summary)
- Taniguchi, K., Hara, M., Sato, Y., Osada, Y., and Suenaga, H., List of marine algae from the Ogatsu coast of Miyagi Prefecture. *Bull. Tohoku Reg. Fish. Res. Lab.*, **47**, 11-20 (1985)
- Taniguchi, K., Igarashi, T., Utoh, H., Nakano, H., and Minami, T., Current status of aquatic organisms and fisheries under warm hydrographic conditions along the coasts of Tohoku and Hokkaido region, Japan. *Kaiyo monthly*, **38**, 157-160 (2006) (in Japanese)
- Taniguchi, K., Kito, H., and Akiyama, K., Marine algae from the Miyako coast of Iwate Prefecture. *Bull. Tohoku Reg. Fish. Res. Lab.*, **41**, 141-147 (1979) (in Japanese, with English summary)
- Yoshida, T., "*Marine Algae of Japan*" ed. by T. Yoshida, Uchida Roukakuho, Tokyo, pp. 1-1222 (1998) (in Japanese)