

# The Ortho-Plastic Approach to Soft Tissue Management in Trauma

Daniel J. Jordan<sup>1</sup>, Marco Malahias<sup>2</sup>, Wasim Khan<sup>3</sup> and Sandip Hindocha<sup>\*,4</sup>

<sup>1</sup>Plastic Surgery Unit, The Christie NHS Foundation Trust, Manchester, UK

<sup>2</sup>Plastic Surgery Department, Good Hope Hospital, West Midlands, UK

<sup>3</sup>Royal National Orthopaedic Hospital, London, UK

<sup>4</sup>Plastic Surgery Unit, Whiston Hospital, Liverpool, UK

**Abstract:** Fractures with associated soft tissue injuries, or those termed ‘open,’ are not uncommon. There has been much discussion regarding their management, with the guidance from the combined British Orthopaedic Association and British Association and Aesthetic Surgeons teams widely accepted as the gold level of therapy. We aim to discuss the current evidence about the initial management of this group of injuries, taking a journey from arrival in the accident and emergency department through to the point of definitive closure. Other modes of therapy are also reviewed.

**Keywords:** Debridement; open fracture; soft-tissue.

## 1. INTRODUCTION

Fractures are a commonly seen injury in the accident and emergency department. Open and closed fractures differentiate themselves due to the amount of damage to the soft tissues enveloping the bone. Any injury to the bone will damage the surrounding soft tissues to a degree, but where there is a communication to the outside, through a wound, the injury is termed as an open, or compound, fracture. Closed (simple) fractures are those where the bone fracture is not linked to the outside environment.

The communication with the environment can be formed in several ways. High trauma injuries often result in an inside to outside injury where the bone tears through the soft tissues. In comparison, a high velocity penetrating injury, such as in a gun shot wound, will cause injury to the soft tissue and then the underlying bone, dragging material from the environment into the wound. The latter is also seen in blast injuries of war and animal bites, and can often be associated with a greater soft tissue to bone ratio injury. The main concern regarding open fractures is that they permit contamination by the outside world, increasing the morbidity and outcome associated with the bony injury.

The main morbidity results from deep infection, non-union of the fracture, osteomyelitis and amputation [2], with fracture healing being dependant, to a significant extent, on the soft tissue envelope quality surrounding the injured bone, and its blood supply [3].

## 2. EPIDEMIOLOGY

Open fractures studies have often been studied in regional rather than national terms, and its epidemiology

varies depending on war and peace situations. It is thought overall incidence of open fractures is 3.2% of all fractures or 11.5 per 100,000 people. Anatomically it is thought around 3.3% of upper limb, 3.7% lower limb and 0.3% of limb girdle, fractures are open. Up to 21% of tibial fractures are open, mainly due to the thinner surrounding soft tissue envelope as compared to other bones [4].

Open fractures have been seen to affect the population in a bimodal form. Often a younger group, generally male, caused by high energy injuries [mainly relating to sport and road traffic accidents] and an elderly peak, involving low energy injuries [often the osteoporotic female] caused by for instance a fall from standing [4].

Open fractures, despite improved management strategies, continues to involve late infection rate ranging up to 25% [5, 6]. Worldwide agreement on mandatory management follows along intravenous antibiotic use, emergency wound debridement and copious irrigation [6-13], although the timing and choice of these treatments is less conclusive.

The desired outcome in the management of open injuries is not merely salvage in more extreme injuries, but a limb which is functional, painless and aesthetically pleasing and this should be remembered at all stages of management.

## 3. THE MANAGEMENT OF OPEN FRACTURES

### 3.1. Initial Management

Open fractures can present in many scenarios, varying from isolated injuries to multiple traumatic injuries to the patient [4].

Although the more minor open fractures, for example of the upper limb digits may appear to be inconsequential, a quick and safe approach to examining the patient must be adhered to so as to not miss more serious problems. In respect to this, the more traumatic open fractures often act as a distraction to the treating physician masking a more

\*Address correspondence to this author at the Department of Plastic Surgery, Whiston Hospital, Liverpool, UK L35 5DR, UK;  
Tel: + 44(0)1244366265; Fax: +44(0)1244366265;  
E-mail: [hindocha2001@yahoo.com](mailto:hindocha2001@yahoo.com)

immediate threat to both the unconscious and conscious patient. For this reason, the internationally recognised Advanced Trauma Life Support (ATLS) protocols [14] are best adhered to [1].

The first aspect of the ATLS principles is to stabilise the cervical spine and assess the airway. The unconfused patient that walks in with an open injury to the arm usually allows these areas to be considered without formal need, as well as hinting to an adequate breathing and circulatory state. In more severe or multiple trauma cases these areas must be assessed in order and treated appropriately. Radiological scans at this time should be undertaken and include any joints above and below any queried bony injuries [1].

Preliminary assessment of open fracture/s may be commenced once life threatening injuries have been managed appropriately. Initial splinting of the fracture will help with both haemorrhage and pain relief, but prior to this the wound should be examined and the limb status documented [1]. The wound should be examined in a careful and systematic approach, noting size, approximate depth and any active bleeding. This includes a circumferential assessment of the limbs as the soft tissue injury may be initially hidden from sight. Concerns at this time regarding distal vascularity should be addressed early in the therapy. Sensation and motor function should also be grossly examined as to confirm injury pre or post therapy, but often nerves are highly tolerant of stretching injuries and there viability is often unaltered [15, 16].

Direct pressure, and as a last resort, application of a tourniquet usually resolves gross bleeding, but its use is currently only indicated in lethal haemorrhage [17].

At this stage, guidance suggests only gross contamination should be removed and if able to, photography of the wound for records [1]. Finally dressing of the wound ought to take place. Photography of the wound allows easy review of the wound without causing further distress to the patient with repeated dressing changes.

There is currently debate regarding the optimum dressing to cover the wound, with the main debate involving the addition of a povidone-iodine or antibacterial dressing or a simple saline soaked gauze alternative. The former, although with the aim of reducing any present and potential bacterial load [18], has been stated as being a potential source of further soft tissue and cell damage due to the cytotoxic properties of the soak [19]. Rogers *et al.* [20] showed no statistical difference in infection rate between saline and povidone-iodine based dressings, and so at present saline is suggested in guidelines by the BOA/BAPRAS [1].

After gross contamination is removed from the wound, provisional cleaning of the wound with both exploration and irrigation should not be undertaken unless done so by the orthopaedic or plastic teams involved [1]. Antibiotic administration should start as early as possible with appropriate tetanus prophylaxis administered. In the next part, we aim to discuss the current evidence for the above guidance as well as the key steps in the initial debridement process.

Once the patient is stable, and the open fracture is stabilised with appropriate splinting helping to ease both

haemorrhage and pain, the management of the wound can take priority. As stated the current guidance is that the next stage of management is early wound debridement and this needs to be preferably undertaken at a specialist centre if in respect to soft tissues, there is any query regarding tension free closure of the wound, muscular or vascular injury and degloving-type injuries [1].

#### 4. WOUND MANAGEMENT

The key parts of wound management involve firstly grading or attempting to classify the wound to assess the degree of surgical input, and by what teams is required. End morbidity studies have shown that the key factors in wound management involve antibiotics and surgical debridement, the latter involving timing, irrigation of the wound and actual surgical technique. These areas will now be discussed.

##### 4.1. Wound Assessment and Classification

Open fractures are common injuries, and often will present to a hospital where the definitive management is unable to be performed. For this reason, classification of the injury is needed for a concise, reproducible and comparable description of the wound that can be explained to the specialist unit most often over the phone. This classification should also be simple to perform, accurate as well as helping both the initial and specialist unit decide a management plan, based on predictive outcomes related to the injury's score. There are multiple classification systems in use for open fractures but since 1976 one system [13] has been felt to be the most simple and reproducible system available. The Gustilo and Anderson classification also suitably predicts prognostic outcome of the injury, taking into account the bone and soft tissue injuries as well as contamination of the wound. The system was based on diaphyseal long bone injuries and so is less functional for metaphyseal, intra-articular, as well as small bone injuries. The Gustilo and Anderson grading system unfortunately has been questioned as to whether it gives poor inter-observer reliability but as stated is the current mainstay of classification [1, 21, 22]. Gustilo modified his scale further in 1984 to take into account the bone exposure and consequent soft tissue coverage as well as vascular injury of the more severe grade III injuries and the scale is shown in Table 1 [6].

The Gustilo classification system must be used with adjuncts of information given. These include the mechanism and force involved in the injury, and other concomitant injuries or co-morbidities [23].

As well as the Gustilo and Anderson classification, many other indexes have been designed to firstly categorize the injury, and also predict whether or not salvage of the limb is feasible.

Other systems in use for the grading of an injury include the Tscherné Classification, which later evolved to form the Hannover Fracture Scale. Likewise the AO classification is used to describe the degree of bony and soft tissue injury and may be more useful for audit and data collection as it is a more comprehensive scale [1, 24].

The Byrd -Spicer classification is another less used scale [25]. It has a large degree of inter-observer variance, but scores include the force of energy and any presence of

**Table 1. Gustilo-Anderson classification of open fracture injuries.**

Gustilo Type	Definition
I	Open simple fracture, Clean wound, Wound <1cm in length
II	Open simple fracture, Wound >1cm in length without extensive tissue damage including flaps or avulsion type injury
III	High energy injuries resulting in Open segmental/multi-fragmental fracture or bone loss associated with extensive soft tissue laceration, damage or loss. This includes severely contaminated wounds, including farmyard related, any vascular involvement, severe crush injuries and fractures that have been open for over 8 hours pre-treatment
IIIA	As above with adequate periosteal coverage of the fractured bone despite soft tissue damage/loss
IIIB	As above with extensive tissue loss and periosteal stripping. Associated with massive contamination
IIIC	As above associated with an arterial injury requiring repair irrespective of degree of soft tissue injury

devitalised tissue, although it is more vascular injury weighted.

Salvage of more serious injuries, avoiding amputation both early and late, has caused several classifications to be suggested. The majority of these are based on battlefield injuries. Unfortunately each has their own positives and flaws ranging from complexity, reproducibility and predictive outcome [23].

Regarding the need for early or late amputation of the limb, several scores have been designed to assess the benefit of limb salvage *versus* amputation and include the MESI, MESS, PSI, LSI and NISSA indexes (Table 2).

Bosse *et al.* [26] examined the usefulness of the above scores as well as the Hanover fracture scale and found all lack sensitivity, with the MESS appearing to have a reasonable specificity in limb salvage. They found lower

**Table 2. Open fracture injury scores.**

Score	Unabbreviated	Main Indices for Scoring	Predictive Outcome [26]
MESI	Mangled Extremity Syndrome Index [27]	<ul style="list-style-type: none"> <li>Injury Severity Score (ISS)</li> <li>Tegmentum Injury</li> <li>Vascular Injury</li> <li>Fracture Type</li> <li>Bone Loss</li> <li>Wait Time</li> <li>Age</li> <li>Pre-existent Disease</li> </ul>	>20 = amputation
MESS	Mangled Extremity Severity Score [28]	<ul style="list-style-type: none"> <li>Type injury</li> <li>Limb ischemia</li> <li>Presence of shock</li> <li>Age</li> <li>Score is doubled for Ischaemia &gt; 6 hours</li> </ul>	>7 amputation [29]
PSI	Predictive Salvage Index [30]	<ul style="list-style-type: none"> <li>Level of arterial injury</li> <li>Degree of bone injury</li> <li>Degree of muscle injury</li> <li>Interval from injury until arrival in operation room</li> </ul>	>8 amputation
LSI	Limb Salvage Index [31]	<ul style="list-style-type: none"> <li>Artery injury</li> <li>Nerve Injury</li> <li>Bone Injury</li> <li>Skin Injury</li> <li>Muscle Injury</li> <li>Deep Vein Injury</li> <li>Warm Ischaemia Time</li> </ul>	>6 likely amputation
NISSA	See Indices [initialled] [32]	<ul style="list-style-type: none"> <li>Nerve injury</li> <li>Ischemia</li> <li>Soft tissue contamination</li> <li>Skeletal injury</li> <li>Shock</li> <li>Age</li> </ul>	<7 salvage and > 11 amputate

numbers correlated well with salvage, but that higher numbers should not equal amputation. These scores do allow the categorisation of injuries, as no two are identical, and should be used in aiding clinical judgement rather than being a set decision making tool. Ultimately, the patient must be assessed in terms of an individual with personal, psychological and social circumstances as well as their critical physical needs being considered.

There is also some debate whether the classification of an open wound should be done post debridement, [33] and also whether the amount of tissue contamination should be taken into account, especially as this and the size of the wound do not always correlate. These debates are the reason for the numerous attempts to classify all wounds into groups, and these should really only be used for guidance rather than the rule. BOA/BAPRAS guidance states post debridement assessment for wound grading [1].

In this paper injuries will be talked about in terms of Gustilo's classification unless stated.

#### 4.2. Antibiotic Use

The choice of antibiotic has been under debate since studies into open fractures began.

Debate ranges from choice of antibiotic, the way it is administered and even to whether or not antibiotics are even indicated in the first instance.

It is however, well established that antibiotic use is a requirement in the open fracture [1, 6, 13, 34].

Initially the antibiotic regime was given as prophylaxis as injury to the soft tissue envelope of the bone pre-disposes to both wound infection as well as late osteomyelitis, although the higher rates of infection seen in these injuries without antibiotics means it is now thought of as therapeutic [34]. Excision of non viable bone and soft tissues allows any source of bacteria to be removed and aided by the high vascularity of bone allows systemic administration to reach the area and thus be a suitable choice. It is generally accepted that a combination of debridement and antibiotics, as opposed to either one course on its own reduces the risk of future complications. Both these therapies also appear to be only of benefit while there is still non-viable tissue and until the normal fracture healing process has begun, and so serial debridement may be required.

It has been shown that open fractures managed only by debridement may result in infection rates as high as 13.9%, while the addition of antibiotics reduces this rate to 2.3% [34].

A Cochrane study suggested the role of prophylactic antibiotics in reducing infection rates by 59% [35].

The wound can be contaminated at the time of injury, particularly in outside to inside injuries, as well as whilst in the hospital, with Cooney *et al.* showing close ties with the bacterial cultures taken from both the patients nose or skin as well as that from the air surrounding the wound [36, 37]. Despite this, pre-debridement cultures do not appear to correlate with either later infections of the wound or systemic sepsis [5] and this association only mildly improves with post debridement cultures [38]. The general consensus is that if a wound appears infected, then at this point there is

a higher prediction of link to quantitative positive culture results [39, 40]. It has been seen that there is no correlation between the time elapsed between injury and debridement and quantitative bacterial counts [39, 41, 42].

Studies have shown benefits of antibiotic therapy in open fractures, when compared to administration of a normal saline placebo [43]. Gustilo [44] also pointed to a mixture of gram positive, gram negative and mixed bacterial growths from these injuries on arrival of the patient to hospital.

The cephalosporin family has long been the source of antibiotics in the treatment of open fractures due to their good broad spectrum activity against both gram positive and negative organisms. Bischoff *et al.* [45] showed that Cefuroxime was secreted into open fracture wounds increasing the antibacterial activity of the wound secretion, and that this reached a highest peak 4 hours after intravenous administration. This, alongside the paper by Gustilo [44] showing cephalosporin's administered within 3 hours reduced morbidity, points towards the use of antibiotics early in the management of these injuries.

In addition to this the use of a local, or topical, antibiotic therapy has been researched. Neomycin, Bacitracin and Polymyxin have all been assessed with varying benefits suggested, and often only in animal studies [46-49].

An 'antibiotic bead pouch' appears to aid reduction of the bacterial load of the wound, as well as reducing consequent infection rates and osteomyelitis [50-53].

Aminoglycoside impregnated Polymethylmethacrylate (PMMA) beads appear to be the most researched in this field. The benefits of the beads include stopping the re-growth of soft tissue interposing the bone ends as well as creating a space in the healing envelope for later bone grafting/management. As well as this it reduces any systemic toxicity, and reduces secondary wound contamination [54]. And these are now recognised as a useful adjunct in the open fracture therapy, especially in higher grade injuries where bone loss is significant [1].

Studies have equally shown that Type I and II injuries require one pre and two post surgery doses of antibiotic (in this case Cephalosporins again) due to mainly gram positive infection, whilst grade III benefit from a longer course, 48-72 hours post definitive soft tissue coverage, due to an increased incidence of gram negative organisms [55]. The addition of Penicillin G as a third antimicrobial agent is highly recommended for open fractures which have been exposed to soil or a farm environment and in injuries with a considerable crush component or vascular compromise] [56, 57].

Several doses of Teicoplanin has also been shown to help reduce the incidence of infection following open fractures, although results seem to be mainly beneficial in soft tissue injuries with Type II fractures, and the study did not include grade III injuries [58].

Longer courses of antibiotics have not been shown to aid benefit, and especially once definitive coverage of the injury has taken place [13, 56, 59-62].

More recent studies have shown that in grade IIIB infections, the benefit of a one off dose of Teicoplanin and

Gentamycin at the time of definitive soft tissue closure would help reduce the risk of future deep surgical site infection. The reason behind this being the increase of nosocomial organisms present at wound closure as opposed to skin flora, present at time of injury, which are covered by the more broad-spectrum antibiotics [63].

The amount of soft tissue injury does appear to correlate with the infection rates of the wound, and so higher graded injuries appear to require more extensive debridement and antibiotic duration. Infection rates for grade I injuries appear to be up to 2%, rising to 2-7% in grade II injuries. Grade III injuries appear to have an overall incidence of 10-25% of infection rates, with this sitting at 7% for grade IIIA, 10-50% in grade IIIB and 25-50% in grade IIIC [2, 5, 6, 13, 62].

This low rate in grade I fractures is the area which questions whether there is any additional benefit of antibiotics to the thoroughly cleaned wound [64].

The current BOA/BAPRAS guidance takes many of the above points [1]. It states antibiotics should be started as soon as achievable, preferably before 3 hours have elapsed since injury. Co-amoxiclav (1.2 g 8 hourly) is now first choice as it has a similar spectrum of therapy to the cephalosporins (Cefuroxime 1.5 g 8 hourly) but is currently less associated with complications such as *Clostridium Difficile* related Colitis. Clindamycin is a non penicillin related alternative for patients with history of allergic reactions (600 mg 6 hourly).

This should be continued for a maximum of 72 hours, or until soft tissue coverage is achieved, with a dose of whichever antibiotic as well as Gentamycin (1.5 mg/kg) being administered at time of induction of the first debridement.

A further dose of Gentamycin should be given, with either a single dose of Teicoplanin (800 mg) or Vancomycin (1g), at the time of skeletal fixation.

Regarding tetanus, Rhee *et al.* suggested after reviewing the literature that tetanus vaccine should be administered in traumatic wounds in the form of tetanus toxoid if the last booster was given more than 10 years prior or if history is not reliable or available, and as tetanus immunoglobulin in patients with incomplete primary immunization or to patients for whom it has been longer than 10 years since their last booster dose [65].

### 4.3. Surgical Management

It has been suggested that in high energy tissue injuries, surgical debridement is the most important prophylactic measure in reducing infection risk [66] and the first procedure may well determine the long term outcome of the injury and patient [67]. This has been evident in the opinion that inadequate debridement may in fact cause an iatrogenic injury and raise later morbidity rates and long term outcome [68, 69].

This theory has been challenged, as for the need of antibiotics, in low energy injuries [70], where it has been considered grade I injuries could be treated as closed injuries, although currently this should be the exception rather than the rule, at least until further evidence is shown. Grade II and above type injuries are often unstable or

involve displacement and will almost definitely need some aspect of surgical therapy.

Debridement is one of the key stages of open fracture management; its basics running through the fact that nonviable tissues and foreign material in the wound both hinder the host defence mechanisms and enhance bacterial growth [71]. Failure to excise the devitalised tissue often causes increased morbidity, as the amount of devitalised tissue contributes to the amount of present bacteria and is often linked to the degree of energy released into the tissues by the injury. Hence why the larger injuries appear to have higher infection rates and require more radical debridement.

#### 4.3.1. Vascular Injury

Vascular injury occurs in 1-4% of open fractures that present in non-war situations, and often requires rapid management, often with the use of shunts or vein interposition especially with the risk of infection in these injuries [72]. These are best analysed in the suspicion of a vascular injury using a Doppler probe, with signs of ischaemia, absent pulses and neurological signs, or a bony injury known to have high arterial damage association [73].

Angiography can be used to confirm, particularly upper limb ischaemia, as this may define limb salvage *versus* amputation [73], although it can delay therapy, and clinical judgement in light of findings and the level and configuration of the fracture can strongly guide in decision making [1]. Vascular compromise should be recognised immediately [loss of distal pulses]. Restoration of circulation should take place under six hours, although muscle death has been seen to occur as soon as 3 hours post injury [1].

#### 4.3.2. Time to Debridement

Thorough the literature there is described a 6 hour rule in which time surgical debridement should take place. However, there does not seem to be any agreed study or text where this rule arose from although numerous studies report the potential benefits, the majority experimental [74-76].

Early debridement has been seen to significantly reduce later infection rates in war related open fractures [77, 78] as well as within civilian populations [79]. Kindsfater [80] showed increased risk in debridement after 5 hours post-injury. This study is not as conclusive as other studies as several variables were not taken into account and numerous dissimilarities between the treatment groups was seen.

Naique *et al.* [81] found that there was no significant difference in deep infections between pre 6 and post 6 debridement but that there was a higher percentage of infections in the second group.

Likewise, a number of studies have questioned the 6 hour rule, with delayed debridement [often within 24 hours] showing no significant outcome difference when compared to pre-6 hour post injury debridement [62, 82-84].

Webb [85] and Harley [86] showed no difference between the time of debridement and end clinical or functional outcome in type 3 fractures specifically, whilst a delay from injury to presentation, such as in more remote regions has been shown to have good outcomes as long as the main principles of satisfactory debridement, fracture stabilisation and soft tissue management [87].

Some studies have shown that the specific timing of debridement relates no correlation to quantitative bacterial counts [42, 88] whilst studies in children have shown no benefit in urgent debridement of the wound under 6 hours post injury, as long as intravenous antibiotics have been started [89]. In this study procedures occurring after the 7<sup>th</sup> hour had similar long term complications to those operated on earlier, and the complication risk being more dependent on the grade of injury.

This is reflected in the suggestion that all open wounds should be explored in normal working hours, but within 24 hours of injury [1].

Immediate exploration is indicated in wounds grossly contaminated, devascularised distally, or those with evidence of compartment syndrome. The multiply injured patient should also be considered in this group [1].

#### 4.3.3. Irrigation

Irrigation is used as an adjunct to aid the reduction in any bacterial load of a wound. Again an area of debate, many surgeons have shown benefit of using free flowing solutions and high pressure pulsatile lavage in cleaning the wound [90]. However, it also been suggested that high pressure pulsatile lavage can damage the viable soft tissues of the wound, including the healing bone [91], as well as driving foreign materials further into the wound [92, 93].

High pressure pulsatile lavage may be useful in removing large particulate debris and foreign matter. However, it may lead to a illogical rebound in the bacterial count of the wound, possibly due to the above factors [94].

Guidance by Crowley [90], based on literature available, suggested the use of normal saline without adjunctive solutions through low pressure irrigation methods, and is matched with current guideline from the BOA/BAPRAS [1].

Soap has been used as an adjunct, similar to povidone-iodine to help reduce any bacterial load of the wound. Conroy [47] and Anglen [48] showed no great benefit when comparing soap vs antibiotic and antiseptic solutions, and its use is not widespread due to concerns regarding potential cytotoxic properties. Likewise the use of additives, such as bacitracin, castile soap, and benzalkonium chloride, which do reduce the initial bacterial load, also appear to cause a later rebound bacterial count [94].

Tap water and normal saline solutions appear to have no great difference in terms of benefit in reducing bacterial load *in vitro* [95], although tap water consistently will grow pseudomonas when cultured in optimum conditions.

#### 4.4. Surgical Debridement

Primary debridement should take place in a suitably sterile location. The fractured limb may benefit from a proximally based tourniquet especially when massive active blood loss is occurring or expected, although its use should be minimised to maximise tissue salvage, and prevent further ischaemia. Tourniquet use has been shown to increase the incidence of wound infection, more than likely due to worsening tissue hypoxia and acidosis [71, 96], but the benefits of a clearer working field, not covered in blood, may be of more benefit initially [1].

Cleansing of the wound and limb skin with soap and then a chlorhexidine scrub should occur, with povidone-iodine as an option. The use of hydrogen peroxide has been recommended to cleanse the area [97]. All foreign matter, as well as tissue whose viability is undetermined should be excised including muscle, bone, and the skin edges of the wound. This can aid exploration of the wound, although the finer structures of the limbs, nerves and tendons, are often found to be more resilient and should not be excised unless there non-viability is undoubted e.g. detached [15, 16]. If further wound opening, rather than excision is required, the fasciotomy lines should be used [1].

The actual amount of tissue to be excised is still a controversial point, with the only agreement being all non-viable tissue excised, although the definition of non-viable is not clear cut. Necrotic tissue should be radically excised [97], but tissue whose viability is questioned is a finer point, and may benefit from serial review of the wound and further debridement. This is key in degloving injuries where the amount of injury can be under-estimated [1].

Systematic exploration of each level of the wound, from superficial to deep, allows a reduction in any non-viable tissue being missed [1].

Wound edge debridement is often dependant on the surgeons experience [98, 99] and is often guided by the four C's which can be found in Table 3 [100,101].

**Table 3. Excision of devitalised tissue 'The 4 C's'.**

<b>Colour</b>	Red (not pale or brown/black)
<b>Consistency</b>	Non-waxy
<b>Contractility</b>	On being pinched
<b>Capillary Bleeding</b>	On being incised/cut

Prophylactic fasciotomy should be performed in vascular injuries associated with open fractures and significant crush injuries. Acute compartment syndrome is a severe threat to limb survival and distal limb ischaemia. An open fracture does not equate to an open compartment and it is key to remember that these patients can still develop this complication [96] and a low threshold for performing the procedure should be followed [102].

This risk is exaggerated in the patient who can tell the surgeon symptoms are worsened and inter-compartmental pressures may be mandatory in the unconscious or anaesthetised patient with a swollen limb [1].

There is traditional agreement that it is better to leave a traumatic wound open after the initial debridement [10, 103] although there is now a shift to early closure which may reduce osteomyelitis and non-union rates, although these factors are dependant on a satisfactory wound debridement in the first instance [104-106].

Repeat, or serial, debridement should take place between 24 and 48 hours and be performed under general anaesthesia allowing a full examination of the wound. Once the wound is deemed satisfactorily clean, closure of the wound is advised with whichever technique is best indicated.

The presence of granulation tissue at later debridement is an encouraging sign showing that a wound is capable of mounting a healing response on both local and systemic levels. It may help with tissue loss in terms of defect although long term wise it aids little with respect to flap coverage and ultimate function, potentially even causing functional impairment. As well as this, the formation of granulation tissue over the surface of necrotic muscle can obscure the true extent of any necrosis as well as harbouring bacteria. Its debridement should be considered throughout review of the wound [107].

#### 4.4.1. Surgical Debridement Adjuncts

Bone that is present in the wound but potentially viable has also been reapplied to the injured site as a bone graft, and salvage techniques include washing in povidone-iodine, autoclaving and gentamycin soak with good results in a case study [108]. Currently bone that fails the 'tug test' should be excised [1].

Alongside this, several papers have assessed the use of laser Doppler flowmetry in the assessment of bone viability, with mixed results [109, 110].

Studies have also show relationships with the amount of oxygen in the tissues surrounding a fracture, stating higher concentrations around the injury site as compared to both the non-injured opposite limb and region of non-injury on the injured limb. This change is most evident during the first 4 days post injury and returns to normal by day 10, and may be able to be used as a predictor of complications [111, 112].

New studies are also looking at addition of Interleukin-12 as a way of reducing infection, particularly with nanocoating of any bone fixator used [113], as well as autologous platelet-rich plasma (PRP) gel [114].

## 5. POST EMERGENCY DEBRIDEMENT

The options available after emergency debridement are to leave the wound open, to temporarily cover or definitively cover. Negative Pressure Wound Therapy (NWPT) has revolutionised this area is now commonly used as a temporary dressing aiding in reducing both morbidity and the need for further surgery.

### 5.1. Negative Pressure Wound Therapy (NPWT)

NWPT has gained universal appeal and appear to help in both the reduction of wound bacterial load, and the amount of tissue required for definitive coverage. It is suggested that NPWT increases local blood flow, reduces oedema and promotes healthy granulation tissue formation [115].

It has been documented in even helping reduce compartment syndrome pain and has even saved a limb from amputation [116,117].

The scope for NPWT use is changing year on year, with the use of dermal regeneration templates and cryo-preserved dermis, [118,119] but it is important NPWT is used as an adjunct for delayed wound closure and not used as an alternative to this or debridement [1, 120-122].

NWPT should be used in any wound that is left open following debridement until definitive surgery is performed [1].

## 5.2. Bone Fixation and Soft Tissue Coverage

Definitive therapy in terms of bone fixation as well as soft tissue coverage reduces overall infection rates as well as long term morbidity. Adequate reduction and realignment of the limb allows normalising of landmarks and there is much debate over internal *versus* external fixation. Whichever is chosen should minimise further soft tissue trauma, allow further debridement or assessment of the wound and closure as require. External fixation has been the preferred choice as it is quick and less invasive, although it does have additional risks such as joint stiffness and pin-site infections [123-126].

Delayed definitive soft tissue coverage is now the accepted choice worldwide, with immediate wound closure the exception. The definition of early *versus* delayed coverage however, is not a set standard. The urgency of soft tissue cover, especially in grade III injuries suggested by Godina [97] in 1986 [within 72 hours of injury] has been shown to be less required as debridement, antibiotic therapy and wound care has improved [127]. Although the 'fix and flap' technique where immediate fracture fixation, debridement and flap coverage has been observed with results not dissimilar to delayed procedures [115,128].

Definitive soft tissue coverage should be aimed for within 7 days, and studies have shown this to improve long term morbidity in terms of infection rates, bone non-union time and rate of union in comparison with further delay [129], although this should be done dependant on the adequately skilled surgical team availability, the patient's condition and the ability to adequately consent the patient for long term morbidity. This is also dependant on the grade of injury, as grade I and even II injuries may be closed primarily [130].

If amputation is indicated, studies have shown the use of the amputated limb as a donor site for the new flap is possible and helps to reduce the remaining burden on the patient, and this if often best performed at first debridement by the suitable orthoplastic team [131].

## 6. PATIENT FACTORS

It has been commented that the injured patient must be assessed in terms not just relating to an open fracture wound. Outcome of open fractures has been linked to several pre-injury morbidities and this should be highlighted and taken into account before a definitive therapy is chosen.

The management protocol described above unfortunately does not always improve morbidity in the open fracture patient. Patient condition is a key factor, with diabetes increasing the risk of infection and subsequently changing the definitive management, often resulting in amputation to the more distal injury [132].

The general nutritional state of the patient is important, as open fractures with or without multiple traumas increases the physiological burden on the body, and poor nutrition and physical stores will reduce the host immunity as well as increase the healing process time. Smoking has a similar input, often resulting in increased infections, delayed union rates and definitive cover failure [133,134].

Likewise, HIV infection has been shown to increase the risk of infection and may alter the ultimate decision

regarding type and timing of soft tissue and bone fixation [135].

## CONCLUSION

Open fractures continue to be a source of debate. Their occurrence will often be seen, more so in wartime situations, but also in urban and rural situations. Although each injury is often different in site and cause they should all be managed in a clear, structured way so that the potential life-altering complications can be kept to a minimum. This structured approach, currently is best described by the BOA/BAPRAS guideline which are well defined by the two surgical associations most present in their management. These guidelines should be followed in the majority of injuries including children who should be treated, in terms of debridement, as in the recommendations for adults [1]. The basics suggested (assessment, antibiotics and debridement) appear to be the mainstay of therapy although there is the possibility of new suggestions and adjuncts which may revolutionise the therapy of open fractures, such as the now common addition of NWPT. It must be remembered that each of these injuries although may fit into a certain classification, are related to an individual and their pre-injury morbidity as well as social, psychological and end physical needs will always play an important part in their management.

## CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

## ACKNOWLEDGEMENTS

Declared none.

## REFERENCES

- [1] Standards for the management of open fractures of the lower limb, a short guide. Guidelines Ed 1, Sept 2009. British Orthopaedic Association, British Association of Plastic Reconstructive and Aesthetic Surgeons. <http://www.bapras.org.uk/page.asp?id=724> accessed July 15, 2012
- [2] Rittmann WW, Schibli M, Matter P, Allgower M. Open fractures. Long-term results in 200 consecutive cases. *Clin Orthop Relat Res* 1979; 138: 132-40.
- [3] Harry LE, Sandison A, Pearse MF, Paleolog EM, Nanchahal J. Comparison of the vascularity of fasciocutaneous tissue and muscle for coverage of open tibial fractures. *Plast Reconstr Surg* 2009; 124(4): 1211-9.
- [4] Court-Brown CM, Rimmer S, Prakash U, McQueen MM. The epidemiology of open long bone fractures. *Injury* 1998; 29(7): 529-34.
- [5] Gustilo RB, Gruninger RP, Davis T. Classification of type III (severe) open fractures relative to treatment and results. *Orthopedics* 1987; 10: 1781-8.
- [6] Gustilo RB, Mendoza RM, Williams DN. Problems in management of type III (severe) open fractures: a new classification of type III open fractures. *J Trauma* 1984; 24: 742-6.
- [7] Turen CH, DiStasio AJ. Treatment of grade IIIB and grade IIIC open tibial fractures. *Orthop Clin North Am* 1994; 25: 561-71.
- [8] Esterhai JL Jr, Queenan J. Management of soft tissue wounds associated with type III open fractures. *Orthop Clin North Am* 1991; 22: 427-32.
- [9] Burgess AR, Poka A, Brumback RJ, Bosse MJ. Management of open grade III tibial fractures. *Orthop Clin North Am* 1987; 18: 85-93.
- [10] Gustilo RB, Merkow RL, Templeman D. The management of open fractures. *J Bone Joint Surg Am* 1990; 72: 299-304.
- [11] Brumback RJ, Ellison PS Jr, Poka A, Lakatos R, Bathon GH, Burgess AR. Intramedullary nailing of open fractures of femoral shaft. *J Bone Joint Surg Am* 1989; 71: 1324-31.
- [12] Patzakis MJ, Wilkins J, Moore TM. Use of antibiotics in open tibial fractures. *Clin Orthop Relat Res* 1983; 178: 31-5.
- [13] Gustilo RB, Anderson JT. Prevention of infection in the treatment of one thousand and twenty-five open fractures of long bones: retrospective and prospective analyses. *J Bone Joint Surg Am* 1976; 58: 453-8.
- [14] American College of Surgeons Committee on Trauma: Resources for Optimal Care of the Injured Patient. Available at <http://www.facs.org/trauma/atls/index.html> [Accessed July 10, 2012].
- [15] Bumbasarevic M, Lesic A, Mitkovic M, Bumbasarevic V. Treatment of blast injuries of the extremity. *J Am Acad Orthop Surg* 2006; 14: 577-81.
- [16] Has B, Kvolik S, Kristek J, Habek D. Intact radial and median nerve after open third degree distal fracture of the humerus. *Coll Antropol* 2006; 30(2): 447-50.
- [17] Kragh JF, O'Neill ML, Beebe DF, Fox CJ, Beekeley AC, Cain JS, Parsons DL, Mabry RL, Blackburne LH. Survey of the indications for use of emergency tourniquets. *J Spec Oper Med* 2011; 11(1): 30-8.
- [18] Gilmore OJ, Sanderson PJ. Prophylactic interparietal povidone-iodine in abdominal surgery. *Br J Surg* 1975 62: 792-9.
- [19] Lineaweaver W, McMorris S, Soucy D, Howard R. Cellular and bacterial toxicities of topical antimicrobials. *Plast Reconstr Surg* 1985; 75(3): 394-6
- [20] Rogers DM, Blouin GS, O'Leary JP. Povidone-iodine wound irrigation and wound sepsis. *Surg Gynecol Obstet* 1983; 157: 426-30.
- [21] Brumback RJ, Jones AL. Interobserver agreement in the classification of open fractures of the tibia: the results of a survey of two hundred and forty-five orthopaedic surgeons. *J Bone Joint Surg (Am)* 1994; 76-A: 1162-6.
- [22] Cross WW, Swiontkowski M. Treatment principles in the management of open fractures. *Indian J Orthop* 2008; 42(4): 377-86.
- [23] Durrant CA, Mackay SP. Orthoplastic classification systems: the good, the bad, and the ungainly. *Ann Plast Surg* 2011; 66(1): 9-12.
- [24] Ruedi TP, Buckley RE, Moran CG. *AO Principles of Fracture Management, Vol 1-Principles*. 2<sup>nd</sup> Exp Ed. AO Publishing 2007; pp. 91-111.
- [25] Byrd HS, Spicer TE, Cierny G. 3rd Management of open tibial fractures. *Plast Reconstr Surg* 1985; 76(5): 719-28.
- [26] Bosse MJ, MacKenzie EJ, Kellam JF, et al. A prospective evaluation of the clinical utility of the lower-extremity injury-severity scores. *J Bone Joint Surg Am* 2001; 83(1): 3-14.
- [27] Gregory RT, Gould RJ, Pecllet M, et al. The mangled extremity syndrome (M.E.S.): a severity grading system for multisystem injury of the extremity. *J Trauma* 1985; 25(12): 1147-50.
- [28] Johansen K, Daines M, Howey T, et al. Objective criteria accurately predicts amputation following lower extremity trauma. *J Trauma* 1990; 30: 568-73.
- [29] Helfet DL, Howey T, Sanders R, Johansen. Limb salvage versus amputation. Preliminary results of the Mangled Extremity Severity Score. *Clin Orthop Relat Res* 1990; (256): 80-6.
- [30] Howe Jr HR, Poole Jr GV, Hansen Jr ST, et al. Salvage of lower extremities following combined orthopaedic and vascular trauma: A predictive salvage index. *Am Surg* 1987; 53: 205-8.
- [31] Russell WL, Sailors DM, Whittle TB, et al. Limb salvage versus traumatic amputation: A decision based on a seven-part predictive index. *Ann Surg* 1991; 213: 473-81.
- [32] McNamara MG, Heckman JD, Corley FG. Severe open fractures of the lower extremity: A retrospective evaluation of the Mangled Extremity Severity Score (MESS). *J Orthop Trauma* 1994; 8: 81-7.
- [33] Zalavras CG, Marcus RE, Levin LS, Patzakis MJ. Management of open fractures and subsequent complications. *J Bone Joint Surg Am* 2007. 89; 883-95.
- [34] Patzakis MJ, Harvey JP Jr, Ivler D. The role of antibiotics in the management of open fractures. *J Bone Joint Surg Am* 1974; 56: 532-41.
- [35] Gosselin RA, Roberts I, Gillespie WJ. Antibiotics for preventing infection in open limb fractures. *Cochrane Database Syst Rev* 1: 2004; CD003764.
- [36] Sochen JE. Orthopedic wounds. *Am J Surg* 1994; 167(1A): 52S-4; discussion 54S-5.
- [37] Cooney WP 3rd, Fitzgerald RH Jr, Dobyms JH, Washington JA 2nd. Quantitative wound cultures in upper extremity trauma. *J Trauma* 1982; 22: 112-7.
- [38] Lee J. Efficacy of cultures in the management of open fractures. *Clin Orthop Relat Res* 1997; 339: 71-5.



- [39] D'Souza A, Rajagopalan N, Amaravati RS. The use of qualitative cultures for detecting infection in open tibial fractures. *J Orthop Surg* 2008; 16(2): 175-8.
- [40] Sen RK, Murthy NR, Gill SS, Nagi ON. Bacterial load in tissues and its predictive value for infection in open fractures. *J Orthop Surg (Hong Kong)* 2000; 8: 1-5.
- [41] Moore TJ, Mauney C, Barron J. The use of quantitative bacterial counts in open fractures. *Clin Orthop Relat Res* 1989; (248): 227-30.
- [42] Wilkins J, Patzakis M. Choice and duration of antibiotics in open fractures. *Orthop Clin North Am* 1991; 22: 433-7.
- [43] Bergman BR. Antibiotic prophylaxis in open and closed fractures: a controlled clinical trial. *Acta Orthop Scand* 1982; 53(1): 57-62.
- [44] Gustilo RB. Use of antimicrobials in the management of open fractures. *Arch Surg* 1979; 114(7): 805-8.
- [45] Bischoff M, Beck A, Frei P, Bischoff G. Pharmacokinetics of cefuroxime in traumatic wound secretion and antibacterial activity under vacuum therapy. *J Chemother* 2010; 22(2): 92-7.
- [46] Rosenstein BD, Wilson FC, Funderburk CH. The use of bacitracin irrigation to prevent infection in postoperative skeletal wounds: An experimental study. *J Bone Joint Surg Am* 1989; 71: 427-30.
- [47] Conroy BP, Anglen JO, Simpson WA, *et al.* Comparison of castle soap, benzalkonium chloride, and bacitracin as irrigation solutions for complex contaminated orthopaedic wounds. *J Orthop Trauma* 1989; 13: 332-7.
- [48] Anglen JO. Comparison of soap and antibiotic solutions for irrigation of lower-limb open fracture wounds. A prospective, randomized study. *J Bone Joint Surg Am* 2005; 87(7): 1415-22.
- [49] Anglen JO. Wound irrigation in musculoskeletal injury. *J Acad Orthop Surg* 2001; 9: 219-26.
- [50] Seligson D, Ostermann PA, Henry SL, Wolley T. The management of open fractures associated with arterial injury requiring vascular repair. *J Trauma* 1994; 37(6): 938-40.
- [51] Seligson D, Henry SL. Treatment of compound fractures. *Am J Surg* 1991; 161(6): 693-701.
- [52] Prasam ML, Zych G, Ostermann PA. Am Wound management for severe open fractures: use of antibiotic bead pouches and vacuum-assisted closure. *J Orthop (Belle Mead NJ)* 2009; 38(11): 559-63.
- [53] Bowyer GW. Antibiotic impregnated beads in open fractures. A report on the technique and possible applications in military surgery. *J R Army Med Corps* 1993; 139(3): 100-4.
- [54] Zalavras CG, Patzakis MJ, Holtom PD, Sherman R. Management of open fractures. *Infect Dis Clin North Am* 2005; 19(4): 915-29.
- [55] Antrum RM, Solomkin JS. A review of antibiotic prophylaxis for open fractures. *Orthop Rev* 1987; 16(4): 246-54.
- [56] Tscherne H. The management of open fractures. In: Tscherne H, Gotzen L, eds. *Fractures with soft tissue injuries*. Berlin: Springer-Verlag 1984; pp. 10-3257.
- [57] Sanders R, Swiontkowski M, Nunley J, Spiegel P. The management of fractures with soft-tissue disruptions. *J Bone Joint Surg Am* 1993; 75-A: 778-89.
- [58] Hughes SP, Strachan E, Miles R, Williams AH. Teicoplanin in open fractures: a preliminary report. *Eur J Surg Suppl* 1992; (567): 15-7.
- [59] Patzakis MJ, Wilkins J, Moore TM. Considerations in reducing the infection rate in open tibial fractures. *Clin Orthop* 1983; 178: 36-41.
- [60] Olson SA. Open fractures of the tibial shaft: current treatment. *J Bone Joint Surg Am* 1996; 78-A: 1428-37.
- [61] Lawrence RM, Hoepflich PD, Huston AC, Benson DR, Riggins RS. Quantitative microbiology of traumatic orthopaedic wounds. *J Clin Microbiol* 1978; 8(6): 673-5.
- [62] Patzakis MJ, Wilkins J. Factors influencing infection rate in open fracture wounds. *Clin Orthop Relat Res* 1989; 243: 36-40.
- [63] Glass GE, Barrett SP, Sanderson F, Pearse MF, Nanchahal. The microbiological basis for a revised antibiotic regimen in high-energy tibial fractures: preventing deep infections by nosocomial organisms. *J Plast Reconstr Aesthet Surg* 2011; 64(3): 375-80.
- [64] Stevenson J, McMaughton G, Riley J. The use of prophylactic flucloxacillin in treatment of open fractures of the distal phalanx within an accident and emergency department: a double-blind randomized placebo-controlled trial. *J Hand Surg Br* 2003; 28(5): 388-94.
- [65] Rhee P, Nunley MK, Demetriades D, Velmahos G, Doucet JJ. Tetanus and trauma: a review and recommendations. *J Trauma* 2005; 58(5): 1082-8.
- [66] Covey DC. Blast and fragment injuries of the musculoskeletal system. *J Bone Joint Surg Am* 2002; 84: 1221-34.
- [67] Lerner A, Reis D, Soudry M. Severe injuries to the limbs. Staged treatment. Springer. Berlin Heidelberg 2007.
- [68] Swiontkowsky M. Criteria for bone debridement in massive lower limb trauma. *Clin Orthop* 1989; 243: 41-7.
- [69] Bielawski J, Sygnatowicz. Is posttraumatic osteomyelitis of iatrogenic nature? *Chir Narzadow Ruchu Ortop Pol* 1998; 63(4): 353-61.
- [70] Yang EC, Eisler J. Treatment of isolated type I open fractures: Is emergent operative debridement necessary? *Clin Orthop Relat Res* 2003; 410: 289-94.
- [71] Zalavras CG, Patzakis MJ. Open fractures: evaluation and management. *J Am Acad Orthop Surg* 2003; 11: 212-9.
- [72] Gabel G, Pyrc J, Hintersheer I, Zwipp H, Saeger HD, Bergert H. Arterial injuries combined with open fractures--management and therapy. *Zentralbl Chir* 2009; 134(4): 292-7.
- [73] Schlickewei W, Kuner EH, Mullaji AB, Gotze B. Upper and lower limb fractures with concomitant arterial injury. *J Bone Joint Surg Br* 2003; 74(2): 181-8.
- [74] Gregory P, Sanders R. The management of severe fractures of the lower extremities. *Clin Orthop* 1995; 318: 95-105.
- [75] Robson MC, Duke WF, Krizek TJ. Rapid bacterial screening in the treatment of civilian wounds. *J Surg Red* 1973; 14: 426-30.
- [76] Friedrich P. Die Aseptische Versorgung frischer Wunden. *Arch Klin Chir* 1898; 57: 240-88.
- [77] Jacob E, Erpekling JM, Murphy KP. A retrospective analysis of open fractures sustained by U.S. Military personnel during operation Just Cause. *Mil Med* 1992; 157: 552-6.
- [78] Lerner A, Fodor L, Soudry M; Is staged external fixation a valuable strategy for war injuries to the limbs? *Curr Orthop Relat Res* 2006; 448: 217-24.
- [79] Yokoyama K, Itoman M, Uchino M, Fukushima K, Nitta H, KoJima Y. Immediate *versus* delayed intramedullary nailing for open fractures of the tibial shaft: a multivariate analysis of factors affecting deep infection and fracture healing. *Indian J Orthop* 2008; 42(4): 410-9.
- [80] Kindsfater K, Jonassen EA. Osteomyelitis in grade II and grade III open tibia fractures with late debridement. *J Orthop Trauma* 1995; 9: 121-7.
- [81] Naique SB, Pearse M, Nanchahal J. Management of severe open tibial fractures: the need for combined orthopaedic and plastic surgical treatment in specialist centres. *J Bone Joint Surg Br* 2006; 88(3): 351-7.
- [82] Singh J, Rambani R, Hashim Z, Raman R, Sharma HK. The relationship between time to surgical debridement and incidence of infection in grade III open fractures. *Strategies Trauma Limb Reconstr* 2012; 7(1): 33-7.
- [83] Bednar DA, Parikh J. Effect of time delay from injury to primary management on the incidence of deep infection after open fractures of lower extremities caused by blunt trauma in adults. *J Orthop Trauma* 1993; 7: 532-5.
- [84] Pollak AN, Jones AL, Castillo RC, Bosse MJ, *et al.* LEAP Study Group. The relationship between time to surgical debridement and incidence of infection after open high-energy lower extremity trauma. *J Bone Joint Surg Am* 2010; 92(1): 7-15.
- [85] Webb LX, Boss MJ, Castillo RC, Macme zie EJ. LEAP Study Group. Analysis of surgeon-controlled variables in the treatment of limb-threatening type-III open tibial diaphyseal fractures. *J Bone Joint Surg Am* 2007; 89(5): 923-8.
- [86] Harley BJ, Beaupre LA, Jones CA, *et al.* The effect of time to definitive treatment on the rate of non-union and infection in open fractures. *J Orthop Trauma* 2002; 16: 484-90.
- [87] Ashford RU, Mehta JA, Cripps R. Delayed presentation is no barrier to satisfactory outcome in the management of open tibial fractures. *Injury* 2004; 35(4): 411-6.
- [88] Moore TJ, Mauney C, Barron J. The use of quantitative bacterial counts in open fractures. *Clin Orthop Relat Res* 1989; 248: 227-30.
- [89] Skaggs DL, Friend L, Alman B, *et al.* The effect of surgical delay on acute infection following 554 open fractures in children. *J Bone Jt Surg Am* 2005; 87(1): 8-12.
- [90] Crowley DJ, Kanakaris NK, Giannoudis PV. Irrigation of the wounds in open fractures. *J Orthop Trauma* 2007; 89(5): 580-5.
- [91] Dirschl DR, Duff GP, Dahners LE, Edin M, Rahn BA, Miclau T. High pressure pulsatile lavage irrigation of intraarticular fractures: effects on fracture healing. *Orthop Trauma* 1998; 12(7): 460-3.
- [92] Hassinger SM, Harding G. High-pressure pulsatile lavage propagates bacteria into soft tissue. *Clin Orthop Relat Res* 2005; 439: 27-31.
- [93] Draeger RW, Dahners LE. Traumatic wound debridement. A comparison of irrigation methods. *J Orthop Trauma* 2006; 20: 83-8.
- [94] Owens BD, White DW, Wenke JC. Comparison of irrigation solutions and devices in a contaminated musculoskeletal wound survival model. *J Bone Joint Surg Am* 2009; 91(1): 92-8.

- [95] Gaines RJ, DeMaio M, Peters D, Hasty J, Blanks J. Management of contaminated open fractures: a comparison of two types of irrigation in a porcine model. *J Trauma Acute Care Surg* 2012; 72(3): 733-6.
- [96] Blick SS, Brumback RJ, Poka A, et al. Compartment syndrome in open tibial fractures. *J Bone Joint Surg Am* 1986; 68: 1348-53.
- [97] Godina M. Early microsurgical reconstruction of complex trauma of the extremities. *Plast Reconstr Surg* 1986; 78: 285-92.
- [98] Bartlett C. Clinical update gunshot wound ballistics. *Clin Orthop* 2003; 408: 28-57.
- [99] Volgas D, Stannard J, Alonso J. Current orthopaedic treatment of ballistic injuries. *Injury* 2005; 36: 380-6.
- [100] Bartlett C, Helfet D, Hausman M, Strauss E. Ballistics and gunshot wounds: Effects on musculoskeletal tissues. *J Am Acad Orthop Surg* 2000; 8: 21-36.
- [101] Bowyer GW. Management of small fragment wounds: Experience from the Afghan border. *J Trauma* 1996; 40: S170-2.
- [102] Moed BR, Fakhouri AJ. Compartment syndrome after low-velocity gunshot wounds to the forearm. *J Orthop Trauma* 1991; 5: 134-7.
- [103] Weitz-Marshall A, Bosse M. Timing of closure of open fractures. *J Am Acad Orthop Surg* 2002; 10: 379-84.
- [104] Fischer MD, Gustilo RB, Varecka TF. The timing of flap coverage, bone-grafting, and intramedullary nailing in patients who have a fracture of the tibial shaft with extensive soft-tissue injury. *J Bone Joint Surg Am* 1991; 73: 1316-22.
- [105] Gopal S, MaJumder S, Batchelor AG, Knight SL, De Boer P, Smith RM. Fix and flap: the radical orthopaedic and plastic treatment of severe open fractures of the tibia. *J Bone Joint Surg Br* 2000; 82: 959-66.
- [106] Cierny G 3rd, Byrd HS, Jones RE. Primary versus delayed soft tissue coverage for severe open tibial fractures: a comparison of results. *Clin Orthop* 1983; 178: 54-63.
- [107] Baechler MF, Groth AT, Nesti LJ, Martin BD. Soft Tissue Management of War Wounds to the Foot and Ankle. *Foot Ankle Clin* 2010; 15(1): 113-38.
- [108] Kumar P, Shrestha D, Bajaeacharya S. Replacement of an extruded segment of radius after autoclaving and sterilising with gentamicin. *J Hand Surg Br* 2006; 31(6): 616-8.
- [109] Hinsley DE, Hobbs CM, Watkins PE. The role of laser Doppler flowmetry in assessing the viability of bone fragments in an open fracture. *Injury* 2002; 33(5): 435-8.
- [110] Swiontkowski MF. Criteria for bone debridement in massive lower limb trauma. *Clin Orthop Relat Res* 1989; (243): 41-7.
- [111] Seekamp A, Ziegler M, Gunderoth M, Regel G. Transcutaneous po2 measurement in follow-up of severe soft tissue injuries of open fractures. *Zentralbl Chir* 1995; 120(1): 16-21; discussion 22-3.
- [112] Robla J, Zych GA, Matos LA. Assessment of soft tissue injury in open tibial shaft fractures by transcutaneous oximetry. *Clin Orthop Relat Res* 1994; (304): 222-8.
- [113] Li B, McKeague AL. Emerging ideas: Interleukin-12 nanocoatings prevent open fracture-associated infections. *Clin Orthop Relat Res* 2011; 469(11): 3262-5.
- [114] Kazakos K, Lyras DN, Verettas D, Tilkeridis K, Tryfonidis. The use of autologous PRP gel as an aid in the management of acute trauma wounds. *Injury* 2009; 40(8): 801-5.
- [115] Stannard JP, Singanamala N, Volgas DA. Fix and flap in the era of vacuum suction devices: What do we know in terms of evidence based medicine? *Injury* 2010; 41(8): 780-6.
- [116] Brem MH, Blanke M, Olk A, Schmidt J, Mueller O, Hennig FF, Gusinde J. The vacuum-assisted closure (V.A.C.) and instillation dressing: limb salvage after 3 degrees open fracture with massive bone and soft tissue defect and superinfection. *Unfallchirurg* 2008; 111(2): 122-5.
- [117] Kneser U, Leffler M, Bach AD, Kopp J, Horch RE. Vacuum assisted closure (V.A.C.) therapy is an essential tool for treatment of complex defect injuries of the upper extremity. *Zentralbl Chir* 2006; 131(Suppl 1): S7-12.
- [118] Barnett TM, Shilt JS. Use of vacuum-assisted closure and a dermal regeneration template as an alternative to flap reconstruction in pediatric grade IIIB open lower-extremity injuries. *Am J Orthop (Belle Mead NJ)* 2009; 38(6): 301-5.
- [119] Brandi C, Grimaldi L, Nisi G, et al. Treatment with vacuum-assisted closure and cryo-preserved homologous de-epidermalised dermis of complex traumas to the lower limbs with loss of substance, and bones and tendons exposure. *J Plast Reconstr Asht Surg* 2008; 61(12): 1507-11.
- [120] Bhattacharyya T, Mehta P, Smith M, Pomahac B. Routine use of wound vacuum-assisted closure does not allow coverage delay for open tibia fractures. *Plast Reconstr Surg* 2008; 121(4): 1263-6.
- [121] Liu DS, Sofiadllis F, Ashton M, Macgill K, Webb A. Early soft tissue coverage and negative pressure wound therapy optimises patient outcomes in lower limb trauma. *Injury* 2012; 43(6): 772-8.
- [122] Hou Z, Irgit K, Strohecker KA, et al. Delayed flap reconstruction with vacuum-assisted closure management of the open IIIB tibial fracture. *J Trauma* 2011; 71(6): 1705-8.
- [123] Roberts CS, Pape HC, Jones AL, Malkani AL, Rodriguez JL, Giannoudis PV. Damage control orthopaedics: Evolving concepts in the treatment of patients who have sustained orthopaedic trauma. *Instr Course Lect* 2005; 54: 447-62.
- [124] Tuttle MS, Smith WR, Williams AE, et al. Safety and efficacy of damage control external fixation versus early definitive stabilization for femoral shaft fractures in the multiple-injured patient. *J Trauma* 2009; 67(3): 602-5.
- [125] Taeger G, Ruchholtz S, Waydhas C, Lewan U, Schmidt B, Nast-Kolb D. Damage control orthopedics in patients with multiple injuries is effective, time saving, and safe. *J Trauma* 2005; 59(2): 409-16.
- [126] Harwood PJ, Giannoudis PV, Probst C, Krettek C, Pape HC. The risk of local infective complications after damage control procedures for femoral shaft fracture. *J Orthop Trauma* 2006; 20(3): 181-9.
- [127] Karanas YL, Nigriny J, Chang J. The timing of microsurgical reconstruction in lower extremity trauma. *Microsurgery* 2008; 28(8): 632-4.
- [128] Gopal S, Giannoudis PV, Murray A, Matthews SJ, Smith RM. The functional outcome of severe, open tibial fractures managed with early fixation and flap coverage. *J Bone Joint Surg Br* 2004; 86(6): 861-7.
- [129] Li L, Yang LQ, Wang H. The effect of soft tissue repairing at different times on the union of type III b tibial fracture. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi* 2000; 14(6): 355-7.
- [130] DeLong WG Jr, Born CT, Wei SY, Petrik ME, Ponzio R, Schwab CW. Aggressive treatment of 119 open fracture wounds. *J Trauma* 1999; 46(6): 1049-54.
- [131] Yang W, Wu X, Yang D, Jiang C. Free tissue transplantation from amputated limbs for covering raw surface of stumps. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi* 2012; 26(2): 215-8.
- [132] White CB, Turner NS, Lee GC, Haidukewych GJ. Open ankle fractures in patients with diabetes mellitus. *Clin Orthop Relat Res* 2003; (414): 37-44.
- [133] Sorensen LT. Wound healing and infection in surgery: the pathophysiological impact of smoking, smoking cessation, and nicotine replacement therapy: a systematic review. *Ann Surg* 2012; 255(6): 1069-79.
- [134] Hoogendoorn JM, Simmermacher RK, Schellekens PP, van der Werken C. Adverse effects if smoking on healing of bones and soft tissues. *Unfallchirurg* 2002; 105(1): 76-81.
- [135] Harrison WJ. Open tibia fractures in HIV positive patients. *Malawi Med J* 2009; 21(4): 174-5.