

LE MATEMATICHE  
Vol. LII (1997) – Fasc. I, pp. 25–39

## ON FRACTIONAL DEVIATION OPERATORS

CARLOS C. PEÑA

The so called *fractional deviation operators* are introduced. This class of integral transforms appears naturally from the study of iteration of fractional integrals of Riemann-Liouville type. Since B. Ross' formulation on fractional iteration process, among other problems selected by T. Osler [5] toward 1974, several authors have been working on this subject. In particular, are worth mentioning contributions of B. Rubin [7] that allowed an intrinsic connection between fractional integrals with different limits of integration (Love's question, see [5] also) and the corresponding Ross' problem for Chen fractional integrals, handled by A. Nahushev [4] and M. Salahitdinov, with broad applications to non local boundary value problems. In this article we consider *deviation operators* as integral transforms, their connection with operators of Rubin type and mapping properties between classical and weighted Lebesgue spaces.

### 1. Preliminaries.

**Definition 1.** Let  $c, x \in \mathbb{R}$ ,  $c \leq x$ ,  $f \in L_{loc}^1 [c, +\infty]$ ,  $\gamma \in \mathbb{R}^+$ . By the Riemann-Liouville transform of  $f$  of order  $\gamma$  and lower limit of integration  $c$  we mean the function  $I_{c+}^\gamma f$  given as

$$I_{c+}^\gamma f(x) = \int_c^x \frac{(x-t)^{\gamma-1}}{\Gamma(\gamma)} f(t) dt.$$

---

Entrato in Redazione il 12 novembre 1996.

1997 AMS Subject Classification: 26A33.

**Proposition 1.** <sup>(1)</sup> *With the above notation:*

a) *The R - L transform is well defined and*

$$I_{c+}^{\gamma} : L_{loc}^1 [c, +\infty] \rightarrow L_{loc}^1 [c, +\infty].$$

b) *For  $\gamma_1, \gamma_2 \in R^+$  the following semigroup property holds*

$$I_{c+}^{\gamma_1} \circ I_{c+}^{\gamma_2} = I_{c+}^{\gamma_1 + \gamma_2}.$$

**Definition 2.** *Let  $a, b, \gamma$  be real numbers,  $a < b, \gamma \in R^+, f \in L^1 [a, b]$ . We also denote*

$$\begin{aligned} {}_{a,b}I_{+}^{\gamma} f(y) &= \int_a^b \frac{(y-t)^{\gamma-1}}{\Gamma(\gamma)} f(t) dt, \quad y > b, \\ {}_{a,b}I_{-}^{\gamma} f(x) &= \int_a^b \frac{(t-x)^{\gamma-1}}{\Gamma(\gamma)} f(t) dt, \quad x < a. \end{aligned}$$

**Definition 3.** *Let  $a < c < b, \alpha, \beta \in R^+$ . By the fractional deviation operator  ${}_{a,b}D_{c+}^{\alpha,\beta}$  of R - L class  $(a, b, \alpha, \beta)$  and lower limit  $c$  we mean the map*

$${}_{a,b}D_{c+}^{\alpha,\beta} : L_{loc}^1 [a, +\infty] \rightarrow L_{loc}^1 [b, +\infty],$$

$${}_{a,b}D_{c+}^{\alpha,\beta} \doteq I_{b+}^{\beta} \circ I_{a+}^{\alpha} - I_{c+}^{\alpha+\beta}.$$

**Definition 4.** *Let  $\alpha, \beta \in R^+, x^1 < x^2 < x^3$ . We'll write*

$$\begin{aligned} F_{\alpha,\beta}(x^1, x^2, x^3) &= F_{\alpha,\beta}^{x^2}(x^1, x^3) \\ &\doteq \int_{x^2}^{x^3} (x - x^1)^{\alpha-1} (x^3 - x)^{\beta-1} dx. \end{aligned}$$

---

<sup>(1)</sup> See [3].

It is easy to see that  $F_{\alpha,\beta}$  is well defined and in terms of the incomplete beta function

$$B_{\alpha,\beta}(x) = \int_x^{+\infty} s^{\alpha-1}(1+s)^{-\alpha-\beta} ds,$$

where  $x > 0$ , the following formulae hold:

$$(1) \quad B_{\alpha,\beta}(0+) = Be(\alpha, \beta),$$

$$(2) \quad B_{\alpha,\beta}(x) + B_{\beta,\alpha}(x^{-1}) = Be(\alpha, \beta),$$

$$(3) \quad F_{\alpha,\beta}(x^1, x^2, x^3) = (x^3 - x^1)^{\alpha+\beta-1} B_{\alpha,\beta}\left(\frac{x^2 - x^1}{x^3 - x^2}\right),$$

$$(4) \quad F_{\alpha,\beta}(px^1, px^2, px^3) = p^{\alpha+\beta-1} F_{\alpha,\beta}(x^1, x^2, x^3), \text{ } p\text{-positive,}$$

$$(5) \quad F_{\alpha,\beta}(x^1, x^2, x^3) = F_{\alpha,\beta}^0(x^1 - x^2, x^3 - x^2).$$

**Definition 5.** Let  $a < b$ ,  $\alpha, \beta \in R^+$ . We introduce the operators

$${}_{a,b}E_+^{\alpha,\beta} : L^1[a, b] \rightarrow L_{loc}^1[b, +\infty],$$

$${}_{a,b}E_+^{\alpha,\beta} f(y) \doteq \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b F_{\alpha,\beta}^b(u, y) f(u) du, \quad f \in L^1[a, b], \quad b < y,$$

and

$${}_{a,b}E_-^{\alpha,\beta} : L^1[a, b] \rightarrow L_{loc}^1[-\infty, a],$$

$${}_{a,b}E_-^{\alpha,\beta} f(x) \doteq \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b F_{\alpha,\beta}^a(x, v) f(v) dv, \quad f \in L^1[a, b], \quad x < a.$$

**Definition 6.** Let  $-\infty \leq a < b < +\infty$ ,  $\gamma \in R^+$ ,  $f \in L^1[a, b]$ . By a Rubin type transform  $R_{a,b}^\gamma f$  of  $f$  we mean the function given as

$$R_{a,b}^\gamma f(x) = \frac{\sin(\pi\gamma)}{\pi} \int_a^b \left(\frac{b-t}{x-b}\right)^\gamma \frac{f(t)}{x-t} dt, \quad x > b.$$

Finally, the author wishes to thank the referee for his helpful suggestions.

## 2. On deviation operators.

**Theorem 1.** *Let  $a < c < b$ ,  $\alpha, \beta, \gamma \in \mathbb{R}^+$ ,  $f \in L^1_{loc}[a, +\infty]$ . Then*

$$(6) \quad {}_{a,b}D_{c+}^{\alpha,\beta} f = {}_{a,b}E_+^{\alpha,\beta} f - {}_{c,b}I_+^{\alpha+\beta} f.$$

$$(7) \quad {}_{a,b}E_+^{\alpha,\beta} f + \left( {}_{-b,-a}E_-^{\beta,\alpha} f^\sim \right)^\sim = {}_{a,b}I_+^{\alpha+\beta} f,$$

where  $f^\sim(x) = f(-x)$  and  $x \leq a$ .

$$(8) \quad ({}_{a,b}I_-^\gamma f)^\sim = {}_{-b,-a}I_+^\gamma f^\sim.$$

$$(9) \quad \left( {}_{a,b}D_{c+}^{\alpha,\beta} f \right)^\sim = {}_{-c,-a}I_-^{\alpha+\beta} f^\sim - {}_{-b,-a}E_-^{\beta,\alpha} f^\sim.$$

*In general, the above identities must be interpreted in the almost everywhere sense.*

*Proof.* With fixed  $d > b \geq e \geq a$  we obtain

$$(10) \quad \|I_{e+}^\gamma\|_{L^1[b,d]} \leq \frac{(d-e)^\gamma}{\Gamma(\gamma+1)} \|f\|_{L^1[e,d]}$$

and therefore

$$(11) \quad \begin{aligned} \|{}_{a,b}D_{c+}^{\alpha,\beta} f\|_{L^1[b,d]} &\leq \\ &\leq \left[ \frac{(d-a)^\alpha}{\Gamma(\alpha+1)} \cdot \frac{(d-b)^\beta}{\Gamma(\beta+1)} + \frac{(d-c)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \right] \|f\|_{L^1[a,d]}. \end{aligned}$$

On the other hand:

$$(12) \quad \|{}_{c,b}I_+^\gamma f\|_{L^1[b,d]} \leq \frac{(d-c)^\gamma}{\Gamma(\gamma+1)} \|f\|_{L^1[c,b]},$$

$$(13) \quad \|{}_{a,b}E_+^{\alpha,\beta} f\|_{L^1[b,d]} \leq \frac{(d-a)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \|f\|_{L^1[a,b]}.$$

From the above relations we deduce that  ${}_{a,b}D_{c+}^{\alpha,\beta}$  and  ${}_{a,b}E_+^{\alpha,\beta} - {}_{c,b}I_+^{\alpha+\beta}$  maps  $L^1[a, d]$  linearly and continuously into  $L^1[b, d]$  for each  $d > b$ . It is

now expedient to observe that  $I_{a+}^\alpha g \in C[a, d]$  whenever  $g \in C[a, d]$  ([3], Ch. 1, Corollary 2). By identifying  $I_{a+}^\alpha g$  with its restriction over  $[b, d]$  we obtain  $I_{b+}^\beta(I_{a+}^\alpha g) \in C[b, d]$ . Analogously  $I_{c+}^{\alpha+\beta} g \in C[b, d]$  by identifying  $g$  with its restriction to  $[c, d]$ . Now  ${}_{a,b}D_{c+}^{\alpha,\beta} g \in C[b, d]$  and from a direct application of Fubini's Theorem we obtain (6). In the general case, given  $f \in L^1[a, d]$  and a positive number  $\zeta$  we may write  $f = g + h$ ,  $g \in C[a, d]$  and  $\|h\|_{L^1[a,d]} \leq \zeta$ . Using (11) – (13) we have

$$\begin{aligned}
& \|{}_{a,b}D_{c+}^{\alpha,\beta} f - ({}_{a,b}E_+^{\alpha,\beta} - {}_{c,b}I_+^{\alpha+\beta}) f\|_{L^1[b,d]} = \\
& = \|{}_{a,b}D_{c+}^{\alpha,\beta} h - ({}_{a,b}E_+^{\alpha,\beta} - {}_{c,b}I_+^{\alpha+\beta}) h\|_{L^1[b,d]} \\
& \leq \|{}_{a,b}D_{c+}^{\alpha,\beta} h\|_{L^1[b,d]} + \|{}_{a,b}E_+^{\alpha,\beta} h\|_{L^1[b,d]} + \|{}_{c,b}I_+^{\alpha+\beta} h\|_{L^1[b,d]} \\
& \leq \left[ \frac{(d-a)^\alpha}{\Gamma(\alpha+1)} \cdot \frac{(d-b)^\beta}{\Gamma(\beta+1)} + \frac{(d-c)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \right] \|h\|_{L^1[a,d]} + \\
& \quad + \frac{(d-a)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \|h\|_{L^1[a,b]} + \frac{(d-c)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \|h\|_{L^1[c,b]} \\
& \leq \left[ \frac{(d-a)^\alpha}{\Gamma(\alpha+1)} \cdot \frac{(d-b)^\beta}{\Gamma(\beta+1)} + \frac{(d-c)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} + \right. \\
& \quad \left. + \frac{(d-a)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} + \frac{(d-c)^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \right] \zeta.
\end{aligned}$$

Since  $\zeta$  was arbitrary equality (6) became valid in the  $\|\cdot\|_{L^1[b,d]}$ -norm for each  $d > b$  and hence it is also valid in the a.e. sense. The other identities may be proved in a similar way.  $\square$

**Theorem 2.** Let  $a, b, \gamma \in \mathbb{R}$ ,  $a < b$ ,  $0 < \gamma < 1$ ,  $f \in L^1[a, b]$ . Then

$$(14) \quad I_{b+}^\gamma R_{a,b}^\gamma f(y) = {}_{a,b}I_+^\gamma f(y) \quad a.e. y > b.$$

*Proof.* In general, Rubin type transforms satisfy the following translation rule

$$(15) \quad \tau_{-b} R_{a-b,0}^\gamma \tau_b = R_{a,b}^\gamma,$$

and so we shall consider  $R_{c,0}^\gamma$  with  $-\infty \leq c < 0$ . If we write

$$(16) \quad \kappa(t, x) = \left(\frac{t}{x}\right)^\gamma \frac{1}{x+t}$$

with both  $x$  and  $t$  positive, then  $\kappa$  is a measurable homogeneous function of degree  $-1$ , and if  $1 < p < \infty$ :

$$(17) \quad \int_0^{+\infty} |\kappa(t, 1)| t^{-1/p} dt = \int_0^{+\infty} |\kappa(1, x)| x^{-1/p'} dx = \chi,$$

where  $\chi$  would be finite if  $\gamma < 1/p$ . In this case, from the Hardy - Littlewood - Pólya Theorem on boundedness of homogeneous operators [2], Theorem 319, the Rubin transform

$$(18) \quad R_{-\infty, 0}^\gamma f(x) = \frac{\sin(\pi\gamma)}{\pi} \int_0^{+\infty} \left(\frac{t}{x}\right)^\gamma \frac{f(-t)}{x+t} dt, \quad x\text{-positive},$$

becomes continuous between  $L^p[-\infty, 0]$  and  $L^p[0, +\infty]$  with

$$(19) \quad \|R_{-\infty, 0}^\gamma f\|_{L^p[0, +\infty]} \leq \frac{\sin(\pi\gamma)}{\sin(\pi(\gamma + 1/p'))} \|f\|_{L^p[-\infty, 0]}.$$

It is now easy to see that if  $1 \leq p < 1/\gamma$  then  $R_{c, 0}^\gamma$ ,  $c$ -negative, maps linear and continuously  $L^p[c, 0]$  on  $L^p[0, +\infty]$  with

$$(20) \quad \|R_{c, 0}^\gamma f\|_{L^p[0, +\infty]} \leq \frac{\sin(\pi\gamma)}{\sin(\pi(\gamma + 1/p'))} \|f\|_{L^p[c, 0]}.$$

Moreover, under this conditions  $R_{c, 0}^\gamma f(x)$  will be defined and finite for every  $x$ -positive with the limit  $c$  finite or infinite.

Now for a given  $f \in L^p[c, 0]$  we write

$$\begin{aligned} I_{0+}^\gamma R_{c, 0}^\gamma f(y) &= \int_0^y \frac{(y-x)^{\gamma-1}}{\Gamma(\gamma)} \left[ \frac{\sin(\pi\gamma)}{\pi} \int_c^0 \left(\frac{|t|}{x}\right)^\gamma \frac{f(t)}{x+|t|} dt \right] dx \\ &= \frac{\sin(\pi\gamma)}{\pi\Gamma(\gamma)} \int_c^0 f(t) |t|^\gamma \left[ \int_0^y \frac{(y-x)^{\gamma-1} x^{-\gamma}}{x+|t|} dx \right] dt. \end{aligned}$$

The application of Fubini's Theorem is justified because the double integral is still absolutely convergent if  $c = -\infty$  and fractional integrals of order  $\gamma$  on the whole real axis are defined for  $L^p$ -functions if  $1 \leq p < 1/\gamma$ . Finally the inner integral may be evaluated after the change of variable  $z = \frac{-t(y-x)}{(x-t)y}$ , and so we obtain (14).  $\square$

**Theorem 3.** Let  $a < c < b$ ,  $\alpha, \beta \in R^+$ . The following formulae hold

$$(21) \quad {}_{a,b}I_+^{\alpha+\beta} = I_{b+}^\beta \circ {}_{a,b}I_+^\alpha + {}_{a,b}I_+^\beta \circ I_{a+}^\alpha,$$

$$(22) \quad {}_{a,b}E_+^{\alpha,\beta} = I_{b+}^\beta \circ {}_{a,b}I_+^\alpha,$$

$$(23) \quad {}_{a,b}D_{c+}^{\alpha,\beta} = I_{b+}^\beta \circ {}_{a,c}I_+^\alpha - {}_{c,b}I_+^\beta \circ I_{c+}^\alpha,$$

$$(24) \quad {}_{a,b}D_{a+}^{\alpha,\beta} = -{}_{a,b}I_+^\beta \circ I_{a+}^\alpha,$$

$$(25) \quad {}_{a,b}D_{b+}^{\alpha,\beta} = I_{b+}^\beta \circ {}_{a,b}I_+^\alpha,$$

$$(26) \quad {}_{a,b}D_{b+}^{\alpha,\beta} - {}_{a,b}D_{c+}^{\alpha,\beta} = {}_{c,b}I_+^\beta \circ I_{c+}^\alpha.$$

**Corollary 1.** If  $a < c < b$ ,  $\alpha, \beta \in R$  and  $0 < \alpha < \alpha + \beta < 1$ , then

$${}_{a,b}D_{c+}^{\alpha,\beta} = I_{b+}^{\alpha+\beta} \left( R_{a,b}^\alpha - R_{c,b}^{\alpha+\beta} \right).$$

### 3. On some boundedness conditions.

**Proposition 2.** Let  $\alpha, \beta \in R^+$ . There exist positive constants  $c_1, c_2$  such that for every positive  $x$  we have the inequality

$$(27) \quad c_1(1+x)^{-\beta} \leq B_{\alpha,\beta}(x) \leq c_2(1+x)^{-\beta}.$$

Moreover, we may take  $c_1 = \min \{ \beta^{-1}, Be(\alpha, \beta) \}$ ,  $c_2 = \max \{ \beta^{-1}, Be(\alpha, \beta) \}$ .

*Proof.* Given a positive  $x$  and assuming  $\alpha \geq 1$  we have

$$(28) \quad B_{\alpha,\beta}(x) \leq \frac{(1+x)^{-\beta}}{\beta}.$$

In particular, if  $x \rightarrow 0+$  in (28) there result  $Be(\alpha, \beta) \leq 1/\beta$  and on one hand (27) holds. If  $0 < \alpha < 1$  we write

$$(29) \quad B_{\alpha,\beta}(x) \geq \frac{(1+x)^{-\beta}}{\beta}$$

and making  $x \rightarrow 0+$  in (29) we obtain  $Be(\alpha, \beta) \geq 1/\beta$ . We now introduce the functions of the non negative real variable  $x$

$$f(x) = (1+x)^\beta B_{\alpha,\beta}(x)$$

and

$$g(x) = B_{\alpha,\beta}(x) - \frac{x^{\alpha-1}(1+x)^{1-\alpha-\beta}}{\beta}.$$

In particular,  $f(x) \equiv 1/\beta$  if  $\alpha = 1$  and our claim follows. On the other hand we may write

$$\begin{aligned} f'(x) &= \beta(1+x)^{\beta-1}g(x), \\ g'(x) &= \frac{1-\alpha}{\beta}x^{\alpha-2}(1+x)^{-\alpha-\beta}, \end{aligned}$$

i.e.  $g$  is monotone increasing or decreasing according as  $0 < \alpha \leq 1$  or  $\alpha \geq 1$  respectively. But  $g(+\infty) = 0$ , i.e.  $f$  becomes a monotone decreasing function in the first case and a monotone increasing one in the second. Moreover,  $f(0) = Be(\alpha, \beta)$  and  $f(+\infty) = 1/\beta$  and hence both estimates follow.  $\square$

**Remark 1.** We'll consider formally the following expressions

$$\begin{aligned} [a] \quad & \sup_{\kappa>0} \kappa^{1-\alpha} \left( \int_a^c \frac{w_1(x)^{-p'/p} dx}{(c-x+\kappa)^{(1-\alpha)p'}} \right)^{\frac{1}{p'}} \left( \int_b^{+\infty} \frac{(y-b)^{\beta q} w_2(y) dy}{(y-c+\kappa)^{(1-\alpha)q}} \right)^{\frac{1}{q}}, \\ [b] \quad & \sup_{\kappa>0} \kappa^{1-\beta} \left( \int_c^b \frac{(b-x)^{\alpha p'} w_1(x)^{-p'/p} dx}{(b-x+\kappa)^{(1-\beta)p'}} \right)^{\frac{1}{p'}} \left( \int_b^{+\infty} \frac{w_2(y) dy}{(y-b+\kappa)^{(1-\beta)q}} \right)^{\frac{1}{q}}, \\ [c] \quad & \sup_{\kappa>0} \left( \int_c^b \left( \frac{b-x+\kappa}{b-x} \right)^{(1-\beta)p'} (b-x)^{\alpha p'} w_1(x)^{-p'/p} dx \right)^{\frac{1}{p'}} \\ & \quad \cdot \left( \int_b^{+\infty} \frac{w_2(y) dy}{(y-b+\kappa)^{(1-\beta)q}} \right)^{\frac{1}{q}}, \\ [d] \quad & \sup_{\kappa>0} \left( \int_a^c \left( \frac{c-x+\kappa}{c-x} \right)^{(1-\alpha)p'} w_1(x)^{-p'/p} dx \right)^{\frac{1}{p'}} \\ & \quad \cdot \left( \int_b^{+\infty} \frac{(y-b)^{\beta q} w_2(y) dy}{(y-c+\kappa)^{(1-\alpha)q}} \right)^{\frac{1}{q}}, \\ [e] \quad & \left[ \int_a^c \left( \int_b^{+\infty} \frac{(y-b)^{\beta q} w_2(y) dy}{(y-x)^{(1-\alpha)q}} \right)^{r/q} \right. \\ & \quad \cdot \left. \left( \int_a^c \frac{w_1(z)^{1-p'} dz}{\left(1 + \frac{c-z}{c-x}\right)^{(1-\alpha)p'}} \right)^{r/q'} w_1(x)^{1-p'} dx \right]^{1/r}, \\ [f] \quad & \left[ \int_c^b \left( \int_b^{+\infty} \frac{w_2(y) dy}{(y-x)^{(1-\beta)q}} \right)^{r/q} \right. \end{aligned}$$



$$\left( \int_c^b \frac{(b-z)^{\alpha p'} w_1(z)^{1-p'} dz}{\left(1 + \frac{b-z}{b-x}\right)^{(1-\beta)p'}} \right)^{r/q'} (b-x)^{\alpha p'} w_1(x)^{1-p'} dx \Big]^{1/r}.$$

**Remark 2.** Here and throughout the paper, when any of  $p$ ,  $q$ , or  $p'$  is  $\infty$  integrals such as those  $[a] - [f]$  have the current interpretations and as usual  $w_1(x)^{-p'/p} = (w_1(x)^{-1})^{p'/p}$ . Thus for example

$$\left( \int_a^c \frac{w_1(x)^{-p'/p} dx}{(c-x+\kappa)^{(1-\alpha)p'}} \right)^{\frac{1}{p'}} = \operatorname{ess\,sup}_{a \leq x \leq c} \frac{w_1(x)^{-1} dx}{(c-x+\kappa)^{1-\alpha}}, \quad (p = 1)$$

and

$$w_1(x)^{-p'/p} = \begin{cases} 1 & \text{if } 0 < w_1(x) < \infty \\ \infty & \text{if } w_1(x) = 0 \\ 0 & \text{if } w_1(x) = \infty \end{cases} \quad (p = \infty).$$

Moreover, products of the form  $0 \cdot \infty$  are taken to be zero.

**Theorem 4.** Let  $-\infty < a < c < b < +\infty$ ,  $\alpha, \beta \in R^+$ ,  $w_1, w_2$  two weights defined on  $[a, b]$  and  $[b, +\infty]$  respectively and  $1 \leq p, q \leq +\infty$ . Let:

- (i)  $0 < \alpha, \beta \leq 1, 1 \leq p \leq q \leq +\infty$ ;
- (ii)  $0 < \alpha \leq 1 < \beta, 1 \leq p \leq q \leq +\infty, 1 < q$ ;
- (iii)  $0 < \beta \leq 1 < \alpha, 1 \leq p \leq q \leq +\infty, 1 < q$ ;
- (iv)  $1 < \alpha, 1 < \beta, 1 \leq p \leq +\infty, 1 < q$ ;
- (v)  $0 < \alpha, \beta < 1, 1 < q < p < +\infty$ ;
- (vi)  $0 < \beta < 1 \leq \alpha, 1 < q < p < +\infty$ ;
- (vii)  $0 < \alpha < 1 \leq \beta, 1 < q < p < +\infty$ ;

the general linear deviation operator  ${}_{a,b}D_{c+}^{\alpha,\beta}$  will be bounded among the weighted Lebesgue spaces  $L_{w_1(x)dx}^p[a, b]$  and  $L_{w_2(y)dy}^q[b, +\infty]$  <sup>(2)</sup> if and only if, with the notation of the above remark, the following respective conditions hold:

- In (i), the expressions  $[a]$  and  $[b]$  are finite.
- In (ii), the expressions  $[a]$  and  $[c]$  are finite.
- In (iii), the expressions  $[b]$  and  $[d]$  are finite.
- In (iv), the expressions  $[c]$  and  $[d]$  are finite.
- In (v), with  $r = 1/q - 1/p$ ,  $[e]$  and  $[f]$  are finite.
- In (vi), with  $r = 1/q - 1/p$ ,  $[d]$  and  $[f]$  are finite.
- In (vii), with  $r = 1/q - 1/p$ ,  $[c]$  and  $[e]$  are finite.

---

<sup>(2)</sup> As usual, in this case we write  ${}_{a,b}D_{c+}^{\alpha,\beta} \in [L_{w_1(x)dx}^p[a, b], L_{w_2(y)dy}^q[b, +\infty]]$ .

*Proof.* Given  $f \in L^1_{loc}[a, +\infty]$ , from (2) and (6) we have

$$\begin{aligned} {}_{a,b}D_{c+}^{\alpha,\beta} f(y) &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \left[ \int_a^c (y-x)^{\alpha+\beta-1} B_{\alpha,\beta} \left( \frac{b-x}{y-b} \right) f(x) dx - \right. \\ &\quad \left. - \int_c^b (y-x)^{\alpha+\beta-1} B_{\beta,\alpha} \left( \frac{y-b}{b-x} \right) f(x) dx \right] \end{aligned}$$

and hence we write

$$(30) \quad {}_{a,b}D_{c+}^{\alpha,\beta} f = {}_{a,b}D_{1;c+}^{\alpha,\beta} f + {}_{a,b}D_{2;c+}^{\alpha,\beta} f,$$

with

$${}_{a,b}D_{1;c+}^{\alpha,\beta} f(y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^c (y-x)^{\alpha+\beta-1} B_{\alpha,\beta} \left( \frac{b-x}{y-b} \right) f(x) dx$$

and

$${}_{a,b}D_{2;c+}^{\alpha,\beta} f(y) = -\frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_c^b (y-x)^{\alpha+\beta-1} B_{\beta,\alpha} \left( \frac{y-b}{b-x} \right) f(x) dx.$$

Since

$$(31) \quad {}_{a,b}D_{1;c+}^{\alpha,\beta} = {}_{a,b}D_{c+}^{\alpha,\beta} \chi_{[a,c]} \quad \text{and} \quad {}_{a,b}D_{2;c+}^{\alpha,\beta} = -{}_{a,b}D_{c+}^{\alpha,\beta} \chi_{[c,b]}$$

it is immediate that  ${}_{a,b}D_{c+}^{\alpha,\beta} \in \left[ L^p_{w_1(x)dx}[a, b], L^q_{w_2(y)dy}[b, +\infty] \right]$  iff

$$(32) \quad \begin{aligned} {}_{a,b}D_{1;c+}^{\alpha,\beta} &\in \left[ L^p_{w_1(x)dx}[a, c], L^q_{w_2(y)dy}[b, +\infty] \right] \quad \text{and} \\ {}_{a,b}D_{2;c+}^{\alpha,\beta} &\in \left[ L^p_{w_1(x)dx}[c, b], L^q_{w_2(y)dy}[b, +\infty] \right]. \end{aligned}$$

But by Prop. 1 we deduce that (32) will hold iff

$$(33) \quad \begin{aligned} {}_{a,c}I_+^{\alpha} &\in \left[ L^p_{w_1(x)dx}[a, c], L^q_{(y-b)^{\beta q} w_2(y)dy}[b, +\infty] \right] \quad \text{and} \\ {}_{c,b}I_+^{\beta} &\in \left[ L^p_{(b-x)^{-\alpha p} w_1(x)dx}[c, b], L^q_{w_2(y)dy}[b, +\infty] \right]. \end{aligned}$$

Moreover the following inequalities hold

$$(34) \quad c_1 \frac{(y-b)^\beta}{\Gamma(\beta)} I_{a,c}^\alpha f(y) \leq_{a,b} D_{1;c+}^{\alpha,\beta} f(y) \leq c_2 \frac{(y-b)^\beta}{\Gamma(\beta)} I_{a,c}^\alpha f(y),$$

$$(35) \quad \left| {}_{a,b}D_{2;c+}^{\alpha,\beta} f(y) \right| \leq c_2 {}_{c,b}I_+^\beta \left[ \frac{(b-y)^\alpha}{\Gamma(\alpha)} |f(y)| \right]$$

and

$$(36) \quad \left| {}_{c,b}I_+^\beta \left[ \frac{(b-y)^\alpha}{\Gamma(\alpha)} f(y) \right] \right| \leq_{c,b} I_+^\beta \left[ \frac{(b-y)^\alpha}{\Gamma(\alpha)} |f(y)| \right] \leq \frac{1}{c_1} \left| {}_{a,b}D_{2;c+}^{\alpha,\beta} f(y) \right|$$

where by  $c_1$  and  $c_2$  we denote the constants determined in Prop. 2.

In particular we may write

$$(37) \quad {}_{a,c}I_+^\alpha = \frac{1}{\Gamma(\alpha)} \tau_{-c} S_{1-\alpha} (\tau_c)^\sim,$$

where  $S_{1-\alpha}$  denote the usual Stieltjes transform. From the corresponding representations of  ${}_{a,c}I_+^\alpha$  and  ${}_{c,b}I_+^\beta$  as composition of maps

$$(38) \quad \begin{array}{ccc} L_{w_1(x)dx}^p[a, c] & & L_{(b-x)^{-\alpha p} w_1(x)dx}^p[c, b] \\ \downarrow (\tau_c)^\sim & & \downarrow (\tau_b)^\sim \\ L_{w_1(c-x)dx}^p[0, c-a] & & L_{x^{-\alpha p} w_1(b-x)dx}^p[0, b-c] \\ \downarrow S_{1-\alpha} & \text{and} & \downarrow S_{1-\beta} \\ L_{(y-b+c)^{\beta q} w_2(y+c)dy}^q[b-c, +\infty] & & L_{w_2(y+b)dy}^q[0, +\infty] \\ \downarrow \tau_{-c} & & \downarrow \tau_{-b} \\ L_{(y-b)^{\beta q} w_2(y)dy}^q[b, +\infty] & & L_{w_2(y)dy}^q[b, +\infty] \end{array}$$

it is clear that (33) will hold iff

$$(39) \quad S_{1-\alpha} \in \left[ L_{w_1(c-x)dx}^p[0, c-a], L_{(y-b+c)^{\beta q} w_2(y+c)dy}^q[b-c, +\infty] \right]$$

and

$$(40) \quad S_{1-\beta} \in \left[ L_{x^{-\alpha p} w_1(b-x)dx}^p[0, b-c], L_{w_2(y+b)dy}^q[0, +\infty] \right].$$

We've therefore reduced our matter to the study of boundedness conditions of the general Stieltjes transform. Therefore known results of K. Andersen [1] and G. Sinnamon [8] in this direction go on.

**Remark 3.** V. D. Stepanov [9] obtained necessary and sufficient conditions for the boundedness from weighted Lebesgue real spaces of Volterra convolution operators of the form

$$Kf(x) = \int_0^x k(x-y) f(y) dy,$$

where  $k(x)$  is a non negative non decreasing kernel satisfying the inequality  $k(x+y) \leq D(k(x) + k(y))$  for all  $x, y \in \mathbb{R}^+$ . In particular, Stepanov research will allow us to relate  ${}_{a,b}D_{c+}^{\alpha,\beta}$  with  $I_{c+}^\gamma$  when  $\gamma \geq 1$ ,  $\gamma = \alpha + \beta$ , i.e. precisely when the corresponding Riemann - Liouville kernel  $k_\gamma(x) = x^{\gamma-1}/\Gamma(\gamma)$  satisfies Stepanov conditions. With our notation we have

$${}_{a,b}I_{c+}^\gamma = \tau_{-a} \circ I_{0+}^\gamma \circ \tau_a \circ \chi_{[a,b]}$$

and it is of interest for us to consider

$$\begin{array}{ccc} L_{w_1(x)dx}^p[a, b] & \xrightarrow{{}_{a,b}I_{c+}^\gamma} & L_{w_2(y)dy}^q[b, +\infty]. \\ \downarrow \tau_a & & \uparrow \tau_{-a} \\ L_{\tau_a w_1(x)dx}^p[0, b-a] & \xrightarrow{I_{0+}^\gamma} & L_{\tau_a w_2(y)dy}^q[b-a, +\infty]. \end{array}$$

Following Stepanov research, for  $t > 0$ , we must consider the numbers

$$(41) \quad A_0(t) = \left[ \int_a^{\min\{t+a,b\}} w_1(x)^{-p'/p} dx \right]^{1/p'} \cdot \left[ \int_{\max\{t+a,b\}}^\infty (y-a-t)^{q(\gamma-1)} w_2(y) dy \right]^{1/q},$$

$$(42) \quad A_1(t) = \left[ \int_a^{\min\{t+a,b\}} (t+a-x)^{p'(\gamma-1)} w_1(x)^{-p'/p} dx \right]^{1/p'} \cdot \left[ \int_{\max\{t+a,b\}}^\infty w_2(y) dy \right]^{1/q},$$

and

$$(43) \quad A_0 = \sup_{t>0} A_0(t), \quad A_1 = \sup_{t>0} A_1(t), \quad A = \max\{A_0, A_1\}.$$

Now, for the boundedness of the operator

$$L_{w_1(x)dx}^p[a, b] \xrightarrow{{}_{a,b}I_{c+}^\gamma} L_{w_2(y)dy}^q[b, +\infty]$$

it is necessary and sufficient that the number  $A$  be finite (see [9], Th. 1).

**Remark 4.** Following Remark 3, we may prove that, in general, our deviation operators are small compared to the operators whose deviation they measure. For instance, let us consider the case  $0 \leq \alpha, \beta \leq 1$ ,  $1 \leq \gamma$ ,  $1 < p \leq q < \infty$ . We assume that  ${}_{a,b}I_+^{\alpha,\beta} \in \left[ L_{w_1(x)dx}^p[a, b], L_{w_2(y)dy}^q[b, +\infty] \right]$ , and for a fixed  $\kappa > 0$  we write

$$\begin{aligned} A &\geq A_0 \\ &\geq A_0(b-a) \\ &= \left[ \int_a^b w_1(x)^{-p'/p} dx \right]^{1/p'} \left[ \int_b^\infty (y-b)^{q(\gamma-1)} w_2(y) dy \right]^{1/q} \\ &\geq \kappa^{1-\alpha} \left[ \int_a^c \frac{w_1(x)^{-p'/p} dx}{(c-x+\kappa)^{(1-\alpha)p'}} \right]^{1/p'} \left[ \int_b^{+\infty} \frac{(y-b)^{\beta q} w_2(y) dy}{(y-c+\kappa)^{(1-\alpha)q}} \right]^{1/q} \end{aligned}$$

and we deduce that

$$\sup_{\kappa>0} \kappa^{1-\alpha} \left( \int_a^c \frac{w_1(x)^{-p'/p} dx}{(c-x+\kappa)^{(1-\alpha)p'}} \right)^{\frac{1}{p'}} \left( \int_b^{+\infty} \frac{(y-b)^{\beta q} w_2(y) dy}{(y-c+\kappa)^{(1-\alpha)q}} \right)^{\frac{1}{q}} < \infty.$$

Analogously

$$\begin{aligned} A &\geq A_1 \\ &\geq A_1(b-a) \\ &= \left[ \int_a^b (b-x)^{p'(\gamma-1)} w_1(x)^{-p'/p} dx \right]^{1/p'} \left[ \int_b^\infty w_2(y) dy \right]^{1/q} \\ &\geq \kappa^{1-\beta} \left[ \int_b^{+\infty} \frac{w_2(y) dy}{(y-b+\kappa)^{(1-\beta)q}} \right]^{1/q} \\ &\quad \cdot \left[ \int_c^b \frac{(b-x)^{\alpha p'}}{(b-x+\kappa)^{(1-\beta)p'}} w_1(x)^{-p'/p} dx \right]^{1/p'} \end{aligned}$$

and now

$$\begin{aligned} \sup_{\kappa>0} \kappa^{1-\beta} \left( \int_c^b \frac{(b-x)^{\alpha p'} w_1(x)^{-p'/p} dx}{(b-x+\kappa)^{(1-\beta)p'}} \right)^{\frac{1}{p'}} \\ \cdot \left( \int_b^{+\infty} \frac{w_2(y) dy}{(y-b+\kappa)^{(1-\beta)q}} \right)^{\frac{1}{q}} < \infty. \end{aligned}$$

From Theorem 4 we deduce that  ${}_{a,b}D_{c+}^{\alpha,\beta} \in \left[ L_{w_1(x)dx}^p[a, b], L_{w_2(y)dy}^q[b, +\infty] \right]$  if  ${}_{a,b}I_+^{\alpha,\beta} \in \left[ L_{w_1(x)dx}^p[a, b], L_{w_2(y)dy}^q[b, +\infty] \right]$ .

Nevertheless we may find weighted functions  $w_1, w_2$  for which  ${}_{a,b}D_{c+}^{\alpha,\beta} \in \left[ L_{w_1(x)dx}^p[a, b], L_{w_2(y)dy}^q[b, +\infty] \right]$  but  ${}_{a,b}I_+^{\alpha,\beta} \notin \left[ L_{w_1(x)dx}^p[a, b], L_{w_2(y)dy}^q[b, +\infty] \right]$ . For instance, for  $a \leq x \leq b \leq y$  we'll write

$$\begin{aligned} w_1(x) &= (b-x)^\sigma, \quad p/p' \leq \sigma < (\alpha + 1/p')p, \\ w_2(y) &= (y-b)^\lambda e^{-y}, \quad \lambda > -1 + (1-\beta)q. \end{aligned}$$

We observe that the number  $A_0(b-a) = +\infty$  in (41), because  $1 - \sigma p'/p \leq 0$ . Hence  $A = +\infty$  in (43) and by the Stepanov condition  ${}_{a,b}I_+^{\alpha,\beta}$  is not bounded. On the other hand,

$$\begin{aligned} \sup_{\kappa>0} \kappa^{1-\alpha} \left( \int_a^c \frac{w_1(x)^{-p'/p} dx}{(c-x+\kappa)^{(1-\alpha)p'}} \right)^{\frac{1}{p'}} \left( \int_b^{+\infty} \frac{(y-b)^{\beta q} w_2(y) dy}{(y-c+\kappa)^{(1-\alpha)q}} \right)^{\frac{1}{q}} &\leq \\ &\leq (b-c)^{\alpha-1} \left( \int_a^c (b-x)^{-\sigma p'/p} dx \right)^{\frac{1}{p'}} \left( \int_b^{+\infty} (y-b)^{\beta q + \lambda} e^{-y} dy \right)^{\frac{1}{q}} \end{aligned}$$

and

$$\begin{aligned} \sup_{\kappa>0} \kappa^{1-\beta} \left( \int_c^b \frac{(b-x)^{\alpha p'} w_1(x)^{-p'/p} dx}{(b-x+\kappa)^{(1-\beta)p'}} \right)^{\frac{1}{p'}} \left( \int_b^{+\infty} \frac{w_2(y) dy}{(y-b+\kappa)^{(1-\beta)q}} \right)^{\frac{1}{q}} &\leq \\ &\leq \left( \int_c^b (b-x)^{(\alpha-\sigma/p)p'} dx \right)^{\frac{1}{p'}} \left( \int_b^{+\infty} (y-b)^{\lambda+(\beta-1)q} e^{-y} dy \right)^{\frac{1}{q}}. \end{aligned}$$

On using Theorem 4 (i) we obtain  ${}_{a,b}D_{c+}^{\alpha,\beta} \in \left[ L_{w_1(x)dx}^p[a, b], L_{w_2(y)dy}^q[b, +\infty] \right]$ .

## REFERENCES

- [1] K.F. Andersen, *Weighted Inequalities for the Stieltjes Transformation and Hilbert's Double Series*, Proc. Royal Soc. Edinburgh, 86A (1980), pp. 75–84.
- [2] G.H. Hardy - J.E. Littlewood - G.Pólya, *Inequalities*, Cambridge Univ. Press, 2nd. ed., 1952.
- [3] A.A. Kilbas - S.G. Samko - O.I. Marichev, *Fract. Integrals and derivatives*, Gordon and Breach Science Publ., N. Y., 1993.
- [4] A.M. Nahushev - M.S. Salahitdinov, *The Law of Composition of Operators for Fractional Integrodifferentiation with Various Origins*, (Russian). Dokl. Acad. Nauk SSSR, 299 (1988), pp. 1313–1316.
- [5] T.J. Osler, *Open Questions For Research*, in Proc. Intern. Conf. Fract. Calculus and Its Appl, N. H., 1974, B. Ross (ed.), Lect. Notes in Math., 457, pp. 376–381.
- [6] C.C. Peña , *On Some Classes of Operators Derived From R - L Fract. Differentiation*, Boletim da Sociedade Paranaense de Matematica. Nova Serie, 15 (1 - 2) (1996), pp. 83–96.
- [7] B.S. Rubin, *On Spaces of Fract. Integrals and Straightline Contours*, (Russian). Izv. Akad. Nauk Armyan. SSR, Ser. Mat., 7 (1972), pp. 373–386.
- [8] G. Sinnamon, *A Note on Stieltjes Transformation*, Proc. Royal Soc. Edinburgh, 110A (1988), pp. 73–78.
- [9] V.D. Stepanov, *Weighted Inequalities for a Class of Volterra Convolution Operators*, J. London Math. Soc., 45 (1992), pp. 232–242.

*Universidad Nacional del Centro de la Provincia de Bs. As.,  
Facultad de Cs. Exactas,  
Dpto. de Matemática,  
Paraje Arroyo Seco,  
Campus Universitario,  
Tandil (C.P. 7000) (ARGENTINA)*