

LE MATEMATICHE

Vol. LXVI (2011) – Fasc. I, pp. 65–77

doi: 10.4418/2011.66.1.6

## SUBORDINATION AND SUPERORDINATION PROPERTIES FOR ANALYTIC FUNCTIONS INVOLVING WRIGHT'S FUNCTIONS

G. MURUGUSUNDARAMOORTHY - N. MAGESH - R. K. RAINA

In the present investigation, we obtain some subordination and superordination results for the Hadamard product of certain normalized analytic functions in the open unit disk involving the linear operator introduced in [J. Dziok and R. K. Raina, *Demonstratio Math.*, 37 (3) (2004), 533–542]. Several consequences of the results are presented. It is also pointed out that one of the main results (Theorem 2.8 below) provides a corrected form of the proof stated in two recent known results.

### 1. Introduction

Let  $\mathcal{H}$  denote the class of analytic functions in  $\mathbb{U} := \{z : |z| < 1\}$ , and  $\mathcal{H}[a, n]$  be the subclass of  $\mathcal{H}$  comprising of functions of the form  $f(z) = a + a_n z^n + a_{n+1} z^{n+1} + \dots$ , and  $\mathcal{A}$  be another subclass of  $\mathcal{H}$  which consists of functions of the form

$$f(z) = z + a_2 z^2 + \dots \quad (1)$$

Suppose  $p, h \in \mathcal{H}$  and let  $\phi(r, s, t; z) : \mathbb{C}^3 \times \mathbb{U} \rightarrow \mathbb{C}$ . If  $p$  and  $\phi(p(z), zp'(z), z^2 p''(z); z)$  are univalent and if  $p$  satisfies the second order superordination

$$h(z) \prec \phi(p(z), zp'(z), z^2 p''(z); z), \quad (2)$$

---

Entrato in redazione: 11 marzo 2011

*AMS 2010 Subject Classification:* 30C45, 30C80.

*Keywords:* Univalent functions, subordination, superordination, Hadamard product(convolution), Wright's generalized hypergeometric functions.

then  $p$  is a solution of the differential superordination (2). (If  $f$  is subordinate to  $F$ , then  $F$  is superordinate to  $f$ .) An analytic function  $q$  is called a *subordinant* if  $q \prec p$  for all  $p$  satisfying (2). A univalent subordinant  $\tilde{q}$  that satisfies  $q \prec \tilde{q}$  for all subordinants  $q$  of (2) is said to be the best subordinant. Recently Miller and Mocanu[15] obtained conditions on  $h$ ,  $q$  and  $\phi$  for which the following implication holds:

$$h(z) \prec \phi(p(z), zp'(z), z^2 p''(z); z) \Rightarrow q(z) \prec p(z).$$

For two functions  $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$  and  $g(z) = z + \sum_{k=2}^{\infty} b_k z^k$ , the Hadamard product (or convolution) of  $f$  and  $g$  is defined by

$$(f * g)(z) := z + \sum_{k=2}^{\infty} a_k b_k z^k =: (g * f)(z).$$

For positive real parameters  $\alpha_1, A_1, \dots, \alpha_l, A_l$  and  $\beta_1, B_1, \dots, \beta_m, B_m$  ( $l, m \in \mathbb{N} = 1, 2, 3, \dots$ ) such that

$$1 + \sum_{i=1}^m B_i - \sum_{j=1}^l A_j \geq 0, \quad (3)$$

the Wright's generalization [24] given by

$${}_l\Psi_m[(\alpha_1, A_1), \dots, (\alpha_l, A_l); (\beta_1, B_1), \dots, (\beta_m, B_m); z] = {}_l\Psi_m[(\alpha_i, A_i)_{1,l}; (\beta_j, B_j)_{1,m}; z]$$

of the generalized hypergeometric function  ${}_pF_q(\alpha_1, \dots, \alpha_p; \beta_1, \dots, \beta_q; z)$  is defined by

$${}_l\Psi_m[(\alpha_t, A_t)_{1,l}; (\beta_t, B_t)_{1,m}; z] = \sum_{k=0}^{\infty} \left\{ \prod_{t=0}^l \Gamma(\alpha_t + kA_t) \right\} \left\{ \prod_{t=0}^m \Gamma(\beta_t + kB_t) \right\}^{-1} \frac{z^k}{k!} \quad (z \in \mathbb{U}).$$

If  $A_t = 1$  ( $t = 1, \dots, l$ ) and  $B_t = 1$  ( $t = 1, \dots, m$ ), we have the relationship :

$$\begin{aligned} \Omega {}_l\Psi_m[(\alpha_t, 1)_{1,l}; (\beta_t, 1)_{1,m}; z] &\equiv {}_lF_m(\alpha_1, \dots, \alpha_l; \beta_1, \dots, \beta_m; z) \\ &= \sum_{k=0}^{\infty} \frac{(\alpha_1)_k \dots (\alpha_l)_k}{(\beta_1)_k \dots (\beta_m)_k} \frac{z^k}{k!}, \end{aligned} \quad (4)$$

where  ${}_lF_m$  is the generalized hypergeometric function (see, for example the details in [19]) such that  $l \leq m + 1$  ( $l, m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$ ),  $z \in \mathbb{U}$ ;  $\mathbb{N}$  denotes the set of all positive integers,  $(\lambda)_n = \lambda(\lambda + 1)\dots(\lambda + n - 1)$  is the Pochhammer symbol, and  $\Omega$  is given by

$$\Omega = \left( \prod_{t=0}^l \Gamma(\alpha_t) \right)^{-1} \left( \prod_{t=0}^m \Gamma(\beta_t) \right). \quad (5)$$

By using the generalized hypergeometric function (4), Dziok and Srivastava [7] introduced a linear operator which was subsequently extended by Dziok and Raina [8] by using the Wright's generalized hypergeometric function (defined above). For the purpose of this paper, we recall the Dziok-Raina linear operator as follows:

$$\Theta[(\alpha_t, A_t)_{1,l}; (\beta_t, B_t)_{1,m}] : \mathcal{A} \rightarrow \mathcal{A}$$

is a linear operator which is defined (in terms of the convolution) by

$$\Theta[(\alpha_t, A_t)_{1,l}; (\beta_t, B_t)_{1,m}]f(z) := z {}_l\Psi_m[(\alpha_t, A_t)_{1,l}; (\beta_t, B_t)_{1,m}; z] * f(z).$$

We observe for the function  $f(z)$  of the form (1) that

$$\Theta[(\alpha_t, A_t)_{1,l}; (\beta_t, B_t)_{1,m}]f(z) = z + \sum_{k=2}^{\infty} \sigma_k(\alpha_1) a_k z^k, \quad (6)$$

where  $\sigma_k(\alpha_1)$  is defined by

$$\sigma_k(\alpha_1) = \frac{\Omega \Gamma(\alpha_1 + A_1(k-1)) \dots \Gamma(\alpha_l + A_l(k-1))}{(k-1)! \Gamma(\beta_1 + B_1(k-1)) \dots \Gamma(\beta_m + B_m(k-1))} \quad (7)$$

and  $\Omega$  is given by (5).

For notational convenience, we write

$$\Theta[\alpha_1]f(z) = \Theta[(\alpha_1, A_1), \dots, (\alpha_l, A_l); (\beta_1, B_1), \dots, (\beta_m, B_m)]f(z), \quad (8)$$

which in view of (6) gives (see [8])

$$z A_1 (\Theta[\alpha_1]f(z))' = \alpha_1 \Theta[\alpha_1 + 1]f(z) - (\alpha_1 - A_1) \Theta[\alpha_1]f(z). \quad (9)$$

The Dziok-Raina operator (6) has been studied recently in the theory of analytic functions by many authors and one may refer to [9], [10] for the latest work on the subject. In view of the relationship (4), the linear operator (6) includes (as its special cases) various other linear operators introduced and studied by Bernardi [1], Carlson and Shaffer [4], Cho-Kwon-Srivastava [5], Choi-Saigo-Srivastava [6], Libera [11], Livingston [12], Ruscheweyh [21] and Srivastava-Owa [23].

In [3], Bulboacă (see also [2]) considered certain classes of first order differential subordinations as well as subordination-preserving integral operators by using the results of Miller and Mocanu [15]. Further, using the results in [2] and [15], Shanmugam *et al.* [22] obtained sufficient conditions for the normalized analytic function  $f(z)$  to satisfy

$$q_1(z) \prec \frac{f(z)}{z f'(z)} \prec q_2(z) \text{ and } q_1(z) \prec \frac{z^2 f'(z)}{\{f(z)\}^2} \prec q_2(z).$$

Recently, Murugusundaramoorthy and Magesh in their papers [13], [16], [17] and [18] obtained certain sufficient conditions for the normalized analytic function  $f(z)$  to satisfy

$$q_1(z) \prec \left( \frac{H_m^l[\alpha_1]f(z)}{(z)} \right)^\delta \prec q_2(z), \quad q_1(z) \prec \frac{(f * \Phi)(z)}{(f * \Psi)(z)} \prec q_2(z)$$

and

$$q_1(z) \prec \frac{zf'(z)}{f(z)} \prec q_2(z), \quad q_1(z) \prec \frac{f(z)}{zf'(z)} \prec q_2(z),$$

where  $q_1, q_2$  are given univalent functions in  $\mathbb{U}$  with  $q_1(0) = 1$  and  $q_2(0) = 1$ .

The main aim of the present paper is to find sufficient conditions for the Hadamard product of a fixed normalized analytic function  $f(z)$  with the normalized analytic functions  $\phi(z)$  and  $\psi(z)$  in the open unit disk  $\mathbb{U}$ , such that  $(f * \Psi)(z) \neq 0$ , involving the Dziok-Raina operator [8] which satisfy

$$q_1(z) \prec \frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \prec q_2(z),$$

where  $q_1, q_2$  are given univalent functions in  $\mathbb{U}$ , and  $\Phi(z) = z + \sum_{n=2}^{\infty} \lambda_n z^n$ ,  $\Psi(z) = z + \sum_{n=2}^{\infty} \mu_n z^n$  are analytic functions in  $\mathbb{U}$  with  $\lambda_n \geq 0$ ,  $\mu_n \geq 0$  ( $\lambda_n \geq \mu_n$ ). Several results are deduced as worthwhile consequences of the main results. In particular, we mention two such special cases of one of our main results which provide the corrected proof details of certain recent results established in [17] and [18].

## 2. Subordination and superordination results

In our present investigation, we require following key lemmas.

**Lemma 2.1.** [14, p.132, Theorem 3.4h] *Let  $q$  be univalent in the unit disk  $\mathbb{U}$  and let  $\theta$  and  $\phi$  be analytic in a domain  $\mathbb{D}$  containing  $q(\mathbb{U})$ , with  $\phi(w) \neq 0$ , when  $w \in q(\mathbb{U})$ . Set  $Q(z) := zq'(z)\phi(q(z))$  and  $h(z) := \theta(q(z)) + Q(z)$ . Suppose that  $Q(z)$  is starlike univalent in  $\mathbb{U}$  and  $\text{Re} \left\{ \frac{zh'(z)}{Q(z)} \right\} > 0$  for  $z \in \mathbb{U}$ . If  $p$  is analytic in  $\mathbb{U}$ , with  $p(0) = q(0)$ ,  $p(\mathbb{U}) \subset \mathbb{D}$  and*

$$\theta(p(z)) + zp'(z)\phi(p(z)) \prec \theta(q(z)) + zq'(z)\phi(q(z)), \quad (10)$$

*then  $p(z) \prec q(z)$  and  $q$  is the best dominant.*

**Lemma 2.2.** [3, p.289, Corollary 3.2] *Let  $q$  be convex univalent in the unit disk  $\mathbb{U}$  and  $\vartheta$  and  $\varphi$  be analytic in a domain  $\mathbb{D}$  containing  $q(\mathbb{U})$ . Suppose that,  $Re \{ \vartheta'(q(z))/\varphi(q(z)) \} > 0$  for  $z \in \mathbb{U}$  and  $\psi(z) = zq'(z)\varphi(q(z))$  is star-like univalent in  $\mathbb{U}$ . If  $p(z) \in \mathbb{H}[q(0), 1] \cap \mathcal{Q}$ , with  $p(\mathbb{U}) \subseteq \mathbb{D}$ , and  $\vartheta(p(z)) + zp'(z)\varphi(p(z))$  is univalent in  $\mathbb{U}$  and*

$$\vartheta(q(z)) + zq'(z)\varphi(q(z)) \prec \vartheta(p(z)) + zp'(z)\varphi(p(z)), \tag{11}$$

then  $q(z) \prec p(z)$  and  $q$  is the best subordinated.

Making use of Lemma 2.1, we first prove the following theorem.

**Theorem 2.3.** *Let  $\Phi, \Psi \in \mathcal{A}$ ,  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , and assume that*

$$Re \left\{ \frac{\gamma_3}{\gamma_4} + \frac{2\gamma_2}{\gamma_4}q(z) + \left( 1 + \frac{zq''(z)}{q'(z)} \right) \right\} > 0 \quad (z \in \mathbb{U}). \tag{12}$$

If  $f \in \mathcal{A}$  satisfies

$$\Delta^{(\gamma)_1^{\dagger}}(f; \Phi, \Psi) = \Delta(f, \Phi, \Psi, \gamma_1, \gamma_2, \gamma_3, \gamma_4) \prec \gamma_1 + \gamma_2q^2(z) + \gamma_3q(z) + \gamma_4zq'(z), \tag{13}$$

where

$$\Delta^{(\gamma)_1^{\dagger}}(f; \Phi, \Psi) := \begin{cases} \gamma_1 + \gamma_2 \left( \frac{\Theta[\alpha_1](f*\Phi)(z)}{\Theta[\alpha_1+1](f*\Psi)(z)} \right)^2 + \gamma_3 \frac{\Theta[\alpha_1](f*\Phi)(z)}{\Theta[\alpha_1+1](f*\Psi)(z)} \\ + \frac{\gamma_4}{A_1} \left( \alpha_1 \frac{\Theta[\alpha_1+1](f*\Phi)(z)}{\Theta[\alpha_1](f*\Phi)(z)} - (\alpha_1 + 1) \frac{\Theta[\alpha_1+2](f*\Psi)(z)}{\Theta[\alpha_1+1](f*\Psi)(z)} + 1 \right) \\ \times \left( \frac{\Theta[\alpha_1](f*\Phi)(z)}{\Theta[\alpha_1+1](f*\Psi)(z)} \right), \end{cases} \tag{14}$$

then

$$\frac{\Theta[\alpha_1](f*\Phi)(z)}{\Theta[\alpha_1+1](f*\Psi)(z)} \prec q(z)$$

and  $q$  is the best dominant.

*Proof.* Define the function  $p$  by

$$p(z) := \frac{\Theta[\alpha_1](f*\Phi)(z)}{\Theta[\alpha_1+1](f*\Psi)(z)} \quad (z \in \mathbb{U}). \tag{15}$$

We observe (in view of (6)) that the function  $p(z)$  is analytic in  $\mathbb{U}$  with  $p(0) = 1$ . Differentiating (15) with respect to  $z$ , and making use of (9), we obtain after elementary calculations the following:

$$\begin{aligned} & \gamma_1 + \gamma_2 \left( \frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \right)^2 + \gamma_3 \frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \\ & + \frac{\gamma_4}{A_1} \left( \alpha_1 \frac{\Theta[\alpha_1 + 1](f * \Phi)(z)}{\Theta[\alpha_1](f * \Phi)(z)} - (\alpha_1 + 1) \frac{\Theta[\alpha_1 + 2](f * \Psi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} + 1 \right) \\ & \times \left( \frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \right) = \gamma_1 + \gamma_2 p^2(z) + \gamma_3 p(z) + \gamma_4 z p'(z). \end{aligned} \quad (16)$$

Using (16) in (13), we get

$$\gamma_1 + \gamma_2 p^2(z) + \gamma_3 p(z) + \gamma_4 z p'(z) < \gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z), \quad (17)$$

and by setting  $\theta(w) := \gamma_1 + \gamma_2 w^2 + \gamma_3 w$  and  $\phi(w) := \gamma_4$  ( $w = q(z)$ ), it can be easily observed that both the functions  $\theta(w)$  and  $\phi(w)$  are analytic in  $\mathbb{C} \setminus \{0\}$  and  $\phi(w) \neq 0$ . Also, we notice that

$$Q(z) := z q'(z) \phi(q(z)) = \gamma_4 z q'(z)$$

and

$$h(z) := \theta(q(z)) + Q(z) = \gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z).$$

It is clear that  $Q(z)$  is starlike univalent in  $\mathbb{U}$  and

$$\operatorname{Re} \left\{ \frac{z h'(z)}{Q(z)} \right\} = \operatorname{Re} \left\{ \frac{\gamma_3}{\gamma_4} + \frac{2\gamma_2}{\gamma_4} q(z) + \left( 1 + \frac{z q''(z)}{q'(z)} \right) \right\} > 0.$$

In view of the hypothesis of Theorem 2.3, the result now follows by an application of Lemma 2.1.  $\square$

By choosing  $A_t = 1$  ( $t = 1, \dots, l$ ) and  $B_t = 1$  ( $t = 1, \dots, m$ ),  $l = 2$ ,  $m = 1$ ,  $\alpha_1 = 1$ ,  $\alpha_2 = 1$  and  $\beta_1 = 1$  in Theorem 2.3, we state the following corollary.

**Corollary 2.4.** *Let  $\Phi, \Psi \in \mathcal{A}$ ,  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , and suppose that (12) holds true. If  $f \in \mathcal{A}$  satisfies*

$$\begin{aligned} & \gamma_1 + \gamma_2 \left( \frac{(f * \Phi)(z)}{z(f * \Psi)'(z)} \right)^2 \\ & + \gamma_4 \frac{(f * \Phi)(z)}{z(f * \Psi)'(z)} \left[ \frac{z(f * \Phi)'(z)}{(f * \Phi)(z)} - \frac{z(f * \Psi)''(z)}{(f * \Psi)'(z)} + \frac{\gamma_3}{\gamma_4} - 1 \right] \\ & < \gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z), \end{aligned}$$

then  $\frac{(f * \Phi)(z)}{z(f * \Psi)'(z)} < q(z)$  and  $q$  is the best dominant.

Corollary 2.4 in the special case when  $\Psi(z) = \Phi(z) = \frac{z}{1-z}$ , yields the following interesting result.

**Corollary 2.5.** *Let  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , and suppose that (12) holds true. If  $f \in \mathcal{A}$  satisfies*

$$\begin{aligned} & \gamma_1 + \gamma_2 \left( \frac{f(z)}{zf'(z)} \right)^2 + \gamma_4 \frac{f(z)}{zf'(z)} \left[ \frac{zf'(z)}{f(z)} - \frac{zf''(z)}{f'(z)} + \frac{\gamma_3}{\gamma_4} - 1 \right] \prec \\ & \gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z), \end{aligned}$$

then  $\frac{f(z)}{zf'(z)} \prec q(z)$  and  $q$  is the best dominant.

**Remark 2.6.** For  $\gamma_1 = \gamma_2 = 0$  and  $\gamma_3 = 1$ , Corollary 2.5 yields the known result due to Shanmugam *et.al* [22].

If we choose  $q(z)$  as a bilinear function, viz.  $q(z) = \frac{1+Az}{1+Bz}$  ( $-1 \leq B < A \leq 1$ ) in Theorem 2.3, we get the following:

**Corollary 2.7.** *Let  $\Phi, \Psi \in \mathcal{A}$ ,  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , assume that (12) holds true. If  $f \in \mathcal{A}$  and*

$$\Delta^{(\gamma)_1^4}(f; \Phi, \Psi) \prec \gamma_1 + \gamma_2 \left( \frac{1+Az}{1+Bz} \right)^2 + \gamma_3 \frac{1+Az}{1+Bz} + \gamma_4 \frac{(A-B)z}{(1+Bz)^2},$$

then

$$\frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \prec \frac{1+Az}{1+Bz},$$

where  $\Delta^{(\gamma)_1^4}(f; \Phi, \Psi)$  is given by (14), and  $\frac{1+Az}{1+Bz}$  is the best dominant.

Next, by applying Lemma 2.2, we prove the following.

**Theorem 2.8.** *Let  $\Phi, \Psi \in \mathcal{A}$ ,  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , and assume that*

$$Re \left\{ \frac{\gamma_3}{\gamma_4} + \frac{2\gamma_2}{\gamma_4} q(z) \right\} \geq 0. \tag{18}$$

Suppose that  $f \in \mathcal{A}$ ,  $\frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \in \mathbb{H}[q(0), 1] \cap \mathcal{Q}$ , and let  $\Delta^{(\gamma)_1^4}(f; \Phi, \Psi)$  be univalent in  $\mathbb{U}$  and

$$\gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z) \prec \Delta^{(\gamma)_1^4}(f; \Phi, \Psi), \tag{19}$$

where  $\Delta^{(\gamma)_1^4}(f; \Phi, \Psi)$  is given by (14), then

$$q(z) \prec \frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)}$$

and  $q$  is the best subordinated.

*Proof.* Let the function  $p(z)$  be defined by (15) (which is analytic in the unit disk  $\mathbb{U}$  with  $p(0) = 1$ ). Following the same steps as in proof of Theorem 2.3, and using (19), we infer that

$$\gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z) \prec \gamma_1 + \gamma_2 p^2(z) + \gamma_3 p(z) + \gamma_4 z p'(z).$$

By setting  $\vartheta(w) = \gamma_1 + \gamma_2 w^2 + \gamma_3 w$  and  $\phi(w) = \gamma_4$ , it is easily observed that  $\vartheta(w)$  is analytic in  $\mathbb{C}$ . Also,  $\phi(w)$  is analytic in  $\mathbb{C} \setminus \{0\}$  and that  $\phi(w) \neq 0$ . In view of the result [20, p.159, Theorem 6.2], if we let

$$L(z, t) = \gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 t z q'(z) = a_1(t)z + \dots, \quad (20)$$

then on differentiating (20) with respect to  $z$  and  $t$ , we get

$$\frac{\partial L(z, t)}{\partial z} = 2\gamma_2 q(z)q'(z) + \gamma_3 q'(z) + t\gamma_4 z q''(z) + t\gamma_4 q'(z) = a_1(t) + \dots$$

and  $\frac{\partial L(z, t)}{\partial t} = \gamma_4 z q'(z)$ . Also,  $\frac{\partial L(0, t)}{\partial z} = \gamma_4 q'(0) \left[ \frac{\gamma_3}{\gamma_4} + \frac{2\gamma_2}{\gamma_4} q(0) + t \right]$ .

From the univalence of  $q$  we have  $q'(0) \neq 0$  and  $q(0) = 1$ , it follows that  $a_1(t) \neq 0$  for  $t \geq 0$  and  $\lim_{t \rightarrow \infty} |a_1(t)| = +\infty$ , and simple computation now yields

$$\operatorname{Re} \left\{ z \frac{\frac{\partial L(z, t)}{\partial z}}{\frac{\partial L(z, t)}{\partial t}} \right\} = \operatorname{Re} \left\{ \frac{\gamma_3}{\gamma_4} + \frac{2\gamma_2}{\gamma_4} q(z) + t \left( 1 + \frac{z q''(z)}{q'(z)} \right) \right\}.$$

Using the fact that  $q$  is a convex univalent function in  $\mathbb{U}$  and  $\gamma_4 \neq 0$ , we conclude that

$$\operatorname{Re} \left\{ z \frac{\frac{\partial L(z, t)}{\partial z}}{\frac{\partial L(z, t)}{\partial t}} \right\} > 0 \text{ if } \operatorname{Re} \left\{ \frac{\gamma_3}{\gamma_4} + \frac{2\gamma_2}{\gamma_4} q(z) \right\} > 0 \quad (z \in \mathbb{U}, t \geq 0)$$

and Theorem 2.8, follows by applying Lemma 2.2.  $\square$

**Remark 2.9.** We mention here that a recent known result [18, p.90, Theorem 2.12] evidently follows from the above Theorem 2.8 by reducing the Dziok-Raina operator (8) to the corresponding Dziok-Srivastava operator (defined in [7]) in view of the parametric substitutions stated in (4). It may also be noted that Theorem 2.8 gives a *corrected form of the proof derivation* of two known recent results [17, p.123, Theorem 3.1] and [18, p.90, Theorem 2.12].



When  $A_t = 1$  ( $t = 1, \dots, l$ ) and  $B_t = 1$  ( $t = 1, \dots, m$ ),  $l = 2$ ,  $m = 1$ ,  $\alpha_1 = 1$ ,  $\alpha_2 = 1$  and  $\beta_1 = 1$  in Theorem 2.8, we derive the following corollary.

**Corollary 2.10.** *Let  $\Phi, \Psi \in \mathcal{A}$ ,  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , and assume that (18) holds true.*

*If  $f \in \mathcal{A}$ ,  $\frac{(f*\Phi)(z)}{z(f*\Psi)'(z)} \in \mathbb{H}[q(0), 1] \cap \mathcal{Q}$  let  $\gamma_1 + \gamma_2 \left( \frac{(f*\Phi)(z)}{z(f*\Psi)'(z)} \right)^2 + \gamma_4 \frac{(f*\Phi)(z)}{z(f*\Psi)'(z)} \left[ \frac{z(f*\Phi)'(z)}{(f*\Phi)(z)} - \frac{z(f*\Psi)''(z)}{(f*\Psi)'(z)} + \frac{\gamma_3}{\gamma_4} - 1 \right]$  be univalent in  $\mathbb{U}$  and*

$$\begin{aligned} & \gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z) \prec \gamma_1 + \gamma_2 \left( \frac{(f*\Phi)(z)}{z(f*\Psi)'(z)} \right)^2 \\ & + \gamma_4 \frac{(f*\Phi)(z)}{z(f*\Psi)'(z)} \left[ \frac{z(f*\Phi)'(z)}{(f*\Phi)(z)} - \frac{z(f*\Psi)''(z)}{(f*\Psi)'(z)} + \frac{\gamma_3}{\gamma_4} - 1 \right]. \end{aligned}$$

*Then  $q(z) \prec \frac{(f*\Phi)(z)}{z(f*\Psi)'(z)}$  and  $q$  is the best subordinator.*

By fixing  $\Phi(z) = \Psi(z) = \frac{z}{1-z}$  in Corollary 2.10, we obtain the following corollary.

**Corollary 2.11.** *Let  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , and (18) holds true. If  $f \in \mathcal{A}$ ,  $\frac{f(z)}{zf'(z)} \in \mathbb{H}[q(0), 1] \cap \mathcal{Q}$ . Let  $\gamma_1 + \gamma_2 \left( \frac{f(z)}{zf'(z)} \right)^2 + \gamma_4 \frac{f(z)}{zf'(z)} \left[ \frac{zf'(z)}{f(z)} - \frac{zf''(z)}{f'(z)} + \frac{\gamma_3}{\gamma_4} - 1 \right]$  be univalent in  $\mathbb{U}$  and*

$$\begin{aligned} & \gamma_1 + \gamma_2 q^2(z) + \gamma_3 q(z) + \gamma_4 z q'(z) \\ & \prec \gamma_1 + \gamma_2 \left( \frac{f(z)}{zf'(z)} \right)^2 + \gamma_4 \frac{f(z)}{zf'(z)} \left[ \frac{zf'(z)}{f(z)} - \frac{zf''(z)}{f'(z)} + \frac{\gamma_3}{\gamma_4} - 1 \right], \end{aligned}$$

*then  $q(z) \prec \frac{f(z)}{zf'(z)}$  and  $q$  is the best subordinator.*

If we choose  $q(z) = (1 + Az)/(1 + Bz)$  ( $-1 \leq B < A \leq 1$ ) in Theorem 2.8, then we obtain the following corollary.

**Corollary 2.12.** *Let  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ),  $q$  be convex univalent with  $q(0) = 1$ , and assume that (18) holds true. If  $f \in \mathbb{A}$ ,  $\frac{\Theta[\alpha_1](f*\Phi)(z)}{\Theta[\alpha_1+1](f*\Psi)(z)} \in \mathbb{H}[q(0), 1] \cap \mathcal{Q}$ . Let  $\Delta^{(\gamma)_1^\dagger}(f; \Phi, \Psi)$  be univalent in  $\mathbb{U}$  and*

$$\gamma_1 + \gamma_2 \left( \frac{1 + Az}{1 + Bz} \right)^2 + \gamma_3 \frac{1 + Az}{1 + Bz} + \gamma_4 \frac{(A - B)z}{(1 + Bz)^2} \prec \Delta^{(\gamma)_1^\dagger}(f; \Phi, \Psi),$$

then

$$\frac{1 + Az}{1 + Bz} \prec \frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)}$$

and  $\frac{1+Az}{1+Bz}$  is the best subordinate.

### 3. Composition Results

Combining the two results given by Theorem 2.3 (involving differential subordination) and Theorem 2.8 (involving differential superordination), we have the following sandwich theorem.

**Theorem 3.1.** Let  $q_1$  and  $q_2$  be convex univalent in  $\mathbb{U}$ ,  $\gamma_i \in \mathbb{C}$  ( $i = 1, \dots, 4$ ) ( $\gamma_4 \neq 0$ ), and let  $q_2$  satisfy (12) and  $q_1$  satisfy (18). For  $f, \Phi, \Psi \in \mathcal{A}$ , let  $\frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \in \mathbb{H}[1, 1] \cap Q$  and  $\Delta^{(\gamma)}_1(f; \Phi, \Psi)$  defined by (14) be univalent in  $\mathbb{U}$  satisfying

$$\begin{aligned} \gamma_1 + \gamma_2 q_1^2(z) + \gamma_3 q_1(z) + \gamma_4 z q_1'(z) &\prec \Delta^{(\gamma)}_1(f; \Phi, \Psi) \\ &\prec \gamma_1 + \gamma_2 q_2^2(z) + \gamma_3 q_2(z) + \gamma_4 z q_2'(z), \end{aligned}$$

then

$$q_1(z) \prec \frac{\Theta[\alpha_1](f * \Phi)(z)}{\Theta[\alpha_1 + 1](f * \Psi)(z)} \prec q_2(z)$$

and  $q_1, q_2$  are, respectively, the best subordinate and best dominant.

On putting  $q_1(z) = \frac{1+A_1z}{1+B_1z}$  ( $-1 \leq B_1 < A_1 \leq 1$ ) and  $q_2(z) = \frac{1+A_2z}{1+B_2z}$  ( $-1 \leq B_2 < A_2 \leq 1$ ) in Theorem 3.1, we obtain the following result.

**Corollary 3.2.** For  $f, \Phi, \Psi \in \mathcal{A}$ , let  $\frac{(f * \Phi)(z)}{z(f * \Psi)'(z)} \in \mathcal{H}[1, 1] \cap Q$  and  $\Delta^{(\gamma)}_1(f; \Phi, \Psi)$  defined by (14) be univalent in  $\mathbb{U}$  satisfying

$$\begin{aligned} \gamma_1 + \gamma_2 \left( \frac{1 + A_1 z}{1 + B_1 z} \right)^2 + \gamma_3 \frac{1 + A_1 z}{1 + B_1 z} + \gamma_4 \frac{(A_1 - B_1)z}{(1 + B_1 z)^2} &\prec \Delta^{(\gamma)}_1(f; \Phi, \Psi) \\ &\prec \gamma_1 + \gamma_2 \left( \frac{1 + A_2 z}{1 + B_2 z} \right)^2 + \gamma_3 \frac{1 + A_2 z}{1 + B_2 z} + \gamma_4 \frac{(A_2 - B_2)z}{(1 + B_2 z)^2}, \end{aligned}$$

then

$$\frac{1 + A_1 z}{1 + B_1 z} \prec \frac{(f * \Phi)(z)}{z(f * \Psi)'(z)} \prec \frac{1 + A_2 z}{1 + B_2 z}$$

and  $\frac{1+A_1z}{1+B_1z}, \frac{1+A_2z}{1+B_2z}$  are, respectively, the best subordinate and best dominant.

**Remark 3.3.** We conclude this paper by remarking that the various results can be derived from Theorems 2.3, 2.8 and 3.1 for the different choices of the arbitrary functions  $\Phi(z)$  and  $\Psi(z)$ . In fact, by appropriately selecting the arbitrary parameters  $A_t, B_t, l, m, \alpha_1, \dots, \alpha_l, \beta_1, \dots, \beta_m$ , involved in the operator (6), we observe that the generalized form of the linear operator of Dziok and Raina [8] reduced to well known operators like the Dziok-Srivastava linear operator, Hohlov linear operator, Saitoh generalized linear operator, Carlson-Shaffer linear operator, Ruscheweyh derivative operator (as well as its generalized versions), Bernardi-Libera-Livingston operator, Cho-Kwon-Srivastava operator, Choi-Saigo-Srivastava operator and Srivastava-Owa fractional derivative operator. These consequences are easy to deduce, and we make here no attempt to mention these obvious special cases of Theorems 2.3, 2.8 and 3.1.

### Acknowledgments

The authors would like to record their sincere thanks to the referee(s) for their valuable comments and insightful suggestions.

### REFERENCES

- [1] S. D. Bernardi, *Convex and starlike univalent functions*, Trans. Amer. Math. Soc. 135 (1969), 429–446.
- [2] T. Bulboacă, *A class of superordination-preserving integral operators*, Indag. Math. New Ser. 13 (3) (2002), 301–311.
- [3] T. Bulboacă, *Classes of first order differential superordinations*, Demonstratio Math. 35 (2) (2002), 287–292.
- [4] B. C. Carlson - S. B. Shaffer, *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal. 15 (1984), 737–745.
- [5] N. E. Cho - O. H. Kwon - H. M. Srivastava, *Inclusion and argument properties for certain subclasses of multivalent functions associated with a family of linear operators*, J. Math. Anal. Appl. 292 (2004), 470–483.
- [6] J. H. Choi - M. Saigo - H. M. Srivastava, *Some inclusion properties of a certain family of integral operators*, J. Math. Anal. Appl. 276 (2002), 432–445.
- [7] J. Dziok - H. M. Srivastava, *Certain subclasses of analytic functions associated with the generalized hypergeometric function*, Intergral Transform Spec. Funct. 14 (2003), 7–18.
- [8] J. Dziok - R. K. Raina, *Families of analytic functions associated with the Wright's generalized hypergeometric function*, Demonstratio Math. 37 (3) (2004), 533–542.

- [9] J. Dziok - R. K. Raina, *Some results based on first order differential subordination with the Wright's generalized hypergeometric function*, Comment. Math. Univ. St. Pauli. 58 (2) (2009), 87–94.
- [10] J. Dziok - R. K. Raina, *Extremal problems in the class of analytic functions associated with the Wright's function*, Adv. Pure. Appl. Math. 1 (2010), 1–12.
- [11] R. J. Libera, *Some classes of regular univalent functions*, Proc. Amer. Math. Soc. 16 (1965), 755–758.
- [12] A. E. Livingston, *On the radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc. 17 (1966), 352–357.
- [13] N. Magesh - G. Murugusundaramoorthy, *Differential Subordinations and superordinations for Comprehensive class of analytic functions*, SUT J. Math. 44 (2) (2008), 237–255.
- [14] S. S. Miller - P. T. Mocanu, *Differential Subordinations: Theory and Applications*, Marcel Dekker Inc., New York, 2000.
- [15] S. S. Miller - P. T. Mocanu, *Subordinants of differential superordinations*, Complex Variables 48 (10) (2003), 815–826.
- [16] G. Murugusundaramoorthy - N. Magesh, *Differential subordinations and superordinations for analytic functions defined by Dziok-Srivastava linear operator*, J. Inequal. Pure Appl. Math. 7 (4) (2006), 1–9.
- [17] G. Murugusundaramoorthy - N. Magesh, *Differential Sandwich theorems for analytic functions defined by Hadamard product*, Annales Univ. M. Curie-Sklodowska Sec.A 59 (2007), 117–127.
- [18] G. Murugusundaramoorthy - N. Magesh, *Differential subordinations and superordinations for analytic functions defined by convolution structure*, Studia Univ. Babes-Bolyai Math. 54 (20) (2009), 83–96.
- [19] S. Owa - H. M. Srivastava, *Univalent and starlike generalized hypergeometric functions*, Canada J. Math. 39 (5) (1987), 1057–1077.
- [20] Ch. Pommerenke, *Univalent functions*, Vanderhoeck and Ruprecht, Göttingen, 1975.
- [21] St. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc. 49 (1975), 109–115.
- [22] T. N. Shanmugam - V. Ravichandran - S. Sivasubramanian, *Differential sandwich theorems for some subclasses of analytic functions*, Aust. J. Math. Anal. Appl. 3 (1) (2006), 1–11.
- [23] H. M. Srivastava - S. Owa, *Some characterization and distortion theorems involving fractional calculus, generalized hypergeometric functions, Hadamard products, linear operators and certain subclasses of analytic functions*, Nagoya Math. J. 106 (1987), 1–28.
- [24] E. M. Wright, *The asymptotic expansion of the generalized hypergeometric function*, Proc. London. Math. Soc. 46 (1946), 389–408.

*G. MURUGUSUNDARAMOORTHY*

*School of Advanced Sciences,  
VIT University, Vellore - 632014,  
Tamilnadu, India  
e-mail: gmsmoorthy@yahoo.com*

*N. MAGESH*

*Department of Mathematics,  
Government Arts College (Men),  
Krishnagiri - 635001, Tamilnadu, India  
e-mail: nmagi\_2000@yahoo.co.in*

*R. K. RAINA*

*M. P. University of Agriculture and Technology,  
Udaipur-313001, Rajasthan, India  
Present address: 10/11 Ganpati Vihar, Opposite Sector 5  
Udaipur 313002, Rajasthan, India  
e-mail: rkrainna\_7@hotmail.com*