

LE MATEMATICHE

Vol. LXVII (2012) – Fasc. II, pp. 77–92

doi: 10.4418/2012.67.2.7

## SOME DIFFERENTIAL SUBORDINATION AND SUPERORDINATION PROPERTIES OF SYMMETRIC ANALYTIC FUNCTIONS INVOLVING NOOR INTEGRAL OPERATOR

ALI MUHAMMAD - AMJADULLAH KHATTAK

In this paper, we obtain some interesting properties of differential subordination and superordination for the classes of symmetric functions analytic in the unit disc, by applying Noor integral operator. We investigate several sandwich theorems on the basis of this theory.

### 1. Introduction

Let  $H(E)$  denote the class of analytic functions in the open unit disc  $E = \{z \mid z \in \mathbb{C}, |z| < 1\}$  and let  $H[a, 1]$  denote the subclass of the functions  $f \in H(E)$  of the form

$$f(z) = a + a_1z + a_2z^2 + \dots \quad a \in \mathbb{C}.$$

Also, let  $\mathcal{A}$  be the subclass of functions  $f \in H(E)$  of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k. \quad (1)$$

---

Entrato in redazione: 22 dicembre 2011

AMS 2010 Subject Classification: 30C45, 30C10.

Keywords: Convex functions, Symmetric functions, Differential subordination and superordination, Noor integral operator.

If  $f$  and  $g$  are analytic in  $E$ , we say that  $f$  is subordinate to  $g$ , written  $f \prec g$  or  $f(z) \prec g(z)$ , if there exists a Schwarz function  $w$  in  $E$  such that  $f(z) = g(w(z))$ .

Suppose that  $h$  and  $k$  are two analytic functions in  $E$ , let

$$\varphi(r, s, t; z) : \mathbb{C}^3 \times E \longrightarrow \mathbb{C}.$$

If  $h$  and  $\varphi(h(z), zh'(z), z^2h''(z); z)$  are univalent functions in  $E$  and if  $h$  satisfies the second order superordination

$$k(z) \prec \varphi(h(z), zh'(z), z^2h''(z); z), \quad (2)$$

then  $k$  is said to be a solution of the differential superordination (2). A function  $q \in H(E)$  is called a subordinated to (2) if  $q(z) \prec h(z)$  for all the functions  $h$  satisfying (2).

A univalent subordinated  $\tilde{q}$  that satisfies  $q(z) \prec \tilde{q}(z)$  for all of the subordinants  $q$  of (2), is said to be the best subordinated.

Recently, Miller and Mocanu [10] obtained sufficient conditions on the functions  $k, q$  and  $\varphi$  for which the following implication holds:

$$k(z) \prec \varphi(h(z), zh'(z), z^2h''(z); z) \implies q(z) \prec h(z).$$

Using these results, the authors in [2] considered certain classes of first-order differential superordinations, see also [5], as well as superordination-preserving integral operators [4]. Aouf et al. [2, 3], obtained sufficient conditions for certain normalized analytic functions  $f$  to satisfy

$$q_1(z) \prec \frac{zf'(z)}{f(z)} \prec q_2(z),$$

where  $q_1$  and  $q_2$  are given univalent normalized functions in  $E$ . Very recently Shanmugam et al. ([19,20]) obtained the so called sandwich results for certain classes of analytic functions. For interested readers we refer to the work done by the authors [1, 2, 13, 14, 18].

In [17], Sakaguchi defined the class of starlike functions with respect to symmetrical points as follows:

Let  $f \in \mathcal{A}$ . Then  $f$  is said to be starlike with respect to symmetrical points in  $E$  if, and only if,

$$\operatorname{Re} \frac{zf'(z)}{f(z) - f(-z)} > 0, z \in E.$$

Obviously, it forms a subclass of close-to-convex functions and hence univalent. Moreover, this class includes the class of convex functions and odd starlike functions with respect to the origin, see [17].

Let  $f \in \mathcal{A}$ . Denote by  $D^\alpha : \mathcal{A} \rightarrow \mathcal{A}$  the operator defined by

$$D^\alpha f(z) = \frac{z}{(1-z)^{\alpha+1}} * f(z) \quad (\alpha > -1)$$

or equivalently,

$$D^n f(z) = \frac{z(z^{n-1}f(z))^{(n)}}{n!}, \quad n \in \mathbb{N}_0 = \{0, 1, 2, \dots\},$$

where the symbol  $(*)$  stands for the Hadamard product (or Convolution). We note that  $D^0 f(z) = f(z)$  and  $D^1 f(z) = zf'(z)$ . The operator  $D^n f$  is called the Ruscheweyh derivative of  $n$ th order of  $f$ , see [16]. Analogous to  $D^n f$ , Noor [11] and Noor and Noor [12] defined an integral operator  $I_n : \mathcal{A} \rightarrow \mathcal{A}$  as follows.

Let  $f_n(z) = \frac{z}{(1-z)^{n+1}}$ ,  $n \in \mathbb{N}_0$ , and let  $f_n^{(\dagger)}$  be defined such that

$$f_n(z) * f_n^{(\dagger)}(z) = \frac{z}{(1-z)^2}.$$

Then

$$I_n f(z) = f_n(z) * f_n^{(\dagger)}(z) = \left[ \frac{z}{(1-z)^{n+1}} \right]^{(\dagger)} * f(z). \tag{3}$$

From (3) it is easy to verify that

$$z(I_{n+1}f(z))' = (n+1)I_n f(z) - nI_{n+1}f(z). \tag{4}$$

We note that  $I_0 f(z) = zf'(z)$  and  $I_1 f(z) = f(z)$ . The operator  $I_n f(z)$  defined by (3) is called the Noor Integral operator of  $n$ th order of  $f$ , see [7]. Moreover, Liu [7] introduced some new subclasses of strongly starlike functions defined by using the Noor integral operator and studied their properties. Liu and Noor [8] investigated some interesting properties of the Noor integral operator.

## 2. Preliminary Results

**Definition 2.1** ([10]). Let  $Q$  be the set of all functions  $f$  that are analytic and injective on  $\bar{E} \setminus U(f)$ , where

$$U(f) = \left\{ \zeta \in \partial E : \lim_{z \rightarrow \zeta} f(z) = \infty \right\},$$

and are such that  $f'(\zeta) \neq 0$  for  $\zeta \in \partial E \setminus U(f)$ .

To establish our main results we need the following Lemmas.

**Lemma 2.2** (Miller and Mocanu [9]). *Let  $q$  be univalent in the unit disc  $E$ , and let  $\theta$  and  $\varphi$  be analytic in a domain  $D$  containing  $q(E)$ , with  $\varphi(w) \neq 0$  when  $w \in q(E)$ . Set  $Q(z) = zq'(z)\varphi(q(z))$ ,  $h(z) = \theta(q(z) + Q(z))$  and suppose that*

(i)  $Q$  is a starlike function in  $E$ ,

(ii)  $\operatorname{Re} \frac{zh'(z)}{Q(z)} > 0$ ,  $z \in E$ .

If  $p$  is analytic in  $E$  with  $p(0) = q(0)$ ,  $p(E) \subseteq D$  and

$$\theta(p(z)) + zp'(z)\varphi(p(z)) \prec \theta(q(z)) + zq'(z)\varphi(q(z)), \quad (5)$$

then  $p(z) \prec q(z)$ , and  $q$  is the best dominant of (5).

**Lemma 2.3** (Shanmugam et al. [20]). *Let  $\mu, \gamma \in \mathbb{C}$  with  $\gamma \neq 0$ , and let  $q$  be a convex function in  $E$  with*

$$\operatorname{Re} \left( 1 + \frac{zq''(z)}{q'(z)} \right) > \max \left\{ 0; -\operatorname{Re} \frac{\mu}{\gamma} \right\}, \quad z \in E.$$

If  $p$  is analytic in  $E$  and

$$\mu p(z) + \gamma zp'(z) \prec \mu q(z) + \gamma zq'(z), \quad (6)$$

then  $p(z) \prec q(z)$ , and  $q$  is the best dominant of (6).

**Lemma 2.4** (Bulboacă [6]). *Let  $q$  be a univalent function in the unit disc  $E$  and let  $\theta$  and  $\varphi$  be analytic in a domain  $D$  containing  $q(E)$ . Suppose that*

(i)  $\operatorname{Re} \frac{\theta'(q(z))}{\varphi(q(z))} > 0$  for  $z \in E$ ,

(ii)  $h(z) = zq'(z)\varphi(q(z))$  is starlike in  $E$ .

If  $p \in H[q(0), 1] \cap Q$  with  $p(E) \subseteq D$ ,  $\theta(p(z) + zp'(z))\varphi(p(z))$  is univalent in  $E$  and

$$\theta(q(z) + zq'(z)\varphi(q(z))) \prec \theta(p(z) + zp'(z)\varphi(p(z))), \quad (7)$$

then  $q(z) \prec p(z)$  and  $q$  is the best subdominant of (7).

**Lemma 2.5** (Miller and Mocanu [10]). *Let  $q$  be convex in  $E$  and let  $\gamma \in \mathbb{C}$ , with  $\operatorname{Re} \gamma > 0$ . If  $p \in H[q(0), 1] \cap Q$  and  $p(z) + \gamma zp'(z)$  is univalent in  $E$ , then*

$$q(z) + \gamma zq'(z) \prec p(z) + \gamma zp'(z), \quad (8)$$

implies  $q(z) \prec p(z)$ , and  $q$  is the best subdominant of (8).

**Lemma 2.6** (Royster [11]). *The function  $q(z) = \frac{1}{(1-z)^{2ab}}$  is univalent in  $E$  if and only if  $|2ab - 1| \leq 1$  or  $|2ab + 1| \leq 1$ .*

### 3. Main Results

**Theorem 3.1.** *Let  $q$  be univalent in  $E$ , with  $q(0) = 1$ , and suppose that*

$$\operatorname{Re} \left( 1 + \frac{zq''(z)}{q(z)} \right) > \max \left\{ 0; -(n+1)\operatorname{Re} \frac{1}{\lambda} \right\}, \quad n \in \mathbb{N}_0, z \in E, \quad (9)$$

where  $\lambda \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$ . If  $f \in \mathcal{A}$  satisfies the subordination

$$(1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right) \prec q(z) + \frac{\lambda}{(n+1)} zq'(z), \quad (10)$$

then

$$\left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) \prec q(z),$$

and  $q$  is the best dominant of (10).

*Proof.* Set

$$\left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) = h(z),$$

where  $h(z)$  is analytic in  $E$  with  $h(0) = 1$ .

A simple computation along with (4) shows that

$$(1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right) = h(z) + \frac{\lambda}{(n+1)} zh'(z),$$

hence the subordination (10) is equivalent to

$$h(z) + \frac{\lambda}{(n+1)} zh'(z) \prec q(z) + \frac{\lambda}{(n+1)} zq'(z).$$

Combining this last relation together with Lemma 2.3 for special case  $\gamma = \frac{\lambda}{(n+1)}$  and  $\mu = 1$ , we obtain our result.  $\square$

Taking  $q(z) = \frac{1+Az}{1+Bz}$  in Theorem 3.1, where  $-1 \leq B < A \leq 1$ , the condition (9) reduces to

$$\operatorname{Re} \frac{1-Bz}{1+Bz} > \max \left\{ 0; -(n+1)\operatorname{Re} \frac{1}{\lambda} \right\}, \quad n \in \mathbb{N}_0 \quad z \in E. \quad (11)$$

It is easy to verify that the function  $\varphi(\zeta) = \frac{(1-\zeta)}{(1+\zeta)}$ ,  $|\zeta| < B$ , is convex in  $E$ , and since  $\varphi(\bar{\zeta}) = \overline{\varphi(\zeta)}$  for all  $|\zeta| < |B|$ , it follows that  $\varphi(E)$  is a convex domain symmetric with respect to the real axis, hence

$$\inf \left\{ \operatorname{Re} \frac{1 - Bz}{1 + Bz} : z \in E \right\} = \frac{1 - |B|}{1 + |B|} > 0. \quad (12)$$

Then, the inequality (11) is equivalent to

$$(n+1) \operatorname{Re} \frac{1}{\lambda} \geq \frac{|B| - 1}{|B| + 1},$$

hence we have the following result.

**Corollary 3.2.** Let  $\lambda \in \mathbb{C}^*$ ,  $n \in \mathbb{N}_0$ ,  $-1 \leq B < A \leq 1$  with

$$\max \left\{ 0; -(n+1) \operatorname{Re} \frac{1}{\lambda} \right\} \leq \frac{1 - |B|}{1 + |B|}.$$

If  $f \in \mathcal{A}$ , and

$$\begin{aligned} (1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right) \\ \prec \frac{1 + Az}{1 + Bz} + \frac{\lambda}{(n+1)} \frac{(A-B)z}{(1+Bz)^2}, \end{aligned} \quad (13)$$

then

$$\left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) \prec \frac{1 + Az}{1 + Bz},$$

and  $\frac{1 + Az}{1 + Bz}$  is the best dominant of (13).

**Example 3.3.** Let  $A = 1$ ,  $B = -1$ ,  $n \in \mathbb{N}_0$ ,  $\lambda \in \mathbb{C}^*$  with  $(n+1) \operatorname{Re} \frac{1}{\lambda} \geq 0$ . If  $f \in \mathcal{A}$  and

$$\begin{aligned} (1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right) \\ \prec \frac{1+z}{1-z} + \frac{\lambda}{(n+1)} \frac{2z}{(1-z)^2}, \end{aligned} \quad (14)$$

then

$$\left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) \prec \frac{1+z}{1-z},$$

and  $\frac{1+z}{1-z}$  is the best dominant of (14).

**Theorem 3.4.** *Let  $q$  be univalent in  $E$ , with  $q(0) = 1$  and  $q(z) \neq 0$  for all  $z \in E$ . Let  $\gamma, \mu \in \mathbb{C}^*$  and  $\nu, \eta \in \mathbb{C}$ , with  $\nu + \eta \neq 0$ . Let  $f \in \mathcal{A}$  and suppose that  $f$  and  $q$  satisfy the following conditions:*

$$\frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(\nu + \eta)z} \neq 0, \quad n \in \mathbb{N}_0, \quad z \in E, \quad (15)$$

and

$$\operatorname{Re} \left( 1 + \frac{zq''(z)}{q'(z)} - \frac{zq'(z)}{q(z)} \right) > 0, \quad z \in E. \quad (16)$$

If

$$1 + \gamma \mu \left[ \frac{\nu z(I_n f(z) - I_n f(-z))' + \eta z(I_{n+1} f(z) - I_{n+1} f(-z))'}{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))} - 1 \right] \prec 1 + \gamma \frac{zq'(z)}{q(z)}, \quad (17)$$

then

$$\left[ \frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(\nu + \eta)z} \right]^\mu \prec q(z),$$

and  $q$  is the best dominant of (17). The power is the principal one.

*Proof.* We begin by setting

$$\left[ \frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(\nu + \eta)z} \right]^\mu = h(z), \quad z \in E, \quad (18)$$

where  $h(z)$  is analytic in  $E$  with  $h(0) = 1$ . Differentiating Equation (18) logarithmically with respect to  $z$ , we have

$$\mu \left[ \frac{\nu z(I_n f(z) - I_n f(-z))' + \eta z(I_{n+1} f(z) - I_{n+1} f(-z))'}{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))} - 1 \right] = \frac{zh'(z)}{h(z)}.$$

To prove our result we use Lemma 2.2. Consider in this lemma

$$\theta(w) = 1 \text{ and } \varphi(w) = \frac{\gamma}{w},$$

then  $\theta$  is analytic in  $\mathbb{C}$  and  $\varphi(w) \neq 0$  is analytic in  $\mathbb{C}^*$ . Also, if we let

$$Q(z) = zq'(z)\varphi(q(z)) = \gamma \frac{zq'(z)}{q(z)},$$

and

$$g(z) = \theta(q(z)) + Q(z) = 1 + \gamma \frac{zq'(z)}{q(z)},$$

then, since  $Q(0) = 0$  and  $Q'(0) \neq 0$ , the assumption (16) would yield that  $Q$  is a starlike function in  $E$ . From (16), we have

$$\operatorname{Re} \frac{zg'(z)}{Q(z)} = \operatorname{Re} \left( 1 + \frac{zq''(z)}{q'(z)} - \frac{zq'(z)}{q(z)} \right) > 0, \quad z \in E,$$

and by using Lemma 2.2 we deduce that the subordination (17) implies that  $h(z) \prec q(z)$ , and the function  $q$  is the best dominant of (17).  $\square$

In particular,  $\nu = 0$ ,  $\eta = \gamma = 1$  and  $q(z) = \frac{1+Az}{1+Bz}$  in the above Theorem 3.4, it is easy to see that the assumption (16) holds whenever  $-1 \leq A < B \leq 1$ , which leads to the next result:

**Corollary 3.5.** *Let  $-1 \leq A < B \leq 1$ ,  $n \in \mathbb{N}_0$ , and  $\mu \in \mathbb{C}^*$ . Let  $f \in \mathcal{A}$  and suppose that  $\frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \neq 0$ ,  $z \in E$ . If*

$$1 + \mu \left[ \frac{z(I_{n+1}f(z) - I_{n+1}f(-z))'}{(I_{n+1}f(z) - I_{n+1}f(-z))} - 1 \right] \prec 1 + \frac{(A-B)z}{(1+Az)(1+Bz)}, \quad (19)$$

then

$$\left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right)^\mu \prec \frac{1+Az}{1+Bz},$$

and  $\frac{1+Az}{1+Bz}$  is the best dominant of (19). The power is the principal one.

Putting  $\nu = n = 0$ ,  $\eta = 1$ ,  $\gamma = \frac{1}{ab}$ ,  $a, b \in \mathbb{C}^*$ ,  $\mu = a$ , and  $q(z) = \frac{1}{(1-z)^{2ab}}$  in Theorem 3.4, then combining this together with Lemma 2.6, we have the next result, see [1].

**Corollary 3.6.** (see [1]) *Let  $a, b \in \mathbb{C}^*$  such that  $|2ab - 1| \leq 1$  or  $|2ab + 1| \leq 1$ . Let  $f \in \mathcal{A}$  and let  $\frac{f(z) - f(-z)}{2z} \neq 0$  for all  $z \in E$ . If*

$$1 + \frac{1}{b} \left( \frac{z(f'(z) + f'(-z))}{(f(z) - f(-z))} - 1 \right) \prec \frac{1+z}{1-z},$$

then

$$\left( \frac{f(z) - f(-z)}{2z} \right)^a \prec \frac{1}{(1-z)^{2ab}}, \quad (20)$$

and  $\frac{1}{(1-z)^{2ab}}$  is the best dominant of (20). The power is the principal one.

Putting  $\nu = n = 0$ ,  $\eta = \gamma = 1$  and  $q(z) = (1+Bz)^{\frac{\mu(A-B)}{B}}$ ,  $-1 \leq B < A \leq 1$ ,  $B \neq 0$  in Theorem 3.4, and using Lemma 2.6, we have the next result.



**Corollary 3.7.** [1]. Let  $-1 \leq B < A \leq 1$  with  $B \neq 0$ , and suppose that  $\left| \frac{\mu(A-B)}{B-1} \right| \leq 1$  or  $\left| \frac{\mu(A-B)}{B+1} \right| \leq 1$ . Let  $f \in \mathcal{A}$  such that  $\frac{(f(z)-f(-z))}{2z} \neq 0$  for all  $z \in E$ , and let  $\mu \in \mathbb{C}^*$ . If

$$1 + \mu \left( \frac{z(f'(z) + f'(-z))}{(f(z) - f(-z))} - 1 \right) \prec \frac{1 + [B + \mu(A - B)]z}{1 + Bz}, \tag{21}$$

then

$$\left( \frac{(f(z) - f(-z))}{2z} \right)^\mu \prec (1 + Bz)^{\frac{\mu(A-B)}{B}},$$

and  $(1 + Bz)^{\frac{\mu(A-B)}{B}}$  is the best dominant of (21). Here the power is the principal one.

By taking  $\nu = n = 0$ ,  $\eta = 1$ ,  $\gamma = \frac{e^{i\lambda}}{ab \cos \lambda}$ ,  $a, b \in \mathbb{C}^*$ ,  $|\lambda| < \frac{\pi}{2}$ ,  $\mu = a$  and  $q(z) = \frac{1}{(1-z)^{2ab \cos \lambda e^{-i\lambda}}}$  in Theorem 3.4, we obtain the following result.

**Corollary 3.8** ([1]). Let  $a, b \in \mathbb{C}^*$ ,  $|\lambda| < \frac{\pi}{2}$ , suppose  $|ab \cos \lambda e^{-i\lambda} - 1| \leq 1$  or  $|ab \cos \lambda e^{-i\lambda} + 1| \leq 1$ . Let  $f \in \mathcal{A}$  such that  $\frac{(f(z)-f(-z))}{2z} \neq 0$  for all  $z \in E$ . If

$$1 + \frac{e^{i\lambda}}{b \cos \lambda} \left( \frac{z(f'(z) + f'(-z))}{(f(z) - f(-z))} - 1 \right) \prec \frac{1+z}{1-z}, \tag{22}$$

then

$$\left( \frac{(f(z) - f(-z))}{2z} \right)^a \prec \frac{1}{(1-z)^{2ab \cos \lambda e^{-i\lambda}}},$$

and  $\frac{1}{(1-z)^{2ab \cos \lambda e^{-i\lambda}}}$  is the best dominant of (22). The power is the principal one.

**Theorem 3.9.** Let  $q$  be univalent in  $E$  with  $q(0) = 1$ , let  $\mu, \gamma \in \mathbb{C}^*$ , and let  $\mu, \gamma \in \mathbb{C}^*$  and let  $\delta, \Omega, \nu, \eta \in \mathbb{C}$  with  $\nu + \eta \neq 0$ . Let  $f \in \mathcal{A}$  and suppose that  $f$  and  $q$  satisfy the next two conditions:

$$\frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(\nu + \eta)z} \neq 0, \quad n \in \mathbb{N}_0, \quad z \in E, \tag{23}$$

and

$$\operatorname{Re} \left( 1 + \frac{z q''(z)}{q'(z)} \right) > \max \left\{ 0; -\operatorname{Re} \frac{\delta}{\gamma} \right\}, \quad z \in E. \tag{24}$$

If

$$\begin{aligned} \psi(z) \equiv & \left[ \frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(\nu + \eta)z} \right]^\mu \\ & \left[ \delta + \gamma \mu \left( \frac{\nu z(I_n f(z) - I_n f(-z))' + \eta z(I_{n+1} f(z) - I_{n+1} f(-z))'}{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))} - 1 \right) \right] + \Omega, \end{aligned} \tag{25}$$

and

$$\psi(z) \prec \delta q(z) + \gamma z q'(z) + \Omega, \quad (26)$$

then

$$\left[ \frac{v(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(v + \eta)z} \right]^\mu \prec q(z),$$

and  $q$  is the best dominant of (26). All the powers are the principal ones.

*Proof.* We begin by setting

$$\left[ \frac{v(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(v + \eta)z} \right]^\mu = h(z). \quad (27)$$

Then  $h(z)$  is analytic in  $E$  with  $h(0) = 1$ . Logarithmic differentiating of (27) yields

$$\mu \left( \frac{vz(I_n f(z) - I_n f(-z))' + \eta z(I_{n+1} f(z) - I_{n+1} f(-z))'}{v(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))} - 1 \right) = \frac{zh'(z)}{h(z)},$$

and hence

$$\mu h(z) \left( \frac{vz(I_n f(z) - I_n f(-z))' + \eta z(I_{n+1} f(z) - I_{n+1} f(-z))'}{v(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))} - 1 \right) = zh'(z).$$

Let us consider the functions:

$$\begin{aligned} \theta(w) &= \delta w + \Omega, \quad \varphi(w) = \gamma, \quad w \in \mathbb{C}, \\ Q(z) &= zq'(z)\varphi(q(z)) = \gamma zq'(z), \quad z \in E, \end{aligned}$$

and

$$g(z) = \theta(q(z)) + Q(z) = \delta q(z) + \gamma zq'(z) + \Omega, \quad z \in E.$$

From the assumption (24) we see that  $Q$  is starlike in  $E$  and, that

$$\operatorname{Re} \frac{zg'(z)}{Q(z)} = \operatorname{Re} \left( \frac{\delta}{\gamma} + 1 + \frac{zq''(z)}{q'(z)} \right) > 0, \quad z \in E,$$

thus, by applying Lemma 2.2, this completes the proof.  $\square$

Taking  $q(z) = \frac{(1+Az)}{(1+Bz)}$  in Corollary 3.7, where  $-1 \leq B < A \leq 1$  and according to (12), the condition (24) becomes

$$\max \left\{ 0; -\operatorname{Re} \frac{\delta}{\gamma} \right\} \leq \frac{1 - |B|}{1 + |B|}.$$

Hence, for the special case  $v = 1 = \gamma$ ,  $\eta = 0$ , we have the following result:

**Corollary 3.10.** *Let  $-1 \leq B < A \leq 1$  and let  $\delta \in \mathbb{C}$  with*

$$\max \{0; -\operatorname{Re} \delta\} \leq \frac{1 - |B|}{1 + |B|}.$$

*Let  $f \in \mathcal{A}$  and suppose that*

$$\frac{(I_n f(z) - I_n f(-z))}{2z} \neq 0, \quad n \in \mathbb{N}_0, z \in E$$

*and let  $\mu \in \mathbb{C}^*$ . If*

$$\begin{aligned} \left[ \frac{(I_n f(z) - I_n f(-z))}{2z} \right]^\mu & \left[ \delta + \mu \left( \frac{z(I_{n+1} f(z) - I_{n+1} f(-z))'}{(I_{n+1} f(z) - I_{n+1} f(-z))} - 1 \right) \right] + \Omega \\ & \prec \delta \frac{1 + Az}{1 + Bz} + \Omega + \frac{z(A - B)}{(1 + Bz)^2}, \end{aligned} \quad (28)$$

*then*

$$\left( \frac{(I_n f(z) - I_n f(-z))}{2z} \right)^\mu \prec \frac{1 + Az}{1 + Bz},$$

*and  $\frac{1+Az}{1+Bz}$  is the best dominant of (28). All the powers are the principal ones.*

Taking  $\gamma = \eta = 1, v = n = 0$  and  $q(z) = \frac{1+z}{1-z}$  in Corollary 3.7, we obtain the next result.

**Corollary 3.11** ([1]). *Let  $f \in \mathcal{A}$  such that  $\frac{f(z) - f(-z)}{2z} \neq 0$  for all  $z \in E$ , and let  $\mu \in \mathbb{C}^*$ . If*

$$\begin{aligned} \left[ \frac{f(z) - f(-z)}{2z} \right]^\mu & \left[ \delta + \mu \left( \frac{z(f'(z) + f'(-z))}{f(z) - f(-z)} - 1 \right) \right] + \Omega \\ & \prec \delta \frac{1+z}{1-z} + \Omega + \frac{2z}{(1-z)^2}, \end{aligned} \quad (29)$$

*then*

$$\left[ \frac{f(z) - f(-z)}{2z} \right]^\mu \prec \frac{1+z}{1-z},$$

*and  $\frac{1+z}{1-z}$  is the best dominant of (29). All the powers are the principal ones.*

#### 4. Superordination and Sandwich results

**Theorem 4.1.** *Let  $q$  be convex in  $E$  with  $q(0) = 1, n \in \mathbb{N}_0$ , let  $\lambda \in \mathbb{C}^*$  with  $(n+1)\operatorname{Re} \lambda > 0$ . Let  $f \in \mathcal{A}$  and suppose that  $\frac{I_{n+1} f(z) - I_{n+1} f(-z)}{2z} \in H[q(0), 1] \cap \mathcal{Q}$ . If the function*

$$(1 - \lambda) \left( \frac{I_{n+1} f(z) - I_{n+1} f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right)$$

is univalent in the unit disc  $E$  and

$$q(z) + \frac{\lambda}{(n+1)}zq'(z) < (1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right), \quad (30)$$

then

$$q(z) < \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right),$$

and  $q$  is the best subordinator of (30).

*Proof.* Set

$$\left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) = h(z), \quad z \in E.$$

Then  $h(z)$  is analytic in  $E$  with  $h(0) = 1$ . Taking logarithmic differentiation with respect  $z$ , we have

$$z \left( \frac{(I_{n+1}f(z))' - (I_{n+1}f(-z))'}{I_{n+1}f(z) - I_{n+1}f(-z)} \right) - 1 = \frac{zh'(z)}{h(z)}. \quad (31)$$

A simple computation along with identity (4) shows that

$$\begin{aligned} h(z) + \frac{\lambda}{(n+1)}zh'(z) &= (1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right), \end{aligned}$$

and now, by using Lemma 2.5, we obtain the desired result.  $\square$

Taking  $q(z) = \frac{1+Az}{1+Bz}$  in Theorem 4.1, where  $-1 \leq B < A \leq 1$ , we obtain the next result.

**Corollary 4.2.** *Let  $q$  be convex in  $E$  with  $q(0) = 1$ ,  $n \in \mathbb{N}_0$ , let  $\lambda \in \mathbb{C}^*$  with  $(n+1)\operatorname{Re}\lambda > 0$ . Let  $f \in \mathcal{A}$  and suppose that  $\frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \in H[q(0), 1] \cap \mathcal{Q}$ . If the function*

$$(1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right)$$

is univalent in the unit disc  $E$ , and

$$\begin{aligned} \frac{1+Az}{1+Bz} + \frac{\lambda(A-B)z}{(1+Bz)^2} < (1-\lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right), \quad (32) \end{aligned}$$

then

$$\frac{1 + Az}{1 + Bz} \prec \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right),$$

and  $\frac{1+Az}{1+Bz}$  is the best subordinant of (32), where  $-1 \leq B < A \leq 1$ .

Using the same techniques as we used in Theorem 3.9, and then applying Lemma 2.4, we have the following theorem.

**Theorem 4.3.** *Let  $q$  be convex in  $E$  with  $q(0) = 1$ , let  $\mu, \gamma \in \mathbb{C}^*$ , and let  $\delta, \Omega, v, \eta \in \mathbb{C}$  with  $v + \eta \neq 0$  and  $\operatorname{Re} \frac{\delta}{\gamma} > 0$ . Let  $f \in \mathcal{A}$  and suppose that  $f$  satisfies the following conditions:*

$$\frac{v(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(v + \eta)z} \neq 0, \quad n \in \mathbb{N}_0, z \in E,$$

and

$$\left[ \frac{v(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(v + \eta)z} \right]^\mu \in H[q(0), 1] \cap \mathcal{Q}.$$

If the function  $\psi$  given by equation (25) is univalent in  $E$ , and

$$\delta q(z) + \gamma z q'(z) + \Omega \prec \psi(z), \tag{33}$$

then

$$q(z) \prec \left[ \frac{v(I_n f(z) - I_n f(-z)) + \eta(I_{n+1} f(z) - I_{n+1} f(-z))}{(v + \eta)z} \right]^\mu,$$

and  $q$  is the best subordinate of (33) (all powers are the principal ones).

Note that by combining Theorem 3.1 with Theorem 4.1 and Corollary 3.10 with Theorem 4.3, we have, respectively, the following two sandwich theorems:

**Theorem 4.4.** *Let  $q_1$  and  $q_2$  be two convex functions in  $E$  with  $q_1(0) = q_2(0) = 1$ ,  $n \in \mathbb{N}_0$ , let  $\lambda \in \mathbb{C}^*$  with  $(n + 1)\operatorname{Re} \lambda > 0$ . Let  $f \in \mathcal{A}$  and suppose that  $\frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \in H[q(0), 1] \cap \mathcal{Q}$ . If the function*

$$(1 - \lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right)$$

is univalent in the unit disc  $E$ , and

$$q_1(z) + \frac{\lambda}{(n + 1)} z q_1'(z) \prec (1 - \lambda) \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) + \lambda \left( \frac{I_n f(z) - I_n f(-z)}{2z} \right) \prec q_2(z) + \frac{\lambda}{(n + 1)} z q_2'(z), \tag{34}$$

then

$$q_1(z) \prec \left( \frac{I_{n+1}f(z) - I_{n+1}f(-z)}{2z} \right) \prec q_2(z),$$

and  $q_1$  and  $q_2$  are, respectively, the best subordinate and the best dominant of (34).

**Theorem 4.5.** Let  $q_1$  and  $q_2$  be two convex functions in  $E$  with  $q_1(0) = q_2(0) = 1$ ,  $n \in \mathbb{N}_0$ , let  $\mu, \gamma \in \mathbb{C}^*$ , and let  $\delta, \Omega, \nu, \eta \in \mathbb{C}$  with  $\nu + \eta \neq 0$  and  $\operatorname{Re} \frac{\delta}{\gamma} > 0$ . Let  $f \in \mathcal{A}$  satisfy the following conditions:

$$\left[ \frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1}f(z) - I_{n+1}f(-z))}{(\nu + \eta)z} \right] \neq 0, z \in E,$$

and

$$\left[ \frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1}f(z) - I_{n+1}f(-z))}{(\nu + \eta)z} \right]^\mu \in H[q(0), 1] \cap \mathcal{Q}.$$

If the function  $\psi$  given by (25) is univalent in  $E$  and

$$\delta q_1(z) + \gamma z q_1'(z) + \Omega \prec \psi(z) \prec \delta q_2(z) + \gamma z q_2'(z) + \Omega, \quad (35)$$

then

$$q_1(z) \prec \left[ \frac{\nu(I_n f(z) - I_n f(-z)) + \eta(I_{n+1}f(z) - I_{n+1}f(-z))}{(\nu + \eta)z} \right]^\mu \prec q_2(z),$$

and  $q_1$  and  $q_2$  are, respectively, the best subordinate and the best dominant of (35) (all the powers are the principal ones).

## Acknowledgements

We are grateful for the help to the Editorial Board and Referees.

## REFERENCES

- [1] A. Muhammad, *Some differential subordination and superordination properties of symmetric functions*, Rend. Sem. Mat. Univ. Poletic. Torino (to appear).
- [2] M. K. Aouf - T. Bulboacă, *Subordination and superordination properties of multivalent functions defined by certain integral operator*, J. Franklin Inst, 347 (2010), 641–653.

- [3] M. K. Aouf - F. M. Al-Oboudi - M. M. Haidan, *On some results for  $\lambda$ -spirallike and  $\lambda$ -Robertson functions of complex order*, Publ. Inst. Math. Belgrade 77 (91) (2005), 93–98.
- [4] T. Bulboacă, *A class of superordination preserving integral operators*, Indag. Math. (New Ser.) 13 (3) (2002), 301–311.
- [5] T. Bulboacă, *Classes of first-order differential subordination*, Demonstratio Math. 35 (2) (2002), 287–292.
- [6] T. Bulboacă, *Differential subordination and superordinations. Recent Results*, House of Scientific Book Publ, Cluj-Napoca, 2005.
- [7] J. L. Liu, *The Noor integral and strongly starlike functions*, J. Math. Anal. Appl. 261 (2001), 441–447.
- [8] J. L. Liu - K. I. Noor, *Some properties of Noor integral operator*, J. Nat. Geom. 21 (2002), 81–90.
- [9] S. S. Miller - P. T. Mocanu, *Differential subordination Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics vol. 225, Marcel Dekker Inc., New York, Basel, 2000.
- [10] S. S. Miller - P. T. Mocanu, *Subordinations of differential superordinations*, Complex Variables 48 (10) (2003), 815–826.
- [11] K. I. Noor, *On new classes of integral operators*, J. Natur. Geom. 16 (1999), 71–80.
- [12] K. I. Noor - M. A. Noor, *On integral operators*, J. Math. Anal. Appl. 238 (1999), 341–352.
- [13] M. Obradovic - M. K. Aouf - S. Owa, *On some results for starlike functions of complex order*, Pub. Inst. Math. Belgrade 46 (60) (1989), 79–85.
- [14] M. Obradovic - S. Owa, *On certain properties for some classes of starlike functions*, J. Math. Anal. Appl. 145 (2) (1990), 357–364.
- [15] W. C. Royster, *On the univalence of a certain integral*, Michigan Math. J. 12 (1965), 385–387.
- [16] S. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc. 49 (1975), 109–115.
- [17] K. Sakaguchi, *On a certain univalent mapping*, J. Math. Soc. Japan, 11 (1959), 72–75.
- [18] S. Shams - S. R. Kulkarni - J. M. Jahangiri, *Subordination properties of  $P$ -valent functions defined by integral operators*, Int. J. Math. Math. Sci. (2006), Article ID 94572, 1–3.
- [19] T. N. Shanmugam - C. Ramachandran - M. Darus - S. Sivasubbramanian, *Differential sandwich theorems for some subclasses of analytic functions involving a linear operator*, Acta Math. Univ. Comenianae 74 (2) (2007), 287–294.
- [20] T. N. Shanmugam - S. Sivasubbramanian - H. M. Srivastava, *On sandwich theorems for some classes of analytic functions*, Int. J. Math. Math. Sci. (2006), Article ID 29684 1–13.

*ALI MUHAMMAD*  
*Department of Basic Sciences*  
*University of Engineering and Technology*  
*Peshawar, Pakistan*  
*e-mail: ali7887@gmail.com*

*AMJADULLAH KHATTAK*  
*Department of Electrical and Electronic Engineering*  
*University of Engineering and Technology*  
*Peshawar, Pakistan*  
*e-mail: amjad67@gmail.com*