

LE MATEMATICHE

Vol. LXVIII (2013) – Fasc. I, pp. 87–98

doi: 10.4418/2013.68.1.8

COMMON FIXED POINT THEOREMS FOR WEAKLY COMPATIBLE MAPPINGS IN FUZZY METRIC SPACES

S. CHAUHAN - S. BHATNAGAR - S. RADENović

The aim of this paper is to prove a common fixed point theorem for a pair of weakly compatible mappings in fuzzy metric space by using the (CLR_g) property. An example is also furnished which demonstrates the validity of our main result. As an application to our main result, we present a fixed point theorem for two finite families of self mappings in fuzzy metric space by using the notion of pairwise commuting. Our results improve the results of Sedghi, Shobe and Aliouche [31].

1. Introduction

The concept of a fuzzy set is investigated by Zadeh [40] in his seminal paper. In 1975, Kramosil and Michalek [14] introduced the concept of fuzzy metric space, which opened an avenue for further development of analysis in such spaces. Further, George and Veeramani [8] modified the concept of fuzzy metric space introduced by Kramosil and Michalek [14] with a view to obtain a Hausdorff topology which has very important applications in quantum particle physics, particularly in connection with both string and ε^∞ theory (see, [21–23]). Fuzzy set theory also has applications in applied sciences such as neural network theory, stability theory, mathematical programming, modeling theory,

Entrato in redazione: 21 maggio 2012

AMS 2010 Subject Classification: 54H25, 47H10.

Keywords: Fuzzy metric space, Weakly compatible mappings, (E.A) property, (CLR_g) property, Fixed point.

engineering sciences, medical sciences (medical genetics, nervous system), image processing, control theory, communication etc. Consequently in due course of time some metric fixed point results were generalized to fuzzy metric spaces by various authors viz Grabiec [9], Cho [6, 7], Subrahmanyam [38] and Vasuki [39].

In 2002, Aamri and El-Moutawakil [2] defined the notion of (E.A) property for self mappings which contained the class of non-compatible mappings in metric spaces. It was pointed out that (E.A) property allows replacing the completeness requirement of the space with a more natural condition of closedness of the range as well as relaxes the completeness of the whole space, continuity of one or more mappings and containment of the range of one mapping into the range of other which is utilized to construct the sequence of joint iterates. Many authors have proved common fixed point theorems in fuzzy metric spaces for different contractive conditions. For details, we refer to [5, 10, 11, 15–17, 20, 24–26, 29, 30, 32–34, 36, 37]. Recently, Sintunavarat and Kumam [35] defined the notion of (CLRg) property in fuzzy metric spaces and improved the results of Miheţ [18] without any requirement of the closedness of the subspace.

In this paper, we prove a common fixed point theorem for a pair of weakly compatible mappings by using (CLRg) property in fuzzy metric space. We also present a common fixed point theorem for two finite families of self mappings in fuzzy metric space by using the notion of pairwise commuting due to Imdad, Ali and Tanveer [12]. Our results improve the results of Sedghi, Shobe and Aliouche [31].

2. Preliminaries

Definition 2.1. [28] A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is a continuous t-norm if it satisfies the following conditions:

1. $*$ is associative and commutative,
2. $*$ is continuous,
3. $a * 1 = a$ for all $a \in [0, 1]$,
4. $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$ for all $a, b, c, d \in [0, 1]$.

Examples of continuous t-norms are $a * b = ab$ and $a * b = \min\{a, b\}$.

Definition 2.2. [8] A 3-tuple $(X, M, *)$ is said to be a fuzzy metric space if X is an arbitrary set, $*$ is a continuous t-norm and M is a fuzzy set on $X^2 \times (0, \infty)$ satisfying the following conditions: for all $x, y, z \in X, t, s > 0$,

1. $M(x, y, t) > 0$,
2. $M(x, y, t) = 1$ if and only if $x = y$,
3. $M(x, y, t) = M(y, x, t)$,
4. $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$,
5. $M(x, y, \cdot) : [0, \infty) \rightarrow [0, 1]$ is continuous.

Then M is called a fuzzy metric on X . Then $M(x, y, t)$ denotes the degree of nearness between x and y with respect to t .

Let $(X, M, *)$ be a fuzzy metric space. For $t > 0$, the open ball $\mathcal{B}(x, r, t)$ with center $x \in X$ and radius $0 < r < 1$ is defined by

$$\mathcal{B}(x, r, t) = \{y \in X : M(x, y, t) > 1 - r\}.$$

Now let $(X, M, *)$ be a fuzzy metric space and τ the set of all $A \subset X$ with $x \in A$ if and only if there exist $t > 0$ and $0 < r < 1$ such that $\mathcal{B}(x, r, t) \subset A$. Then τ is a topology on X induced by the fuzzy metric M .

In the following example (see [8]), we know that every metric induces a fuzzy metric:

Example 2.3. Let (X, d) be a metric space. Denote $a * b = ab$ (or $a * b = \min\{a, b\}$) for all $a, b \in [0, 1]$ and let M_d be fuzzy sets on $X^2 \times (0, \infty)$ defined as follows:

$$M_d(x, y, t) = \frac{t}{t + d(x, y)}.$$

Then $(X, M_d, *)$ is a fuzzy metric space and the fuzzy metric M induced by the metric d is often referred to as the standard fuzzy metric.

Lemma 2.4. [9] *Let $(X, M, *)$ be a fuzzy metric space. Then $M(x, y, t)$ is non-decreasing for all $x, y \in X$.*

Definition 2.5. [13] Two self mappings f and g of a non-empty set X are said to be weakly compatible (or coincidentally commuting) if they commute at their coincidence points, i.e. if $fz = gz$ some $z \in X$, then $fgz = gfgz$.

Remark 2.6. [13] Two compatible self mappings are weakly compatible, but the converse is not true. Therefore the concept of weak compatibility is more general than that of compatibility.

Definition 2.7. [3] A pair of self mappings f and g of a fuzzy metric space $(X, M, *)$ is said to satisfy the (E.A) property, if there exists a sequence $\{x_n\}$ in X for some $z \in X$ such that

$$\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = z.$$

Remark 2.8. It is noted that weak compatibility and (E.A) property are independent to each other (see [27], Example 2.1, Example 2.2).

Definition 2.9. [3] Two self mappings f and g of a fuzzy metric space $(X, M, *)$ are non-compatible if and only if there exists at least one sequence $\{x_n\}$ in X such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = z$ for some $z \in X$, but for some $t > 0$, $\lim_{n \rightarrow \infty} M(fgx_n, gfx_n, t)$ is either less than 1 or nonexistent.

Remark 2.10. From Definition 2.9, it is easy to see that any non-compatible self mappings of a fuzzy metric space $(X, M, *)$ satisfy the (E.A) property. But two mappings satisfying the (E.A) property need not be non-compatible (see [27], Remark 4.8).

Definition 2.11. [35] A pair of self mappings f and g of a fuzzy metric space $(X, M, *)$ is said to satisfy the (CLRg) property if there exists a sequence $\{x_n\}$ in X such that

$$\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = gu,$$

for some $u \in X$.

Inspired by Sintunavarat and Kumam [35], we show examples of self mappings f and g which are satisfying the (CLRg) property.

Example 2.12. Let $(X, M, *)$ be a fuzzy metric space with $X = [0, \infty)$ and

$$M(x, y, t) = \begin{cases} \frac{t}{t + |x - y|}, & \text{if } t > 0; \\ 0, & \text{if } t = 0. \end{cases}$$

for all $x, y \in X$. Define self mappings f and g on X by $f(x) = x + 3$ and $g(x) = 4x$ for all $x \in X$. Let a sequence $\{x_n\} = \{1 + \frac{1}{n}\}_{n \in \mathbb{N}}$ in X , we have

$$\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = 4 = g(1) \in X,$$

which shows that f and g satisfy the (CLRg) property.

Example 2.13. The conclusion of Example 2.12 remains true if the self mappings f and g is defined on X by $f(x) = \frac{x}{7}$ and $g(x) = \frac{x}{8}$ for all $x \in X$. Let a sequence $\{x_n\} = \{\frac{1}{n}\}_{n \in \mathbb{N}}$ in X . Since

$$\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = 0 = g(0) \in X,$$

therefore f and g satisfy the (CLRg) property.

Definition 2.14. [12] Two families of self mappings $\{f_i\}_{i=1}^m$ and $\{g_k\}_{k=1}^n$ are said to be pairwise commuting if

1. $f_i f_j = f_j f_i$ for all $i, j \in \{1, 2, \dots, m\}$,
2. $g_k g_l = g_l g_k$ for all $k, l \in \{1, 2, \dots, n\}$,
3. $f_i g_k = g_k f_i$ for all $i \in \{1, 2, \dots, m\}$ and $k \in \{1, 2, \dots, n\}$.

Throughout this paper, $(X, M, *)$ is considered to be a fuzzy metric space with condition $\lim_{t \rightarrow \infty} M(x, y, t) = 1$ for all $x, y \in X$.

3. Results

In 2010, Sedghi, Shobe and Aliouche [31] proved a common fixed point theorem for a pair of weakly compatible mappings with (E.A) property in fuzzy metric space by using the following function:

Let Φ is a set of all increasing and continuous functions $\phi : (0, 1] \rightarrow (0, 1]$, such that $\phi(t) > t$ for every $t \in (0, 1)$.

Example 3.1. [31] Let $\phi : (0, 1] \rightarrow (0, 1]$ defined by $\phi(t) = t^{\frac{1}{2}}$.

Theorem 3.2. (Theorem 1, [31]) Let $(X, M, *)$ be a fuzzy metric space and f and g be self mappings of X satisfying the following conditions:

1. $f(X) \subseteq g(X)$ and $f(X)$ or $g(X)$ is a closed subset of X ,
- 2.

$$M(fx, fy, t) \geq \phi \left(\min \left\{ \begin{array}{l} M(gx, gy, t), \\ \sup_{t_1+t_2=\frac{2}{k}t} \min \left\{ \begin{array}{l} M(gx, fx, t_1), \\ M(gy, fy, t_2) \end{array} \right\}, \\ \sup_{t_3+t_4=\frac{2}{k}t} \max \left\{ \begin{array}{l} M(gx, fy, t_3), \\ M(gy, fx, t_4) \end{array} \right\} \end{array} \right\} \right) \quad (1)$$

for all $x, y \in X$, $t > 0$ and for some $1 \leq k < 2$. Suppose that the pair (f, g) satisfies the (E.A) property and (f, g) is weakly compatible. Then f and g have a unique common fixed point in X .

Now we prove our main result:

Theorem 3.3. Let $(X, M, *)$ be a fuzzy metric space, where $*$ is a continuous t -norm. Further let f, g be mappings from X into itself and satisfying the inequality (1) of Theorem 3.2. If the pair (f, g) satisfies the (CLRg) property then f and g have a unique common fixed point provided the pair (f, g) is weakly compatible.

Proof. Since the pair (f, g) satisfies the (CLRg) property, there exists a sequence $\{x_n\}$ in X such that

$$\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = gu,$$

for some $u \in X$. Now we assert that $fu = gu$. Let, on the contrary, $fu \neq gu$, then there exists $t_0 > 0$ such that

$$M\left(fu, gu, \frac{2}{k}t_0\right) > M(fu, gu, t_0). \quad (2)$$

To support the claim, let it be untrue. Then we have

$$M\left(fu, gu, \frac{2}{k}t\right) > M(fu, gu, t), \text{ for all } t > 0.$$

Repeatedly using this equality, we obtain

$$M(fu, gu, t) = M\left(fu, gu, \frac{2}{k}t\right) = \dots = M\left(fu, gu, \left(\frac{2}{k}\right)^n t\right) \rightarrow 1,$$

as $n \rightarrow \infty$. This shows that $M(fu, gu, t) = 1$ for all $t > 0$ which contradicts $fu \neq gu$ and hence (2) is proved. On using inequality (1), with $x = x_n$, $y = u$, we get

$$M(fx_n, fu, t_0) \geq \phi \left(\min \left\{ \begin{array}{l} M(gx_n, gu, t_0), \\ \sup_{t_1+t_2=\frac{2}{k}t_0} \min \left\{ \begin{array}{l} M(gx_n, fx_n, t_1), \\ M(gu, fu, t_2) \end{array} \right\}, \\ \sup_{t_3+t_4=\frac{2}{k}t_0} \max \left\{ \begin{array}{l} M(gx_n, fu, t_3), \\ M(gu, fx_n, t_4) \end{array} \right\} \end{array} \right\} \right)$$

for all $\varepsilon \in (0, \frac{2}{k}t_0)$. As $n \rightarrow \infty$, it follows that

$$\begin{aligned} M(gu, fu, t_0) &\geq \phi \left(\min \left\{ \begin{array}{l} M(gu, gu, t_0), \\ \min \{M(gu, gu, \varepsilon), M(gu, fu, \frac{2}{k}t_0 - \varepsilon)\}, \\ \max \{M(gu, fu, \frac{2}{k}t_0 - \varepsilon), M(gu, gu, \varepsilon)\} \end{array} \right\} \right) \\ &= \phi \left(M\left(gu, fu, \frac{2}{k}t_0 - \varepsilon\right) \right) > M\left(gu, fu, \frac{2}{k}t_0 - \varepsilon\right), \end{aligned}$$

as $\varepsilon \rightarrow 0$, we have

$$M(gu, fu, t_0) \geq M\left(gu, fu, \frac{2}{k}t_0\right),$$

which contradicts (2), we have $gu = fu$. Next, we let $z = fu = gu$. Since the pair (f, g) is weakly compatible, $fgu = gfu$ which implies that $fz = fgu = gfu =$

gz . Now we show that $z = fz$. Suppose that $z \neq fz$, then on using (1) with $x = z$, $y = u$, we get, for some $t_0 > 0$,

$$M(fz, fu, t_0) \geq \phi \left(\min \left\{ \begin{array}{l} M(gz, gu, t_0), \\ \sup_{t_1+t_2=\frac{2}{k}t_0} \min \left\{ \begin{array}{l} M(gz, fz, t_1), \\ M(gu, fu, t_2) \end{array} \right\}, \\ \sup_{t_3+t_4=\frac{2}{k}t_0} \max \left\{ \begin{array}{l} M(gz, fu, t_3), \\ M(gu, fz, t_4) \end{array} \right\} \end{array} \right\} \right), \\ M(fz, z, t_0) \geq \phi \left(\min \left\{ \begin{array}{l} M(fz, z, t_0), \\ \min \{M(fz, fz, \varepsilon), M(z, z, \frac{2}{k}t_0 - \varepsilon)\}, \\ \max \{M(fz, z, \varepsilon), M(z, fz, \frac{2}{k}t_0 - \varepsilon)\} \end{array} \right\} \right),$$

for all $\varepsilon \in (0, \frac{2}{k}t_0)$. As $\varepsilon \rightarrow 0$, we have

$$M(fz, z, t_0) \geq \phi \left(\min \left\{ M(fz, z, t_0), M\left(z, fz, \frac{2}{k}t_0\right) \right\} \right) \\ = \phi(M(fz, z, t_0)) > M(fz, z, t_0),$$

which is a contradiction. Hence $fz = gz = z$. Therefore z is a common fixed point of f and g .

Uniqueness: Let $w (\neq z)$ be another common fixed point of f and g . On using inequality (1) with $x = z$, $y = w$, we get, for some $t_0 > 0$,

$$M(fz, fw, t_0) \geq \phi \left(\min \left\{ \begin{array}{l} M(gz, gw, t_0), \\ \sup_{t_1+t_2=\frac{2}{k}t_0} \min \left\{ \begin{array}{l} M(gz, fz, t_1), \\ M(gw, fw, t_2) \end{array} \right\}, \\ \sup_{t_3+t_4=\frac{2}{k}t_0} \max \left\{ \begin{array}{l} M(gz, fw, t_3), \\ M(gw, fz, t_4) \end{array} \right\} \end{array} \right\} \right), \\ M(z, w, t_0) \geq \phi \left(\min \left\{ \begin{array}{l} M(z, w, t_0), \\ \min \{M(z, z, \varepsilon), M(w, w, \frac{2}{k}t_0 - \varepsilon)\}, \\ \max \{M(z, w, \varepsilon), M(w, z, \frac{2}{k}t_0 - \varepsilon)\} \end{array} \right\} \right),$$

for all $\varepsilon \in (0, \frac{2}{k}t_0)$. As $\varepsilon \rightarrow 0$, we have

$$M(z, w, t_0) \geq \phi \left(\min \left\{ M(z, w, t_0), M\left(w, z, \frac{2}{k}t_0\right) \right\} \right) \\ = \phi(M(z, w, t_0)) > M(z, w, t_0),$$

which is a contradiction. Therefore $Bz = z = Tz$. It implies that f and g have a unique a common fixed point. \square

Remark 3.4. Theorem 3.3 improves the main result of Sedghi, Shobe and Aliouche ([31], Theorem 1) without any requirement on containment of ranges amongst the involved mappings and closedness of one or more subspaces.

The following examples illustrates Theorem 3.3.

Example 3.5. Let $(X, M, *)$ be a fuzzy metric space, where $X = [3, 19)$, with t -norm $*$ is defined by $a * b = ab$ for all $a, b \in [0, 1]$ and

$$M(x, y, t) = \begin{cases} \frac{t}{t+|x-y|}, & \text{if } t > 0; \\ 0, & \text{if } t = 0. \end{cases}$$

for all $x, y \in X$. Let the function $\phi : (0, 1] \rightarrow (0, 1]$ defined by $\phi(t) = t^{\frac{1}{2}}$. Define the self mappings f and g by

$$f(x) = \begin{cases} 3, & \text{if } x \in \{3\} \cup (5, 19); \\ 12, & \text{if } x \in (3, 5]. \end{cases} \quad g(x) = \begin{cases} 3, & \text{if } x = 3; \\ 11, & \text{if } x \in (3, 5]; \\ \frac{x+1}{2}, & \text{if } x \in (5, 19). \end{cases}$$

Taking $\{x_n\} = \{5 + \frac{1}{n}\}_{n \in \mathbb{N}}$ or $\{x_n\} = \{3\}$, it is clear that the pair (f, g) satisfies the (CLRg) property.

$$\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = 3 = g(3) \in X.$$

It is noted that $f(X) = \{3, 12\} \not\subseteq [3, 10) \cup \{11\} = g(X)$. Thus, all the conditions of Theorem 3.3 are satisfied and 3 is a unique common fixed point of the pair (f, g) . Also, all the involved mappings are even discontinuous at their unique common fixed point 3. Here, it may be pointed out that $g(X)$ is not a closed subspace of X .

Now we utilize the notion of commuting pairwise due to Imdad, Ali and Tanveer [10] and extend Theorem 3.3 to two finite families of self mappings in fuzzy metric space.

Corollary 3.6. Let $(X, M, *)$ be a fuzzy metric space, where $*$ is a continuous t -norm. Further let $\{f_1, f_2, \dots, f_p\}$ and $\{g_1, g_2, \dots, g_q\}$ be two finite families of mappings from X into itself such that $f = f_1 f_2 \dots f_p$ and $g = g_1 g_2 \dots g_q$ and satisfy inequality (1) of Theorem 3.2. Suppose that the pair (f, g) shares the (CLRg) property.

Moreover, if the family $\{f_i\}_{i=1}^p$ commutes pairwise with the family $\{g_i\}_{i=1}^q$, then (for all $i \in \{1, 2, \dots, p\}$ and $j \in \{1, 2, \dots, q\}$) f_i and g_j have a unique common fixed point.

Proof. The proof of this theorem is similar to that of Theorem 3.1 contained in Imdad, Ali and Tanveer [12], hence details are avoided. \square

Remark 3.7. Corollary 3.6 improve the result of Sedghi, Shobe and Aliouche ([31], Theorem 2).

By setting $f_1 = f_2 = \dots = f_p = f$ and $g_1 = g_2 = \dots = g_q = g$ in Corollary 3.6, we deduce the following:

Corollary 3.8. *Let $(X, M, *)$ be a fuzzy metric space, where $*$ is a continuous t -norm. Further let f and g be mappings from X into itself such that the pair (f^p, g^q) satisfies the (CLRg) property such that*

$$M(f^p x, f^p y, t) \geq \phi \left(\min \left\{ \begin{array}{l} M(g^q x, g^q y, t), \\ \sup_{t_1+t_2=\frac{2}{k}t} \min \left\{ \begin{array}{l} M(g^q x, f^p x, t_1), \\ M(g^q y, f^p y, t_2) \end{array} \right\}, \\ \sup_{t_3+t_4=\frac{2}{k}t} \max \left\{ \begin{array}{l} M(g^q x, f^p y, t_3), \\ M(g^q y, f^p x, t_4) \end{array} \right\} \end{array} \right\} \right) \quad (3)$$

holds for all $x, y \in X$, $t > 0$, for some $1 \leq k < 2$ and p, q are fixed positive integers. Then f and g have a unique common fixed point provided that the pair (f^p, g^q) commutes pairwise.

Open problem.

Can the above mentioned theorems be proved for intuitionistic fuzzy metric spaces?

Acknowledgment

The authors would like to express their sincere thanks to Professor Dorel Mihet for his paper [18]. The third author is also thankful to the Ministry of Science and Technological Development of Serbia.

REFERENCES

- [1] I. Aalam - S. Kumar - B.D. Pant, *A common fixed point theorem in fuzzy metric space*, Bull. Math. Anal. Appl. 2 (4) (2010), 76–82.
- [2] M. Aamri - D. El Moutawakil, *Some new common fixed point theorems under strict contractive conditions*, J. Math. Anal. Appl. 270 (1) (2002), 181–188.
- [3] M. Abbas - I. Altun - D. Gopal, *Common fixed point theorems for non compatible mappings in fuzzy metric spaces*, Bull. Math. Anal. Appl. 1 (2) (2009), 47–56.

- [4] S. S. Chang - Y. J. Cho - B. S. Lee - J. S. Jung - S. M. Kang, *Coincidence point theorems and minimization theorems in fuzzy metric spaces*, Fuzzy Sets and Systems 88 (1) (1997), 119–127.
- [5] S. Chauhan - B. D. Pant, *Common fixed point theorems in fuzzy metric spaces*, Bull. Allahabad Math. Soc. 27 (2012), in press.
- [6] Y. J. Cho, *Fixed points for compatible mappings of type (A)*, Math. Japon. 38 (3) (1993), 497–508.
- [7] Y. J. Cho, *Fixed points in fuzzy metric spaces*, J. Fuzzy Math. 5 (4) (1997), 949–962.
- [8] A. George - P. Veeramani, *On some result in fuzzy metric space*, Fuzzy Sets and Systems 64 (1994), 395–399.
- [9] M. Grabiec, *Fixed points in fuzzy metric spaces*, Fuzzy Sets and System 27 (3) (1988), 385–389.
- [10] M. Imdad - J. Ali, *Some common fixed point theorems in fuzzy metric spaces*, Math. Commun. 11 (2) (2006), 153–163.
- [11] M. Imdad - J. Ali, *A general fixed point theorem in fuzzy metric spaces via an implicit function*, J. Appl. Math. Inform. 26 (3-4) (2008), 591–603.
- [12] M. Imdad - J. Ali - M. Tanveer, *Coincidence and common fixed point theorems for nonlinear contractions in Menger PM spaces*, Chaos, Solitons Fractals 42 (5) (2009), 3121–3129.
- [13] G. Jungck - B. E. Rhoades, *Fixed points for set valued functions without continuity*, Indian J. Pure Appl. Math. 29 (3) (1998), 227–238.
- [14] I. Kramosil - J. Michalek, *Fuzzy metric and statistical metric spaces*, Kybernetika (Prague) 11 (5) (1975), 336–344.
- [15] S. Kumar, *Fixed point theorems for weakly compatible maps under E.A. property in fuzzy metric spaces*, J. Appl. Math. Inform. 29 (1-2) (2011), 395–405. MR2798597
- [16] S. Kumar - S. Chauhan, *Common fixed point theorems using implicit relation and property (E.A) in fuzzy metric spaces*, Ann. Fuzzy Math. Inform. (2012), in press.
- [17] S. Kumar - B. Fisher, *A Common fixed point theorem in fuzzy metric space using property (E.A.) and implicit relation*, Thai J. Math. 8 (3) (2010), 439–446.
- [18] D. Mihet, *Fixed point theorems in fuzzy metric spaces using property E.A.*, Non-linear Anal. 73 (7) (2010), 2184–2188.
- [19] S. N. Mishra - N. Sharma - S. L. Singh, *Common fixed points of maps on fuzzy metric spaces*, Int. J. Math. Math. Sci. 17 (2) (1994), 253–258.
- [20] P. P. Murthy - S. Kumar - K. Tas, *Common fixed points of self maps satisfying an integral type contractive condition in fuzzy metric spaces*, Math. Commun. 15 (2) (2010), 521–537.
- [21] M. S. El Naschie, *On the uncertainty of Cantorian geometry and two-slit experiment*, Chaos Solitons Fractals 9 (3) (1998), 517–529.

- [22] M. S. El Naschie, *A review of E-infinity theory and the mass spectrum of high energy particle physics*, Chaos Solitons Fractals 19 (2004), 209–236.
- [23] M. S. El Naschie, *A review of applications and results of E-infinity theory*, Int. J. Nonlinear Sci. Numer. Simul. 8 (2007), 11–20.
- [24] D. O'Regan - M. Abbas, *Necessary and sufficient conditions for common fixed point theorems in fuzzy metric space*, Demonstratio Math. 42 (4) (2009), 887–900.
- [25] B. D. Pant - S. Chauhan, *Common fixed point theorems for two pairs of weakly compatible mappings in Menger spaces and fuzzy metric spaces*, Vasile Alecsandri Univ. Bacău Sci. Stud. Res. Ser. Math. Inform. 21 (2) (2011), 81–96.
- [26] V. Pant - R. P. Pant, *Fixed points in fuzzy metric space for noncompatible maps*, Soochow J. Math. 33 (4) (2007), 647–655.
- [27] H. K. Pathak - R. R. López - R., K. Verma, *A common fixed point theorem using implicit relation and property (E.A) in metric spaces*, Filomat 21 (2) (2007), 211–234.
- [28] B. Schweizer - A. Sklar, *Probabilistic Metric Spaces*, North-Holland Series in Probability and Applied Mathematics. North-Holland Publishing Co., New York 1983.
- [29] S. Sedghi - C. Alaca - N. Shobe, *On fixed points of weakly commuting mappings with property (E.A)*, J. Adv. Stud. Topol. 3 (3) (2012), 11–17.
- [30] S. Sedghi - N. Shobe, *Common fixed point theorems for weakly compatible mappings satisfying contractive condition of integral type*, J. Adv. Res. Appl. Math. 3 (4) (2011), 67–78.
- [31] S. Sedghi - N. Shobe - A. Aliouche, *A common fixed point theorem for weakly compatible mappings in fuzzy metric spaces*, Gen. Math. 18 (3) (2010), 3–12.
- [32] Y. Shen - D. Qiu - W. Chen, *Fixed point theorems in fuzzy metric spaces*, Appl. Math. Lett. 25 (2) (2012), 138–141.
- [33] S. L. Singh - A. Tomar, *Fixed point theorems in FM-spaces*, J. Fuzzy Math. 12 (4) (2004), 845–859.
- [34] W. Sintunavarat - Y. J. Cho - P. Kumam, *Coupled coincidence point theorems for contractions without commutative condition in intuitionistic fuzzy normed spaces*, Fixed Point Theory Appl. Vol. 2011, 2011.
- [35] W. Sintunavarat - P. Kumam, *Common fixed point theorems for a pair of weakly compatible mappings in fuzzy metric spaces*, J. Appl. Math. Vol. 2011, Article ID 637958, 14 pages, 2011.
- [36] W. Sintunavarat - P. Kumam, *Fixed point theorems for a generalized intuitionistic fuzzy contraction in intuitionistic fuzzy metric spaces*, Thai J. Math. 10 (1) (2012), 123–135.
- [37] W. Sintunavarat - P. Kumam, *Common fixed points for R-weakly commuting in fuzzy metric spaces*, Annali dell'Università di Ferr., in press.
- [38] P. V. Subrahmanyam, *A common fixed point theorem in fuzzy metric spaces*, Inform. Sci. 83 (3-4) (1995), 109–112.

- [39] R. Vasuki, *Common fixed points for R-weakly commuting maps in fuzzy metric spaces*, Indian J. Pure Appl. Math. 30 (4) (1999), 419–423.
- [40] L. A. Zadeh, *Fuzzy sets*, Inform. Control 8 (1965), 338–353.

SUNNY CHAUHAN

Near Nehru Training Centre

H. No: 274, Nai Basti B-14

Bijnor, 246701, Uttar Pradesh, India.

e-mail: sun.gkv@gmail.com

SHUBHAM BHATNAGAR

Near Jain Temple, Shambha Bazar

Bijnor-246701, Uttar Pradesh, India.

STOJAN RADENOVIC

University of Belgrade

Faculty of Mechanical Engineering

Kraljice Marije 16, 11120 Beograd, Serbia.

e-mail: radens@beotel.net